



Increase in chemokine CXCL1 by ER β ligand treatment is a key mediator in promoting axon myelination

Hawra Karim^{a,b}, Sung Hoon Kim^c, Andrew S. Lapato^{a,b}, Norio Yasui^c, John A. Katzenellenbogen^c, and Seema K. Tiwari-Woodruff^{a,b,1}

^aDivision of Biomedical Sciences, University of California, Riverside School of Medicine, Riverside, CA 92521; ^bCenter for Glial-Neuronal Interactions, University of California, Riverside School of Medicine, CA 92506; and ^cDepartment of Chemistry, University of Illinois at Urbana-Champaign, Urbana, IL 61801

Edited by Bruce S. McEwen, The Rockefeller University, New York, NY, and approved April 27, 2018 (received for review December 19, 2017)

Estrogen receptor β (ER β) ligands promote remyelination in mouse models of multiple sclerosis. Recent work using experimental autoimmune encephalomyelitis (EAE) has shown that ER β ligands induce axon remyelination, but impact peripheral inflammation to varying degrees. To identify if ER β ligands initiate a common immune mechanism in remyelination, central and peripheral immunity and pathology in mice given ER β ligands at peak EAE were assessed. All ER β ligands induced differential expression of cytokines and chemokines, but increased levels of CXCL1 in the periphery and in astrocytes. Oligodendrocyte CXCR2 binds CXCL1 and has been implicated in normal myelination. In addition, despite extensive immune cell accumulation in the CNS, all ER β ligands promoted extensive remyelination in mice at peak EAE. This finding highlights a component of the mechanism by which ER β ligands mediate remyelination. Hence, interplay between the immune system and central nervous system may be responsible for the remyelinating effects of ER β ligands. Our findings of potential neuroprotective benefits arising from the presence of CXCL1 could have implications for improved therapies for multiple sclerosis.

immunomodulation | oligodendrocytes | remyelination | multiple sclerosis | CXCL1

Multiple sclerosis (MS) is an autoimmune central nervous system (CNS) demyelinating disease (1). Experimental autoimmune encephalomyelitis (EAE) is a widely utilized, T cell-mediated mouse model that shares key pathological features of MS, including development of inflammatory lesions, demyelination, axon pathology, and gliosis (2). In both MS and EAE, proinflammatory cytokines produced by autoreactive T helper (Th) cells induce oligodendrocyte (OL) apoptosis, resulting in demyelination and neurodegeneration (3–5).

Current approved therapies for relapsing-remitting MS prevent relapses and slow disability progression primarily by reducing inflammation. However, they fail to reverse axonal pathology or restore myelination (6). No directly neuroprotective or remyelinating agents are currently available (7).

Recently, estrogens have emerged as attractive candidates for MS therapy that fulfill this need. Estrogenic effects are primarily mediated by ligand-activated nuclear estrogen receptor (ER) α and β isoforms. Nonselective endogenous estrogens produced during pregnancy reduce MS and EAE relapses and severity through primarily ER α -mediated suppression of inflammation (8–10). Furthermore, treatment of nonpregnant female MS patients with pregnancy levels of the placenta-derived estrogen hormone estradiol reduces circulating CD4⁺ T cell numbers and proinflammatory TNF- α and IFN- γ , while increasing antiinflammatory IL-5 and IL-10 (11). In EAE mice, selective ER α ligands are broadly antiinflammatory and decrease leukocyte infiltration into the CNS (12). Unfortunately, estrogenic signaling through ER α is also feminizing in males and associated with proliferative effects that increase cancer risk, limiting the clinical usefulness of endogenous estrogens (13). In contrast, selective ER β ligands offer many of estrogen's benefits without these

deleterious side effects (see Fig. 1 for structure and selectivity of ER β vs. ER α ligands).

Treatment with highly selective ER β ligands modulates autoimmunity, improves neurological deficits, and increases myelination in mice with EAE, despite ongoing infiltration of autoreactive leukocytes (12, 14–16). For example, diarylpropionitrile (DPN), which displays >70-fold ER β selectivity over ER α , ameliorates late-stage EAE clinical disease (12, 14, 17, 18), but does not affect production of IFN- γ , TNF- α , or IL-6 by peripheral mononuclear cells or reduce CNS infiltration of CD45⁺ leukocytes (12, 18). Notably, while overall immunity was not significantly altered, the frequency of antigen-presenting dendritic cells within the CNS was reduced by DPN, which also impeded their production of TNF- α (18).

In addition to their immunological effects, ER β agonists also mediate therapeutic effects through action on CNS populations. In both EAE and cuprizone models of MS, DPN or chloroindazole (IndCl) treatment increased callosal OLs and myelination, while simultaneously improving functional measures (12, 15, 17). In contrast to DPN, either prophylactic or therapeutic treatment with IndCl reduced leukocyte infiltration into the CNS of EAE mice (15), decreased peripheral Th1 and Th17 cytokine production (15), and suppressed expression of IL-1 β , IL-6, and IL-23 by microglia (19). DPN and IndCl treatment during EAE increased brain-derived neurotrophic factor levels and activated the PI3K/Akt/mammalian target of rapamycin pathway, which is involved in OL progenitor (OPC) proliferation and differentiation (15). The nonsteroidal agonist WAY-202041 (WAY) exhibits >200-fold selectivity for ER β over ER α , and also decreases inflammation and shows promise for the treatment of MS (20). Our recent results show that WAY has similar neuroprotective effects to DPN and IndCl in chronic EAE.

Significance

CXCL1 is a major neutrophil chemoattractant that binds to the chemokine receptor, CXCR2, on neutrophils and oligodendrocytes. Estrogen receptor β ligand treatment in a mouse model of multiple sclerosis induces an increase in peripheral and brain CXCL1 levels that correlate with an increase in axon remyelination. Oligodendrocyte progenitor recruitment and differentiation by CXCL1, in combination with attenuated IFN- γ production reducing apoptosis, may account for at least one avenue whereby estrogen receptor β ligands exert their clinical benefits.

Author contributions: J.A.K. and S.K.T.-W. designed research; H.K., S.K., A.S.L., and N.Y. performed research; H.K., J.A.K., and S.K.T.-W. analyzed data; and H.K., A.S.L., and S.K.T.-W. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

¹To whom correspondence should be addressed. Email: seema.tiwari-woodruff@medsch.ucr.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1721732115/-DCSupplemental.

Published online May 29, 2018.




Fig. 1. Structure and selectivity of ER β ligands: ER β vs. ER α .

Previously, ER β ligand-mediated immune effects have been assessed after peak disease when demyelination and CNS inflammation has plateaued (12, 15). To understand how ER β ligands modify EAE pathogenesis and to expose a potential common immunomodulatory role of ER β ligands, leukocyte populations and secreted cytokines/chemokines were analyzed from mice treated with one of several ER β ligands from EAE onset to peak disease. In the present study, we show that therapeutic ER β ligands enhance peripheral and CNS concentrations of the C-X-C motif chemokine, CXCL1, with similar remyelinating and neuroprotective properties.

Results

Effect of Therapeutic ER β Ligand Treatment on EAE Clinical Score and Uterine Weight. A persistent disease course starting at day 15–16 with chronic motor deficits was observed in vehicle-treated EAE mice. Therapeutic IndCl and WAY treatment administered at peak disease decreased clinical disease severity over time (15) (Fig. 2*A* and *SI Appendix*, Table S2). Not previously reported, therapeutic WAY treatment induced an increase in myelin density, increase in the OL population, and decrease in “g-ratio” of white-matter tracts (*SI Appendix*, Fig. S2), similar to EAE groups treated with DPN (17) or IndCl (15).

To address therapeutic response of different ER β ligand treatment on peak EAE pathology and immune response, treatment was started at day 8 and suspended on day 21 (Fig. 2*B* and *SI Appendix*, Table S3). Estradiol pretreatment (pre-E2) completely abrogated clinical disease (15). All other treatment groups displayed onset of clinical disease beginning 7–10 d post-induction, with peak disease (defined as 2 to 3 sequential days of maximal clinical score) occurring between days 18 and 21 (Fig. 2*A* and *B*). All therapeutic ER β ligands administered at onset of disease did not attenuate clinical symptoms by day 21 (Fig. 2*B*). Pre-E2 treatment and not ER β ligands increased uterine weight by nearly fourfold (Fig. 2*C*), as previously observed (15).

Improved Axon Myelination and No Effect on Leukocyte Infiltration and Astrogliosis. Reduced axon numbers and myelination at peak EAE has been reported (17, 21). Similarly, vehicle-treated mice at peak disease (Fig. 2*B*) displayed extensive demyelination, indicated by reduced myelin basic protein (MBP⁺) staining intensity, proximal to CD45⁺ (a pan-leukocyte marker) perivascular cuffs and leukocyte infiltrates that were not observed in normal animals (Fig. 3*A*). Significant loss of MBP⁺ and NF200⁺ axons was observed in the vehicle-treated mice, while ER β ligand treatment rescued both myelin and axons (Fig. 3*B* and *C*).

There was an increase in CD45⁺ leukocyte infiltration in the white matter of vehicle-treated and ER β ligand-treated mice, whereas pre-E2 treatment had minimal CD45⁺ immunoreactivity (Fig. 4*A* and *C*). Similarly, immunoreactivity for the myeloid cell marker ionized calcium-binding adaptor molecule 1 Iba1⁺ and the astrocyte marker glial fibrillary acidic protein (GFAP) was significantly elevated in vehicle- and ER β ligand-treated mice (Fig. 4). Pre-E2 abrogated this increase (Fig. 4*D* and *E*).

No Change in CNS-Infiltrating Leukocyte Populations. To profile CNS infiltrating leukocytes after ER β ligand treatment, mononuclear cells were isolated at peak EAE disease and analyzed

by flow cytometry. Vehicle-treated mice showed elevated CD11b⁺CD45^{hi} activated microglia/macrophage frequency (gate R2) (Fig. 5*A* and *C*). Pre-E2 significantly reduced CD11b⁺CD45^{hi} frequency (Fig. 5*C*), while ER β ligand and vehicle treated groups displayed similar frequency of activated microglia/macrophages (Fig. 5*C*). Similarly, increased CD4⁺IFN- γ ⁺ Th1 and CD4⁺IL-17⁺ Th17 frequencies were observed in vehicle-treated mice but not pre-E2 (Fig. 5*D* and *E* and *SI Appendix*, Fig. S3). ER β ligands did not affect Th1 or Th17 frequency (Fig. 5*D* and *E*).

Modification of Cytokine and Chemokine Production by Peripheral Leukocytes.

To characterize the effects of ER β ligands on the peripheral immune system, splenocytes were harvested from mice 21 d after EAE induction and assessed by flow cytometry. Splenic CD11b⁺CD45⁺ macrophage/monocyte composition was attenuated by pre-E2, but not ER β ligand treatment (Fig. 6*A* and *B*). Th1 frequency was significantly reduced in pre-E2 and ER β ligand treated groups vs. vehicle (Fig. 6*C*), while Th17 frequency was not affected (Fig. 6*D* and *SI Appendix*, Fig. S3).

Cytokine and chemokine expression was analyzed to assess the role of leukocyte effector function in ER β ligand derived benefits during peak disease (22).

Proinflammatory cytokines. The Th1 cytokine IFN- γ was significantly up-regulated in vehicle-treated mice vs. normal. In contrast, pre-E2, post-IndCl, WAY, and DPN decreased IFN- γ concentration relative to vehicle (Fig. 7*A*, *i*). IL-2 was significantly increased in vehicle-treated mice vs. normal and was attenuated only by pre-E2 treatment (Fig. 7*A*, *ii*). Vehicle-treated mice produced higher levels of TNF- α and IL-17 vs. normal (Fig. 7*A*, *iii* and *iv*). Pre-E2, IndCl, WAY, and DPN produced similar levels of TNF- α and IL-17 as vehicle treatment (Fig. 7*A*, *iii* and *iv*). Increased IL-1 β was observed in vehicle-treated and pre-E2 groups vs. normal. Interestingly, all ER β ligand groups exhibited greater IL-1 β production vs. vehicle (Fig. 7*A*, *v*).




Fig. 2. Therapeutic ER β ligand treatment decreases EAE severity with no effect on uterine weight. (A) Treatment with ER β ligands, IndCl (post-EAE+IndCl; purple), and WAY (post-EAE+WAY; orange) began on day 21 until day 40. Onset of clinical disease occurred between days 17 and 19 in vehicle-treated EAE mice (post-EAE+vehicle; red), with peak severity at day 21. Treatment with IndCl and WAY significantly decreased disease severity vs. vehicle beginning by day 24 through 40 (IndCl-purple stars; WAY-orange stars). Differences in EAE clinical scores were determined by two-way unbalanced ANOVA with Dunnett's multiple comparisons test (48). $n = 8$ –10 mice per group; * $P < 0.05$, *** $P < 0.01$, **** $P < 0.001$, **** $P < 0.0001$. (B) Therapeutic treatment with ER β ligands, IndCl, WAY and DPN (post-EAE+DPN; blue) began on day 8 until day 21. Prophylactic E2 (pre-EAE+E2; green) prevented onset of clinical disease. Vehicle-treated EAE mice displayed onset of clinical disease symptoms between days 7 and 10 with disease severity peaking around day 17. During peak disease, IndCl, WAY, and DPN treatment did not affect EAE clinical scores. (C) Assessment of postperfusion uterus to body weight. Only pre-E2-treated female mice exhibited a 4x increase in uterus:body weight ratio. $n = 8$ –10 mice per group.




Fig. 3. Therapeutic ER β ligand treatment improves myelination during peak EAE disease. (A) Representative 10 \times magnification images of thoracic spinal cords collected at peak disease (day 21). Experimental groups shown in Fig. 2B, were immunostained for MBP (red), CD45 (green), and DAPI (blue), with 4 \times normal (i) and vehicle (ii) images to show pathology. (B) Magnified 40 \times images showing ventral column immunostained with MBP (red) and NF200 (green). (C) Quantification of MBP intensity and NF200 $^+$ axon numbers revealed significant decreases in only vehicle treated groups. $n = 5$ mice per group; * $P < 0.05$, ** $P < 0.01$.

Antiinflammatory cytokines. Skewing immune response toward production of Th2 cytokines, such as IL-4, IL-5, and IL-10, with parallel suppression of Th1 cytokines ameliorates EAE (23). IL-4 was up-regulated only in vehicle-treated mice (Fig. 7B, i). Similarly, neither IL-5 nor IL-10 was changed by E2 or ER β ligand treatment (Fig. 7B, ii and iii).

Chemokines. CXCL1 was significantly up-regulated by ER β ligands relative to vehicle (Fig. 7C, i). No change was observed in pre-E2 groups. CXCL9 and CXCL10, closely related IFN- γ -inducible T cell and monocyte chemoattractants, were increased in vehicle-treated mice vs. normal. CXCL9 levels were similar in all treatment groups compared with vehicle; however, pre-E2 and IndCl treatment significantly reduced CXCL10 levels, with no change by WAY or DPN treatment (Fig. 7C, ii and iii).

ER β Ligands Up-Regulate CXCL1. Due to its role in OPC recruitment and up-regulation by splenic leukocytes in response to ER β ligand treatment, CXCL1 expression was examined in white matter from mice at peak EAE by immunohistochemistry (IHC). In the CNS, astrocytes are a primary source of CXCL1 during white-matter development (24) and neuroinflammation (25). Compared with vehicle, IndCl- and DPN-treated mice exhibited increased CXCL1 $^+$ immunoreactivity that appeared to overlap with GFAP (Fig. 8A–C) (26). In addition, CXCL1 immunoreactivity colocalized with IL-1 β and GFAP in astrocytes during ER β ligand treatment (SI Appendix, Fig. S4). In addition, supernatant from primary astrocyte cultures stimulated with IL-1 β contained increased levels of CXCL1. The same supernatant when added to OL cultures induced an increase in MBP expression (SI Appendix, Fig. S5).

CXCR2 receptor is critical for developmental OPC positioning and differentiation in the CNS (27). OL primary cultures in the presence of CXCR2 antagonist, SB 225002 and IL-1 β astrocyte-conditioned media showed decreased differentiation and enhanced OL apoptosis (SI Appendix, Fig. S5). To determine the distribution of CXCR2 during peak EAE disease in response to ER β ligand treatment, CXCR2 expression by OPCs/OLs was assessed by colabeling with the OL lineage-specific transcription factor, Olig2 in spinal cords. Olig2 $^+$ nuclei were comparable between all groups evaluated and CXCR2 was

detected in a subset of Olig2 $^+$ nuclei (Fig. 8B and C). The fraction of CXCR2 $^+$ Olig2 $^+$ cells increased in all treatment groups (Fig. 8E), consistent with previous reports that OPC/OL CXCR2 is up-regulated in MS (28).




Fig. 4. Treatment with ER β ligands do not exhibit significant changes in CNS immune cell populations during peak EAE disease. (A and B) Representative 40 \times magnification images of thoracic spinal cords collected at peak disease (day 21) from experimental groups shown in Fig. 2B were immunostained for astrocytes (GFAP, green), leukocytes (CD45, red), microglia/macrophages (Iba1, red), and DAPI (blue). (C–E) Quantification of normalized intensities of CD45, GFAP, and Iba1. Vehicle-treated mice exhibited increased CD45, Iba1, and GFAP intensity that was decreased only with pre-E2 treatment. $n = 5$ mice per group; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Fig. 5. Therapeutic ER β ligand treatment effect on CNS inflammatory immune cell numbers during peak EAE disease. (A) Representative plots of CD11b (y axis) and CD45 (x axis) distribution in the CNS from vehicle, pre-E2, and therapeutic ER β ligand treated mice collected from peak EAE mice. In the CNS, resting microglia (R1) are represented by CD11b⁺CD45^{lo} populations, while activated microglia/macrophages (R2) are represented by CD11b⁺CD45^{hi}. (B and C) Quantification of resting (CD11b⁺CD45^{lo}; R1) and activated microglia/macrophages (CD11b⁺CD45^{hi}; R2) from prophylactic E2 and therapeutic ER β ligands treated EAE mice. Pre-E2 decreased activated macrophages compared with vehicle. (D and E) Th1 (CD4⁺IFN- γ ⁺) and Th17 (CD4⁺IL-17⁺) frequency in the CNS was decreased by pre-E2 treatment. ER β ligands did not affect Th1 or Th17 frequency. $n = 8-10$ mice per group; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Discussion

In this study, we characterized the therapeutic impact of three ER β ligands in a clinically relevant mouse model of MS. Our results suggest that the protective effects of ER β ligands on remyelination and neuroprotection may be mediated by changes in cytokine and chemokine production. Comparisons among the ER β ligands tested show improved myelination. Consistent with previous studies (12, 15), this occurred with little impact on the composition or polarization of CNS or splenic leukocytes, but involved reduction of peripheral IFN- γ production and increased CXCL1 expression in the CNS and spleen. Individual ER β ligands displayed differential effects on the production of the IFN- γ -inducible CXCL10, with IndCl being the only compound tested to suppress production aside from pretreatment with E2. These data indicate that individual ER β ligands may have various effects on leukocyte activity in EAE, while also pointing to a potential immunological mechanism common to ER β -mediated remyelination. Specifically, three findings that were observed with all ER β ligands tested stand out: (i) increased myelination, (ii) reduced IFN- γ , and (iii) increased CXCL1. These data suggest that ER β ligands may facilitate an environment within the inflamed CNS where OPCs are recruited to the site of ongoing demyelination by astrocytic CXCL1 and allowed to differentiate into myelinating OLs through attenuation of cytotoxic IFN- γ production (SI Appendix, Fig. S6).

Although the ligands used in this study were all selective for ER β , their activity in the peripheral and CNS may differ (12, 15, 19, 29). For example, IndCl, but not DPN or WAY, was effective at repressing inflammatory response gene products in microglia and astrocytes (19). The differential immune effects observed by treatment with the ER β ligands tested in the present study may be due to how these compounds recruit cofactors. As an example, IndCl promotes entry of ER β into the C-terminal binding protein-dependent transrepression pathway while DPN does not (19). However, while transcriptional cofactor recruitment differs, these ER β ligands converge in their ability to induce OPC/OL survival and myelination during demyelinating disease (15, 30). Importantly, this was shown to be mediated by direct action on OLs as conditional knockout of ER β from these cells abrogated the clinical benefits of DPN (30). Quiescent microglia are distributed throughout the homeostatic CNS and are characterized by coexpression of the myeloid cell integrin/complement receptor CD11b and low levels of CD45. In the present study, unlike pre-E2, no ER β ligand tested abrogated the rise in activated

microglia/macrophages recovered from the CNS or spleen of vehicle-treated mice.

Unexpectedly, ER β ligands had divergent effects on peripheral and CNS T lymphocyte populations. Pre-E2 reduced the frequency of Th1 cells in splenic and CNS mononuclear cells. In contrast, pre-E2 reduced CNS-infiltrating Th17 cells, but had no effect on splenic lymphocytes. Similarly, ER β ligand treatment decreased Th1 cell frequency, as well as proinflammatory cytokine IFN- γ in the periphery, with no change in Th1 populations in the CNS. This finding may reflect CNS infiltration at disease onset or reduced Th1 differentiation resulting from decreased IFN- γ concentrations. Another ER β ligand, 4-(2-phenyl-5,7-bis[trifluoromethyl]pyrazolo[1,5-a]pyrimidin-3-yl)phenol (PHTPP), was shown to suppress Th17 response by acting directly on CD4⁺ T cells (16) while IndCl altered CD4⁺ T cell ER β expression in vitro (16). We did not observe a suppression of Th17 frequency or peripheral IL-17 secretion with ER β ligands. Additional study is required to assay the effect of ER β ligands on CD8⁺ populations; however, E2 has been shown to increase their activity in a viral of lung infection (31).

Proinflammatory cytokines, such as IFN- γ , drive EAE by potentiating antigen presentation and secretion of additional cytokines and chemokines (22). Of potential importance to the mechanism whereby ER β ligands promote remyelination, splenocytes from ER β ligand treatment produced less IFN- γ in response to MOG₃₅₋₅₅ peptide. IFN- γ elicits OL apoptosis in vitro (32) and ectopic CNS expression results in OL death and impaired myelination (33). These data indicate that IFN- γ -induced death of OLs/OPCs may represent a potential etiology for remyelination failure in EAE and MS. However, IFN- γ may possess pleiotropic effects on OLs during demyelinating disease (34). Thus, ER β ligand-mediated peripheral IFN- γ reduction may play a key role in supporting remyelination by slowing Th1 differentiation, while lowered levels of CNS IFN- γ may improve OL survival.

Interestingly, IL-1 β , which is associated with various inflammatory and demyelinating disorders (35) and responsive to E2 signaling (36), was significantly increased by ER β ligand treatments. While IL-1 β is cytotoxic to mature OLs in vitro, neither demyelination nor OL loss during cuprizone demyelination are attenuated in mice lacking IL-1 β (35). However, remyelination fails due to lack of IL-1 β -mediated expression of insulin growth factor-1 (37), pointing toward a complex role for this cytokine. Our results suggest that IL-1 β in the presence of the ER β ligands may be associated with myelin repair by promoting mature OL differentiation and remyelination during EAE.

Chemokines and their cognate receptors play a critical role in the recruitment and trafficking of leukocytes in the context of




Fig. 6. Effect of therapeutic ER β ligand treatment on splenic immune cell numbers during peak EAE. (A) Representative plots of splenic CD11b (y axis) and CD45 (x axis) distribution in vehicle, prophylactic E2, and therapeutic ER β ligand treated mice collected at peak disease. Activated monocytes/macrophages (R2) are represented by CD11b⁺CD45^{hi}. (B) Quantification of activated microglia/macrophage frequency was decreased only in pre-E2-treated mice. (C and D) ER β ligands decreased Th1 frequency. No change was observed in Th17 frequency in any group vs. vehicle. $n = 8-10$ mice per group; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Fig. 7. Therapeutic ER β ligand IndCl treatment effects on cytokine and chemokine production by peripheral immune cells during peak EAE disease. (A) Splenocytes collected from vehicle-treated mice at peak disease stimulated with MOG_{35–55} for 48 h showed increased IFN- γ , IL-2, TNF- α , IL-17, and IL-1 β vs. normal. (i) Pre-E2 and therapeutic ER β ligand treatment decreased IFN- γ . (ii) Only pre-E2 groups decreased IL-2 vs. vehicle. (iii and iv) No change was observed in TNF- α and IL-17 in any treatment group. (v) ER β ligands increased IL-1 β levels vs. vehicle. (B, i–iii) Vehicle treated mice exhibited increased levels of anti-inflammatory cytokine IL-4 vs. normal. No change was observed in IL-4, IL-5, or IL-10 production in any treatment groups. (C) Vehicle treated mice exhibited increased levels of CXCL9 and CXCL10 with no increase in CXCL1 levels. (i) ER β ligands, but not preE2, increased CXCL1. (ii) Similar levels of CXCL9 was observed in all treatment groups. (iii) Only pre-E2 and therapeutic IndCl groups reduced CXCL10 vs. vehicle. $n = 5–8$ mice per group; * $P < 0.05$, ** $P < 0.01$.

disease conditions. ER β ligands increased CXCL1 and decreased CXCL10. Abundant CXCL10 has been detected at early time points in the development of EAE and MS lesions (38) and promotes neuronal and OL apoptosis (39). Here, only IndCl decreased CXCL10 levels.

CXCL1, a major neutrophil chemoattractant expressed by astrocytes and microglia, affects chemoattraction through its receptor CXCR2, which is present on diverse cellular populations, including leukocytes, keratinocytes, dermal fibroblasts, neutrophils, and OLs (40, 41). E2 has been shown to be a negative regulator of CXCL1/CXCR2 signaling pathways (42–45) through predominantly ER α -dependent pathways (42). However, CXCL1-mediated functions extend beyond chemoattraction. In MS, both CXCL1⁺ astrocytes and CXCR2⁺ OLs (46) have been detected in lesions (46), indicating that it may play a role in recruitment of OPCs to demyelinating sites. CXCL1 overexpression by astrocytes attenuates EAE disease severity (47), while CXCR2 deficient mice exhibit reduced white-matter volume and myelin proteins in the spinal cord (24). Furthermore, CXCR2 signaling promotes OPC survival by increasing levels of the antiapoptotic protein, Bcl-2 (39). Together, these results indicate a potential role for astrocyte derived CXCL1 in promoting OL survival in ER β ligand treated EAE mice.

Our finding that ER β ligand therapy promotes increased astrocytic CXCL1 production may represent an additional facet of how these compounds induce remyelination. OPC recruitment and differentiation by CXCL1, in combination with attenuated IFN- γ production reducing OL apoptosis, may account for at least one avenue whereby ER β ligands exert their clinical benefits. In summary, our data show that ER β ligand neuroprotection/remyelination may be partly mediated by skewing the proinflammatory

phenotype to a protective phenotype. Our results demonstrate that the interplay between CNS- and immune-derived signals is central to the induction and regulation of neuroinflammatory diseases such as MS. The possibility that ER β ligands modulate the cytokine and chemokine milieu to potentially promote repair/remyelination opens up exciting therapeutic options.

Materials and Methods

Treatment. Drugs were dissolved in vehicle (10% ethanol and 90% Miglyol 812N) at the following concentrations and administered subcutaneously daily; IndCl 5 mg/kg/d, WAY 1 or 10 mg/kg/d, and DPN 8 mg/kg/d (Fig. 1); 2.5 mg/kg/d E2 was used as a positive control. Treatment was initiated at postimmunization day 0, 8, or 21.

EAE. Active EAE was induced in 8-wk-old female C57BL/6 mice, as previously described (14, 48) (SI Appendix, Fig. S1). Animals were maintained according to the University of California, Riverside Office of Research Integrity and the Institutional Animal Care and Use Committee.




Fig. 8. Modulation of CXCL1 and its receptor CXCR2 in the CNS from ER β ligand-treated EAE mice. (A) Representative 40 \times images of thoracic spinal cord ventral column at peak disease (Fig. 2B) immunostained for CXCL1 (red), GFAP (green), and DAPI (blue). (A and B) White dashed boxes from merged images were split and enlarged 3 \times . Arrows indicate costaining between CXCL1 and GFAP. (B) Magnified 40 \times images of CXCR2 (red) and OPCs/OLs (Olig2, green), arrows indicate CXCR2⁺Olig2⁺ cells. (C) ER β ligands increased CXCL1 staining intensity vs. vehicle. (D and E) Olig2⁺ cell numbers were comparable among groups, while the fraction of Olig2⁺CXCR2⁺ OPCs/OLs were increased in groups with EAE. $n = 5–8$ mice per group; * $P < 0.05$, *** $P < 0.001$, **** $P < 0.0001$.

Mononuclear Cell Isolation. Splenocytes were isolated and stimulated in vitro using 25 $\mu\text{g}/\text{mL}$ MOG_{35–55} (30, 49). Supernatants were collected after 48 h and cytokine levels were determined by Luminex. CNS mononuclear cells were isolated per the protocol in ref. 50.

Flow Cytometry and ELISA. Leukocytes were stained with antibodies (SI Appendix, Table S1) and flow cytometry data were acquired. CXCL1 concentrations supernatant were measured using the murine CXCL1 ELISA kit.

Cell Culture. Primary astrocytes and OPCs were isolated as described previously (51). Astrocytes were treated with IL-1 β or media alone for 48 h then supernatant was applied to primary OLs \pm CXCR2 antagonist SB 225002.

IHC and Analysis. IHC was performed on formalin-fixed thoracic spinal cord sections and primary cell cultures with different antibodies (SI Appendix, Table S1). Images were quantified using unbiased stereology (49) and ImageJ software.

Statistical Analysis. All data were analyzed in GraphPad Prism 6. Differences in EAE clinical scores were determined by two-way unbalanced ANOVA with Dunnett's multiple comparisons test (48). All others were analyzed by ordinary one-way ANOVA with Dunnett's multiple comparisons test or Kruskal-Wallis with Dunn's multiple comparisons test. Data are presented as mean \pm SEM; $n = 2$ –3 independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Data Availability. All data generated or analyzed during this study are included in herein and in the SI Appendix.

- Trapp BD, Nave KA (2008) Multiple sclerosis: An immune or neurodegenerative disorder? *Annu Rev Neurosci* 31:247–269.
- Constantinescu CS, Farooqi N, O'Brien K, Gran B (2011) Experimental autoimmune encephalomyelitis (EAE) as a model for multiple sclerosis (MS). *Br J Pharmacol* 164: 1079–1106.
- Fletcher JM, Lalor SJ, Sweeney CM, Tubridy N, Mills KH (2010) T cells in multiple sclerosis and experimental autoimmune encephalomyelitis. *Clin Exp Immunol* 162: 1–11.
- McFarland HF, Martin R (2007) Multiple sclerosis: A complicated picture of autoimmunity. *Nat Immunol* 8:913–919.
- Peterson LK, Fujinami RS (2007) Inflammation, demyelination, neurodegeneration and neuroprotection in the pathogenesis of multiple sclerosis. *J Neuroimmunol* 184: 37–44.
- Villoslada P (2016) Neuroprotective therapies for multiple sclerosis and other demyelinating diseases. *Mult Sci Dem Dis* 1:1–11.
- Münzel EJ, Williams A (2013) Promoting remyelination in multiple sclerosis—recent advances. *Drugs* 73:2017–2029, and erratum (2014) 74:157.
- Offner H, Polanczyk M (2006) A potential role for estrogen in experimental autoimmune encephalomyelitis and multiple sclerosis. *Ann N Y Acad Sci* 1089:343–372.
- Morales LBJ, et al. (2006) Treatment with an estrogen receptor alpha ligand is neuroprotective in experimental autoimmune encephalomyelitis. *J Neurosci* 26:6823–6833.
- Lélu K, et al. (2011) Estrogen receptor α signaling in T lymphocytes is required for estradiol-mediated inhibition of Th1 and Th17 cell differentiation and protection against experimental autoimmune encephalomyelitis. *J Immunol* 187:2386–2393.
- Sicotte NL, et al. (2002) Treatment of multiple sclerosis with the pregnancy hormone estradiol. *Ann Neurol* 52:421–428.
- Tiwari-Woodruff S, Morales LB, Lee R, Voskuhl RR (2007) Differential neuroprotective and antiinflammatory effects of estrogen receptor (ER) α and ER β ligand treatment. *Proc Natl Acad Sci USA* 104:14813–14818.
- Banks E, Canfell K (2009) Invited commentary: Hormone therapy risks and benefits—The Women's Health Initiative findings and the postmenopausal estrogen timing hypothesis. *Am J Epidemiol* 170:24–28.
- Kumar S, et al. (2013) Estrogen receptor β ligand therapy activates PI3K/Akt/mTOR signaling in oligodendrocytes and promotes remyelination in a mouse model of multiple sclerosis. *Neurobiol Dis* 56:131–144.
- Moore SM, et al. (2014) Multiple functional therapeutic effects of the estrogen receptor β agonist indazole-Cl in a mouse model of multiple sclerosis. *Proc Natl Acad Sci USA* 111:18061–18066.
- Aggelakopoulou M, Kourepini E, Paschalidis N, Panoutsakopoulou V (2016) ER β in CD4+ T cells is crucial for ligand-mediated suppression of central nervous system autoimmunity. *J Immunol* 196:4947–4956.
- Crawford DK, et al. (2010) Oestrogen receptor beta ligand: A novel treatment to enhance endogenous functional remyelination. *Brain* 133:2999–3016.
- Du S, Sandoval F, Trinh P, Umeda E, Voskuhl R (2011) Estrogen receptor- β ligand treatment modulates dendritic cells in the target organ during autoimmune demyelinating disease. *Eur J Immunol* 41:140–150.
- Saijo K, Collier JG, Li AC, Katzenellenbogen JA, Glass CK (2011) An ADIOL-ER β -CtBP transrepression pathway negatively regulates microglia-mediated inflammation. *Cell* 145:584–595.
- Harris HA, et al. (2003) Evaluation of an estrogen receptor-beta agonist in animal models of human disease. *Endocrinology* 144:4241–4249.
- Mangiardi M, et al. (2011) An animal model of cortical and callosal pathology in multiple sclerosis. *Brain Pathol* 21:263–278.
- Imitola J, Chitnis T, Khoury SJ (2005) Cytokines in multiple sclerosis: From bench to bedside. *Pharmacol Ther* 106:163–177.
- Moss RB, et al. (2004) Th1/Th2 cells in inflammatory disease states: Therapeutic implications. *Expert Opin Biol Ther* 4:1887–1896.
- Padovani-Claudio DA, Liu L, Ransohoff RM, Miller RH (2006) Alterations in the oligodendrocyte lineage, myelin, and white matter in adult mice lacking the chemokine receptor CXCR2. *Glia* 54:471–483.
- Hennessy E, Griffin EW, Cunningham C (2015) Astrocytes are primed by chronic neurodegeneration to produce exaggerated chemokine and cell infiltration responses to acute stimulation with the cytokines IL-1 β and TNF- α . *J Neurosci* 35: 8411–8422.
- Filipovic R, Zecevic N (2008) The effect of CXCL1 on human fetal oligodendrocyte progenitor cells. *Glia* 56:1–15.
- Robinson S, Tani M, Strieter RM, Ransohoff RM, Miller RH (1998) The chemokine growth-regulated oncogene-alpha promotes spinal cord oligodendrocyte precursor proliferation. *J Neurosci* 18:10457–10463.
- Omari KMJ, John GR, Sealton SC, Raine CS (2005) CXC chemokine receptors on human oligodendrocytes: Implications for multiple sclerosis. *Brain* 128:1003–1015.
- Minutolo F, Macchia M, Katzenellenbogen BS, Katzenellenbogen JA (2011) Estrogen receptor β ligands: Recent advances and biomedical applications. *Med Res Rev* 31: 364–442.
- Khalaj AJ, et al. (2013) Estrogen receptor (ER) β expression in oligodendrocytes is required for attenuation of clinical disease by an ER β ligand. *Proc Natl Acad Sci USA* 110:19125–19130.
- Robinson DP, Hall OJ, Nilles TL, Bream JH, Klein SL (2014) 17 β -estradiol protects females against influenza by recruiting neutrophils and increasing virus-specific CD8 T cell responses in the lungs. *J Virol* 88:4711–4720.
- Vartanian T, Li Y, Zhao M, Stefansson K (1995) Interferon-gamma-induced oligodendrocyte cell death: Implications for the pathogenesis of multiple sclerosis. *Mol Med* 1:732–743.
- Balabanov R, Strand K, Kemper A, Lee JY, Popko B (2006) Suppressor of cytokine signaling 1 expression protects oligodendrocytes from the deleterious effects of interferon-gamma. *J Neurosci* 26:5143–5152.
- Gao X, et al. (2000) Interferon-gamma protects against cuprizone-induced demyelination. *Mol Cell Neurosci* 16:338–349.
- Mason JL, Suzuki K, Chaplin DD, Matsushima GK (2001) Interleukin-1beta promotes repair of the CNS. *J Neurosci* 21:7046–7052.
- Brown CM, Mulcahey TA, Filipek NC, Wise PM (2010) Production of proinflammatory cytokines and chemokines during neuroinflammation: Novel roles for estrogen receptors alpha and beta. *Endocrinology* 151:4916–4925.
- Mason JL, Xuan S, Dragatsis I, Efstratiadis A, Goldman JE (2003) Insulin-like growth factor (IGF) signaling through type 1 IGF receptor plays an important role in remyelination. *J Neurosci* 23:7710–7718.
- Sørensen TL, et al. (2002) Multiple sclerosis: A study of CXCL10 and CXCR3 co-localization in the inflamed central nervous system. *J Neuroimmunol* 127:59–68.
- Tirotta E, Ransohoff RM, Lane TE (2011) CXCR2 signaling protects oligodendrocyte progenitor cells from IFN- γ /CXCL10-mediated apoptosis. *Glia* 59:1518–1528.
- Nguyen D, Stangel M (2001) Expression of the chemokine receptors CXCR1 and CXCR2 in rat oligodendroglial cells. *Brain Res Dev Brain Res* 128:77–81.
- Filipovic R, Jakovcevic I, Zecevic N (2003) GRO-alpha and CXCR2 in the human fetal brain and multiple sclerosis lesions. *Dev Neurosci* 25:279–290.
- Hickey DK, Fahey JV, Wira CR (2013) Estrogen receptor α antagonists mediate changes in CCL20 and CXCL1 secretions in the murine female reproductive tract. *Am J Reprod Immunol* 69:159–167.
- Lei ZB, et al. (2003) Effect of estradiol on chemokine receptor CXCR2 expression in rats: Implications for atherosclerosis. *Acta Pharmacol Sin* 24:670–674.
- Lasarte S, et al. (2016) Sex hormones coordinate neutrophil immunity in the vagina by controlling chemokine gradients. *J Infect Dis* 213:476–484.
- Haddad SN, Wira CR (2014) Estradiol regulation of constitutive and keratinocyte growth factor-induced CCL20 and CXCL1 secretion by mouse uterine epithelial cells. *Am J Reprod Immunol* 72:34–44.
- Omari KMJ, John G, Lango R, Raine CS (2006) Role for CXCR2 and CXCL1 on glia in multiple sclerosis. *Glia* 53:24–31.
- Omari KM, Lutz SE, Santambrogio L, Lira SA, Raine CS (2009) Neuroprotection and remyelination after autoimmune demyelination in mice that inducibly overexpress CXCL1. *Am J Pathol* 174:164–176.
- Hasselmann JPC, Karim H, Khalaj AJ, Ghosh S, Tiwari-Woodruff SK (2017) Consistent induction of chronic experimental autoimmune encephalomyelitis in C57BL/6 mice for the longitudinal study of pathology and repair. *J Neurosci Methods* 284:71–84.
- Moore S, et al. (2013) Therapeutic laquinimod treatment decreases inflammation, initiates axon remyelination, and improves motor deficit in a mouse model of multiple sclerosis. *Brain Behav* 3:664–682.
- Pino PA, Cardona AE (2011) Isolation of brain and spinal cord mononuclear cells using percoll gradients. *J Vis Exp*, 10.3791/2348.
- Tiwari-Woodruff S, et al. (2006) K+ channel KV3.1 associates with OSP/claudin-11 and regulates oligodendrocyte development. *Am J Physiol Cell Physiol* 291:C687–C698.