NEWS

uegjournal WILEY

UEG position paper: Obesity and digestive health

THE SCALE OF THE ISSUE

The obesity epidemic is one of the most significant public health challenges in Europe today. Obesity rates have more than doubled in the last 40 years—over half of the EU's adult population and nearly one in three children in the World Health Organisation European region are overweight or obese.

OBESITY AND CHRONIC DIGESTIVE DISEASES

Obesity is a major risk factor for chronic digestive diseases, placing a substantial strain on European health systems.

Childhood obesity

- Childhood obesity is a significant risk factor for digestive and metabolic diseases later in life¹
- Non-alcoholic fatty liver disease (NAFLD) is a consequence of adolescent obesity, with the risk of developing liver cirrhosis^{2,3}

Obesity and functional digestive disorders

- Four in 10 adults suffer from one or more disorders of gut-brain interaction (DGBI),⁴ and people with obesity are even more likely to suffer DGBI symptoms⁵
- There is a strong association between obesity and gastroesophageal reflux disease⁶

Obesity and inflammatory bowel disease

- Twenty percent of paediatric⁷ and 40% of adult⁸ patients with IBD are overweight or obese
- A western-type diet is thought to contribute to obesity and IBD⁹
- Obesity and/or visceral adiposity may increase the risk for developing surgical complications, hospitalisation and post-operative recurrence in patients with IBD¹⁰⁻¹⁴

Obesity and liver disease

- NAFLD is common in people who are obese or morbidly obese (prevalence of 70%–95%)¹⁵
- NAFLD is a leading cause of liver-related mortality in Europe, predicted to become the leading cause of end-stage liver disease unless urgent action is taken^{16,17}

Obesity and digestive cancers

- Obesity is an established risk factor for multiple types of cancer,¹⁸ including pancreatic,^{19,20} liver,^{21,22} and colorectal²³ cancer
- In 2019, 4.6% of all cancer deaths were attributable to obesity²⁴

PREVENTION AND TREATMENT

The increase in people who are overweight and obese across Europe is alarming and has been exacerbated by the COVID-19 pandemic.²⁵ There is an urgent need for preventive and therapeutic strategies at both individual and public health levels.

Lifestyle-related prevention

- Promoting healthy lifestyle choices, such as following a Mediterranean diet and reducing alcohol consumption
- Tackling the over-consumption of ultra-processed foods, often high in calories, salt, sugar, and saturated fats
- Promoting physical activity to improve overall health and maintain a healthy body weight

Pharmacological therapies

- Despite the high prevalence of NAFLD in Europe, no effective pharmacological treatment is currently approved for use
- Existing therapies mainly focus on metabolic disorders associated with NAFLD^{26,27}

Endoscopic and metabolic therapies

 Patients with obesity who receive intragastric balloons or endoscopic sleeve gastroplasty and lifestyle intervention show a higher mean total bodyweight loss compared to those who receive lifestyle intervention only^{28,29}

The role of bariatric surgery

- Bariatric surgery should be considered in patients with morbid obesity
- Bariatric surgery is associated with greater and faster weight loss and, in most cases, with improved long-term results compared with conventional therapy. This has the potential to reduce all-cause mortality and the development of obesity-related

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

^{© 2022} The Authors. United European Gastroenterology Journal published by Wiley Periodicals LLC. on behalf of United European Gastroenterology.

diseases.³⁰ Additionally, this decreases the incidence of obesityrelated cancer and cancer-related mortality^{31,32}

 It is important that bariatric surgery be accompanied by structured behavioural therapies, nutritional programmes and lifestyle changes to reduce calorie intake and increase exercise

UEG RECOMMENDATIONS

In accordance with World Health Organisation guidelines, UEG recommends the following policy interventions:

- Effective prevention strategies targeting food, including food reformulation, mandatory EU-wide front-of-pack nutrition labelling, food marketing restrictions (especially to protect children), taxation of unhealthy food high in fats, sugar and salt, increased availability and affordability of healthy foods to influence consumer choices.
- Coherent action throughout EU and member states and involvement of all relevant sectors to create environments for people and communities that are conducive to limiting the production and consumption of unhealthy foods.
- Population prevention efforts must target all age groups throughout the life course, with a particular focus on children as early life interventions can have a substantial impact.
- 4. Mass media awareness campaigns and educational programmes in schools, with a focus on improving overall health outcomes rather than on weight loss alone and the inclusion of healthy school meals and integration of food science into the school curriculum.
- 5. Consistent uptake of clinical practice guidelines for healthcare to deliver effective, evidence-informed obesity care and improve the quality of healthcare and patient outcomes.
- 6. **Research and development investment** to provide innovative prevention and treatment options.

Patrizia Burra¹ D Marianna Arvanitakis^{2,3} Jorge Amil Dias⁴ Michael Bretthauer^{5,6} Ana Dugic^{7,8} Daniel Hartmann⁹ Patrick Michl¹⁰ Thomas Seufferlein¹¹ Joana Torres^{12,13} Hans Törnblom¹⁴ D Monique E. van Leerdam¹⁵ Shira Zelber-Sagi^{16,17} Andreea Botos¹⁸

¹Multivisceral Transplant Unit, Department of Surgery, Oncology and Gastroenterology, Padua University Hospital, Padua, Italy

- ²Department of Gastroenterology, Hepatopancreatology and GI Oncology, Erasme University Hospital, Université Libre de Bruxelles, Brussels, Belgium
- ³Nutrition Team, Erasme University Hospital, Université Libre de
 - Bruxelles, Brussels, Belgium
- ⁴Department of Pediatrics, Hospital S. João, Porto, Portugal ⁵Department of Transplantation Medicine, Clinical Effectiveness Research Group, Oslo University Hospital, Oslo, Norway
- ⁶Clinical Effectiveness Research Group, Institute of Health and
 - Society, University of Oslo, Oslo, Norway
- ⁷Department of Gastroenterology, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Medizincampus Oberfranken, Bayreuth, Germany
 - ⁸Department of Medicine, Karolinska Institutet, Stockholm, Sweden
 - ⁹Department of Surgery, Klinikum rechts der Isar Technische Universität Munich, Munich, Germany
 - ¹⁰Department of Internal Medicine I, Martin-Luther University Halle-Wittenberg, Halle, Germany
- ¹¹Department of Internal Medicine I, Ulm University Hospital, Ulm, Germany
 - ¹²Division of Gastroenterology, Hospital Beatriz Ângelo, Loures, Portugal
- ¹³Faculdade de Medicina, Universidade de Lisboa, Lisboa, Portugal ¹⁴Department of Molecular and Clinical Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg,
 - Gothenburg, Sweden
- ¹⁵Department of Gastroenterology, Netherlands Cancer Institute, Amsterdam, The Netherlands
- ¹⁶School of Public Health, Faculty of Social Welfare and Health Sciences, University of Haifa, Haifa, Israel
- ¹⁷Department of Gastroenterology, Tel Aviv Medical Center, Tel Aviv, Israel
 - ¹⁸United European Gastroenterology (UEG), Vienna, Austria

Correspondence

Patrizia Burra, Multivisceral Transplant Unit, Department of Surgery, Oncology and Gastroenterology, Padua University Hospital, Padua, Italy.

Email: burra@unipd.it

ORCID

Patrizia Burra b https://orcid.org/0000-0002-8791-191X Hans Törnblom b https://orcid.org/0000-0003-2117-9874

REFERENCES

- Barton M. Childhood obesity: a life-long health risk. Acta Pharmacol Sin. 2012;33(2):189–93. https://doi.org/10.1038/aps.2011.204
- Kumar S, Kelly AS. Review of childhood obesity: from epidemiology, etiology, and comorbidities to clinical assessment and treatment. Mayo Clin Proc. 2017;92(2):251–65. https://doi.org/10.1016/j. mayocp.2016.09.017
- Shaunak M, Byrne CD, Davis N, Afolabi P, Faust SN, Davies JH. Nonalcoholic fatty liver disease and childhood obesity. Arch Dis Child.

2021;106(1):3-8. https://doi.org/10.1136/archdischild-2019-31 8063

- Sperber AD, Bangdiwala SI, Drossman DA, Ghoshal UC, Simren M, Tack J, et al. Worldwide prevalence and burden of functional gastrointestinal disorders, results of Rome Foundation Global Study. Gastroenterology. 2021;160(1):99–114.e3. https://doi.org/10.1053/ j.gastro.2020.04.014
- Aro P, Ronkainen J, Talley NJ, Storskrubb T, Bolling-Sternevald E, Agréus L. Body mass index and chronic unexplained gastrointestinal symptoms: an adult endoscopic population based study. Gut. 2005;54(10):1377–83. https://doi.org/10.1136/gut.2004.057497
- Singh S, Sharma AN, Murad MH, Buttar NS, El-Serag HB, Katzka DA, et al. Central adiposity is associated with increased risk of esophageal inflammation, metaplasia, and adenocarcinoma: a systematic review and meta-analysis. Clin Gastroenterol Hepatol. 2013;11(11):1399–412.e7. https://doi.org/10.1016/j.cgh.2013. 05.009
- Nic Suibhne T, Raftery TC, McMahon O, Walsh C, O'Morain C, O'Sullivan M. High prevalence of over weight and obesity in adults with Crohn's disease: associations with disease and lifestyle factors. J Crohns Colitis. 2013;7(7):e241–8. https://doi.org/10.1016/j.cro hns.2012.09.009
- Long MD, Crandall WV, Leibowitz IH, Duffy L, del Rosario F, Kim SC, et al. Prevalence and epidemiology of over weight and obesity in children with inflammatory bowel disease. Inflamm Bowel Dis. 2011;17(10):2162–8. https://doi.org/10.1002/ibd.21585
- Inczefi O, Bacsur P, Resal T, Keresztes C, Molnar T. The influence of nutrition on intestinal permeability and the microbiome in health and disease. Front Nutr. 2022;9:718710. https://doi.org/10.3389/ fnut.2022.718710
- Jiang K, Chen B, Lou D, Zhang M, Shi Y, Dai W, et al. Systematic review and meta-analysis: association between obesity/over weight and surgical complications in IBD. Int J Colorectal Dis. 2022;37(7): 1485–96. https://doi.org/10.1007/s00384-022-04190-y
- Li Y, Zhu W, Gong J, Zhang W, Gu L, Guo Z, et al. Visceral fat area is associated with a high risk for early postoperative recurrence in Crohn's disease. Colorectal Dis. 2015;17(3):225–34. https://doi.org/ 10.1111/codi.12798
- El-Dallal M, Stein DJ, Raita Y, Feuerstein JD. The impact of obesity on hospitalized patients with ulcerative colitis. Ann Gastroenterol. 2021;34(2):196–201. https://doi.org/10.20524/aog.2021.0592
- Grillot J, D'Engremont C, Parmentier AL, Lakkis Z, Piton G, Cazaux D, et al. Sarcopenia and visceral obesity assessed by computed tomography are associated with adverse outcomes in patients with Crohn's disease. Clin Nutr. 2020;39(10):3024–30. https://doi.org/ 10.1016/j.clnu.2020.01.001
- Gu P, Chhabra A, Chittajallu P, Chang C, Mendez D, Gilman A, et al. Visceral adipose tissue volumetrics inform odds of treatment response and risk of subsequent surger y in IBD patients starting antitumor necrosis factor therapy. Inflamm Bowel Dis. 2022;28(5): 657–66. https://doi.org/10.1093/ibd/izab167
- Henry L, Paik J, Younossi ZM. Review article: the epidemiologic burden of non-alcoholic fatty liver disease across the world. Aliment Pharmacol Ther. 2022;56(6):942–56. https://doi.org/10.1111/apt. 17158
- Pais R, Barritt AS, Calmus Y, Scatton O, Runge T, Lebray P, et al. NAFLD and liver transplantation: current burden and expected challenges. J Hepatol. 2016;65(6):1245–57. https://doi.org/10.1016/ j.jhep.2016.07.033
- Schattenberg JM, Lazarus JV, Newsome PN, Serfaty L, Aghemo A, Augustin S, et al. Disease burden and economic impact of diagnosed non-alcoholic steatohepatitis in five European countries in 2018: a

cost-of-illness analysis. Liver Int. 2021;41(6):1227-42. https://doi. org/10.1111/liv.14825

- Fang Z, Giovannucci EL. The timing of adiposity and changes in the life course on the risk of cancer. Cancer Metastasis Rev. 2022;41(3): 471–89. https://doi.org/10.1007/s10555-022-10054-2
- Michl P, Lohr M, Neoptolemos JP, Capurso G, Rebours V, Malats N, et al. UEG position paper on pancreatic cancer. Bringing pancreatic cancer to the 21st century: prevent, detect, and treat the disease earlier and better. United Eur Gastroenterol J. 2021;9(7):860–71. https://doi.org/10.1002/ueg2.12123
- Stolzenberg-Solomon RZ, Schairer C, Moore S, Hollenbeck A, Silverman DT. Lifetime adiposity and risk of pancreatic cancer in the NIH-A ARP Diet and Health Study cohort. Am J Clin Nutr. 2013; 98(4):1057–65. https://doi.org/10.3945/ajcn.113.058123
- Yang C, Lu Y, Xia H, Liu H, Pan D, Yang X, et al. Excess body weight and the risk of liver cancer: systematic review and a meta-analysis of cohort studies. Nutr Cancer. 2020;72(7):1085–97. https://doi.org/ 10.1080/01635581.2019.1664602
- Sohn W, Lee HW, Lee S, Lim JH, Lee MW, Park CH, et al. Obesity and the risk of primary liver cancer: a systematic review and metaanalysis. Clin Mol Hepatol. 2021;27(1):157–74. https://doi.org/10. 3350/cmh.2020.0176
- Johnson CM, Wei C, Ensor JE, Smolenski DJ, Amos CI, Levin B, et al. Meta-analyses of colorectal cancer risk factors. Cancer Causes Control. 2013;24(6):1207–22. https://doi.org/10.1007/s10552-013-0201-5
- Zhi X, Kuang X, Liu K, Li J. The global burden and temporal trend of cancer attributable to high body mass index: estimates from the Global Burden of Disease Study 2019. Front Nutr. 2022;9:918330. https://doi.org/10.3389/fnut.2022.918330
- 25. Horton R. Offline: COVID-19 is not a pandemic. Lancet. 2020; 396(10255):874. https://doi.org/10.1016/s0140-6736(20)32000-6
- Francque S, Vonghia L. Pharmacological treatment for non-alcoholic fatty liver disease. Adv Ther. 2019;36(5):1052–74. https://doi.org/ 10.1007/s12325-019-00898-6
- Yan H, Wu W, Chang X, Xia M, Ma S, Wang L, et al. Gender differences in the efficacy of pioglitazone treatment in nonalcoholic fatty liver disease patients with abnormal glucose metabolism. Biol Sex Differ. 2021;12(1):1–8. https://doi.org/10.1186/s13293-020-00344-1
- Abu Dayyeh BK, Maselli DB, Rapaka B, Lavin T, Noar M, Hussan H, et al. Adjustable intragastric balloon for treatment of obesity: a multicentre, open-label, randomised clinical trial. Lancet. 2021; 398(10315):1965–73. https://doi.org/10.1016/s0140-6736(21)02 394-1
- Abu Dayyeh BK, Bazerbachi F, Vargas EJ, Sharaiha RZ, Thompson CC, Thaemert BC, et al. Endoscopic sleeve gastroplasty for treatment of class 1 and 2 obesity (MERIT): a prospective, multicentre, randomised trial. Lancet. 2022;400(10350):441–51. https://doi.org/ 10.1016/s0140-6736(22)01280-6
- Wiggins T, Guidozzi N, Welbourn R, Ahmed AR, Markar SR. Association of bariatric surgery with all-cause mortality and incidence of obesity-related disease at a population level: a systematic review and meta-analysis. PLoS Med. 2020;17(7):e1003206. https://doi.org/ 10.1371/journal.pmed.1003206
- Aminian A, Wilson R, Al-Kurd A, Tu C, Milinovich A, Kroh M, et al. Association of bariatric surgery with cancer risk and mortality in adults with obesity. JAMA. 2022;327(24):2423–33. https://doi.org/ 10.1001/jama.2022.9009
- Rustgi VK, Li Y, Gupta K, Minacapelli CD, Bhurwal A, Catalano C, et al. Bariatric surgery reduces cancer risk in adults with nonalcoholic fatty liver disease and severe obesity. Gastroenterology. 2021;161(1): 171–84.e10. https://doi.org/10.1053/j.gastro.2021.03.021