

Structure, Biosynthesis, and Localization of Dipeptidyl Aminopeptidase B, an Integral Membrane Glycoprotein of the Yeast Vacuole

Christopher J. Roberts, Gabriele Pohlig, Joel H. Rothman, and Tom H. Stevens

Institute of Molecular Biology, University of Oregon, Eugene, Oregon 97403

Abstract. We have characterized the structure, biogenesis, and localization of dipeptidyl aminopeptidase B (DPAP B), a membrane protein of the yeast vacuole. An antibody specific for DPAP B recognizes a 120-kD glycoprotein in yeast that behaves like an integral membrane protein in that it is not removed from membranes by high pH Na_2CO_3 treatment. Inspection of the deduced amino acid sequence of DPAP B reveals a hydrophobic domain near the NH_2 terminus that could potentially span a lipid bilayer. The in vitro enzymatic activity and apparent molecular weight of DPAP B are unaffected by the allelic state of *PEP4*, a gene essential for the proteolytic activation of a number of soluble vacuolar hydrolases. DPAP B is synthesized as a glycosylated precursor that is converted to the mature 120-kD species by carbohydrate addition. The precursor

form of DPAP B accumulates in *sec* mutants (Novick, P., C. Field, and R. Schekman. 1980. *Cell*. 21:205–215) that are blocked at the ER (*sec18*) or Golgi apparatus (*sec7*), but not at secretory vesicles (*sec1*). Immunolocalization of DPAP B in wild-type or *sec1* mutant cells shows that the protein resides in the vacuolar membrane. However, it is present in non-vacuolar compartments in *sec18* and *sec7* cells, confirming that the delivery of DPAP B is blocked in these mutants. Interestingly, DPAP B appears to stain the nuclear envelope in a *sec18* mutant, which is consistent with the accumulation of DPAP B in the ER membrane at the restrictive temperature. These results suggest that soluble and membrane-bound vacuolar proteins use the same stages of the secretory pathway for their transport.

THE vacuole of the yeast *Saccharomyces cerevisiae* is considered to be the equivalent of the lysosome of mammalian cells because it contains a large number of hydrolytic enzymes and has an acidic pH (reviewed in Rothman and Stevens, 1988). The biosynthesis and sorting of soluble vacuolar hydrolases, such as carboxypeptidase Y (CPY)¹ and proteinase A, have been examined. The localization of soluble proteins to the vacuole requires the early stages of the secretory pathway; i.e., ER and Golgi apparatus (Stevens et al., 1982). In addition, localization determinants on CPY and proteinase A, as well as genes necessary for the efficient sorting of these proteins, have been identified (Valls et al., 1987; Johnson et al., 1987; Klionsky et al., 1988; Rothman and Stevens, 1986; Bankaitis et al., 1986; Robinson et al., 1988; Rothman et al., 1989).

Much less is known about the sorting of membrane proteins to the vacuole. Studies on mammalian cells indicate that soluble and membrane-bound lysosomal proteins are sorted

by distinct mechanisms (von Figura and Hasilik, 1986). Several lysosomal membrane proteins have been identified (Chen et al., 1985; Lewis et al., 1985; Barriocanal et al., 1986; Lippincott-Schwartz and Fambrough, 1986; Waheed et al., 1988), and of those examined, none possess the mannose-6-phosphate moiety required for sorting soluble lysosomal proteins (von Figura and Hasilik, 1986). Furthermore, several membrane proteins are found at normal levels in the lysosomes of I-cell fibroblasts, which mis-sort soluble lysosomal hydrolases to the cell surface.

Several activities associated with the vacuolar membrane of yeast have been characterized (Rothman and Stevens, 1988), including transport systems for Ca^{2+} (Ohsumi and Anraku, 1981) and basic amino acids (Ohsumi and Anraku, 1983), a protein-translocating ATPase (Uchida et al., 1985; Kane et al., 1989), an α -mannosidase (Opheim, 1978), and dipeptidyl aminopeptidase (DPAP) B (Suarez Rendueles et al., 1981; Julius et al., 1983; Bordallo et al., 1984). Nothing is known about the transport or posttranslational modifications for any of the vacuolar membrane proteins. Analysis of the localization of α -mannosidase and DPAP B in yeast mutants that mis-sort soluble vacuolar proteins suggested that most of these mutants do not mis-sort membrane-bound vacuolar proteins (Bankaitis et al., 1986; Rothman and

J. H. Rothman's present address is MRC Laboratory of Molecular Biology, Hills Road, Cambridge CB2 2QH, United Kingdom.

1. *Abbreviations used in this paper:* CPY, carboxypeptidase Y; DPAP, dipeptidyl aminopeptidase; endo F, endoglycosidase F; N-linked, asparagine-linked.

Stevens, 1986; Robinson et al., 1988; Rothman and Stevens, unpublished results). Thus, soluble and membrane-bound proteins appear to be directed to the vacuole by distinct mechanisms.

As a first step in the study of vacuolar membrane protein sorting and transport, we have characterized the structure, biogenesis, and localization of DPAP B. We show that DPAP B is an integral membrane protein that, like soluble vacuolar proteins, uses the early stages of the secretory pathway for its transport to the vacuole.

Materials and Methods

Strains, Growth Conditions, and Materials

The genotypes of the *Escherichia coli* and *S. cerevisiae* strains used in these studies are listed in Table I. Yeast cultures were grown in minimal medium supplemented with the appropriate nutrients as previously described (Stevens et al., 1986).

Restriction endonucleases and other enzymes used for cloning and sequencing were from New England Biolabs (Beverly, MA), Bethesda Research Laboratories (Gaithersburg, MD), Pharmacia Fine Chemicals (Piscataway, NJ), or Boehringer Mannheim Biochemicals (Indianapolis, IN). ¹²⁵I-protein A was from Amersham Corp. (Arlington Heights, IL) and carrier-free ³⁵S-H₂SO₄ was from ICN Biomedicals Inc. (Irvine, CA). IgG Sorb was from The Enzyme Center (Boston, MA), endoglycosidase F (endo F) was from Boehringer Mannheim Biochemicals, and FITC-conjugated sheep anti-rabbit IgG was from Cappel Laboratories (Malvern, PA). All other reagents used in these experiments were from Sigma Chemical Co. (St. Louis, MO).

Plasmid Constructions, Recombinant DNA Methodology, and DNA Sequence Analysis

Restriction endonuclease and Bal 31 exonuclease digestions and ligations were performed as recommended by the suppliers. Plasmid purification, agarose gel electrophoresis, and DNA-mediated transformation of *E. coli* were performed according to standard procedures (Maniatis et al., 1982). Plasmids were introduced into yeast cells by the lithium acetate transforma-

tion method (Ito et al., 1983). *E. coli* strain MC1061 was used for all plasmid manipulations and JM103 was used for M13 phage work.

DNA sequencing was performed by the chain termination method of Sanger et al. (1977) using M13mpl8 and mpl9 vectors (Yanisch-Perron et al., 1985). In one case, sequencing was primed using a synthetic oligonucleotide corresponding to nucleotides 441-459 of DAP2 (see Fig. 2), prepared at the University of Oregon Biotechnology Laboratory on a DNA synthesizer (model 380B; Applied Biosystems, Inc., Foster City, CA) as described (Ito et al., 1982). DNA and amino acid sequence analyses were performed using the sequence analysis software package (version 4) of the University of Wisconsin Genetics Computer Group.

Plasmid pRG1 was constructed by inserting the 4.7-kb Bam HI-Pst I (blunt-ended by treating with T4 DNA polymerase) DAP2 fragment (Fig. 1), isolated from the original DAP2 plasmid YEpl3-GS13-4 (Julius et al., 1983), into the Bam HI-Sma I sites of pUC12. Plasmid pGP3, a multicopy 2-μm plasmid containing the DAP2 gene, was constructed by inserting the 4.7-kb Bam HI-Pst I (blunt-ended) fragment from pRG1 into the Bam HI-Pvu II sites of YEpl24.

A chromosomal DAP2 deletion was constructed by transposing the genomic copy with the cloned gene into which the LEU2 gene was inserted in place of 1.3 kb of the DAP2 coding region (Rothstein, 1983; Fig. 1). Plasmid pRG1 was digested with BstE II and Kpn I, treated with T4 DNA polymerase to make blunt ends, and ligated with a 2.1-kb Hpa I fragment containing the LEU2 gene. The resulting plasmid (pGP2) was cut with Bam HI and Pst I and used to transform the yeast strain JHRY20-2C (leu2-3, leu2-112). Stable LEU⁺ transformants were purified and screened for the lack of thermolabile dipeptidyl aminopeptidase activity, assayed as described previously (Julius et al., 1983).

Plasmid pCJR6, a multicopy 2-μm yeast vector containing the DPAP B coding region fused to the GALI promoter, was constructed by inserting a 400-bp Eco RI-Hind III GALI promoter fragment (Johnston and Davis, 1984) into the Eco RI and Hind III sites of pSEY8, and subsequently subcloning into the Hind III site of this plasmid (pCJR5) the 2.9 kbp Hind III fragment of DAP2 (Fig. 1), resulting in the fusion of the GALI promoter at position -90 of DAP2.

DPAP B antigen for production of antiserum was made in *E. coli* by fusing a portion of the DAP2 coding region into the *E. coli* expression vector pHSe5, which contains the T4 phage lysozyme gene under the control of the TAC promoter (Muchmore et al., 1988; kindly provided by R. Dahlquist, University of Oregon). Plasmid pRG1 was digested with Stu I, treated briefly with Bal 31 exonuclease, and further digested with Hind III. Fragments of ~1,650 bp were cloned into the Sna BI-Hind III sites of pHSe5,

Table I. Strains Used in This Study

Yeast strain	Genotype	Source or reference
X2180-1AL1	<i>MATa gal2 lys2 mal mel</i>	Derived from X2180
HMSF176	<i>MATa sec18 gal2</i>	Novick et al., 1980
SF294-2B	<i>MATa sec7 gal2</i>	Novick and Schekman, 1983
HMSF1	<i>MATa sec1 gal2</i>	Novick et al., 1980
SEY5186	<i>MATa sec18 ura3-52 leu2-3 leu2-112</i>	Emr et al., 1984
CJRY21-3BΔ1	<i>MATa sec18 ura3-52 leu2-3 leu2-112</i> <i>met14 ade1 ade2-1 trp1 dap2-Δ2::LEU2 GAL⁺</i>	This study
CJRY22-6BΔ1	<i>MATa sec1 ura3-52 leu2-3 leu2-112</i> <i>met14 ade1 ade2-1 dap2-Δ2::LEU2 GAL⁺</i>	This study
CJRY23-2AΔ1	<i>MATa sec7 ura3-52 leu2-3 leu2-112</i> <i>met14 ade1 his3 dap2-Δ2::LEU2 GAL⁺</i>	This study
JHRY20-1AΔ1	<i>MATa ura3-52 leu2-3 leu2-112</i> <i>his3-Δ200 lys2-801 pep4-3 can1 ste13-Δ1::LEU2</i>	Rothman, 1988
JHRY20-2C	<i>MATa ura3-52 leu2-3 leu2-112 his3-Δ200</i>	Rothman, 1988
JHRY20-2CΔ2	<i>MATa ura3-52 leu2-3 leu2-112</i> <i>his3-Δ200 pep4-Δ2::LEU2</i>	Derived from JHRY20-2C
JHRY20-2CΔ3	<i>MATa ura3-52 leu2-3 leu2-112</i> <i>his3-Δ200 dap2-Δ2::LEU2</i>	Derived from JHRY20-2C
<i>E. coli</i>		
MC1061	F ⁻ <i>hsdR hsdM⁺ araD139 (araABOIC-leu)7679</i> (<i>lac</i>)X74 <i>galU galK rpsL</i>	Casadaban and Cohen, 1980
JM103	F ['] Δ(<i>lac pro</i>) <i>thi strA endA sbcB15 supE/F['] traD36</i> <i>proAB⁺ lac⁺ lacZΔM15</i>	Messing, 1983

resulting in the production of a hybrid protein consisting of the NH₂-terminal seven residues of T4 phage lysozyme fused to the COOH-terminal half of DPAP B. Transformants were screened for an isopropyl β -D-thiogalactopyranoside- (IPTG) inducible protein species by SDS-PAGE (Laemmli, 1970) of total *E. coli* cell extracts. A plasmid was isolated (pRG2) which produced a large amount (~5% of total cell protein) of a 55-kD species upon induction with isopropyl β -D-thiogalactopyranoside.

Production of DPAP B Antigen and Generation of DPAP B Antiserum

E. coli cells containing plasmid pRG2 were grown in LB broth plus ampicillin to saturation, then diluted 1:100 in 250 ml of fresh medium, and grown for 1 h at 37°C. Isopropyl β -D-thiogalactopyranoside was added to 1 mM, and incubation was continued for 3 h before collecting the cells by centrifugation. The cells were lysed and the fusion protein was isolated from the insoluble fraction essentially as previously described (Kleid et al., 1981). An SDS-PAGE gel slice containing the fusion protein was resuspended in SDS elution buffer (1% SDS, 50 mM Tris-HCl, pH 8.0, 1 mM EDTA, 1 mM DTT, 100 mM NaCl) in a homogenizer (Dounce; Wheaton Instruments Div., Millville, NJ), incubated for 2 h at 37°C, and centrifuged at 12,000 g for 10 min. The supernatant was saved, and the pellet was resuspended in 5 ml of elution buffer and treated as above. The supernatants were combined, and the protein was precipitated by adding 4 vol of acetone, incubating for 30 min at -80°C, and centrifuging at 15,000 g for 10 min.

A DPAP B-specific antibody was prepared by injecting DPAP B antigen into New Zealand White Rabbits (250 μ g DPAP B antigen per rabbit per injection) essentially as described (Vaitukaitis, 1981). These antibodies were affinity purified as previously described (Stevens et al., 1982). Antiserum to phosphoglycerate kinase was prepared as described (Rothman et al., 1986).

Alkaline Sodium Carbonate Fractionation

Yeast cells containing plasmid pGP3 were grown in minimal medium at 30°C to mid log phase. 12 OD₆₀₀ U of cells (1.2×10^8 cells) were spheroplasted as described previously (Stevens et al., 1986), lysed in 50 mM sodium phosphate, pH 7.5, 2 mM EDTA, and 0.5 mM PMSF, and diluted 100-fold in ice-cold 100 mM sodium carbonate, pH 11.5. An aliquot representing the whole extract fraction was removed, and after 30 min on ice the lysate was centrifuged for 3 h at 100,000 g. The supernatant and whole extract fractions were neutralized with 1 N acetic acid, then precipitated in 5% TCA on ice. Precipitated protein was pelleted, washed twice in diethylether, dried, resuspended in sample buffer, and heated at 70°C until dissolved. The pellet fraction was homogenized in 50 mM Tris-HCl, pH 6.8, and solubilized by addition of SDS to a final concentration of 4%. The sample was incubated at 60°C until complete solubilization had occurred; i.e., the suspension became completely transparent. Sample buffer (without SDS) was added, and the samples were subjected to SDS-PAGE on an 8% polyacrylamide gel. Electrophoresis of the gel onto nitrocellulose was carried out according to Burnette (1981) using a Trans-blot Apparatus from Bio-Rad Laboratories (Richmond, CA). The transferred proteins were detected by probing the nitrocellulose with antibodies to the appropriate antigens and subsequent labeling with 0.3 μ Ci of ¹²⁵I-protein A. The dried nitrocellulose was exposed to film (XAR-5; Eastman Kodak Co., Rochester, NY) at -80°C.

Radiolabeling, Immunoprecipitation, and Endo F Treatment

Yeast cells were grown in minimal media plus 50 μ M Na₂SO₄ (Stevens et al., 1986) to mid log phase, spun down in a clinical centrifuge, washed once and resuspended in sulfate-free media, and incubated at the appropriate labeling temperature for 15–30 min at 8 OD₆₀₀ U/ml. ³⁵S-H₂SO₄ was added (0.5–1.0 mCi) and cells were labeled and then chased with 10 mM Na₂SO₄ (except for the experiments in Fig. 5 B, which were chased with 10 mM Na₂SO₄, 1 mM cysteine, 1 mM methionine, and 0.1 mg/ml cycloheximide). 2 OD₆₀₀ U of cells were then sedimented, converted to spheroplasts, and lysed in 0.1 ml of 1% SDS by heating for 3 min at 100°C. 0.9 ml of immunoprecipitation buffer (Stevens et al., 1986) was added, followed by 0.1 ml of IgG Sorb (prepared as suggested by the manufacturer). The mixture was centrifuged 15 min in a microfuge (model 235B; Fisher Scientific Co., Pittsburgh, PA) and the supernatant was treated with 2 μ l of affinity purified DPAP B antibody and incubated 60 min on ice. The immune complexes were sedimented with IgG Sorb and washed as before

(Stevens et al., 1982), and the final IgG Sorb cell pellet was resuspended in 0.1 ml of 1% SDS and heated 3 min at 100°C. 0.9 ml of immunoprecipitation buffer and 0.1 ml of IgG Sorb was added, and the immunoprecipitation protocol was repeated. Sample buffer was added to the final IgG Sorb pellet and the samples were heated for 3 min at 100°C and centrifuged. Supernatant fractions were transferred to new tubes and aliquots (normalized for the amount of ³⁵S-H₂SO₄ incorporated) were analyzed by SDS-PAGE and fluorography as described previously (Stevens et al., 1986).

Endo F treatment (Elder and Alexander, 1982) was carried out on immunoprecipitated samples of DPAP B essentially as described (Stevens et al., 1986). A time course of endo F treatment was performed by adding 300 mU of endo F to DPAP B immunoprecipitated from SEY5186 cells (containing plasmid pGP3) and withdrawing aliquots at the times indicated in Fig. 6.

Immunofluorescence Microscopy

Yeast cells containing a disruption of the chromosomal *DAP2* locus and transformed with the plasmid pCJR6 were grown in minimal media plus 2% raffinose to an OD₆₀₀ of 1.0, and galactose was added directly to the cultures to a final concentration of 3%. The cultures were then incubated at either 34°C (*SEC⁺*, *sec18*, and *sec1*) or 38°C (*SEC⁺* and *sec7*) for 2 h and the cultures were put on ice. The cells were fixed with formaldehyde, spheroplasted, and prepared for immunofluorescence as described (Kilmartin and Adams, 1984; Adams and Pringle, 1984; Wittenberg et al., 1987). In some cases, fixed spheroplasts were treated with SDS by resuspending in 1 ml of 1.2 M sorbitol/PBS (prepared as described by Adams and Pringle, 1984) plus 5% SDS, mixed gently, centrifuged at 8,000 g, and washed twice in 1.2 M sorbitol. FITC-conjugated sheep anti-rabbit IgG second antibody was used for staining DPAP B antibody. Nuclei were stained with 4',6'-diamidino-2-phenyl-indole (DAPI) as described (Kilmartin and Adams, 1984).

Cells were viewed and photographed using a Zeiss Axioplan Photomicroscope (equipped for epifluorescence at excitation wavelengths appropriate for DAPI and FITC fluorescence) and film and developer (TMAX-400; Eastman Kodak Co.). Photomicrographs are shown at a magnification of 1,000.

Results

The *DAP2* Gene Encodes DPAP B

Yeast cells contain two membrane-associated DPAP activities that differ in their subcellular locations and thermal stabilities (Suarez Rendueles et al., 1981; Julius et al., 1983). DPAP B is a thermolabile enzyme associated with the vacuolar membrane (Bordallo et al., 1984). DPAP A, the *STE13* gene product, is a nonvacuolar, heat-stable enzyme that participates in the proteolytic maturation of the mating pheromone α -factor (Fuller et al., 1988). The *STE13* gene was cloned by complementation of the mating defect of a *MAT α*




Figure 1. Restriction map and nucleotide sequencing strategy of the *DAP2* gene. A linear map of the *DAP2* gene is depicted, with the horizontal arrows indicating direction and extent of sequence determination. The arrow originating in a solid circle corresponds to sequence determined from DNA synthesis primed by a synthetic oligonucleotide. The boxed region denotes the open reading frame, with the predicted initiating methionine codon at the left end. The hatched region shows the portion of *DAP2* coding sequence replaced by *LEU2* sequences in the *dap2-Δ2::LEU2* disruption (see Materials and Methods). Restriction sites that are relevant to the sequence analysis are indicated: B, Bam HI; H, Hind III; A, Acc I; Bs, BstE II; S, Stu I; K, Kpn I; E, Eco RI; Hc, Hinc II; P, Pst I.

-458 BamHI
GGATCCTGTTTGGTGGTGGTTCGTAGCCCTTGCAAGATTTTGAAGAAGTTTATAAGTTTGAGAAACACGGATATTACCATTTCTGAGGGTG

-360 GAAGGTTAAAGAAATGGCCTAATGTTAGCTTCAATTAAGAAGAAGCTGATAATAGCACTTCTACTATATAGCAGGTATGCCACTACCGTTATTGATGGTAAGATCAAAAGTTTAGG

-240 ATTCTAAACGCATAGTTGTAAGTTGATGTATATATATATATATATATATATTAATTAACAATAATATGCTCCGCCCAATTTTCTCCTTCAATACCGCGGAGGCGGTATTGAA

HindIII

-120 GGAAATAGACGAGAAATTCCTTATCAAGAAGCTTCCATCAAGTGACATAAGAAGTGCCGAAATTCGAAGTATCTTTTCAGAGAGTATTTTGCACATACCAATAAGCCAAATTACTC

1 ATG GAA GGT GGC GAA GAA GAA GTT GAG CGC ATT CCT GAT GAA CTT TTC GAT ACA AAA AAG AAG CAT TTG TTA GAT AAG CTC ATA AGG GTC
1 Met Glu Gly Gly Glu Glu Glu Val Glu Arg Ile Pro Asp Glu Leu Phe Asp Thr Lys Lys Lys His Leu Leu Asp Lys Leu Ile Arg Val

AccI

91 GGA ATA ATC CTT GTA CTC CTG ATA TGG GGC ACT GTT TTG TTG CTA AAA AGT ATA CCT CAC CAT TCA AAC ACA CCA GAT TAT CAA GAA CCC
31 Gly Ile Ile Leu Val Leu Leu Ile Trp Gly Thr Val Leu Leu Leu Lys Ser Ile Pro His His Ser Asn Thr Pro Asp Tyr Gln Glu Pro

181 AAC TCT AAT TAC ACC AAT GAT GGG AAA TTA AAG GTG TCG TTT TCT GTT GTA AGA AAC AAT ACA TTT CAT CCC AAA TAT CAC GAG CTG CAA
61 Asn Ser Asn Tyr Thr Asn Asp Gly Lys Leu Lys Val Ser Phe Ser Val Val Arg Asn Asn Thr Phe His Pro Lys Tyr His Glu Leu Gln

AccI

271 TGG ATT AGT GAC AAT AAA ATT GAA AGC AAC GAT CTA GGT CTT TAT GTT ACA TTT ATG AAC GAT AGT TAC GTT GTT AAA TCT GTC TAC GAT
91 Trp Ile Ser Asp Asn Lys Ile Glu Glu Arg Lys Phe Ser Asn Asp Leu Gly Leu Tyr Val Thr Phe Met Asn Asp Ser Tyr Val Val Lys Ser Val Tyr Asp

361 GAC TCA TAT AAT AAC GTT TTA TTA GAG GGA AAA ACT TTC ATT CAT AAC GGT CAA AAC CTC ACT GTG GAG TCA ATA ACT GCG TCT CCC GAC
121 Asp Ser Tyr Asn Asn Val Leu Leu Glu Gly Lys Thr Phe Ile His Asn Gly Gln Asn Leu Thr Val Glu Ser Ile Thr Ala Ser Pro Asp

AccI

451 CTG AAG CGG TTA TTA ATT AGA ACA AAT AGC GTA CAA AAT TGG AGA CAC TCA ACG TTT GGT TCT TAT TTT GTC TAC GAT AAG AGT AGC TCG
151 Leu Lys Arg Leu Leu Ile Arg Thr Asn Ser Val Gln Asn Trp Arg His Ser Thr Phe Gly Ser Tyr Phe Val Tyr Asp Lys Ser Ser Ser

541 TCC TTG AGG AGA TTG GAA ACA GTG GCT CTA GCA ATA TGG TCC CCT AAT TCT AAT AAT ATT GCA TAC GTC CAA GAT AAC AAT ATA TAT ATT
181 Ser Leu Arg Arg Leu Glu Thr Val Ala Leu Ala Ile Trp Ser Asn Ser Asn Ile Ala Tyr Val Gln Asp Asn Asn Ile Tyr Ile

631 TAT TCT GCT ATT TCC AAA AAG ACC ATA CGG GCT GTG ACC AAC GAC GGA AGC TCC TTT CTT TTT AAC GGT AAG CCA GAT TGG GTT TAC GAG
211 Tyr Ser Ala Ile Ser Lys Lys Thr Ile Arg Ala Val Thr Asn Asp Gly Ser Ser Phe Leu Phe Asn Gly Lys Pro Asp Trp Val Tyr Glu

BstEII

721 GAA GAA GTT TTT GAA GAC GAC AAG GCT GCG TGG TGG TCA CCA ACT GGT GAT TAC CTG GCA TTC TTG AAA ATT GAT GAA TCA GAA GTT GGT
241 Glu Glu Val Phe Glu Asp Asp Lys Ala Ala Trp Trp Ser Pro Thr Gly Asp Tyr Leu Ala Phe Leu Lys Ile Asp Glu Ser Glu Val Gly

811 GAG TTT ATC ATT CCA TAT TAT GTC CAA GAT GAA AAG GAT ATA TAC CCC GAA ATG CGC AGT ATC AAG TAT CCA AAA AGT GGC ACG CCA AAT
271 Glu Phe Ile Ile Pro Tyr Trp Lys Thr Val Gln Asp Gly Lys Ser Pro Glu Met Arg Ser Ile Lys Tyr Pro Lys Ser Gly Thr Pro Asn

901 CCT CAT GCA GAG CTA TGG GTT TAC AGT ATG AAA GAT GGA ACA TCG TTC CAT CCA AGA ATA AGT GGA AAT AAA AAA GAT GGA AGT CTG TTA
301 Pro His Ala Glu Leu Trp Val Tyr Ser Met Lys Asp Gly Thr Ser Phe His Pro Arg Ile Ser Gly Asn Lys Lys Asp Gly Ser Leu Leu

991 ATT ACT GAA GTT ACA TGG GTA GGA AAT GGA AAC GTT TTA GTT AAA ACT ACC GAT CGA TCC TCG GAC ATA TTG ACT GTG TTT TTG ATA GAT
331 Ile Thr Glu Val Thr Trp Val Gly Asn Gly Asn Val Leu Val Lys Thr Thr Asp Arg Ser Ser Asp Ile Leu Thr Val Phe Leu Ile Asp

1081 ACA ATT GCC AAA GAC TTC AAA CGT GGT AAG GAA CGA AAG TTT AAC GGA GGA TGG TGG GAG ATT ACT CAT AAT ACT CTG TTT ATT CCG GCA
361 Thr Ile Ala Lys Ser Phe Lys Arg Gly Lys Glu Arg Lys Phe Asn Gly Gly Trp Trp Glu Ile Thr His Asn Thr Leu Phe Ile Pro Ala

StuI

1171 AAT GAA ACA TTT GAT AGG CCT CAT AAT GGT TAT GTT GAT ATT CTT CCG ATT GGT GGT TAC AAT CAT TTG GCT TAT TTC GAA AAT AGT AAT
391 Asn Glu Thr Phe Asp Arg Pro His Asn Gly Tyr Val Asp Ile Leu Pro Ile Gly Gly Tyr Asn His Leu Ala Tyr Phe Glu Asn Ser Asn

1261 AGT TCA CAC TAT AAA ACA TTG ACA GAG GGG AAA TGG GAA GTG GTG AAT GGC CCA CTT GCC TTT GAT TCA ATG GAA AAT CGT CTT TAC TTC
421 Ser Ser His Tyr Lys Thr Leu Thr Glu Gly Lys Trp Glu Val Val Asn Gly Pro Leu Ala Phe Asp Ser Met Glu Asn Arg Leu Tyr Phe

BstEII

1351 ATT TCT ACA CGA AAG AGT TCA ACC GAA CGC CAC GTT TAC TAC ATA GAT TTA CGG TCA CCA AAT GAA ATT ATT GAA GTT ACT GAT ACT TCT
451 Ile Ser Thr Arg Lys Lys Ser Thr Glu Arg His Val Tyr Ser Phe Asn Leu Arg Ser Pro Asn Glu Ile Ile Glu Val Thr Thr Ser

1441 GAG GAT GGT GTC TAT GAT GTG TCC TTT TCT TCT GGT AGA AGG TTT GGT TTA CTC ACC TAT AAA GGA CCA AAA GTT CCA TAT CAA AAA ATT
481 Glu Asp Gly Val Tyr Asp Val Ser Phe Ser Ser Gly Arg Arg Phe Gly Leu Leu Thr Tyr Lys Gly Pro Lys Val Pro Tyr Gln Lys Ile

1531 GTG GAC TTC CAT TCT CGT AAA GCA GAA AAA TGC GAC AAA GGT AAT GTT TTA GGC AAA TCA CTA TAC CAT TTG GAA AAG AAC GAA GTA CTT
511 Val Asp Phe His Ser Arg Lys Ala Glu Lys Cys Asp Lys Gly Asn Val Leu Gly Lys Ser Leu Tyr His Leu Glu Lys Asn Glu Val Leu

KpnI

1621 ACC AAA ATT TTA GAA GAT TAT GCG GTA CCC AGA AAA TCA TTC AGG GAA TTG AAC CTA GGA AAG GAC GAA TTT GGA AAG GAT ATA CTA GTG
541 Thr Lys Ile Leu Glu Asp Tyr Ala Val Pro Arg Lys Ser Phe Arg Glu Lys Leu Gly Lys Asp Glu Phe Gly Lys Ile Leu Val

EcoRI

1711 ACC TCG TAT GAA ATC CTA CCA AAT GAT TTC GAT GAA ACG TTA AGT GAC CAC TAT CCT GTA TTT TTC TTT GCA TAT GGG GGA CCG AAT TCT
571 Asn Ser Tyr Glu Ile Leu Pro Asn Asp Phe Asp Glu Thr Leu Ser Asp His Tyr Pro Val Phe Phe Phe Ala Tyr Gly Gly Pro Asn Ser

1801 CAA CAA GTT GTC AAA ACG TTT TCC GTA GGA TTT AAT GAA GTG GTA GCT TCA CAA TTA AAC GCA ATT GTA GTT GTT GTT GAC GGT CGT GGT
601 Gln Gln Val Val Lys Thr Phe Ser Val Gly Phe Asn Glu Val Val Ala Ser Gln Leu Asn Ala Ile Val Val Val Val Asp Gly Arg Gly

1891 ACT GGC TTC AAA GGT CAA GAC TTT AGA TCC CTT GTT CGC GAT AGG CTC GGT GAT TAC GAG GCC CGC GAC CAA ATA TCT GCG GCT TCC TTA
631 Thr Gly Phe Lys Gly Phe Arg Ser Leu Val Arg Ser Phe Arg Leu Gly Asp Tyr Glu Ala Arg Asp Gln Ile Ser Ala Ala Ser Leu

KpnI

1981 TAT GGT TCT TTA ACT TTT GTT GAT CCG CAA AAG ATT TCC TTA TTT GGT TGG TCA TAC GGG GGG TAC CTG ACA CTA AAA ACT TTG GAG AAA
661 Tyr Gly Ser Leu Thr Phe Val Asp Pro Gln Lys Ile Ser Leu Phe Gly Trp Ser Tyr Gly Gly Tyr Leu Thr Leu Lys Thr Leu Glu Lys

2071 GAT GGC GGA AGA CAT TTC AAA TAC GGG ATG TCA GTT GCG CCA GTA ACC GAC TGG AGA TTT TAC GAT TCT GTT TAT ACT GAG AGG TAC ATG
691 Asp Gly Gly Arg His Phe Lys Tyr Gly Met Ser Val Ala Pro Val Thr Asp Trp Arg Phe Tyr Asp Ser Val Tyr Thr Glu Arg Tyr Met

2161 CAT ACT CCT CAA GAA AAC TTT GAT GGA TAC GTA GAA TCA AGC GTT CAT AAT GTC ACT GCT TTG GCA CAA GCA AAT AGA TTT TTG TTG ATG
721 His Thr Pro Gln Glu Asn Phe Asp Gly Tyr Val Glu Ser Ser Val His Asn Val Thr Ala Leu Ala Gln Ala Asn Arg Phe Leu Leu Met

HincII

2251 CAC GGA ACA GGA GAT GAT AAC GTT CAC TTT CAA AAT TCC CTA AAG TTT CTG GAC CTT TTG GAT CTA AAT GGT GTG GAA AAT TAT GAC GTC
751 His Gly Thr Gly Asp Asp Asn Val His Phe Gln Asn Ser Leu Lys Phe Leu Asp Leu Leu Asp Leu Asn Gly Val Glu Asn Tyr Asp Val

2341 CAC GTC TTT CCT GAC TCA GAT CAT AGT ATA AGA TAC CAT AAT GCG AAT GTA ATC GTT TTT GAC AAG CTA TTG GAT TGG CAA AGC GTG CTT
781 His Val Phe Pro Asp Ser Asp His Ser Ile Arg Tyr His Asn Ala Asn Val Ile Val Phe Asp Lys Leu Leu Asp Trp Gln Ser Val Leu

2431 TCG ATG GGC AAT TTG ACA AAT GAG TTG ACT ATT TAC AGC TCA TCT CAT AGA GAC ATT CAT AAA ACA TTT TCA TAT TTA CAT ACT ATG TAT
811 Ser Met Gly Asn Leu Thr Lys Glu Leu Thr Ile Tyr Ser Ser Ser His Arg Asp Ile His Lys Thr Phe Ser Tyr Leu His Thr Met Tyr

AccI

2521 ATA TAA AATGGTAATATAACAGATTAAATGTCGTAGAAGATGATGTTTACGGTGGCAAAAGCGAGCACGAAATTTGGCAAGGGTATAAGTAATTAAGCAAATTTGGTCAAAT
841 Ile End

AccI

2639 CTATAATCTTCACTAACAACTTGCCCATGGTCTAC

Figure 2. Nucleotide and corresponding amino acid sequence of the *DAP2* gene. Nucleotide residues are numbered in the 5'- to -3' direction, with the A of the predicted ATG initiating methionine given the number 1. The restriction sites used in sequencing are indicated. The TATA-like element is underlined, the amino acids comprising the putative membrane-spanning domain are boxed, and the potential sites of N-linked glycosylation are marked by asterisks.

stel3 mutant (Sprague, G., and Ira Herskowitz, unpublished results; Julius et al., 1983). Another gene, called *STE13'* (also called *DPP2*), was cloned by its ability to complement the α -factor maturation defect of a *stel3* mutant when expressed from a high copy number plasmid (Sprague, G., and I. Herskowitz, unpublished results; Julius et al., 1983). The gene dosage of *DPP2* had a direct effect on the specific activity of DPAP B in extracts of yeast cells (Julius et al., 1983; Sprague, G., unpublished results). Strains transformed with a multicopy plasmid containing *DPP2* on a 4.7-kb Bam HI-Pst I fragment (Fig. 1) showed a 10–20-fold increase in DPAP B activity, whereas strains carrying a disrupted genomic copy of *DPP2* (*dpp2-Δ2::LEU2*, see Materials and Methods; see Fig. 4) showed no measurable DPAP B activity in assays of whole cell extracts (data not shown). Also, a *dpp2-Δ2::LEU2/DPP2* diploid strain had ~50% of the DPAP B activity of a *DPP2/DPP2* diploid (data not shown).

Mutants deficient in DPAP B activity have been previously described (Suarez Rendueles and Wolf, 1987). These mutants, including one with an exceptionally thermolabile DPAP B activity, harbored mutations that fell into a single complementation group, called *DAP2*, and failed to complement the *dpp2-Δ2* mutation. Thus, *DPP2* and *DAP2* define the same locus, which encodes DPAP B. Henceforth, *DAP2* will denote the gene encoding DPAP B. No phenotype other than the lack of DPAP B was observed in strains disrupted at the *DAP2* locus (Sprague, G., unpublished results; data not shown), consistent with the results of Suarez Rendueles and Wolf (1987).

Primary Structure of DPAP B Deduced from the *DAP2* Sequence

We deduced the primary structure of DPAP B by sequencing *DAP2*. Fig. 1 shows the restriction map of *DAP2* and the sequencing strategy for the 3,135-bp Bam HI-Acc I fragment. The complete nucleotide and deduced amino acid sequences are presented in Fig. 2. 459 bp of 5' and 153 bp of 3' flanking sequences are included in addition to the 2,523 bp of coding sequence.

Translation of DPAP B is predicted to initiate at the methionine codon numbered 1 (Fig. 2). The sequenced region contains a long open reading frame coding for a protein of 841 amino acids, with eight canonical sites for asparagine-linked (N-linked) glycosylation. A 24-bp segment located from –210 to –187 relative to the putative translation initiating codon may function as a reiterated TATA-like box for initiation of transcription. An identical sequence is found in a similar location upstream of the yeast *STE3* gene (Hagen et al., 1986). The predicted molecular mass from the deduced amino acid sequence is 96,429 D, which is similar to the apparent molecular mass of DPAP B after removal of N-linked carbohydrate (see below).

Hydropathy analysis of the amino acid sequence (Kyte and Doolittle, 1982; Fig. 3 A) reveals a single hydrophobic domain, beginning 30 residues from the NH₂ terminus, of sufficient length and hydrophobicity to span a lipid bilayer (Adams and Rose, 1985; Davis and Model, 1985). Thus, the predicted topology of DPAP B is that of a type II (or group B; Fig. 3 B) integral membrane protein (Garoff, 1985; Singer et al., 1987), with an NH₂-terminal cytoplasmic do-




Figure 3. Hydrophobicity analysis of the *DAP2* gene product. (A) The mean hydropathic index of successive stretches of 19 amino acid residues is plotted vs. the amino acid number of the middle residue of each stretch. Regions with high positive values are potential membrane spanning domains. (B) Model of the membrane topology of DPAP B predicted from the deduced amino acid sequence. The NH₂- and COOH-terminal ends of the protein are indicated, and the sites for N-linked glycosylation are marked by triangles.

main and a lumenal COOH terminus. The hydrophobic domain presumably functions as both an ER-targeting signal and a membrane anchor, as is true of other type II integral membrane proteins, such as the neuraminidase of influenza virus (Bos et al., 1984) and the transferrin receptor (Zerial et al., 1986). Further support for this disposition of DPAP B in the membrane is the observation that DPAP B is a glycoprotein (see below), and that all sites for N-linked glycosylation are included in the COOH-terminal domain, which therefore must be the lumenal domain.

No significant sequence similarity has been found between DPAP B and any other vacuolar or lysosomal protein for which sequence information is available. However, strong homology has been found between DPAP B and DPAP A, the product of the *STE13* gene (Flanagan, C., and J. Thorner, personal communication; see below).

DPAP B Is a 120-kD Integral Membrane Protein

To facilitate the biochemical characterization of DPAP B, a DPAP B-specific antibody was prepared (see Materials and Methods). Lanes 1 and 2 of Fig. 4 show that the antibody is specific for the *DAP2* gene product in yeast. Isogenic *DAP2* and *dap2-Δ2* cells were labeled with ³⁵S-H₂SO₄ for 30 min and chased for 30 min in the presence of 10 mM Na₂SO₄, and immunoprecipitations were performed from cell extracts. Two species were precipitated from the *DAP2* strain (Fig. 4, lane 2): a major species of ~120 kD and a lesser amount of a 110-kD form. Both of these bands were absent from the *dap2-Δ2* strain (lane 1), indicating that they represent the products of the *DAP2* locus.




Figure 4. Immunoprecipitations of DPAP B. (Lanes 1 and 2) JHRY20-2CΔ3 (Δ) and JHRY20-2C (+) cells were ^{35}S labeled for 30 min and chased 30 min in the presence of Na_2SO_4 . (Lanes 3–10) JHRY20-2C cells were pulse labeled for 5 min and chased in unlabeled Na_2SO_4 , cysteine, methionine, and cycloheximide for the times indicated. Labeled cells were collected, converted to spheroplasts, lysed in 1% SDS, and immunoprecipitated with DPAP B antibody. Immunoprecipitates were analyzed on 6% polyacrylamide SDS gels without (–Endo F) or with (+Endo F) earlier treatment with endo F.

The deduced amino acid sequence predicts that DPAP B is an integral membrane protein. To test this prediction biochemically, yeast spheroplasts were treated with 100 mM sodium carbonate, pH 11.5, and centrifuged at 100,000 g . Under these conditions, only intrinsic membrane proteins pellet with membranes, whereas soluble and peripheral membrane proteins are recovered in the supernatant (Steck and Yu, 1973; Fujiki et al., 1982). Fig. 5 shows western blot analysis of the total, pellet, and supernatant fractions using DPAP B-specific and phosphoglycerate kinase- (PGK) specific antibodies. The DPAP B antigen was found in the membrane fraction, whereas the soluble protein PGK did not sediment with membranes. These data confirm that DPAP B is an integral membrane protein.

DPAP B Is Synthesized as a Glycosylated Precursor

The soluble vacuolar proteins analyzed to date, such as CPY, are synthesized as glycoproteins (Rothman and Stevens, 1988). CPY first appears as a glycosylated precursor which receives further glycosyl residues in the Golgi apparatus before being proteolytically cleaved to the mature form upon delivery to the vacuole (Stevens et al., 1982). To follow the biosynthesis of DPAP B, a culture of *DAP2* cells was pulse labeled for 5 min in ^{35}S - H_2SO_4 and chased for 0, 5, 10, and 30 min (Fig. 4, lanes 3–6). DPAP B initially appeared as two distinct bands of 110 and 113 kD (lane 3) which chased to

the broad 120-kD species seen in lane 2. A small amount of the 110-kD form remained after a 30-min chase (lanes 2 and 6). Longer times of up to 60 min were required to convert this species to the fully mature 120-kD form (data not shown).

Treatment of the immunoprecipitated samples with endo F (Fig. 4, lanes 7–10), which removes N-linked glycosyl residues, showed that the only detectable difference in the species seen in lanes 3–6 is in the amount of N-linked carbohydrate present on DPAP B. At each chase time point, labeled DPAP B migrated as a 96-kD species after endo F treatment, which is very close to the molecular mass predicted from the deduced amino acid sequence. A small amount of a 99-kD species was also detected in lanes 8–10; this appears to correspond to one residual carbohydrate chain remaining on DPAP B since it disappears upon longer incubation with endo F (see Fig. 6). Thus, DPAP B is modified in a manner similar to soluble vacuolar proteins, such as CPY, in that its carbohydrate moieties undergo only modest extension in the Golgi apparatus, and do not receive the extensive outer chain mannose residues found on secreted proteins such as invertase (Esmon et al., 1981; Runge, 1988).

DPAP B Receives Variable Numbers of N-linked Glycosyl Residues

The appearance of two distinct DPAP B species at the earliest times in its biogenesis is similar to that seen with invertase, which is initially synthesized as several discrete species that differ in the number of N-linked glycosyl chains added per protein molecule (Esmon et al., 1981). To determine how many of the eight potential sites for N-linked glycosylation are modified on DPAP B, limited endo F digestion was per-




Figure 5. DPAP B is an integral membrane protein. JHRY20-1AΔ1 cells containing the *DAP2* gene on a multicopy plasmid were converted to spheroplasts, lysed, and diluted into 100 mM sodium carbonate, pH 11.5. A portion of the total extract (Tot) was saved, and the rest was separated into membrane (Mb) and soluble (Sol) fractions by centrifugation at 100,000 g for 3 h. The three fractions were subjected to SDS-PAGE, and the gel was analyzed by immunoblotting with DPAP B and phosphoglycerate kinase antibodies and subsequent incubation with ^{125}I -protein A. An autoradiogram of the immunoblots is shown, with the positions of the protein molecular mass standards indicated.




Figure 6. Time course of endo F digestion. DPAP B was immunoprecipitated from CJRY21-3BΔ1 (*sec18*) cells containing *DAP2* on a multicopy 2- μm plasmid after ^{35}S labeling for 30 min. Samples were digested with endo F for the times indicated (ON, overnight). The different glycosylated species are marked, with the corresponding number of glycosyl residues shown.

formed on the precursor, which was immunoprecipitated from a *sec18* mutant at 34°C (see below). Fig. 6 suggests that the precursor forms seen in lane 3 of Fig. 4 correspond to DPAP B with six and seven glycosyl residues added, although forms with five and eight oligosaccharide chains can also be seen (Fig. 6, 0-min time point). Therefore, DPAP B can be modified at all eight or as few as five sites, but the majority of molecules receives six or seven glycosyl chains.

DPAP B Is Not Affected by the Allelic State of *PEP4*

The *PEP4* gene function is necessary for the activation of a number of vacuolar hydrolases (Jones et al., 1982). Mutations at this locus result in the failure to proteolytically process the large molecular mass, inactive forms of soluble vacuolar proteins such as CPY (Rothman and Stevens, 1988). Suarez Rendueles et al. (1981) showed that total DPAP (DPAP A and DPAP B) activity was similar in a wild-type strain and a *pep4-3* mutant. To examine further the effect of a *pep4* mutation on DPAP B, enzyme assays and immunoprecipitations of DPAP B were performed using isogenic *PEP4* and *pep4-Δ2* strains. No significant effect on the activity of DPAP B was detected in the *pep4-Δ2* strain as compared to the *PEP4* strain (data not shown), confirming that DPAP B is not synthesized as an inactive precursor. This result is consistent with the observation that DPAP B which had accumulated in the ER of *sec18* cells (see below) was enzymatically active (data not shown). Fig. 7 shows that DPAP B immunoprecipitated from isogenic *PEP4* and *pep4-Δ2* strains comigrates before and after digestion with endo F, although a minor breakdown product, not seen in the *pep4-Δ2* strain, was detected in the *PEP4* strain after endo F treatment. These data argue that the biogenesis of DPAP B is independent of the allelic state of the *PEP4* gene.

Early-blocked *sec* Mutants Accumulate the Precursor Form of DPAP B

The temperature-sensitive *sec* mutations define sequential steps in the secretory pathway in yeast, since cells carrying these mutations are blocked for protein secretion at distinct stages (Novick et al., 1980; Novick et al., 1981). The localization of CPY to the vacuole is prevented in *sec* mutants that are blocked at either the ER or Golgi apparatus (*sec18* and *sec7*, respectively; Stevens et al., 1982), and the protein accumulates as an inactive precursor that lacks Golgi-mediated carbohydrate modifications. *sec1* cells, which at 37°C are blocked at a late secretory vesicle stage of the pathway, transport CPY to the vacuole at all temperatures. Thus, secretory and soluble vacuolar proteins diverge after the *sec7*-defined block and before the *sec1*-blocked stage. To determine if the maturation of DPAP B was affected in these mutants, DPAP




Figure 7. Effect of a *pep4* mutation on DPAP B. Strains JHRY20-2C and JHRY20-2CΔ2 were ³⁵S labeled for 15 min and chased for 30 min at 30°C. Labeled cells were collected, and DPAP B was immunoprecipitated and endo F treated as before.




Figure 8. Effect of *sec* mutations on DPAP B. Strains were ³⁵S labeled at their nonpermissive temperatures (X2180, HMSF176, and HMSF1, 34°C; SF294-2B, 38°C) for 15 min, chased for 30 min in the presence of Na₂SO₄ at the same temperature, and DPAP B was immunoprecipitated as before.

B was immunoprecipitated after labeling the cells at the nonpermissive temperature of 34°C (Salminen and Novick, 1987), except for the *sec7* mutant, whose restrictive temperature is 38°C (data not shown). Wild-type cells contained the 120-kD mature form of DPAP B at either 34°C (Fig. 8) or 38°C (data not shown). *sec1* cells also contained mature DPAP B, whereas *sec18* cells accumulated the DPAP B precursor. Thus, the precursor form seen after a short labeling period (Fig. 4, lane 3) is the ER-form of DPAP B. The *sec7* strain accumulated a mixture of core-glycosylated and mature forms of DPAP B. This suggests that the block in transport is at an early stage within the Golgi apparatus such that DPAP B is only partially processed by the mannosyl transferases (Runge, 1988). This phenotype is similar to that seen with invertase, which accumulates as a heterogeneous population of core- and hyper-glycosylated forms in *sec7* cells at the restrictive temperature (Esmon et al., 1981).

The accumulation of the DPAP B precursor in *sec18* and *sec7* cells was thermoreversible: after labeling cells at the restrictive temperature, shifting the cultures to 23°C, and chasing for 3 h in the absence of new protein synthesis, DPAP B was completely converted to the mature 120-kD form (data not shown). This result, which was also observed with CPY accumulated in *sec18* and *sec7* cells (Stevens et al., 1982), suggests that the accumulated DPAP B precursor resides early in the pathway and is able to transit the Golgi apparatus upon shifting to the permissive temperature.

Early-blocked *sec* Mutants Accumulate DPAP B in Nonvacuolar Compartments

To determine whether the accumulation of the DPAP B precursor in *sec18* and *sec7* mutants was correlated with a failure to deliver the protein to the vacuole, DPAP B was localized by indirect immunofluorescence microscopy. For these studies, DPAP B synthesis was placed under the control of the inducible *GAL1* promoter (Johnston and Davis, 1984), allowing analysis of nascent DPAP B synthesis after shifting to the nonpermissive temperature. A multicopy plasmid containing the *GAL1-DAP2* gene fusion was transformed into wild-type and *sec* mutant strains which had been disrupted at the chromosomal *DAP2* locus. Transformed strains were then shifted to the restrictive temperatures in galactose-containing media and incubated for 2 h. This procedure typically resulted in a 4–10-fold induction of DPAP B activity relative to a strain containing a single genomic copy of *DAP2* (data not shown). No DPAP B activity or antigen was detected when these strains were grown in the absence of galactose (data not shown). The cells were fixed in formaldehyde, spheroplasted, and incubated with affinity-purified DPAP B




Figure 9. DPAP B localization by immunofluorescence microscopy. Strains JHRY20-1A (*SEC*⁺), CJRY21-3BΔ1 (*sec18*), CJRY23-2AΔ1 (*sec7*), and CJRY22-6BΔ1 (*sec1*), each containing the *GAL1-DAP2* gene fusion on a multicopy plasmid, were shifted to their respective nonpermissive temperatures and treated for induction of the *GAL1* promoter for 2 h as described in Materials and Methods. The cells were fixed, converted to spheroplasts, and stained with DAPI and affinity-purified DPAP B antibody. Cells were viewed by Nomarski differential interference optics (A, D, G, and J), and under DAPI (B, E, H, and K) and FITC (C, F, I, and L) excitation wavelengths.

antibody, followed by FITC-conjugated sheep anti-goat antibody (Fig. 9). Nomarski optics revealed the position of the vacuole, and the nucleus was localized by DAPI staining. Wild-type cells grown at either 34°C (Fig. 9, A–C) or 38°C (data not shown) exhibited fluorescent labeling of DPAP B around the perimeter of the vacuole, consistent with the fact that DPAP B is a vacuolar membrane protein. A similar labeling pattern was seen in *sec1* cells (Fig. 9, J–L), indicat-

ing that the *sec1* block is later in the secretory pathway than the point at which sorting of vacuolar membrane proteins occurs. The staining pattern seen in *sec18* (Fig. 9, D–F) and *sec7* (Fig. 9, G–I) cells was distinctly nonvacuolar. Interestingly, the *sec18* strain showed DPAP B staining around the nucleus, apparently in the nuclear envelope, which probably reflects the fact that the nuclear envelope and the ER are continuous membranes. These data indicate that transport of

DPAP B to the vacuole is prevented in *sec* mutants that are blocked in protein transport early in the secretory pathway.

Discussion

DPAP B is the first protein of the yeast vacuolar membrane whose structure and biosynthesis has been characterized. Of the vacuolar membrane-associated activities that have been described in yeast, only the proton-translocating ATPase (Uchida et al., 1985; Kane et al., 1989) and α -mannosidase (Yoshihisa et al., 1988) have been extensively characterized biochemically. The proton-translocating ATPase is a multimeric enzyme consisting of both integral and peripheral membrane proteins (Kane et al., 1989). α -Mannosidase, which is frequently used as a marker for the vacuolar membrane, is efficiently removed from membranes by treatment with high-pH sodium carbonate, suggesting that it is peripherally associated with the vacuolar membrane (Yoshihisa et al., 1988). Thus, DPAP B appears to be a more appropriate marker for intrinsic vacuolar membrane proteins.

In a previous biochemical study, DPAP B was reported to be a 40-kD protein (Garcia Alvarez et al., 1985). In the present work we have found that DPAP B is a 120-kD glycoprotein, that this species is absent from a strain with a disruption at *DAP2*, and that the *DAP2* sequence encodes a protein predicted to have a molecular mass of 96,429 D, which corresponds closely to that actually observed for deglycosylated DPAP B. The 40-kD species seen previously may have been a contaminating protein or a proteolytic fragment of DPAP B.

As is the case for the soluble vacuolar protein CPY (Stevens et al., 1982), the delivery of DPAP B to the vacuole is blocked in *sec* mutants that affect the early stages of the secretory pathway. *sec18* (ER-blocked) and *sec7* (Golgi-blocked) cells accumulate DPAP B in nonvacuolar compartments at the restrictive temperature. However, *sec1* cells, which are blocked at a late secretory vesicle stage, transport DPAP B to the vacuole at all temperatures, indicating that DPAP B is transported directly from the Golgi apparatus to the vacuole. This rules out models in which vacuolar membrane proteins are first delivered to the plasma membrane, and suggests that, in yeast, soluble and membrane-bound proteins traverse the same compartments of the secretory pathway en route to the vacuole (Stevens et al., 1982), as is the case for the delivery of proteins to the lysosome in mammalian cells (Griffiths et al., 1988).

The activity and apparent molecular mass of DPAP B are unaffected by a null mutation at the *PEP4* locus. DPAP B is active at all stages of its transport and does not appear to undergo proteolytic cleavage upon delivery to the vacuole. In this respect, DPAP B differs from the soluble vacuolar hydrolases studied thus far, which are synthesized as zymogens and are activated in a *PEP4*-dependent fashion (Rothman and Stevens, 1988). Interestingly, the other vacuolar membrane activities that have been examined, i.e., α -mannosidase (Jones et al., 1982) and the proton-translocating ATPase (Yamashiro, C., and T. Stevens, unpublished results), are also *PEP4* independent.

Overproduction of the soluble vacuolar proteins CPY and proteinase A results in their secretion, presumably due to the saturation of a component of the sorting apparatus (Stevens et al., 1986; Rothman et al., 1986). This observation provided a means of isolating mutations that disrupt the sorting

process, both linked and unlinked to the structural genes of the soluble vacuolar proteins (Valls et al., 1987; Rothman and Stevens, 1986; Bankaitis et al., 1986; Robinson et al., 1988; Rothman et al., 1989). 20-fold overproduction of DPAP B does not result in increased cell surface activity of this enzyme (our unpublished results). However, at this level of expression DPAP B does suppress the defect in α -factor maturation of a *MAT α stel3* mutant (Sprague, G., and I. Herskowitz, unpublished observation; Julius et al., 1983). A reasonable interpretation of this result is that DPAP B is mislocalized into the late secretory pathway when overproduced, allowing it to encounter and process the α -factor precursor to the mature form. Based on this assumption, mutations have been isolated that result in secretion of mature α -factor in a *stel3* background. These mutants do not cause DPAP B to be overproduced, and may define genes that encode components of the sorting apparatus for vacuolar membrane proteins (Pohlig, G., and T. Stevens, unpublished results).

The sorting signal(s) that directs membrane proteins to the vacuole has yet to be defined. The sorting information could reside in any of three distinct domains; i.e., the cytoplasmic, membrane, and/or luminal domains. Moreover, each of these domains has been shown to be important for proper intracellular targeting of particular membrane proteins. The luminal domains of the influenza virus hemagglutinin and vesicular stomatitis virus G protein have been implicated as important for the polarized secretion of these proteins to the apical and basolateral membranes, respectively (Roth et al., 1987; McQueen et al., 1987). Retention of the coronavirus E1 glycoprotein in the Golgi apparatus requires a specific transmembrane region (Machamer and Rose, 1987), and an intact cytoplasmic tail is necessary for endocytosis of the LDL and transferrin receptors (Davis et al., 1986; Rothenberger et al., 1987). Preliminary experiments have shown that the cytoplasmic tail and transmembrane region of DPAP B contain vacuolar sorting information; i.e., fusion of the NH₂-terminal 47 residues of DPAP B to the NH₂ terminus of the cytoplasmic form of invertase results in the delivery of this hybrid protein to the vacuolar membrane (Roberts, C., and T. Stevens, unpublished observation). This approach has been previously used to map the sorting domains of CPY (Johnson et al., 1987) and proteinase A (Klionsky et al.,




Figure 10. Comparison of DPAP B and DPAP A. Models for the disposition of the two proteins in their respective membranes are shown, with their subcellular locations listed at right. The NH₂ and COOH termini of the proteins are indicated, as are the regions of extensive amino acid identity, with the percentage of identical amino acids given.

1988). Further experiments are aimed at the identification of the specific amino acid residues in this region that are necessary for targeting DPAP B to the vacuole.

The DNA sequence of the *STE13* gene, which encodes DPAP A (Julius et al., 1983), has recently been determined (Flanagan, C., and J. Thorner, personal communication). The deduced amino acid sequence predicts a protein of 931 amino acids with striking similarities to DPAP B (Fig. 10). Both proteins are predicted to have type II membrane topologies, with short cytoplasmic tails and ~800 amino acid luminal domains. The luminal domains of these proteins are very similar; in particular, the COOH-terminal 240 amino acids are 48% identical, presumably reflecting the similar enzymatic functions of these proteins in vitro and in vivo (Julius et al., 1983). However, the cytoplasmic and transmembrane domains are unrelated, possibly reflecting the different subcellular addresses of DPAP A and DPAP B. This sorting signal hypothesis can be tested by exchanging these segments of the two proteins, an approach that has been useful in studying the sorting of membrane proteins in polarized cells (Roth et al., 1987; McQueen et al., 1987).

We would like to thank George Sprague for the gift of the plasmid containing the *DAP2* clone and for many useful discussions; Jeremy Thorner for communicating results before publication; Rick Dahlquist for providing the plasmid pHSe5; Ruskin Gould for excellent technical assistance; Harry Howard for expert assistance with the fluorescence microscopy; and Christopher Raymond, Nicholas Davis, Karen Moore, and Patricia Kane for comments on the manuscript.

This work was supported by National Institutes of Health Predoctoral Traineeships to C. J. Roberts and J. H. Rothman, a Deutsche Forschungsgemeinschaft Postdoctoral Fellowship to G. Pohl, and by grants from the National Institutes of Health (PHS 38006) and the Chicago Community Trust Searle Scholars Program to T. H. Stevens.

Received for publication 20 December 1988.

References

- Adams, A. E. M., and J. R. Pringle. 1984. Relationship of actin and tubulin distribution to bud growth in wild-type and morphogenetic-mutant *Saccharomyces cerevisiae*. *J. Cell Biol.* 98:934-945.
- Adams, G. A., and J. K. Rose. 1985. Structural requirements of a membrane-spanning domain for protein anchoring and cell surface transport. *Cell* 41:1007-1015.
- Bankaitis, V. A., L. M. Johnson, and S. D. Emr. 1986. Isolation of yeast mutants defective in protein targeting to the vacuole. *Proc. Natl. Acad. Sci. USA* 83:9075-9079.
- Barriocanal, J. G., J. S. Bonifacio, L. Yuan, and I. V. Sandoval. 1986. Biosynthesis, glycosylation, movement through the Golgi system, and transport to lysosomes by an N-linked carbohydrate-independent mechanism of three lysosomal integral membrane proteins. *J. Biol. Chem.* 261:16755-16763.
- Bordallo, C., J. Schwencke, and M. Suarez Rendueles. 1984. Localization of the thermosensitive X-prolyl dipeptidyl aminopeptidase in the vacuolar membrane of *Saccharomyces cerevisiae*. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 173:199-203.
- Bos, T. J., A. R. Davis, and D. P. Nayak. 1984. NH₂-terminal hydrophobic region of influenza virus neuraminidase provides the signal function in translocation. *Proc. Natl. Acad. Sci. USA* 81:2327-2331.
- Burnette, W. N. 1981. "Western blotting": electrophoretic transfer of proteins from sodium dodecyl sulfate-polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein A. *Anal. Biochem.* 112:195-203.
- Casadaban, M. F., and S. N. Cohen. 1980. Analysis of gene control signals by DNA fusion and cloning in *Escherichia coli*. *J. Mol. Biol.* 138:179-207.
- Chen, J. W., T. L. Murphy, M. C. Willingham, I. Pastan, and J. T. August. 1985. Identification of two lysosomal membrane glycoproteins. *J. Cell Biol.* 101:85-95.
- Davis, N. G., and P. Model. 1985. An artificial anchor domain: hydrophobicity suffices to stop transfer. *Cell* 41:607-614.
- Davis, C. G., M. A. Lehrman, D. W. Russell, R. G. W. Anderson, M. S. Brown, and J. L. Goldstein. 1986. The J. D. mutation in familial hypercholesterolemia: amino acid substitution in cytoplasmic domain impedes internalization of LDL receptors. *Cell* 45:15-24.
- Elder, J. H., and S. Alexander. 1982. Endo- β -N-acetylglucosaminidase F: endoglycosidase from *Flavobacterium meningosepticum* that cleaves both high-mannose and complex glycoproteins. *Proc. Natl. Acad. Sci. USA* 79:4540-4544.
- Emr, S. D., I. Schauer, W. Hansen, P. Esmon, and R. Schekman. 1984. Invertase β -galactosidase hybrid proteins fail to be transported from the endoplasmic reticulum in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 4:2347-2355.
- Esmon, B., P. Novick, and R. Schekman. 1981. Compartmentalized assembly of oligosaccharides on exported glycoprotein in yeast. *Cell* 25:451-460.
- Fujiki, Y., A. L. Hubbard, S. Fowler, and P. B. Lazarow. 1982. Isolation of intracellular membranes by means of sodium carbonate treatment: application to endoplasmic reticulum. *J. Cell Biol.* 93:97-102.
- Fuller, R. S., R. E. Sterne, and J. Thorner. 1988. Enzymes required for yeast prohormone processing. *Annu. Rev. Physiol.* 50:345-362.
- Garcia Alvarez, N., C. Bordallo, S. Gascon, and P. Suarez Rendueles. 1985. Purification and characterization of a thermosensitive X-prolyl dipeptidyl aminopeptidase (dipeptidyl aminopeptidase yscV) from *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 832:119-125.
- Garoff, H. 1985. Using recombinant DNA techniques to study protein targeting in the eucaryotic cell. *Annu. Rev. Cell Biol.* 1:403-445.
- Griffiths, G., B. Hoffack, K. Simons, I. Mellman, and S. Kornfeld. 1988. The mannose 6-phosphate receptor and the biogenesis of lysosomes. *Cell* 52:329-341.
- Hagen, D. C., G. McCaffrey, and G. F. Sprague, Jr. 1986. Evidence the yeast *STE3* gene encodes a receptor for the peptide pheromone a factor: gene sequence and implications for the structure of the presumed receptor. *Proc. Natl. Acad. Sci. USA* 83:1418-1422.
- Ito, H., Y. Fukuda, K. Murata, and A. Kimura. 1983. Transformation of intact yeast cells treated with alkali cations. *J. Bacteriol.* 153:163-168.
- Ito, H., Y. Ike, S. Ikuta, and K. Itakura. 1982. Solid phase synthesis of polynucleotides VI. Further studies on polystyrene copolymers for the solid support. *Nucleic Acids Res.* 10:1755-1769.
- Johnson, L. M., V. A. Bankaitis, and S. D. Emr. 1987. Distinct sequence determinants direct intracellular sorting and modification of a yeast vacuolar proteinase. *Cell* 48:875-885.
- Johnston, M., and R. W. Davis. 1984. Sequences that regulate the divergent *GALI-GALI0* promoter in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 4:1440-1448.
- Jones, E. W., G. S. Zubenko, and R. R. Parker. 1982. *PEP4* gene function is required for expression of several vacuolar hydrolases in *Saccharomyces cerevisiae*. *Genetics* 102:665-677.
- Julius, D., L. Blair, A. Brake, G. F. Sprague, Jr., and J. Thorner. 1983. Yeast α factor is processed from a larger precursor polypeptide: the essential role of a membrane-bound dipeptidyl aminopeptidase. *Cell* 32:839-852.
- Kane, P. M., C. T. Yamashiro, J. H. Rothman, and T. H. Stevens. 1989. Protein sorting in yeast: the role of the vacuolar proton-translocating ATPase. *J. Cell Sci.* In press.
- Kilmartin, J. V., and A. E. M. Adams. 1984. Structural rearrangements of tubulin and actin during the cell cycle of the yeast *Saccharomyces*. *J. Cell Biol.* 98:922-933.
- Kleid, D. G., D. Yansura, B. Small, D. Dowbenko, D. M. Moore, M. J. Grubman, P. D. McKercher, D. O. Morgan, B. H. Robertson, and H. L. Bachrach. 1981. Cloned viral protein vaccine for foot-and-mouth disease: responses in cattle and swine. *Science (Wash. DC)* 214:1125-1129.
- Klionsky, D. J., L. M. Banta, and S. D. Emr. 1988. Intracellular sorting and processing of a yeast vacuolar hydrolase: proteinase A propeptide contains vacuolar targeting information. *Mol. Cell. Biol.* 8:2105-2116.
- Kyte, J., and R. F. Doolittle. 1982. A simple method for displaying the hydrophobic character of a protein. *J. Mol. Biol.* 157:105-132.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (Lond.)* 227:680-685.
- Lewis, V., S. A. Green, M. Marsh, P. Vihko, A. Helenius, and I. Mellman. 1985. Glycoproteins of the lysosomal membrane. *J. Cell Biol.* 100:1839-1847.
- Lippincott-Schwartz, J., and D. Fambrough. 1986. Lysosomal membrane dynamics: structure and interorganelle movement of a major lysosomal membrane protein. *J. Cell Biol.* 102:1593-1605.
- Machamer, C. E., and J. K. Rose. 1987. A specific transmembrane domain of a coronavirus E1 glycoprotein is required for its retention in the Golgi region. *J. Cell Biol.* 105:1205-1214.
- Maniatis, T., E. F. Fritsch, and J. Sambrook. 1982. Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York. 545 pp.
- McQueen, N. L., D. P. Nayak, E. B. Stephens, and R. W. Compans. 1987. Basolateral expression of a chimeric protein in which the transmembrane and cytoplasmic domains of vesicular stomatitis virus G protein have been replaced by those of the influenza virus hemagglutinin. *J. Biol. Chem.* 262:16233-16240.
- Messing, J. 1983. New M13 vectors for cloning. *Methods Enzymol.* 101:20-78.
- Muchmore, D. C., L. P. McIntosh, C. B. Russell, D. E. Anderson, and F. W. Dahlquist. 1988. Expression and ¹⁵N labeling of proteins for proton and nitrogen-15 NMR. *Methods Enzymol.* In press.
- Novick, P., S. Ferro, and R. Schekman. 1981. Order of events in the yeast secretory pathway. *Cell* 25:461-469.
- Novick, P., C. Field, and R. Schekman. 1980. Identification of 23 complemen-

- tation groups required for post-translational events in the yeast secretory pathway. *Cell*. 21:205-215.
- Novick, P., and R. Schekman. 1983. Export of major cell surface proteins is blocked in yeast secretory mutants. *J. Cell Biol.* 96:541-547.
- Ohsumi, Y., and Y. Anraku. 1981. Active transport of basic amino acids driven by a proton motive force in vacuolar membrane vesicles of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 256:2079-2082.
- Ohsumi, Y., and Y. Anraku. 1983. Calcium transport driven by a proton motive force in vacuolar membrane vesicles of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 258:5614-5617.
- Opheim, D. J. 1978. α -D-mannosidase of *Saccharomyces cerevisiae*: characterization and modulation of activity. *Biochim. Biophys. Acta*. 524:121-130.
- Robinson, J. S., D. J. Klionsky, L. M. Banta, and S. D. Emr. 1988. Protein sorting in yeast: isolation of mutants defective in the delivery and processing of multiple vacuolar hydrolases. *Mol. Cell Biol.* 8:4936-4948.
- Roth, M. G., D. Gunderson, N. Patil, and E. Rodriguez-Boulan. 1987. The large external domain is sufficient for the correct sorting of secreted or chimeric influenza virus hemagglutinins in polarized monkey kidney cells. *J. Cell Biol.* 104:769-782.
- Rothenberger, S., B. J. Iacopetta, and L. C. Kuhn. 1987. Endocytosis of the transferrin receptor requires the cytoplasmic domain but not its phosphorylation site. *Cell*. 49:423-431.
- Rothman, J. H. 1988. Protein sorting in yeast. Ph. D. thesis. University of Oregon, Eugene, OR.
- Rothman, J. H., and T. H. Stevens. 1986. Protein sorting in yeast: mutants defective in vacuole biogenesis mislocalize vacuolar proteins into the late secretory pathway. *Cell*. 47:1041-1051.
- Rothman, J. H., and T. H. Stevens. 1988. Protein sorting and biogenesis of the lysosome-like vacuole in yeast. In *Protein Transfer and Organelle Biogenesis*. R. Das and P. Robbins, editors. Academic Press, Inc., San Diego, CA. 159-208.
- Rothman, J. H., C. P. Hunter, L. A. Valls, and T. H. Stevens. 1986. Overproduction-induced mislocalization of a yeast vacuolar protein allows isolation of its structural gene. *Proc. Natl. Acad. Sci. USA*. 83:3248-3252.
- Rothman, J. H., I. Howald, and T. H. Stevens. 1989. Characterization of genes required for protein sorting, vacuolar function, and endocytosis in the yeast *Saccharomyces cerevisiae*. *EMBO (Eur. Mol. Biol. Organ.) J.* In press.
- Rothstein, R. K. 1983. One step gene disruption in yeast. *Methods Enzymol.* 101:202-211.
- Runge, K. W. 1988. Post-translational modification during protein secretion. In *Protein Transfer and Organelle Biogenesis*. R. Das and P. Robbins, editors. Academic Press, Inc., San Diego, CA. 317-362.
- Salminen, A., and P. J. Novick. 1987. A ras-like protein is required for a post-Golgi event in yeast secretion. *Cell*. 49:527-538.
- Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain terminating inhibitors. *Proc. Natl. Acad. Sci. USA*. 74:5463-5467.
- Singer, S. J., P. A. Maher, and M. P. Yaffe. 1987. On the transfer of integral membrane proteins into membranes. *Proc. Natl. Acad. Sci. USA*. 84:1960-1964.
- Steck, T. L., and J. Yu. 1973. Selective solubilization of proteins from red blood cell membranes by protein perturbants. *J. Supramol. Struct.* 1:220-248.
- Stevens, T. H., B. Esmon, and R. Schekman. 1982. Early stages in the yeast secretory pathway are required for transport of carboxypeptidase Y to the vacuole. *Cell*. 30:439-448.
- Stevens, T. H., J. H. Rothman, G. S. Payne, and R. Schekman. 1986. Gene dosage-dependent secretion of yeast vacuolar carboxypeptidase Y. *J. Cell Biol.* 102:1551-1557.
- Suarez Rendueles, M. P., J. Schwencke, N. Garcia Alvarez, and S. Gascon. 1981. A new X-prolyl-dipeptidyl aminopeptidase from yeast associated with a particulate fraction. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 131:296-300.
- Suarez Rendueles, P., and D. H. Wolf. 1987. Identification of the structural gene for dipeptidyl aminopeptidase yscV (*DAP2*) of *Saccharomyces cerevisiae*. *J. Bacteriol.* 169:4041-4048.
- Uchida, E., Y. Ohsumi, and Y. Anraku. 1985. Purification and properties of H^+ -translocating, Mg^{2+} -adenosine triphosphatase from vacuolar membranes of *Saccharomyces cerevisiae*. *J. Biol. Chem.* 260:1090-1095.
- Vaitukaitis, J. L. 1981. Production of antisera with small doses of immunogen: multiple intradermal injections. *Methods Enzymol.* 73:46-52.
- Valls, L. A., C. P. Hunter, J. H. Rothman, and T. H. Stevens. 1987. Protein sorting in yeast: the localization determinant of yeast vacuolar carboxypeptidase Y resides in the propeptide. *Cell*. 48:887-897.
- von Figura, K., and A. Hasilik. 1986. Lysosomal enzymes and their receptors. *Annu. Rev. Biochem.* 55:167-193.
- Waheed, A., S. Gottschalk, A. Hille, C. Krentler, R. Pohlmann, T. Bräulke, H. Hauser, H. Geuze, and K. von Figura. 1988. Human lysosomal acid phosphatase is transported as a transmembrane protein to lysosomes in transfected baby hamster kidney cells. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:2351-2358.
- Wittenberg, C., S. L. Richardson, and S. I. Reed. 1987. Subcellular localization of a protein kinase required for cell cycle initiation in *Saccharomyces cerevisiae*: evidence for an association between the *CDC28* gene product and the insoluble cytoplasmic matrix. *J. Cell Biol.* 105:1527-1538.
- Yanisch-Perron, C., J. Vieira, and J. Messing. 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13 mp19 and pUC19 vectors. *Gene*. 33:103-119.
- Yoshihisa, T., Y. Ohsumi, and Y. Anraku. 1988. Solubilization and purification of α -mannosidase, a marker enzyme of vacuolar membranes in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 263:5158-5163.
- Zerial, M., P. Melancon, C. Schneider, and H. Garoff. 1986. The transmembrane segment of the human transferrin receptor functions as a signal peptide. *EMBO (Eur. Mol. Biol. Organ.) J.* 5:1543-1550.