

Analysis of Suborganellar Fractions from Spinach and Pea Chloroplasts for Calmodulin-binding Proteins

DANIEL M. ROBERTS, RAYMOND E. ZIELINSKI, MICHAEL SCHLEICHER, and D. MARTIN WATTERSON
Laboratory of Cellular and Molecular Physiology, Howard Hughes Medical Institute and Department of Pharmacology, Vanderbilt University, School of Medicine, Nashville, Tennessee 37232. Dr. Schleicher's present address is Max-Planck-Institut für Psychiatrie, D-8033 Martinsried, Munich 40, Federal Republic of Germany.

ABSTRACT Purified chloroplasts from spinach and pea leaves were subfractionated into envelope, thylakoid, and stroma fractions and were analyzed for calmodulin-binding proteins using a ^{125}I -calmodulin gel overlay assay. Calmodulin binding was primarily associated with a major polypeptide (M_r 33,000) in the envelope membrane fraction. In contrast, major calmodulin-binding proteins were not detected in the thylakoid or stroma fractions. Our results provide the first evidence of calmodulin-binding proteins in the chloroplast envelope, and raise the possibility that calmodulin may contribute to the regulation of chloroplast function through its interaction with calmodulin-binding proteins in the chloroplast envelope. In addition, our results combined with those of other investigators support the proposal that subcellular organelles may be a primary site of calmodulin action.

Calcium is required for a number of cellular activities in eucaryotes, and has been implicated as a possible regulatory agent of metabolic activities and growth responses in a variety of plant species (1). Kinetic, thermodynamic, and structural data strongly suggest that the major intracellular receptors involved in calcium-mediated regulation are a class of calcium-binding proteins known as calcium-modulated proteins (for a recent review, see reference 2). Calmodulin is a calcium-modulated protein that has been found in all eucaryotes examined, including higher plants (for reviews, see references 1–3), and appears to be one of the most highly conserved proteins known (1, 4).

Calmodulin has a number of *in vitro* biochemical activities including the ability to stimulate certain enzyme activities and bind to proteins (1–3). Two calmodulin-sensitive enzyme activities have been demonstrated in higher plants: a NAD kinase (5, 6) and a microsomal calcium ATPase (7). A thorough investigation of the properties of calmodulin-sensitive enzymes and calmodulin-binding proteins is necessary before the role of calmodulin in the regulation of cellular processes in plants can be completely understood. An important part of such an investigation is the determination of the subcellular distribution of calmodulin, calmodulin-stimulated enzymes, and calmodulin-binding proteins. The results of a number of subcellular fractionation and immunocytology studies carried out on tissues from vertebrate organisms indicate that, although calmodulin is found predominantly in the freely soluble fraction, small quantities appear to be associated with subcellular organelles (2, 8). In addition, calmodulin-binding

proteins have been demonstrated in isolated organelles including mitochondria from 3T3 cells (9), and nuclear, mitochondrial/lysosomal, and microsomal fractions from chicken embryo fibroblasts (8).

In contrast to the studies on animal systems, information on the subcellular distribution of calmodulin and calmodulin-binding proteins in plant tissues is more limited. Recently, Jarret et al. (10) and Muto (11) have reported the presence of small quantities of calmodulin in the chloroplast fractions of pea and wheat, respectively. Jarrett et al. (10) suggested that this small pool of calmodulin is involved in the light-dependent phosphorylation of NAD by chloroplast NAD kinase. However, Simon et al. (6) demonstrated soluble and chloroplast pools of NAD kinase activity in spinach and found that the soluble enzyme was stimulated by calmodulin in the presence of calcium, whereas the chloroplast enzyme was not. Simon et al. (6) also found small quantities of calmodulin activity in spinach chloroplasts, but suggested that this chloroplast calmodulin may be the result of contamination by cytoplasmic calmodulin. A detailed analysis of chloroplast fractions and subfractions for calmodulin-binding proteins may provide additional insight into the physiological significance of the chloroplast calmodulin pool. In the present study, we examined purified chloroplast subfractions from spinach and pea leaves for calmodulin-binding proteins.

MATERIALS AND METHODS

Spinach leaves were obtained from a local vendor. Pea seeds (*Pisum sativum* var. Early Alaska) were surface sterilized with a 10% (vol/vol) solution of

Chlorox, imbibed overnight, and grown in soil (Terra Lite Metromix 200, W. R. Grace and Co., Baltimore, MD) in a greenhouse under normal day-light. Leaves from 2–3-wk-old seedlings were used for chloroplast preparation.

Purified intact chloroplasts from spinach or pea leaves were prepared as described by Cline et al. (12), using a modification of the silica sol-density gradient method of Morgenthaler et al. (13). Plastids collected from Percoll gradients were washed twice in 5–10 vol of 50 mM HEPES/NaOH, pH 7.5, containing 0.33 M sorbitol. The chloroplasts were lysed by resuspension in 50 mM HEPES/NaOH, pH 7.5, containing 4 mM MgCl₂. Subfractionation of the lysed chloroplasts was achieved using a modification of the discontinuous sucrose gradient centrifugation method of Mendiola-Morgenthaler and Morgenthaler (14). The lysed chloroplast suspension (6 ml, 0.2 mg/ml chlorophyll) was layered on a step gradient consisting of 2 ml of 30% (wt/vol) sucrose and 2.5 ml of 12% (wt/vol) sucrose, and was centrifuged at 21,500 rpm (Beckman SW 41 rotor; Beckman Instruments, Inc., Palo Alto, CA) for 1 h at 4°C. The stroma fraction remains at the top of the gradient and the thylakoid forms a pellet at the bottom. The envelope fraction, which bands at the 30%/12% sucrose interface, was diluted in 50 mM HEPES/NaOH, pH 7.5, and was collected by sedimentation at 35,000 rpm (SW 41 rotor) for 1 h. All fractions were stored at –80°C until analysis.

Calmodulin was purified (15) and radiolabeled with ¹²⁵I using the lactoperoxidase procedure (16) as previously described. Samples were subjected to electrophoresis on polyacrylamide gels using the buffer system of Laemmli (17) and a modified microslab apparatus similar to that described by Matsudaira and Burgess (18). Detection of calmodulin-binding proteins was accomplished using the ¹²⁵I-calmodulin gel binding assay described by Van Eldik and Burgess (8), except that 0.1 mM CaCl₂ was used for the calcium incubations. Crude fractions from chicken gizzard that contain known calmodulin-binding proteins were included in each overlay assay as a positive control. Competition experiments were performed by incubating the gels with unlabeled calmodulin or α-lactalbumin for 30 min prior to the addition of ¹²⁵I-calmodulin.

Protein concentrations were determined by a modification of the procedure of Lowry et al. (19), as described by Markwell et al. (20) or by amino acid analysis. Chlorophyll was determined by the method of Arnon (21).

RESULTS AND DISCUSSION

Chloroplasts were purified by silica sol density gradient centrifugation (13) and were lysed and subfractionated using carefully defined conditions (14) to ensure fraction purity. The polypeptide profiles obtained by SDS gel electrophoresis of the chloroplast subfractions (Fig. 1A) appear comparable to those reported by other investigators (12, 22, 23).

Gel overlay analysis of the chloroplast subfractions from spinach and pea indicates the presence of a major calmodulin-binding protein in the envelope membrane (Fig. 1, B and C). This calmodulin-binding activity is associated with the major Coomassie-Blue-staining polypeptide (apparent *M_r* 33,000) in our envelope preparation (Fig. 1A). In addition to this major binding protein, the envelope fraction contains three minor calmodulin-binding proteins (apparent *M_r* 17,000, 27,000, and 50,000). A minor calmodulin-binding protein (apparent *M_r* 20,000) was also found in the pea thylakoid fraction. Calmodulin-binding proteins were not detected in the thylakoid fraction from spinach chloroplasts or in the stroma fractions from pea or spinach chloroplasts. Although the data shown in Fig. 1 represent the results of one experiment, the results obtained with four separate experiments on four different chloroplast subfraction preparations indicate the *M_r* 33,000 envelope polypeptide is the major calmodulin-binding protein. We were not able to detect the minor calmodulin-binding proteins in all envelope preparations (Fig. 2A).

The association of ¹²⁵I-calmodulin with the major envelope calmodulin-binding protein (*M_r* 33,000) appears to be calcium independent since binding occurs regardless of whether calcium or EDTA is present (Fig. 1, B and C). The calcium-independent interaction of calmodulin with calmodulin-binding proteins in gel overlay analysis is not unique to the calmodulin-binding proteins in the chloroplast envelope. Van Eldik and Burgess (8) found a number of calcium-independent calmodulin-binding proteins in the particulate subcellular fractions from chicken embryo fibroblasts. Our results combined with those of Van Eldik and Burgess (8) suggest that the calcium-independent interaction of calmodulin with particulate calmodulin-binding proteins may be a general phenomenon in plant and animal systems. It is important to recognize that a calcium-independent interaction does not preclude a calcium-dependent regulation of activity.

The binding of ¹²⁵I-calmodulin to the major envelope mem-




FIGURE 1 ¹²⁵I-Calmodulin gel overlay analysis of chloroplast fractions from spinach and pea. A shows a Coomassie-Blue-stained 15% (wt/vol) acrylamide gel and B and C show autoradiograms (10-h exposure). Gel overlay analysis was done in the presence of 0.1 mM CaCl₂ (B) or 5 mM EDTA (C) as described in Materials and Methods. Lane 1: molecular weight standards: bovine serum albumin, *M_r* 67,000; ovalbumin, *M_r* 43,000; carbonic anhydrase, *M_r* 30,000; and soybean trypsin inhibitor, *M_r* 20,000). Lane 2: chloroplast envelope fraction from pea. Lane 3: stroma fraction from pea. Lane 4: thylakoid fraction from pea. Lane 5: envelope fraction from spinach. Lane 6: stroma fraction from spinach. Lane 7: thylakoid fraction from spinach. The amount loaded onto the gels accounts for ~0.15%, 0.15%, and 10% of the stroma, thylakoid, and envelope fractions, respectively. The arrowheads in A indicate the polypeptides which correspond to the major calmodulin-binding polypeptides (apparent *M_r* 33,000) in B and C. The Coomassie-Blue-stained gel in A corresponds to the autoradiogram in B. Nearly identical polypeptide profiles were obtained for the gel corresponding to the autoradiogram in C (data not shown).




FIGURE 2 Inhibition of ^{125}I -calmodulin binding to the major calmodulin-binding protein (M_r 33,000) in spinach chloroplast envelopes. A, B, and C show autoradiograms (10-h exposure). The Coomassie-Blue-stained patterns were indistinguishable from those shown in Fig. 1A. Gel overlay analysis was performed in the presence of 0.1 mM CaCl_2 and in the absence of competing protein (A), or in the presence

of a 1,000-fold molar excess of unlabeled chicken gizzard calmodulin (B) or a 1,000-fold molar excess of unlabeled bovine α -lactalbumin (C). The gels in each panel contain equal quantities of an envelope membrane fraction from spinach chloroplasts.

brane polypeptide (M_r 33,000) is inhibited by pre-incubation with a 1,000-fold molar excess of unlabeled chicken gizzard calmodulin (Fig. 2B). In contrast, the binding of ^{125}I -calmodulin is only weakly inhibited, if at all, by a 1,000-fold molar excess of α -lactalbumin (Fig. 2C), a small, acidic, calcium-binding protein that has no known structural or functional relationship to calmodulin or other calcium-modulated proteins. These observations indicate that the interaction of ^{125}I -calmodulin with the major envelope binding protein is not an artifact of the labeling protocol, and that this binding protein is selective in its interaction with acidic calcium-binding proteins. Although these observations suggest that the interaction of calmodulin with the envelope binding proteins is relatively specific, it must be recognized that calmodulin may not be the endogenous receptor for these binding proteins. For example, calmodulin can substitute for troponin C under *in vitro* conditions as the calcium-sensitizing component of muscle actomyosin ATPase (24, 25). Therefore, on the basis of precedents with animal systems, it is possible to conjecture that a calcium-modulated protein similar to calmodulin may be the physiologically relevant protein.

Our results suggest that the chloroplast envelope may be a primary target of calcium regulation in the chloroplast. The envelope membrane is an integral component of the chloroplast, and participates in several chloroplast processes including the transport of metabolites (26), the uptake and incorporation of cytoplasmically synthesized proteins (27), and a number of metabolic activities, such as fatty acid metabolism (28, 29), galactolipid biosynthesis (30), prenylquinone biosynthesis (31), and carotenoid biosynthesis (32, 33). The calcium dependence of these activities has not been investigated. Simon et al. (6) investigated thylakoid, envelope, and stroma fractions from spinach chloroplasts for calmodulin-stimulated NAD kinase activity. Nearly all chloroplast NAD kinase activity was present in the stroma fraction and was not stimulated by calmodulin.

Data regarding the structure and function of the polypeptides of the chloroplast envelope are limited. Of particular relevance to the present work are the results of Flugge and Heldt (34–36) which suggest that a phosphate translocator activity is associated with the major polypeptide (M_r 29,000) in their envelope preparation. Whether the polypeptide of

Flugge and Heldt corresponds to the major calmodulin-binding protein (M_r 33,000) in our envelope preparation remains to be determined.

The demonstration of calmodulin-binding proteins in the chloroplast envelope is an important step towards identifying the site of calmodulin regulation of chloroplast function. Clearly, more detailed analyses of the interactions between calmodulin and envelope proteins are necessary before any physiological significance can be attributed to these observations. However, as more becomes known regarding the structure and function of the envelope polypeptides, and the effect of calmodulin and calcium on their activities, a more defined role for calmodulin in the regulation of chloroplast processes may emerge.

We would like to thank Wilson Burgess and Linda Van Eldik for their helpful suggestions and their critical review of this manuscript.

This work was supported by National Institutes of Health grant GM 30861 and by National Science Foundation grant PCM 8242875.

Received for publication 27 July 1983, and in revised form 16 August 1983.

REFERENCES

- Schleicher, M., D. B. Iverson, L. J. Van Eldik, and D. M. Watterson. 1982. Calmodulin. In *The Cytoskeleton in Plant Growth and Development*. C. W. Lloyd, editor. Academic Press, Inc., London. 85–106.
- Van Eldik, L. J., J. G. Zenguei, D. R. Marshak, and D. M. Watterson. 1982. Calcium-binding proteins and the molecular basis of calcium action. *Int. Rev. Cytol.* 77:1–61.
- Klee, C. B., and T. C. Vanaman. 1982. Calmodulin. *Adv. Protein Chem.* 35:213–321.
- Watterson, D. M., W. H. Burgess, T. J. Lukas, D. Iverson, D. R. Marshak, M. Schleicher, B. W. Erickson, K.-F. Fok, and L. J. Van Eldik. 1983. Towards a molecular and atomic anatomy of calmodulin and calmodulin binding proteins. *Adv. Cycl. Nucleotide Res.* 16:205–226.
- Anderson, J. M., H. Charbonneau, H. P. Jones, R. O. McCann, and M. J. Cormier. 1980. Characterization of the plant nicotinamide adenine dinucleotide kinase activator protein and its identification as calmodulin. *Biochemistry*. 19:3113–3120.
- Simon, P., P. Dieter, M. Bonzon, H. Greppin, and D. Marme. 1982. Calmodulin-dependent and independent NAD kinase activities from cytoplasmic and chloroplastic fractions of spinach (*Spinacia oleracea* L.). *Plant Cell Reports*. 1:119–122.
- Dieter, P., and D. Marme. 1981. A calmodulin-dependent, microsomal ATPase from corn (*Zea mays*, L.). *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 125:245–248.
- Van Eldik, L. J., and W. H. Burgess. 1983. Analytical subcellular distribution of calmodulin and calmodulin-binding proteins in normal and virus-transformed fibroblasts. *J. Biol. Chem.* 258:4539–4547.
- Pardue, R. L., M. A. Kaetzel, S. H. Hahn, B. R. Brinkley, and J. D. Dedman. 1981. The identification of calmodulin-binding sites on mitochondria in cultured 3T3 cells. *Cell*. 23:533–542.
- Jarrett, H. W., C. J. Brown, C. C. Black, and M. J. Cormier. 1982. Evidence that calmodulin is in the chloroplast of peas and serves a regulatory role in photosynthesis. *J. Biol. Chem.* 257:13795–13804.
- Muto, S. 1982. Distribution of calmodulin in wheat leaf cells. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 147:161–164.
- Cline, K., J. Andrews, B. Mersey, E. H. Newcomb, and K. Keegstra. 1981. Separation and characterization of inner and outer envelope membranes of pea chloroplasts. *Proc. Natl. Acad. Sci. USA*. 78:3595–3599.
- Morgenthaler, J. J., C. A. Price, J. M. Robinson, and M. Gibbs. 1974. Photosynthetic activity of spinach chloroplasts after isopycnic centrifugation in gradients of silica. *Plant Physiol. (Bethesda)*. 54:532–534.
- Mendiola-Morgenthaler, L. R., and J. J. Morgenthaler. 1974. Proteins of the envelope and thylakoid membranes of spinach chloroplasts. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 49:152–155.
- Van Eldik, L. J., and D. M. Watterson. 1979. Characterization of a calcium-modulated protein from transformed chicken fibroblasts. *J. Biol. Chem.* 254:10250–10255.
- Marshak, D. R., D. M. Watterson, and L. J. Van Eldik. 1981. Calcium-dependent interaction of S100b, troponin C, and calmodulin with an immobilized phenothiazine. *Proc. Natl. Acad. Sci. USA*. 78:6793–6797.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (Lond.)*. 227:680–685.
- Matsudaira, P. J., and D. R. Burgess. 1978. SDS microslab linear gradient polyacrylamide gel electrophoresis. *Anal. Biochem.* 87:386–396.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the folin phenol reagent. *J. Biol. Chem.* 193:265–275.
- Markwell, M. K., S. M. Haas, L. L. Bieber, and N. E. Tolbert. 1978. A modification of the Lowry procedure to simplify protein determination in membrane and lipoprotein samples. *Anal. Biochem.* 87:206–210.
- Arnon, D. I. 1949. Copper enzymes in isolated chloroplasts. Polyphenol oxidase in *Beta vulgaris*. *Plant Physiol. (Bethesda)*. 24:1–15.
- Joyard, J., A. Grossman, S. G. Bartlett, R. Douce, and N.-H. Chua. 1982. Characterization of envelope membrane polypeptides from spinach chloroplasts. *J. Biol. Chem.* 257:1095–1101.
- Siegenthaler, P.-A., and T. D. Nguyen. 1983. Proteins and polypeptides of envelope membranes from spinach chloroplasts. I. Isoelectric focusing and sodium dodecyl sulfate polyacrylamide gel electrophoresis separations. *Biochim. Biophys. Acta*. 722:226–233.
- Amphlett, G. W., T. C. Vanaman, and S. V. Perry. 1976. Effect of troponin C-like

- protein from bovine brain (brain modulator protein) on the Mg^{2+} -stimulated ATPase of skeletal muscle actomyosin. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 72:163-168.
25. Dedman, J. R., J. D. Potter, and A. R. Means. 1977. Biological cross-reactivity of rat testis phosphodiesterase activator protein and rabbit skeletal muscle troponin-C. *J. Biol. Chem.* 252:2434-2440.
 26. Douce, R., and J. Joyard. 1979. Structure and function of the plastid envelope. *Adv. Bot. Res.* 7:1-116.
 27. Chua, N.-H., and G. W. Schmidt. 1979. Transport of proteins into mitochondria and chloroplasts. *J. Cell Biol.* 81:461-483.
 28. Joyard, J., and R. Douce. 1977. Site of synthesis of phosphatidic acid and diacylglycerol in spinach chloroplasts. *Biochim. Biophys. Acta.* 486:273-285.
 29. Joyard, J., and P. K. Stumpf. 1981. Synthesis of long chain Acyl-CoA in the chloroplast envelope membranes. *Plant Physiol. (Bethesda).* 67:250-256.
 30. Douce, R., and J. Joyard. 1980. Plant galactolipids. In *The Biochemistry of Plants*. P. K. Stumpf, editor. Academic Press, Inc., New York. 4:321-362.
 31. Soll, J., R. Douce, and G. Schultz. 1980. Site of biosynthesis of α -tocopherol in spinach chloroplasts. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 112:243-246.
 32. Costes, C., C. Burghoffer, J. Joyard, M. Block, and R. Douce. 1979. Occurrence and biosynthesis of violaxanthin in isolated spinach chloroplast envelope. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 103:17-21.
 33. Block, M. A., J. Joyard, and R. Douce. 1980. Site of synthesis of geranylgeraniol derivatives in intact spinach chloroplasts. *Biochim. Biophys. Acta.* 631:210-219.
 34. Flugge, U. I., and H. W. Heldt. 1976. Identification of a protein involved in phosphate transport of chloroplasts. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 68:259-262.
 35. Flugge, U. I., and H. W. Heldt. 1977. Specific labelling of a protein involved in phosphate transport of chloroplasts by pyridoxal-5'-phosphate. *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 82:29-33.
 36. Flugge, U. I., and H. W. Heldt. 1978. Specific labelling of the active site of the phosphate translocator in spinach chloroplasts by 2,4,6-trinitrobenzene sulfone. *Biochem. Biophys. Res. Commun.* 84:37-44.