



Article

Human Serum Proteins and Susceptibility of *Acinetobacter baumannii* to Cefiderocol: Role of Iron Transport

Casin Le ^{1,†}, Camila Pimentel ^{1,†}, Fernando Pasteran ², Marisel R. Tuttobene ^{3,4} , Tomás Subils ⁵, Jenny Escalante ¹, Brent Nishimura ¹, Susana Arriaga ¹, Aimee Carranza ¹, Vyanka Mezcord ¹ , Alejandro J. Vila ^{4,6} , Alejandra Corso ², Luis A. Actis ⁷ , Marcelo E. Tolmasky ¹ , Robert A. Bonomo ^{8,9,10,*} and Maria Soledad Ramírez ^{1,*}

- ¹ Center for Applied Biotechnology Studies, Department of Biological Science, College of Natural Sciences and Mathematics, California State University Fullerton, Fullerton, CA 92831, USA; thanhle1998@csu.fullerton.edu (C.L.); camilapimentel99@csu.fullerton.edu (C.P.); jenni1@csu.fullerton.edu (J.E.); bnish-942@csu.fullerton.edu (B.N.); arriagasusie@gmail.com (S.A.); aimeecarranza1119@gmail.com (A.C.); mezcordvyanka@csu.fullerton.edu (V.M.); mtolmasky@fullerton.edu (M.E.T.)
 - ² National/Regional Reference Laboratory for Antimicrobial Resistance (NRL), Servicio Antimicrobianos, Instituto Nacional de Enfermedades Infecciosas, ANLIS Dr. Carlos G. Malbrán, Buenos Aires C1282, Argentina; fpasteran@anlis.gov.ar (F.P.); acorso@anlis.gov.ar (A.C.)
 - ³ Área Biología Molecular, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario S2000, Argentina; tuttobene@ibr-conicet.gov.ar
 - ⁴ Instituto de Biología Molecular y Celular de Rosario (IBR, CONICET-UNR), Rosario S2000, Argentina; vila@ibr-conicet.gov.ar
 - ⁵ Instituto de Procesos Biotecnológicos y Químicos de Rosario (IPROBYQ, CONICET-UNR), Rosario S2002, Argentina; subils@iprobyq-conicet.gov.ar
 - ⁶ Área Biofísica, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario S2000, Argentina
 - ⁷ Department of Microbiology, Miami University, Oxford, OH 45056, USA; actisla@miamioh.edu
 - ⁸ Departments of Medicine, Pharmacology, Molecular Biology and Microbiology, Biochemistry, Proteomics and Bioinformatics, School of Medicine, Case Western Reserve University, Cleveland, OH 44106, USA
 - ⁹ Research Service and GRECC, Louis Stokes Cleveland Department of Veterans Affairs Medical Center, Cleveland, OH 44106, USA
 - ¹⁰ CWRU-Cleveland VAMC Center for Antimicrobial Resistance and Epidemiology (Case VA CARES), Cleveland, OH 44106, USA
- * Correspondence: robert.bonomo@va.gov (R.A.B.); msramirez@fullerton.edu (M.S.R.); Tel.: +1-216-791-3800 (R.A.B.); Tel.: +1-657-278-4562 (M.S.R.)
- † These authors have contributed equally to this work.



Citation: Le, C.; Pimentel, C.; Pasteran, F.; Tuttobene, M.R.; Subils, T.; Escalante, J.; Nishimura, B.; Arriaga, S.; Carranza, A.; Mezcord, V.; et al. Human Serum Proteins and Susceptibility of *Acinetobacter baumannii* to Cefiderocol: Role of Iron Transport. *Biomedicines* **2022**, *10*, 600. <https://doi.org/10.3390/biomedicines10030600>

Academic Editor: Amedeo Amedei

Received: 16 February 2022

Accepted: 1 March 2022

Published: 3 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Cefiderocol, a recently introduced antibiotic, has a chemical structure that includes a cephalosporin that targets cell wall synthesis and a chlorocatechol siderophore moiety that facilitates cell penetration by active iron transporters. Analysis of the effect that human serum, human serum albumin, and human pleural fluid had on growing *Acinetobacter baumannii* showed that genes related to iron uptake were down-regulated. At the same time, β -lactamase genes were expressed at higher levels. The minimum inhibitory concentrations of this antimicrobial in *A. baumannii* cells growing in the presence of human serum, human serum albumin, or human pleural fluid were higher than those measured when these fluids were absent from the culture medium. These results correlate with increased expression levels of β -lactamase genes and the down-regulation of iron uptake-related genes in cultures containing human serum, human serum albumin, or human pleural fluid. These modifications in gene expression could explain the less-than-ideal clinical response observed in patients with pulmonary or bloodstream *A. baumannii* infections. The exposure of the infecting cells to the host's fluids could cause reduced cefiderocol transport capabilities and increased resistance to β -lactams. The regulation of genes that could impact the *A. baumannii* susceptibility to cefiderocol, or other antibacterials, is an understudied phenomenon that merits further investigation.

Keywords: *Acinetobacter baumannii*; cefiderocol; human fluids; iron; human serum albumin

1. Introduction

Carbapenem-resistant *Acinetobacter baumannii* (CRAB), one of the most feared pathogens in healthcare settings, has been categorized by the Centers for Disease Control and Prevention (CDC) as an “urgent threat” [1–5]. As a consequence of *A. baumannii*'s ability to develop multidrug resistance (MDR), treatment strategies have become extremely limited, with only a few active antibiotics in existence [6–8].

The increasing number of CRAB infections translates into alarmingly high morbidity and mortality [2,9–11]. Consequently, numerous efforts are focusing on finding novel treatment options [12–19]. Cefiderocol (CFDC) was approved by the Food and Drug Administration of the United States (FDA) in November 2019 to treat nosocomial pneumonia and urinary tract infections (Available online: https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/209445s000lbl.pdf (accessed on 14 February 2022)) [20]. CFDC is a hybrid molecule that consists of a cephalosporin component that targets cell wall synthesis and a catechol siderophore moiety that allows cell penetration by active ferric-siderophore transporters [21–25]. This novel synthetic compound uses a “Trojan horse” strategy to improve antibiotic penetration and reach a high concentration at the target site [22,23]. This strategy allows the drug to enter the bacterial cells using active iron transporters. In *A. baumannii*, the most well-characterized iron transporters are *bauA*, *bfnH*, and *fbnN*, which are coupled to a TonB system to translocate the siderophore–iron complex into the periplasm [26]. However, up to 21 putative TonB-dependent outer membrane transporter genes have been identified or predicted in *A. baumannii* genomes, most often associated with putative or confirmed ferric-siderophore and heme uptake genes [27]. The *bauA* gene, which encodes for the major outer membrane receptor of ferric-acinetobactin complexes, was extensively studied in *A. baumannii* and was shown to be highly conserved [28–31]. In addition, PiuA and PirA, and also TonB-dependent receptors, were shown to mediate the uptake of siderophore–iron complexes [32]. These receptors have been associated with the cellular entry of CFDC in Gram-negative pathogens [25,33,34]. CFDC's primary targets are cell wall synthesizing proteins associated with β -lactam activity, such as the penicillin-binding proteins 1 and 3 (PBP1 and PBP3) [35]. Randomized clinical trials, and other small case series studies, showed CFDC as a promising alternative to treat carbapenemase-producing pathogens and other less frequent non-fermenters, such as *Stenotrophomonas maltophilia*, *Burkholderia* spp., and *Achromobacter* spp [7,22,36–41].

Simner et al. explored possible resistance mechanisms to CFDC in 14 isolates of carbapenem-resistant Enterobacterales that exhibited baseline CFDC resistance (minimum inhibitory concentration (MICs) ≥ 4 mg/L) [42]. They identified heterogeneous mechanisms that include the combination of β -lactamase production and permeability defects contributing to elevated CFDC MIC. Particularly, some of the mechanisms identified were the presence of New Delhi metallo- β -lactamase (NDM) carbapenemases, the occurrence of outer membrane porin mutations, and the presence of efflux transporters among the isolates evaluated. In addition, mutations in the *baeS* gene, responsible for encoding a sensor kinase protein of the two-component BaeSR and mutations in *tonB3*, *exbD*, *envZ*, or *ompR*, were identified among the clinical isolates that exhibited elevated CFDC MICs. The down-regulation of specific iron transport receptors in *Pseudomonas aeruginosa*, named Fe-cIRA, is correlated with resistance to CFDC [43]. Similarly, in *A. baumannii*, CFDC resistance was associated with the reduced expression of the TonB-dependent receptor gene *pirA* as well as mutations in the penicillin-binding proteins 3 (PBP₃). The presence of acquired β -lactamases, such as PER, may have also contributed to resistance in some strains [44,45]. So far, studies have not linked β -lactamase activity, nor levels of expression of PBPs, to CFDC resistance.

A. baumannii senses components of human fluids and responds by modifying its transcriptional and phenotypic profiles [46–50]. Human serum albumin (HSA), as well as human pleural fluid (HPF), modulates the expression of genes associated with iron uptake systems, biofilm formation, antibiotic resistance, and DNA acquisition, among others [47,51,52]. In previous studies, we observed that CRAB AB5075 genes associated with iron uptake systems were down-regulated when exposed to HPF and 0.2% HSA [47], while genes associated with β -lactam resistance were up-regulated in the presence of physiological concentrations of HSA and human serum (HS) [51,52]. Understanding the modification in the expression of iron uptake systems and β -lactam resistance genes in *A. baumannii* exposed to biological fluids could allow future strategies to protect antibiotics from a possible deleterious in vivo efficacy. In this work, we describe the effect of HS, HSA, and HPF on representative CRAB strains with different genetic backgrounds, focusing our analysis on the level of expression of iron uptake TonB-dependent receptors and β -lactam resistance genes.

2. Materials and Methods

2.1. Bacterial Strains

The model multidrug-resistant *A. baumannii* strain AB5075 was used [47,52,53]. In addition, *A. baumannii* AMA16 and AB0057 strains were also used [54–57]. The AB5075 strain contains the carbapenem resistance genes *bla*_{OXA-23}, *bla*_{GES-14}, and *bla*_{ADC}. The AMA16 strain presents *bla*_{NDM-1}, *bla*_{PER-7}, and *ISAba125* resistance genes. The AB0057 strain harbors the genes *bla*_{TEM-1}, *bla*_{OXA-23}, and *bla*_{ADC}. This strain was isolated in the Walter Reed Army Medical Center, Bethesda, MD 20814, USA [58]. In addition, all the strains presented conserved resistance genes such as *bla*_{OXA-51-like}, *carO*, *pbp1*, and *pbp3*. Nineteen additional CRAB strains, belonging to different clonal complexes, were included to performed CFDC susceptibility assays [54,55].

2.2. RNA Extraction and Quantitative Reverse Transcription Polymerase Chain Reaction (qRT-PCR)

Overnight cultures of *A. baumannii* strains were diluted 1:10 in fresh LB medium, LB medium containing 3.5% HSA (physiological concentration), containing 4% HPF, or 100% pooled normal HS, and incubated with agitation for 5 h at 37 °C. Pure HSA (Sigma-Aldrich, St. Louis, MO, USA) and pooled normal human serum from a certified vendor (Innovative Research Inc, Novi, MI, USA) were used in the cultures. RNA was extracted from each strain using the Direct-zol RNA Kit (Zymo Research, Irvine, CA, USA), following the manufacturer's instructions. Total RNA extractions were performed in three biological replicates for each condition. The integrity of the RNA samples was verified by agarose gel electrophoresis. The extracted and DNase-treated RNA was used to synthesize cDNA using the manufacturer's protocol provided within the iScript™ Reverse Transcription Supermix for qPCR (Bio-Rad, Hercules, CA, USA). The cDNA concentrations were adjusted to 50 ng/ μ L and qPCR was conducted using the qPCRBIO SyGreen Blue Mix Lo-ROX, following the manufacturer's protocol (PCR Biosystems, Wayne, PA, USA). At least three biological replicates of cDNA were used in triplets and were run using the CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad, Hercules, CA, USA). Transcriptional levels of each sample were normalized to the transcriptional level of *rpoB*. The relative quantification of gene expression was performed using the comparative threshold method $2^{-\Delta\Delta C_t}$. The ratios obtained after normalization were expressed as folds of change compared with cDNA samples isolated from the bacterial cultures in the LB. Asterisks indicate significant differences as determined by ANOVA, followed by Tukey's multiple comparison test ($p < 0.05$), using GraphPad Prism (GraphPad software, San Diego, CA, USA).

2.3. Antimicrobial Susceptibility Testing

Antibiotic susceptibility assays were performed following the procedures recommended by the Clinical and Laboratory Standards Institute (CLSI), Malvern, PA, USA [59]. After OD adjustment, 100 μ L of cells grown in fresh LB medium, LB medium contain-

ing 3.5% HSA, 4% HPF, or 100% HS were inoculated on Mueller–Hinton agar plates, as previously described [47,48,50]. Antimicrobial commercial E-strips (Liofilchem S.r.l., 64026 Roseto degli Abruzzi, Italy) for CFDC was used. Mueller–Hinton agar plates were incubated at 37 °C for 18 h. CLSI breakpoints were used for interpretation [60].

3. Results

3.1. Changes in the Expression Levels of Genes Associated with Iron Uptake Systems in the Presence of Human Fluids

Building upon on our previous observations on the AB5075 strain, we decided to further study the role of HS, HSA, and HPF on CRAB AB0057 and AMA16 strains. The selected strains belong to different clonal complexes and harbor clinically important β -lactamases, such as oxacillinases (OXA-23) and NDM-1, respectively [54,61], which are the most prevalent carbapenemases in *A. baumannii* and Enterobacterales, respectively [4,62,63]. Quantitative RT-PCR (qRT-PCR) assays, to evaluate the expression of the iron uptake TonB-dependent receptor *pirA*, *piuA*, and *bauA*, were carried out using total RNA extracted from cells cultured in lysogeny broth (LB), LB supplemented with 4% HPF and 3.5% HSA, and cells cultured in 100% HS, as previously described [47]. In addition, the analysis of the expression levels of other iron uptake-associated genes, such as *tonB3*, *basE*, *feoA*, *fhuE1*, and *fhuE2*, involved in transport and biosynthesis [64–68], was also performed.

In the CRAB AMA16, the expression of the iron transport-related genes *bauA*, *pirA*, and *piuA* was reduced in all tested conditions evaluated (Figure 1A and Table S1). Similar results were observed for the CRAB AB0057 strain. A reduction in the expression of *bauA*, *pirA*, and *piuA* in the tested condition was also seen (Figure 1B and Table S1). In addition, when the hypervirulent *A. baumannii* AB5075 strain was exposed to 3.5% HSA and the HS, the expression levels of the genes encoding for the above-mentioned TonB-receptor genes were down-regulated (Figure 1C and Table S1). Pimentel et al. previously showed that the levels of expression of AB5075 iron-related genes were down-regulated in cells cultured in iron-rich media [47]; this finding is in agreement with the level of expression of *basE* reported in this work (Figure S1B).

Analysis of available RNA-seq data of AB5075 cultivated in LB vs. 3.5% HSA showed that *fhuE1*, *fhuE2*, *pirA*, and *piuA* genes [47,68] were also down-regulated in the presence of HPF (Figure S1A). Moreover, the transcriptional levels of *fhuE1*, *fhuE2*, *tonB*, and *feoA* genes were further analyzed by qRT-PCR for the three strains (AMA16, AB0057, and AB5075) in the three tested conditions (HPF, HSA, and HS), showing a decrease in the level of expression in all of the strains in all of the tested conditions (Figure S1B), with the exception of an increase in the expression of *feoA* for AMA16 exposed to HPF (Figure S1).

3.2. Modulation of the Expression of β -Lactam Resistance Genes by Human Fluids

We previously observed that genes associated with β -lactam resistance, such as *bla*_{OXA-51-like}, *bla*_{OXA-23}, *pbp1*, and *pbp3*, were up-regulated in *A. baumannii* AB5075, AMA16, and AB0057 when cultured in the presence of 3.5% HSA (51, 52). To further examine if changes in genes associated with β -lactam resistance were affected by HS or HPF in the CRAB strains, qRT-PCR was used to assess the gene expression of *bla*_{OXA-51-like}, *bla*_{OXA-23}, *bla*_{NDM-1}, *pbp1*, *pbp3*, and *carO* in cultures of the strains AMA16 and AB0057 containing HS or HPF.

This transcriptional analysis revealed a down-regulation, with a log₂ fold change of -2.84 and -1.62 , in expression levels of *bla*_{NDM-1} in *A. baumannii* AMA16 cultured in the presence of HS or HPF, respectively (Figure 2A and Table S1). In this strain, the levels of expression of *ISAb125*, *bla*_{PER-7}, *pbp1*, and *pbp3* were up-regulated in the presence of HPF (Table S1). HS also induced the up-regulation of *ISAb125* (Figure 2A). In the case of the AB0057 strain, we observed an increase in the expression of *bla*_{OXA-23}, *bla*_{OXA-51}, *carO*, *bla*_{ADC}, *pbp1*, and *pbp3* in cultures containing HPF. In cultures containing HS, the former four genes were up-regulated (Figure 2B and Table S1). Assessment of the changes in β -lactam resistance-associated *A. baumannii* AB5075 genes in the presence of HPF showed the down-

regulation of the *carO* and *bla*_{GES-14} genes (Figure 2C). The *carO* gene encodes for an outer membrane protein channel that allows the permeation of imipenem in *A. baumannii* [69]. The marked down-regulation of *carO* (more than 4 Log₂ fold change, Table S1) could be contributing to limited CFDC entrance; however, further analysis is needed to establish this. On the other hand, the expression levels of *bla*_{OXA-51}, *pbp1*, *pbp3*, and *bla*_{ADC} were increased (Figure 2C).




Figure 1. Genetic analysis of iron uptake genes of AMA16 (A), AB0057 (B), and AB5075 (C) *A. baumannii* strains and qRT-PCR of genes associated with iron uptake, *pirA*, *piuA*, and *bauA*, expressed in LB, LB supplemented with HPF or with HSA, or cultured in HS. Fold changes were calculated using double Δ Ct analysis. At least three independent samples were used, and four technical replicates were performed from each sample. The LB was used as reference. Statistical significance ($p < 0.05$) was determined by ANOVA followed by Tukey's multiple comparison test: one asterisk: $p < 0.05$; two asterisks: $p < 0.01$; and three asterisks: $p < 0.001$.

The impact of human fluids in CFDC activity was further studied using a panel of twenty CRAB strains, including AB5075, AMA16, and AB0057. Cells cultured in LB, LB supplemented with HPF or 3.5% HSA, were used to determine the MICs of CFDC. Ten strains had baseline resistance to CFDC (Table S1).

An increase in the MIC values was observed in 14 strains when the cells were growing in the presence of HSA or HPF (Table 1 and Figure 2D). We draw attention to the occurrence of heteroresistant colonies, with the inhibition ellipse, in 11 of the tested strains (Table 1 and Figure S2). Most of the strains that exhibited this phenomenon harbored *bla*_{PER-7} [54]. Our results agree with a recent report indicating that PER-like β -lactamases contribute to a decreased susceptibility to CFDC in *A. baumannii* [45]. In addition, they observed that the combination of CFDC with avibactam may inhibit the activity of PER-type β -lactamases [45].




Figure 2. (A–C) Genetic analysis of β -lactamase and PBP genes of AMA16 (A), AB0057 (B), and AB5075 (C) *A. baumannii* strains and qRT-PCR of genes associated with β -lactams resistance expressed in LB, LB supplemented with HPF, or in HS. Fold changes were calculated using double Δ Ct analysis. At least three independent samples were used. LB was used as the reference condition. Statistical significance ($p < 0.05$) was determined by ANOVA followed by Tukey’s multiple comparison test: one asterisk: $p < 0.05$; two asterisks: $p < 0.01$; and three asterisks: $p < 0.001$. (D) Effect of HSA and HPF on the antimicrobial susceptibility of *A. baumannii* representative strains AMA40 and AMA113 grown in LB broth, LB broth plus 3.5% HSA, or HPF, that were used to perform cefiderocol (CFDC) susceptibility. Minimum inhibitory concentration (MIC) on cation-adjusted Mueller–Hinton agar was performed by MTS (Liofilchem S.r.l., 64026 Roseto degli Abruzzi TE, Italy), following the manufacturer’s recommendations.

Table 1. Minimal inhibitory concentrations of cefiderocol (CFDC) for 22 carbapenem-resistant *Acinetobacter baumannii* representative strains performed using CFDC MTS strips (Liofilchem S.r.l., 64026 Roseto degli Abruzzi TE, Italy) on Mueller–Hinton agar (cation adjusted). *A. baumannii* cells were cultured in LB or LB supplemented with 3.5% HSA or HPF, respectively.

Strain	CFDC MICs (mg/L)		
	LB	HPF	3.5% HSA
AB5075	0.5 (S)	1 (S)	2 (S)
ABUH702	0.38 (S)	1.5 (S)	3 (S)
AMA16	>4.5 * (I)	>256 (R)	32 * (R)
AB0057	1 (S)	8 (I)	1.5 (S)
AMA40	0.5 (S)	16 * (R)	3 (S)
AMA41	0.094 (S)	0.5–0.75 (S)	2 (S)
AMA113	0.5 (S)	1.5 (S)	1.5 (S)
AMA181	0.19 (S)	0.19 (S)	0.75 (S)
AMA3	24 (R)	>256 (R)	32 * (R)
AMA4	16 * (R)	48 * (R)	64 * (R)
AMA5	>256 (R)	>256 (R)	16 * (R)
AMA9	32 (R)	48 (R)	16 (R)
AMA14	8 * (I)	16 * (R)	12 (I)
AMA17	>256 (R)	>256 (R)	>256 (R)
AMA18	64 * (R)	16 * (R)	16 * (R)
AMA19	4 (S)	4 (S)	48 * (R)
AMA28	32 * (R)	>256 (R)	32 * (R)
AMA30	64 * (R)	128 * (R)	12 * (I)
AMA31	>256 (R)	>256 (R)	96 * (R)
AMA33	16 * (R)	>256 (R)	>256 (R)

* Intra-colonies are present. S: Susceptible, I: Intermediate, R: Resistant.

4. Discussion

Immediately after the invasion of infecting bacteria, the body responds by reducing free iron levels in the blood and tissues (hypoferremic response) [70–72]. In the face of these nutritional limitations, microorganisms evolve strategies to scavenge sufficient trace elements necessary to support their metabolism and growth. Among them, siderophore-mediated iron uptake systems are widespread among bacterial pathogens and are the systems utilized by CFDC to increase cell entrance.

Previous reports support a link between mutations in iron transporters and a decrease in susceptibility to CFDC [33,42,44]. In particular, PirA and PiuA have been linked with CFDC entry and CFDC resistance when mutations in these genes have been reported [33,34]. Our results show a decrease in the expression of genes coding for the TonB-dependent receptors BauA, PirA, and PiuA, when CRAB strains are exposed to human fluids and HSA. In addition, the increased expression of β -lactam resistance-associated genes, in particular, the increased expression of genes encoding PBPs in the three tested conditions, was also observed. These two observations are coupled with elevated CFDC MICs in 14 CRAB isolates when exposed to HPF and/or HSA.

Supporting our observations, recent studies have shown that the down-regulation of *pirA* and *piuA* in three *A. baumannii* clinical strains is associated with resistance to CFDC [33,44]. In addition, this work showed that the alteration of the hydrophobic isoleucine to the charged asparagine, at position 236 of PBP3, could also contribute to the CFDC resistance [44].

In other Gram-negatives, the relationship between mutations in TonB-dependent catecholate siderophore receptors and CFDC was observed. A recent case report showed the development of resistance to CFDC in a patient with intra-abdominal and bloodstream infections caused by a CR-*Enterobacter cloacae* strain after 21 days of CFDC treatment. Whole-genome sequencing revealed heterogeneous mutations in the *cirA* gene, which encodes a TonB-dependent catecholate siderophore receptor, conferring phenotypic resistance to cefiderocol. *cirA* was the only catecholate receptor present in that clinical isolate,

supporting the role of the mutation of this gene in CFDC resistance [73]. Moreover, in *Pseudomonas aeruginosa*, mutations in *piuD* and *pirR*, major iron transport pathways, and a leucine-to-phenylalanine substitution at amino acid position 147 in the ADC β -lactamase, were associated with CFDC MIC increases [73,74]. Simner et al. reported that substitutions in the region of the AmpC omega loop contribute to increased cefiderocol MICs to *P. aeruginosa* [75].

The present work focuses on the effects of different human bodily fluids on iron uptake and antibiotic resistance genes, some of which were linked with the reduced susceptibility to CFDC. Our results show a down-regulation or up-regulation of the genes associated with iron uptake or β -lactam resistance. These alterations were seen among the different strains and under different conditions. We believe that the sum of all the changes may contribute to the reduced effectiveness of CFDC when studied in vitro.

The exposure of *A. baumannii* to human fluids results in the transcriptional repression of genes coding for proteins involved in TonB-dependent siderophore-mediated iron acquisition. Taking into account these observations and the fact that HPF and HS possess a high concentration of HSA, a protein that has been reported as an important host iron reservoir [76], it is possible to speculate that the reduced efficacy of CFDC in the presence of these fluids is a consequence of the iron-rich environment created by the presence of HSA. In the case of HPF, we cannot discount the possibility that other proteins such as ferritin, which are abundant in HPF, may provide the necessary metal ions through this source, reducing the expression of the catechol iron uptake system and the number of entry sites for cefiderocol, leading to diminished efficacy.

5. Conclusions

Overall, our results show that *A. baumannii* may be unique in its response to human bodily fluids by modulating the expression of iron uptake genes and β -lactam resistance-associated determinants. We hypothesize that the presence of iron-binding proteins in the human fluids is sensed by *A. baumannii*, which in turn responds by down-regulating the expression of iron uptake system genes, impacting CFDC activity. The particular changes in expression of the aforementioned group of genes, which is not evaluated in traditional susceptibility tests, may contribute to increases in MICs. These results also raise questions regarding the expression of other factors that may contribute to challenges in overcoming infections by *A. baumannii*.

The limitations inherent in performing MICs in the presence of bodily fluids (protein binding) are recognized. We are also aware that comparing these analyses does not take into account the pharmacokinetic and pharmacodynamic (PK/PD) parameters that ensure drug efficacy [36,77]. Lastly, we appreciate that these data must also be evaluated in the context of outcomes in animal models of infection [78–80]. Nevertheless, the impact on iron transport mechanisms uncovered in these observations merits further analyses.

Supplementary Materials: The following supporting information can be downloaded at <https://www.mdpi.com/article/10.3390/biomedicines10030600/s1>. Table S1. Comparison of the level of expression of iron associated and β -lactam resistance genes obtained by qRT-PCR in the three CRAB strains. Figure S1. (A) Heatmap outlining the differential expression of genes, *fhuE1*, *fhuE2*, *pirA* and *piuA* associated with iron uptake in presence of 4% HPF or 0.2% HSA. The asterisks represent the DEGs (adjusted *p*-value < 0.05 with \log_2 fold change >1). (B) Differential expression of genes associated with iron-uptake obtained for *A. baumannii* AMA16, AB0057 and AB5075 strains cultured in the presence of HPF, HSA or HS. Figure S2. AMA40 CFDC MICs displaying colonies within the E-test zone and the MICs results of the selected intra colonies (IHC1 and IHC2). CFDC MICs were performed following the CLSI guidelines using CAMHA. The intracolony obtained when AMA40 was exposed to 4% HPF were subculture and the CFDC MIC was performed.

Author Contributions: C.P., C.L., F.P., M.R.T., T.S., R.A.B. and M.S.R. conceived the study and designed the experiments. C.P., C.L., F.P., M.R.T., T.S., J.E., B.N., S.A., A.C. (Aimee Carranza), V.M., A.J.V., A.C. (Alejandra Corso), L.A.A., M.E.T., R.A.B. and M.S.R. performed the experiments and genomics and bioinformatics analyses. C.P., C.L., F.P., M.R.T., T.S., A.J.V., L.A.A., R.A.B., M.E.T.

and M.S.R. analyzed the data and interpreted the results. R.A.B., M.E.T. and M.S.R. contributed reagents/materials/analysis tools. F.P., M.R.T., T.S., L.A.A., R.A.B., M.E.T. and M.S.R. wrote and revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: The authors' work was supported by NIH SC3GM125556 to M.S.R., R01AI100560, R01AI063517, R01AI072219 to R.A.B. and 2R15 AI047115 to M.E.T. This study was supported in part by funds and/or facilities provided by the Cleveland Department of Veterans Affairs, award number 1I01BX001974, to R.A.B. from the Biomedical Laboratory Research and Development Service of the VA Office of Research and Development, and the Geriatric Research Education and Clinical Center VISN 10 to R.A.B., C.P. and J.E. were supported by grant MHRT 2T37MD001368 from the National Institute on Minority Health and Health Disparities, National Institute of Health. S.A. and A.C. (Aimee Carranza) were supported by Project RAISE, U.S. Department of Education HSI-STEM, award number P031C160152. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the Department of Veterans Affairs. M.R.T. and T.S. are recipients of a postdoctoral fellowship from CONICET. A.J.V. is a staff member from CONICET.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data pertaining to the study described in the manuscript is described in the report.

Acknowledgments: We are grateful to Michael R. Jacobs for his critical reading of the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Eichenberger, E.M.; Thaden, J.T. Epidemiology and Mechanisms of Resistance of Extensively Drug Resistant Gram-Negative Bacteria. *Antibiotics* **2019**, *8*, 37. [\[CrossRef\]](#)
2. Holmes, C.L.; Anderson, M.T.; Mobley, H.L.T.; Bachman, M.A. Pathogenesis of Gram-negative bacteremia. *Clin. Microbiol. Rev.* **2021**, *34*, e00234–e20. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Garcia-Ortega, L.; Arch, O.; Perez-Canosa, C.; Lupion, C.; Gonzalez, C.; Rodriguez-Bano, J.; Spanish Study Group of Nosocomial Infections. Control measures for *Acinetobacter baumannii*: A survey of Spanish hospitals. *Enferm. Infecc. Microbiol. Clín.* **2011**, *29*, 36–38. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Ramirez, M.S.; Bonomo, R.A.; Tolmasky, M.E. Carbapenemases: Transforming *Acinetobacter baumannii* into a Yet More Dangerous Menace. *Biomolecules* **2020**, *10*, 720. [\[CrossRef\]](#) [\[PubMed\]](#)
5. CDC. *Antibiotic Resistance Threats in the United States*; US Department of Health and Human Services, CDC: Atlanta, GA, USA, 2019.
6. Karaiskos, I.; Lagou, S.; Pontikis, K.; Rapti, V.; Poulakou, G. The “Old” and the “New” Antibiotics for MDR Gram-Negative Pathogens: For Whom, When, and How. *Front. Public Health* **2019**, *7*, 151. [\[CrossRef\]](#)
7. Bassetti, M.; Echols, R.; Matsunaga, Y.; Ariyasu, M.; Doi, Y.; Ferrer, R.; Lodise, T.P.; Naas, T.; Niki, Y.; Paterson, D.L.; et al. Efficacy and safety of cefiderocol or best available therapy for the treatment of serious infections caused by carbapenem-resistant Gram-negative bacteria (CREDIBLE-CR): A randomised, open-label, multicentre, pathogen-focused, descriptive, phase 3 trial. *Lancet Infect. Dis.* **2021**, *21*, 226–240. [\[CrossRef\]](#)
8. Bassetti, M.; Peghin, M.; Vena, A.; Giacobbe, D.R. Treatment of Infections Due to MDR Gram-Negative Bacteria. *Front. Med.* **2019**, *6*, 74. [\[CrossRef\]](#)
9. Isler, B.; Doi, Y.; Bonomo, R.A.; Paterson, D.L. New treatment options against carbapenem-resistant *Acinetobacter baumannii* infections. *Antimicrob. Agents Chemother.* **2019**, *63*, e01110–e18. [\[CrossRef\]](#)
10. Bonomo, R.A.; Burd, E.M.; Conly, J.; Limbago, B.M.; Poirel, L.; Segre, J.A.; Westblade, L.F. Carbapenemase-Producing Organisms: A Global Scourge. *Clin. Infect. Dis.* **2018**, *66*, 1290–1297. [\[CrossRef\]](#)
11. Spellberg, B.; Bonomo, R.A. The deadly impact of extreme drug resistance in *Acinetobacter baumannii*. *Crit. Care Med.* **2014**, *42*, 1289–1291. [\[CrossRef\]](#)
12. Paterson, D.L.; Isler, B.; Stewart, A. New treatment options for multiresistant gram negatives. *Curr. Opin. Infect. Dis.* **2020**, *33*, 214–223. [\[CrossRef\]](#) [\[PubMed\]](#)
13. McLeod, S.M.; Moussa, S.H.; Hackel, M.A.; Miller, A.A. In Vitro Activity of Sulbactam-Durlobactam against *Acinetobacter baumannii-calcoaceticus* Complex Isolates Collected Globally in 2016 and 2017. *Antimicrob. Agents Chemother.* **2020**, *64*, e02534–e19. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Lomovskaya, O.; Nelson, K.; Rubio-Aparicio, D.; Tsivkovski, R.; Sun, D.; Dudley, M.N. The Impact of Intrinsic Resistance Mechanisms on Potency of QPX7728, a New Ultra-Broad-Spectrum Beta-lactamase Inhibitor of Serine and Metallo Beta-Lactamases in *Enterobacteriaceae*, *Pseudomonas aeruginosa*, and *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **2020**, *65*, e00552–e20.

15. Karaiskos, I.; Galani, I.; Souli, M.; Giamarellou, H. Novel beta-lactam-beta-lactamase inhibitor combinations: Expectations for the treatment of carbapenem-resistant Gram-negative pathogens. *Expert Opin. Drug Metab. Toxicol.* **2019**, *15*, 133–149. [[CrossRef](#)] [[PubMed](#)]
16. Barnes, M.D.; Kumar, V.; Bethel, C.R.; Moussa, S.H.; O'Donnell, J.; Rutter, J.D.; Good, C.E.; Hujer, K.M.; Hujer, A.M.; Marshall, S.H.; et al. Targeting Multidrug-Resistant *Acinetobacter* spp.: Sulbactam and the Diazabicyclooctenone beta-Lactamase Inhibitor ETX2514 as a Novel Therapeutic Agent. *mBio* **2019**, *10*, e00159-e19. [[CrossRef](#)] [[PubMed](#)]
17. Zhanel, G.G.; Lawrence, C.K.; Adam, H.; Schweizer, F.; Zelenitsky, S.; Zhanel, M.; Lagace-Wiens, P.R.S.; Walkty, A.; Denisuik, A.; Golden, A.; et al. Imipenem-Relebactam and Meropenem-Vaborbactam: Two Novel Carbapenem-beta-Lactamase Inhibitor Combinations. *Drugs* **2018**, *78*, 65–98. [[CrossRef](#)] [[PubMed](#)]
18. Wright, H.; Bonomo, R.A.; Paterson, D.L. New agents for the treatment of infections with Gram-negative bacteria: Restoring the miracle or false dawn? *Clin. Microbiol. Infect.* **2017**, *23*, 704–712. [[CrossRef](#)] [[PubMed](#)]
19. Durand-Reville, T.F.; Guler, S.; Comita-Prevoir, J.; Chen, B.; Bifulco, N.; Huynh, H.; Lahiri, S.; Shapiro, A.B.; McLeod, S.M.; Carter, N.M.; et al. ETX2514 is a broad-spectrum beta-lactamase inhibitor for the treatment of drug-resistant Gram-negative bacteria including *Acinetobacter baumannii*. *Nat. Microbiol.* **2017**, *2*, 17104. [[CrossRef](#)]
20. Lee, Y.R.; Yeo, S. Cefiderocol, a New Siderophore Cephalosporin for the Treatment of Complicated Urinary Tract Infections Caused by Multidrug-Resistant Pathogens: Preclinical and Clinical Pharmacokinetics, Pharmacodynamics, Efficacy and Safety. *Clin. Drug Investig.* **2020**, *40*, 901–913. [[CrossRef](#)]
21. Parsels, K.A.; Mastro, K.A.; Steele, J.M.; Thomas, S.J.; Kufel, W.D. Cefiderocol: A novel siderophore cephalosporin for multidrug-resistant Gram-negative bacterial infections. *J. Antimicrob. Chemother.* **2021**, *76*, 1379–1391. [[CrossRef](#)]
22. Bonomo, R.A. Cefiderocol: A Novel Siderophore Cephalosporin Defeating Carbapenem-resistant Pathogens. *Clin. Infect. Dis.* **2019**, *69*, S519–S520. [[CrossRef](#)] [[PubMed](#)]
23. Zhanel, G.G.; Golden, A.R.; Zelenitsky, S.; Wiebe, K.; Lawrence, C.K.; Adam, H.J.; Idowu, T.; Domalaon, R.; Schweizer, F.; Zhanel, M.A.; et al. Cefiderocol: A Siderophore Cephalosporin with Activity Against Carbapenem-Resistant and Multidrug-Resistant Gram-Negative Bacilli. *Drugs* **2019**, *79*, 271–289. [[CrossRef](#)] [[PubMed](#)]
24. Aoki, T.; Yoshizawa, H.; Yamawaki, K.; Yokoo, K.; Sato, J.; Hisakawa, S.; Hasegawa, Y.; Kusano, H.; Sano, M.; Sugimoto, H.; et al. Cefiderocol (S-649266), A new siderophore cephalosporin exhibiting potent activities against *Pseudomonas aeruginosa* and other gram-negative pathogens including multi-drug resistant bacteria: Structure activity relationship. *Eur. J. Med. Chem.* **2018**, *155*, 847–868. [[CrossRef](#)] [[PubMed](#)]
25. Ito, A.; Nishikawa, T.; Matsumoto, S.; Yoshizawa, H.; Sato, T.; Nakamura, R.; Tsuji, M.; Yamano, Y. Siderophore Cephalosporin Cefiderocol Utilizes Ferric Iron Transporter Systems for Antibacterial Activity against *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **2016**, *60*, 7396–7401. [[CrossRef](#)]
26. Sheldon, J.R.; Skaar, E.P. *Acinetobacter baumannii* can use multiple siderophores for iron acquisition, but only acinetobactin is required for virulence. *PLoS Pathog.* **2020**, *16*, e1008995. [[CrossRef](#)] [[PubMed](#)]
27. Antunes, L.C.; Imperi, F.; Towner, K.J.; Visca, P. Genome-assisted identification of putative iron-utilization genes in *Acinetobacter baumannii* and their distribution among a genotypically diverse collection of clinical isolates. *Res. Microbiol.* **2011**, *162*, 279–284. [[CrossRef](#)]
28. Nwugo, C.C.; Arivett, B.A.; Zimble, D.L.; Gaddy, J.A.; Richards, A.M.; Actis, L.A. Effect of ethanol on differential protein production and expression of potential virulence functions in the opportunistic pathogen *Acinetobacter baumannii*. *PLoS ONE* **2012**, *7*, e51936. [[CrossRef](#)]
29. Zimble, D.L.; Penwell, W.F.; Gaddy, J.A.; Menke, S.M.; Tomaras, A.P.; Connerly, P.L.; Actis, L.A. Iron acquisition functions expressed by the human pathogen *Acinetobacter baumannii*. *Biomaterials* **2009**, *22*, 23–32. [[CrossRef](#)]
30. Ramirez, M.S.; Penwell, W.F.; Traglia, G.M.; Zimble, D.L.; Gaddy, J.A.; Nikolaidis, N.; Arivett, B.A.; Adams, M.D.; Bonomo, R.A.; Actis, L.A.; et al. Identification of potential virulence factors in the model strain *Acinetobacter baumannii* A118. *Front. Microbiol.* **2019**, *10*, 1599. [[CrossRef](#)]
31. Moynié, L.; Serra, I.; Scorciapino, M.A.; Oueis, E.; Page, M.G.; Ceccarelli, M.; Naismith, J.H. Preacinetobactin not acinetobactin is essential for iron uptake by the BauA transporter of the pathogen. *eLife* **2018**, *7*, e42270. [[CrossRef](#)]
32. Ghysels, B.; Ochsner, U.; Möllman, U.; Heinisch, L.; Vasil, M.; Cornelis, P.; Matthijs, S. The *Pseudomonas aeruginosa* pirA gene encodes a second receptor for ferrienterobactin and synthetic catecholate analogues. *FEMS Microbiol. Lett.* **2005**, *246*, 167–174. [[CrossRef](#)] [[PubMed](#)]
33. Moynié, L.; Luscher, A.; Rolo, D.; Pletzer, D.; Tortajada, A.; Weingart, H.; Braun, Y.; Page, M.G.; Naismith, J.H.; Kohler, T. Structure and Function of the PiuA and PirA Siderophore-Drug Receptors from *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **2017**, *61*, e02531-e16. [[CrossRef](#)] [[PubMed](#)]
34. Luscher, A.; Moynié, L.; Auguste, P.S.; Bumann, D.; Mazza, L.; Pletzer, D.; Naismith, J.H.; Köhler, T. TonB-Dependent Receptor Repertoire of *Pseudomonas aeruginosa* for Uptake of Siderophore-Drug Conjugates. *Antimicrob. Agents Chemother.* **2018**, *62*, e00097-e18. [[CrossRef](#)] [[PubMed](#)]
35. Sato, T.; Yamawaki, K. Cefiderocol: Discovery, Chemistry, and In Vivo Profiles of a Novel Siderophore Cephalosporin. *Clin. Infect. Dis.* **2019**, *69*, S538–S543. [[CrossRef](#)]
36. McCreary, E.K.; Heil, E.L.; Tamma, P.D. New Perspectives on Antimicrobial Agents: Cefiderocol. *Antimicrob. Agents Chemother.* **2021**, *65*, e02171-e20. [[CrossRef](#)]

37. Wunderink, R.G.; Matsunaga, Y.; Ariyasu, M.; Clevenbergh, P.; Echols, R.; Kaye, K.S.; Kollef, M.; Menon, A.; Pogue, J.M.; Shorr, A.F.; et al. Cefiderocol versus high-dose, extended-infusion meropenem for the treatment of Gram-negative nosocomial pneumonia (APEKS-NP): A randomised, double-blind, phase 3, non-inferiority trial. *Lancet Infect. Dis.* **2021**, *21*, 213–225. [[CrossRef](#)]
38. Jean, S.S.; Gould, I.M.; Lee, W.S.; Hsueh, P.R.; International Society of Antimicrobial Chemotherapy. New Drugs for Multidrug-Resistant Gram-Negative Organisms: Time for Stewardship. *Drugs* **2019**, *79*, 705–714. [[CrossRef](#)]
39. Jacobs, M.R.; Abdelhamed, A.M.; Good, C.E.; Rhoads, D.D.; Hujer, K.M.; Hujer, A.M.; Domitrovic, T.N.; Rudin, S.D.; Richter, S.S.; van Duin, D.; et al. ARGONAUT-I: Activity of Cefiderocol (S-649266), a Siderophore Cephalosporin, against Gram-Negative Bacteria, Including Carbapenem-Resistant Nonfermenters and Enterobacteriaceae with Defined Extended-Spectrum beta-Lactamases and Carbapenemases. *Antimicrob. Agents Chemother.* **2019**, *63*, e01801-e18. [[CrossRef](#)] [[PubMed](#)]
40. Bavaro, D.F.; Belati, A.; Diella, L.; Stufano, M.; Romanelli, F.; Scalone, L.; Stolfa, S.; Ronga, L.; Maurmo, L.; Dell’Aera, M.; et al. Cefiderocol-Based Combination Therapy for “Difficult-to-Treat” Gram-Negative Severe Infections: Real-Life Case Series and Future Perspectives. *Antibiotics* **2021**, *10*, 652. [[CrossRef](#)]
41. Oliva, A.; Ceccarelli, G.; De Angelis, M.; Sacco, F.; Miele, M.C.; Mastroianni, C.M.; Venditti, M. Cefiderocol for compassionate use in the treatment of complicated infections caused by extensively and pan-resistant *Acinetobacter baumannii*. *J. Glob. Antimicrob. Resist.* **2020**, *23*, 292–296. [[CrossRef](#)]
42. Simner, P.J.; Beisken, S.; Bergman, Y.; Ante, M.; Posch, A.E.; Tamma, P.D. Defining Baseline Mechanisms of Cefiderocol Resistance in the *Enterobacterales*. *Microb. Drug Resist.* **2022**, *28*, 161–170. [[CrossRef](#)]
43. Van Delden, C.; Page, M.G.; Köhler, T. Involvement of Fe uptake systems and AmpC β -lactamase in susceptibility to the siderophore monosulfactam BAL30072 in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **2013**, *57*, 2095–2102. [[CrossRef](#)]
44. Malik, S.; Kaminski, M.; Landman, D.; Quale, J. Cefiderocol Resistance in *Acinetobacter baumannii*: Roles of β -Lactamases, Siderophore Receptors, and Penicillin Binding Protein 3. *Antimicrob. Agents Chemother.* **2020**, *64*, e01221-20. [[CrossRef](#)]
45. Poirel, L.; Sadek, M.; Nordmann, P. Contribution of PER-type and NDM-type ss-lactamases to cefiderocol resistance in *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **2021**, *65*, e00877-e21. [[CrossRef](#)]
46. Martinez, J.; Razo-Gutierrez, C.; Le, C.; Courville, R.; Pimentel, C.; Liu, C.; Fung, S.E.; Tuttobene, M.R.; Phan, K.; Vila, A.J.; et al. Cerebrospinal fluid (CSF) augments metabolism and virulence expression factors in *Acinetobacter baumannii*. *Sci. Rep.* **2021**, *11*, 4737. [[CrossRef](#)]
47. Pimentel, C.; Le, C.; Tuttobene, M.R.; Subils, T.; Martinez, J.; Sieira, R.; Papp-Wallace, K.M.; Keppetipola, N.; Bonomo, R.A.; Actis, L.A.; et al. Human Pleural Fluid and Human Serum Albumin Modulate the Behavior of a Hypervirulent and Multidrug-Resistant (MDR) *Acinetobacter baumannii* Representative Strain. *Pathogens* **2021**, *10*, 471. [[CrossRef](#)]
48. Martinez, J.; Fernandez, J.S.; Liu, C.; Hoard, A.; Mendoza, A.; Nakanouchi, J.; Rodman, N.; Courville, R.; Tuttobene, M.R.; Lopez, C.; et al. Human pleural fluid triggers global changes in the transcriptional landscape of *Acinetobacter baumannii* as an adaptive response to stress. *Sci. Rep.* **2019**, *9*, 17251. [[CrossRef](#)]
49. Rodman, N.; Martinez, J.; Fung, S.; Nakanouchi, J.; Myers, A.L.; Harris, C.M.; Dang, E.; Fernandez, J.S.; Liu, C.; Mendoza, A.M.; et al. Human Pleural Fluid Elicits Pyruvate and Phenylalanine Metabolism in *Acinetobacter baumannii* to Enhance Cytotoxicity and Immune Evasion. *Front. Microbiol.* **2019**, *10*, 1581. [[CrossRef](#)]
50. Quinn, B.; Rodman, N.; Jara, E.; Fernandez, J.S.; Martinez, J.; Traglia, G.M.; Montana, S.; Cantera, V.; Place, K.; Bonomo, R.A.; et al. Human serum albumin alters specific genes that can play a role in survival and persistence in *Acinetobacter baumannii*. *Sci. Rep.* **2018**, *8*, 14741. [[CrossRef](#)]
51. Le, C.; Pimentel, C.; Tuttobene, M.R.; Subils, T.; Nishimura, B.; Traglia, G.M.; Perez, F.; Papp-Wallace, K.M.; Bonomo, R.A.; Tolmasky, M.E.; et al. Interplay between meropenem and human serum albumin on expression of carbapenem resistance genes and natural competence in *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **2021**, *65*, e01019-e21. [[CrossRef](#)]
52. Pimentel, C.; Le, C.; Tuttobene, M.R.; Subils, T.; Papp-Wallace, K.M.; Bonomo, R.A.; Tolmasky, M.E.; Ramirez, M.S. Interaction of *Acinetobacter baumannii* with Human Serum Albumin: Does the Host Determine the Outcome? *Antibiotics* **2021**, *10*, 833. [[CrossRef](#)]
53. Jacobs, A.C.; Thompson, M.G.; Black, C.C.; Kessler, J.L.; Clark, L.P.; McQueary, C.N.; Gancz, H.Y.; Corey, B.W.; Moon, J.K.; Si, Y.; et al. AB5075, a Highly Virulent Isolate of *Acinetobacter baumannii*, as a Model Strain for the Evaluation of Pathogenesis and Antimicrobial Treatments. *mBio* **2014**, *5*, e01076-e14. [[CrossRef](#)]
54. Adams, M.D.; Pasteran, F.; Traglia, G.M.; Martinez, J.; Huang, F.; Liu, C.; Fernandez, J.S.; Lopez, C.; Gonzalez, L.J.; Albornoz, E.; et al. Distinct mechanisms of dissemination of NDM-1 metallo- β -lactamase in *Acinetobacter* spp. in Argentina. *Antimicrob. Agents Chemother.* **2020**, *64*, e00324-20. [[CrossRef](#)]
55. Rodgers, D.; Pasteran, F.; Calderon, M.; Jaber, S.; Traglia, G.M.; Albornoz, E.; Corso, A.; Vila, A.J.; Bonomo, R.A.; Adams, M.D.; et al. Characterisation of ST25 NDM-1-producing *Acinetobacter* spp. strains leading the increase in NDM-1 emergence in Argentina. *J. Glob. Antimicrob. Resist.* **2020**, *23*, 108–110. [[CrossRef](#)]
56. Hujer, A.M.; Higgins, P.G.; Rudin, S.D.; Buser, G.L.; Marshall, S.H.; Xanthopoulou, K.; Seifert, H.; Rojas, L.J.; Domitrovic, T.N.; Cassidy, P.M.; et al. Nosocomial outbreak of extensively drug-resistant *Acinetobacter baumannii* isolates containing *bla*OXA-237 carried on a plasmid. *Antimicrob. Agents Chemother.* **2017**, *61*, e00797-e17. [[CrossRef](#)]
57. Adams, M.D.; Chan, E.R.; Molyneaux, N.D.; Bonomo, R.A. Genomewide analysis of divergence of antibiotic resistance determinants in closely related isolates of *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **2010**, *54*, 3569–3577. [[CrossRef](#)]

58. Hujer, K.M.; Hujer, A.M.; Hulten, E.A.; Bajaksouzian, S.; Adams, J.M.; Donskey, C.J.; Ecker, D.J.; Massire, C.; Eshoo, M.W.; Sampath, R.; et al. Analysis of antibiotic resistance genes in multidrug-resistant *Acinetobacter* sp. isolates from military and civilian patients treated at the Walter Reed Army Medical Center. *Antimicrob. Agents Chemother.* **2006**, *50*, 4114–4123. [[CrossRef](#)]
59. CLSI. *Performance Standards for Antimicrobial Susceptibility Testing*, 29th ed.; CLSI Document M100; Clinical Lab Standards Institute: Wayne, PA, USA, 2019.
60. CLSI. *Performance Standards for Antimicrobial Susceptibility Testing*, 30th ed.; CLSI Document M100; Clinical Lab Standards Institute: Wayne, PA, USA, 2020.
61. Adams, M.D.; Goglin, K.; Molyneaux, N.; Hujer, K.M.; Lavender, H.; Jamison, J.J.; MacDonald, I.J.; Martin, K.M.; Russo, T.; Campagnari, A.A.; et al. Comparative genome sequence analysis of multidrug-resistant *Acinetobacter baumannii*. *J. Bacteriol.* **2008**, *190*, 8053–8064. [[CrossRef](#)]
62. Durante-Mangoni, E.; Andini, R.; Zampino, R. Management of carbapenem-resistant Enterobacteriaceae infections. *Clin. Microbiol. Infect.* **2019**, *25*, 943–950. [[CrossRef](#)]
63. Martinez-Martinez, L. Carbapenemases: The never-ending story. *Enferm. Infecc. Microbiol. Clin.* **2019**, *37*, 73–75. [[CrossRef](#)]
64. Moeck, G.S.; Coulton, J.W. TonB-dependent iron acquisition: Mechanisms of siderophore-mediated active transport. *Mol. Microbiol.* **1998**, *28*, 675–681. [[CrossRef](#)]
65. Noinaj, N.; Guillier, M.; Barnard, T.J.; Buchanan, S.K. TonB-dependent transporters: Regulation, structure, and function. *Annu. Rev. Microbiol.* **2010**, *64*, 43–60. [[CrossRef](#)]
66. Lau, C.K.; Krewulak, K.D.; Vogel, H.J. Bacterial ferrous iron transport: The Feo system. *FEMS Microbiol. Rev.* **2016**, *40*, 273–298. [[CrossRef](#)]
67. Mihara, K.; Tanabe, T.; Yamakawa, Y.; Funahashi, T.; Nakao, H.; Narimatsu, S.; Yamamoto, S. Identification and transcriptional organization of a gene cluster involved in biosynthesis and transport of acinetobactin, a siderophore produced by *Acinetobacter baumannii* ATCC 19606T. *Microbiology* **2004**, *150*, 2587–2597. [[CrossRef](#)]
68. Tiwari, V.; Rajeswari, M.R.; Tiwari, M. Proteomic analysis of iron-regulated membrane proteins identify FhuE receptor as a target to inhibit siderophore-mediated iron acquisition in *Acinetobacter baumannii*. *Int. J. Biol. Macromol.* **2018**, *125*, 1156–1167. [[CrossRef](#)]
69. Mussi, M.A.; Limansky, A.S.; Relling, V.; Ravasi, P.; Arakaki, A.; Actis, L.A.; Viale, A.M. Horizontal gene transfer and assortative recombination within the *Acinetobacter baumannii* clinical population provide genetic diversity at the single carO gene, encoding a major outer membrane protein channel. *J. Bacteriol.* **2011**, *193*, 4736–4748. [[CrossRef](#)]
70. Baynes, R.D.; Bezwoda, W.R. Lactoferrin and the inflammatory response. *Adv. Exp. Med. Biol.* **1994**, *357*, 133–141.
71. Fillebeen, C.; Wilkinson, N.; Charlebois, E.; Katsarou, A.; Wagner, J.; Pantopoulos, K. Hepcidin-mediated hypoferremic response to acute inflammation requires a threshold of Bmp6/Hjv/Smad signaling. *Blood* **2018**, *132*, 1829–1841. [[CrossRef](#)]
72. Caldwell, M.; Archibald, F. The effect of the hypoferremic response on iron acquisition by and growth of murine lymphoma cells. *Biochem. Cell Biol.* **1987**, *65*, 651–657. [[CrossRef](#)]
73. Klein, S.; Boutin, S.; Kocer, K.; Fiedler, M.O.; Störzinger, D.; Weigand, M.A.; Tan, B.; Richter, D.; Rupp, C.; Mieth, M.; et al. Rapid development of cefiderocol resistance in carbapenem-resistant *Enterobacter cloacae* during therapy is associated with heterogeneous mutations in the catecholate siderophore receptor cira. *Clin. Infect Dis.* **2021**, ciab511. [[CrossRef](#)]
74. Streling, A.P.; Al Obaidi, M.M.; Lainhart, W.D.; Zangeneh, T.; Khan, A.; Dinh, A.Q.; Hanson, B.; Arias, C.A.; Miller, W.R. Evolution of Cefiderocol Non-Susceptibility in *Pseudomonas aeruginosa* in a Patient without Previous Exposure to the Antibiotic. *Clin. Infect. Dis.* **2021**, *73*, e4472–e4474. [[CrossRef](#)] [[PubMed](#)]
75. Simner, P.J.; Beisken, S.; Bergman, Y.; Posch, A.E.; Cosgrove, S.E.; Tamma, P.D. Cefiderocol Activity against Clinical *Pseudomonas aeruginosa* Isolates Exhibiting Ceftolozane-Tazobactam Resistance. *Open Forum Infect. Dis.* **2021**, *8*, ofab311. [[CrossRef](#)] [[PubMed](#)]
76. Pinsky, M.; Roy, U.; Moshe, S.; Weissman, Z.; Kornitzer, D. Human Serum Albumin Facilitates Heme-Iron Utilization by Fungi. *mBio* **2020**, *11*, e00607–20. [[CrossRef](#)] [[PubMed](#)]
77. König, C.; Both, A.; Rohde, H.; Kluge, S.; Frey, O.R.; Rohr, A.C.; Wichmann, D. Cefiderocol in Critically Ill Patients with Multi-Drug Resistant Pathogens: Real-Life Data on Pharmacokinetics and Microbiological Surveillance. *Antibiotics* **2021**, *10*, 649. [[CrossRef](#)]
78. Kidd, J.M.; Abdelraouf, K.; Nicolau, D.P. Development of Neutropenic Murine Models of Iron Overload and Depletion to Study the Efficacy of Siderophore-Antibiotic Conjugates. *Antimicrob. Agents Chemother.* **2019**, *64*, e01961–e19. [[CrossRef](#)]
79. Kidd, J.M.; Abdelraouf, K.; Nicolau, D.P. Efficacy of Humanized Cefiderocol Exposure Is Unaltered by Host Iron Overload in the Thigh Infection Model. *Antimicrob. Agents Chemother.* **2019**, *64*, e01767–19. [[CrossRef](#)] [[PubMed](#)]
80. Matsumoto, S.; Singley, C.M.; Hoover, J.; Nakamura, R.; Echols, R.; Rittenhouse, S.; Tsuji, M.; Yamano, Y. Efficacy of Cefiderocol against Carbapenem-Resistant Gram-Negative Bacilli in Immunocompetent-Rat Respiratory Tract Infection Models Recreating Human Plasma Pharmacokinetics. *Antimicrob. Agents Chemother.* **2017**, *61*, e00700–17. [[CrossRef](#)] [[PubMed](#)]