



Response to Article “Evaluation of the Anti-Malarial Activity of the Crude Root Extract and Solvent Fraction of *Sesamum indicum* (Fabaceae)” [Letter]

Putri Reno Intan ^{*}, Ariyani Noviantari ^{*}

Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency (BRIN), Cibinong Science Center, Cibinong - Bogor, West Java, Indonesia

^{*}These authors contributed equally to this work

Correspondence: Putri Reno Intan, Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency (BRIN), Genomic Building, Cibinong Science Center, Jalan Raya Bogor Km. 46, Cibinong - Bogor, West Java, 16911, Indonesia, Email putri.reno.intan@brin.go.id

Dear editor

We read, analyzed, and appreciated Fentaw Girmaw and Getachew Ashagrie's study¹ to assess the antimalarial activity of the crude root extract and solvent fraction of *Sesamum indicum* in mice. Based on their methods, the experimental animals used were healthy *Swiss albino* mice with weight (20–35 g) or age (6–8 weeks). However, the sex of the mice used in this study was not clearly stated in the method section.

Here, it is possible to advise using either both male or female mice in the following research. This is because male dies more frequently than females in plasmodium-infected mice, suggesting that the immune response differs between the sexes.² Knowing the varied *P. berghei* infection responses in mice of both sexes and the role of sex hormones in those responses would aid in understanding the physiological mechanisms of plasmodial infections in humans.

Males and females are not equally susceptible to malaria, and this sexual dimorphism may have significant effects on how vaccinations and medications work. On the processes mediating these sexual differences, nevertheless, nothing is known. Sex hormones are what control the primary variations between the sexes.³

In their study, Girmaw and Ashagrie reported that the crude root extract and solvent fractions of *Sesamum indicum* possessed a dose-dependent antimalarial activity and a significant change in other parameters in both models that strengthen the traditional claim.¹ However to further enrich the results obtained and as a basis before conducting an in vivo anti-malarial activity of *Sesamum indicum* test, for future study an in vivo antimalarial activities and toxicological assessment of *Sesamum indicum* (Fabaceae) test can be carried out first.⁴ So that, the classification of the extract can be determined whether it is classified as high extract activity at IC₅₀ <5 mg/mL, promising activity at 5–15 mg/mL, moderate activity at 15–50 mg/mL, and inactivity at > 50 mg/mL.⁵

Acknowledgments

All authors would like to acknowledge Dr. Sunarno and all of the researchers at the Center for Biomedical Research, National Research and Innovation Agency (BRIN) for their ongoing assistance.

Disclosure

There are no conflicts of interest among the authors of this communication.

References

1. Girmaw F, Ashagrie G. Evaluation of the anti-malarial activity of the crude root extract and solvent fraction of *Sesamum indicum* (Fabaceae). *J Exp Pharmacol*. 2023;15:163–175. PMID: 37013163; PMCID: PMC10066629. doi:10.2147/JEP.S407557
2. Klein SL. Hormonal and immunological mechanisms mediating sex differences in parasite infection. *Parasite Immunol*. 2004;26(6–7):247–264. doi:10.1111/j.0141-9838.2004.00710.x
3. Legorreta-Herrera M, Mosqueda-Romo NA, Nava-Castro KE, Morales-Rodríguez AL, Buendía-González FO, Morales-Montor J. Sex hormones modulate the immune response to *Plasmodium berghei* ANKA in CBA/Ca mice. *Parasitol Res*. 2015;114(7):2659–2669. doi:10.1007/s00436-015-4471-6
4. Phuwajaroanpong A, Chaniad P, Horata N, Muangchanburee S, Kaewdana K, Punsawad C. In vitro and in vivo antimalarial activities and toxicological assessment of *pogostemon cablin* (blanco) benth. *J Evid Based Integr Med*. 2020;25:1–8. doi:10.1177/2515690X20978387
5. Lusakibanza M, Mesia G, Tona G, et al. In vitro and in vivo antimalarial and cytotoxic activity of five plants used in congolese traditional medicine. *J Ethnopharmacol*. 2010;129(3):398–402. doi:10.1016/j.jep.2010.04.007

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Journal of Experimental Pharmacology 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Journal of Experimental Pharmacology editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Journal of Experimental Pharmacology

Dovepress

Publish your work in this journal

The Journal of Experimental Pharmacology is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of laboratory and experimental pharmacology. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-experimental-pharmacology-journal>

<https://doi.org/10.2147/JEP.S418029>