

REVIEW

# Tumour-associated carbohydrate antigens in breast cancer

Aurélie Cazet<sup>1,2</sup>, Sylvain Julien<sup>3</sup>, Marie Bobowski<sup>1,2</sup>, Joy Burchell<sup>3</sup> and Philippe Delannoy<sup>1,2\*</sup>

## Abstract

Glycosylation changes that occur in cancer often lead to the expression of tumour-associated carbohydrate antigens. In breast cancer, these antigens are usually associated with a poor prognosis and a reduced overall survival. Cellular models have shown the implication of these antigens in cell adhesion, migration, proliferation and tumour growth. The present review summarizes our current knowledge of glycosylation changes (structures, biosynthesis and occurrence) in breast cancer cell lines and primary tumours, and the consequences on disease progression and aggressiveness. The therapeutic strategies attempted to target tumour-associated carbohydrate antigens in breast cancer are also discussed.

## Introduction

Glycosylation is one of the most important modifications of proteins and lipids. The structure of glycans is highly diversified and their biosynthesis requires specific enzymatic machinery involving a high number of glycosyltransferases. Glycoconjugates of the cell surface are thought to play important roles in a variety of biological functions, such as cell–cell and cell–substrate interactions, bacterial adhesion, membrane organization, cell immunogenicity and protein targeting. For example, sialyl-Lewis<sup>x</sup> (sLe<sup>x</sup>) antigens are the ligands for selectins and are involved in the recruitment of leukocytes to lymphoid tissues and inflammation sites [1].

Glycan structures depend on the cell type, developmental stage and cell differentiation, and are modified in many pathologic states including cancers. The modification of cellular glycosylation is indeed a common phenotypic change of cancer cells that mainly affects the

outer part of glycans, leading to the expression of tumour-associated carbohydrate antigens (TACAs). Increased  $\beta$ 1,6-branching, increased sLe<sup>x</sup> or sialyl-Lewis<sup>a</sup> (sLe<sup>a</sup>) antigens, or the general increase in sialylation are commonly observed in *N*-linked and *O*-linked glycans of carcinoma cells and are associated with grade, invasion, metastasis and poor prognosis. These changes in glycosylation often reflect a deregulation of glycosyltransferase expression at the transcriptional level, and several examples have shown that glycosyltransferase genes, including *ST6GAL1* [2] and *MGAT5* [3], are regulated by oncogenes.

sLe<sup>x</sup>, s-Le<sup>a</sup> and sialyl-Thomsen-nouvelle (sTn) are tumour-associated antigens found in breast cancer [4-6]. When these antigens are detected at the surface of breast carcinoma cells, they are usually associated with a poor prognosis and a reduced overall survival of the patients [7]. These glycosylation changes mostly result from the modification of the expression of glycosyltransferase (GT) genes, and the level of expression of sialyltransferases has been proposed as a prognostic marker for the follow-up of breast cancer patients [8,9].

Several cellular models have been developed in order to explain the mechanisms by which carbohydrate antigens can reinforce breast cancer progression and aggressiveness, showing the implication of TACAs in cell adhesion, migration, proliferation and tumour growth [10,11]. TACAs such as sTn are also targets for breast cancer immunotherapy. Finally, recent data seem to indicate an important role of complex gangliosides in breast cancer progression and metastasis [12,13]. The present review summarizes our current knowledge of glycosylation changes that occur in breast cancer and the consequences of these changes on disease development and aggressiveness.

## Defects in mucin-type *O*-glycosylation

One of the hallmarks of breast adenocarcinomas is the overexpression and aberrant localization of membrane-bound mucins, such as MUC1 and MUC4 [14]. Mucins are characterized by a large number of tandem repeated domains, rich in serine and threonine, which are therefore densely *O*-glycosylated.

\*Correspondence: philippe.delannoy@univ-lille1.fr

<sup>1</sup>Structural and Functional Glycobiology Unit, UMR CNRS 8576, University of Sciences and Technologies of Lille, 59655 Villeneuve d'Ascq, France  
Full list of author information is available at the end of the article

O-glycans expressed by normal breast epithelial cells are composed of Core2 structures extended with a linear polylectosaminic chain, which can be fucosylated or sialylated [15] (Figure 1). Accordingly, a recent study reported a prevalence of Le<sup>x</sup> expression at the apex of normal breast epithelia, where MUC1 is also detected [16].

In contrast, breast cancer cell lines express truncated O-glycans with shorter polylectosaminic chains and a tendency to switch from Core2 O-glycans to sialylated or unsialylated Core1 structures (Figure 1) [17]. It has long been known that breast cancer tissues contain more sialic acid and more sialyltransferase activity [18] than corresponding normal tissues. Breast cancer cells may thus express shorter O-glycans due to premature sialylation of the polylectosaminic chain blocking further extension, or due to truncation of the O-glycans at the core level – owing to the sialylation of the Core1 structure or of the initial GalNAc, leading to the expression of sialyl-Thomsen–Friedenreich (sialyl-T) antigen structures and sTn antigen, respectively (Figure 1).

Alternatively, a defect in GTs that are involved in the biosynthesis of the O-glycan cores can also lead to the incomplete synthesis of the glycans, allowing the expression of O-linked carbohydrate antigens known as Thomsen–Friedenreich (T or TF) antigen (unsialylated Core1) and Thomsen-nouvelle (Tn) antigen (initial unsubstituted GalNAc) (Figure 1).

### Thomsen–Friedenreich antigen

T antigen is the cryptic precursor of Core2 O-glycans, which can be unmasked if the cancer cells lose their ability to synthesize Core2 (Figure 1). T antigen consists of unsialylated Core1 structures (Galβ1-3GalNAcαSer/Thr) initially described on glycoporphins of red blood cells. The galactose residue is crucial for the antigenicity since re-sialylation or β-D-galactosidase treatment abrogates T-antigen detection [19].

Georg Springer was the pioneer of the study of T antigen in breast cancer. Using natural anti-T antisera, he first showed that all breast cancers, but not benign lesions or normal mammary gland, expressed the T antigen [20]. Later studies using monoclonal antibodies demonstrated the presence of T antigen at a low level in normal epithelium or metaplastic lesions of the breast [21].

Lectins such as *Arachis hypogea* agglutinin/peanut lectin (PNA), *Amaranthus caudatus* lectin or *Artocarpus integrifolia*/jacalin have been extensively used to detect T antigen. None of these lectins, however, is exclusively specific for the Galβ1-3GalNAc structure. Both *A. caudatus* agglutinin and *A. integrifolia* agglutinin/jacalin are specific for GalNAc when there is no substitution on C6 of the saccharide (Figure 2). In contrast, substitution on C3 does not impair their binding. Consequently both *A. caudatus* agglutinin and *A. integrifolia* agglutinin/

jacalin can bind to Tn antigen (GalNAcα-Ser/Thr), T antigen (Galβ1-3GalNAcα-Ser/Thr) and sialyl-T antigen (Neu5Acα2-3Galβ1-3GalNAcα-Ser/Thr) [22,23].


PNA is specific for unsubstituted Gal residues [24]. Because of its high reactivity with asialo-glycophorin, PNA has been tagged as the anti-T lectin [24]. PNA was able to react weakly with healthy tissue while anti-T antisera were not, however, suggesting that PNA specificity was slightly wider than the sole T carbohydrate. Indeed, PNA can bind, albeit with lower affinity, to terminal galactose in a variety of glycans including glycolipids such as asialo-G<sub>M1</sub> [23] or N-acetyl-lactosamine (LacNAc) [25]. Studies of T-antigen expression using PNA should therefore be treated cautiously, especially in the absence of complementary analysis of the O-glycosylation of the considered cells or tissues.

In parallel to lectins, mAbs have long been used to investigate T-antigen expression in epithelial cancers [26]. As for lectins, most mAbs are not strictly specific for the sole Galβ1-3GalNAc-Ser/Thr structure – some of them, such as A78-G/A7, cross-reacting with asialo-G<sub>M1</sub>. Other mAbs define site-specific T antigens, such as BW835 with the VTSA motif in MUC1 tandem repeat. The HH8 mAb, as PNA, tolerates further substitution on the C6 of the GalNAc leading to Core2 structures [26] (Figure 2).

These subtle differences of specificity concur with the variability of the detection of T antigen in tumour tissues [26,27], and these discrepancies have fuelled debates regarding the prognosis value of T-antigen expression in breast cancers. PNA staining does not have any prognostic significance [26], while a more recent study using the mAb A78-G/A7 showed a correlation of T-antigen expression with a better prognosis [28], opposite to what is reported for gastrointestinal, lung or cervix cancers.

A recent study has shown that 98% of the disseminated tumour cells in the bone marrow were positively stained by a new anti-T antibody [29], suggesting a role for T antigen in the metastasis process. This putative role has been investigated during the past decade, on the assumption that Galβ1-3GalNAc-R disaccharide could be a good ligand for Galectin-3 [30], which has been shown to promote breast cancer metastasis by mediating adhesion to endothelial cells. Although Galectin-3 seems to be involved in breast cancer metastasis through its interaction with some cancer-associated O-glycans [31], however, the nature of its ligand remains to be clearly elucidated via a comprehensive characterization of the glycosylation pattern required for this interaction.

To summarize, there is strong evidence showing that T antigen is frequently overexpressed in breast cancer, suggesting that the changes in O-glycosylation provide some advantage to the tumour development. Relation to prognosis or metastatic behaviour, however, suffers from



contradictory studies – partially due to the use of different probes – and remains to be unambiguously demonstrated.

### Thomsen-nouvelle antigen







Tn antigen is the cryptic precursor of the T antigen (Core1) that can be unmasked if a cell loses its ability to synthesize the Core1 structure (see Figure 1).

Tn (GalNAc $\alpha$ -R) expression on blood cells is responsible for Tn syndrome, which is a rare autoimmune haematological disorder [32]. Tn syndrome occurs due to mutations in the *COSMC* gene coding the C1 $\beta$ 3GalT-specific molecular chaperone, leading to a drastic decrease of C1 $\beta$ 3GalT (T-synthase) activity [33]. Mutations and loss of heterozygosity of *COSMC* were described in colon cancer and melanoma cancer Tn-positive cells, as well as in tissues from two Tn-positive cervical cancers [34]. Extensive studies of *COSMC* in epithelial cancers,

however, showed that these events were rare and could only partially explain Tn expression in cancers [35].

A range of GalNAc-specific lectins, including *Helix pomiana* lectin (HPA) and plant lectins from *Dolichos biflorus*, *Griffonia simplicifolia*, *Wistaria floribundia*, *Glycine max* soybean, *Vicia villosa* isolectin B4 (VVA) and *Salvia sclarea* agglutinin, can be used to detect Tn antigen [32,36]. These lectins bind to terminal GalNAc and therefore react with Tn antigen, but also with Forssman antigen (GalNAc $\alpha$ 1-3GalNAc $\beta$ -R) and to some extent with blood group A (GalNAc $\alpha$ 1-3[Fuca1-2]Gal $\beta$ -R) or glycolipids [36]. HPA, which has been extensively used to study Tn expression in breast cancer [37], has been shown by some authors to bind T antigen [36]. There is also a range of mAbs available to detect Tn antigen [32]. Some of them, however, can slightly cross-react with sTn.

Tn antigen was found to be expressed in almost 90% of breast cancers while it is barely detected in normal tissue

Structures		Name	ACA and AIA	PNA and HH8
GalNAc $\alpha$ -R		Tn antigen	Yes	No
Neu5Ac $\alpha$ 2-6GalNAc $\alpha$ -R		Sialyl-Tn antigen	No	No
Gal $\beta$ 1-3GalNAc $\alpha$ -R		T or TF antigen Core1	Yes	Yes
Gal $\beta$ 1-3(Neu5Ac $\alpha$ 2-6)GalNAc $\alpha$ -R		Sialyl(6)T	No	Yes
Neu5Ac $\alpha$ 2-3Gal $\beta$ 1-3GalNAc $\alpha$ -R		Sialyl(3)T	Yes	No
Gal $\beta$ 1-4GlcNAc $\beta$ 1-6(Gal $\beta$ 1-3)GalNAc $\alpha$ -R		Core2	-	Yes

**Figure 2. Short O-glycans observed in breast epithelial cells during carcinogenesis.** Structures of short O-glycans observed in breast epithelial cells during carcinogenesis and their binding to *Arachis hypogea* agglutinin/peanut lectin (PNA) lectin or HH8 mAb, and to *Amaranthus caudatus* agglutinin (ACA) or *Artocarpus integrifolia* agglutinin/jacalin (AIA). T or TF, Thomsen–Friedenreich; Tn, Thomsen-nouvelle.

[38]. Association with high grade has been reported for invasive ductal carcinomas (IDC) using a pool of anti-Tn antibodies [39] and for ductal carcinoma *in situ* using VVA and *G. simplicifolia* agglutinin [40]. Using VVA, *G. simplicifolia* agglutinin and 83D4 mAb, other authors reported a tendency to lose Tn expression in grade III tumours [41]. Using either anti-Tn mAb or VVA, Tn was found associated with the Pathological Tumor-Node-Metastasis (pTNM) tumour stage in IDC [39,42]. Tn expression was also associated with decreased overall survival using antibodies [35] or HPA [43] and with increased risks of recurrence using VVA [42] or HPA [44].

Strikingly, Tn expression was always observed in lymph node metastatic cells using either mAb or lectins [39,41] with 100% of Tn-positive metastatic cells, while primary tumours could exhibit various percentages of cell positivity [39,41,42]. Tn expression in primary tumours seems consistently associated with lymph node involvement using mAbs [41,42], HPA [43] or VVA [42], and even with invasion of lymphatic vessels within the

primary tumour [42]. This observation strongly suggests that Tn antigen may play a role in the mechanism involved in lymph node metastasis. Indeed, Danussi and colleagues have recently shown that a newly generated anti-Tn antibody could block Tn-positive MCF-7 breast cancer cell line interaction with lymphatic endothelium of the mammary pad in an animal model [45]. A putative receptor(s) for Tn antigen in lymphatic cells, however, remains to be identified.

One known receptor for Tn antigen is the macrophage galactose-type lectin (MGL), which, in human, is specific for GalNAc [46]. MGL is expressed in immature dendritic cells and macrophages, and is proposed as an immunomodulatory receptor [46]. MGL has been shown to bind Tn antigen carried by MUC1 in colon carcinoma cells [47], and human immature dendritic cells can uptake MUC1-Tn peptide in a MGL-mediated way [48]. The MGL–Tn interaction, however, has not yet been investigated regarding its possible functional involvement in breast cancer development.

### Sialyl-Thomsen-nouvelle antigen

Tn antigen can be sialylated at the C6 position of GalNAc, resulting in the disaccharide Neu5Ac $\alpha$ 2-6GalNAc-R (sTn). STn is absent in normal healthy tissues but can be detected at various frequencies in almost all kinds of carcinomas [49].

ST6GalNAc I is the enzyme catalysing the transfer of sialic acid onto GalNAc *in vivo* [50]. It has been proposed that *COSMC* mutation was necessary to provide the Tn acceptor substrate for ST6GalNAc I to synthesize sTn antigen [34]. Transfection of ST6GalNAc I cDNA, however, has been shown to be sufficient to induce sTn expression in various breast cancer cell lines expressing Core1 and Core2 glycans [11,50,51], proving that ST6GalNAc I can compete with active Core1  $\beta$ 1,3-galactosyltransferase.

Several mAbs have been raised against sTn antigen using various immunogens, leading to subtle differences in the way they recognize sTn [49]. The most commonly used mAbs in breast cancer investigations are B72.3, TKH2 and HB-STn1. B72.3 appeared to be specific for clustered sTn but also cross-reacts with Tn clusters [52,53]. In contrast, TKH2 reacts more strongly to monomeric sTn conjugated to keyhole limpet hemocyanin (STn-KLH) than B72.3 [54].

The reported occurrence of sTn expression in breast cancer is highly variable. It is commonly accepted that at least 25 to 30% of breast cancers are sTn-positive [51]. The reported ranges are 20 to 80% using B72.3 and 30 to 60% using HB-STn1 [49], and correlation with known prognosis markers varies from one study to another. Correlation with a decreased overall survival seems to be the more consistent correlation through the literature. Kinney and colleagues proposed that sTn was an independent predictor of poor prognosis of breast cancers [55]. Miles and colleagues, however, reported a significant correlation of sTn expression with the decreased survival in the restricted node-positive patient group [7], whereas Leivonen and colleagues limited the association to the short-term prognosis [56].

Induction of sTn expression by transfection of the breast cancer cell line with ST6GalNAc I cDNA decreased the adhesion of cells to matrix proteins, and increased mobility and tumour growth in a xenograft model [10,11,50]. These observations are in accordance with a more aggressive phenotype of cancer cell that is usually associated with sTn expression but remain to be elucidated at the molecular level. Of the various sTn-positive proteins present in these transfected cell lines, only four have been identified: MUC1 and CD44 in human [10], and integrin  $\beta_1$  [57] and osteopontin [58] in mouse. All of these molecules have described functions in adhesion and/or migration, and it has been suggested that the modification of their O-glycosylation may influence this function and the mobility of the cancer cells.

### Lewis antigens in breast cancer

#### Structure and biosynthesis of Lewis antigens


The histoblood group Lewis antigens are found in most human epithelial tissues, where they are expressed at the terminal part of glycolipid and glycoprotein glycan chains [59]. These antigens derive from the substitution of type 1 (Gal $\beta$ 1-3GlcNAc) or type 2 (Gal $\beta$ 1-4GlcNAc) disaccharide sequences by fucose and sialic acid residues (Figure 3). Le<sup>a</sup>, Le<sup>b</sup> and sLe<sup>a</sup> derive from type 1 sequences, and Le<sup>x</sup>, Le<sup>y</sup> and sLe<sup>x</sup> derive from type 2. In healthy individuals, type 1 (Le<sup>a</sup> and Le<sup>b</sup>) antigens are widely expressed, while type 2 (Le<sup>x</sup> and Le<sup>y</sup>) antigens are only expressed at relatively low levels. The biosynthesis of these antigens requires specific fucosyltransferases and sialyltransferases (Table 1).

The GlcNAc  $\alpha$ 1,4-fucosylation of type 1 disaccharide generates Le<sup>a</sup> antigen, whereas the  $\alpha$ 1,3-fucosylation of type 2 gives Le<sup>x</sup> (Figure 3). The fucosyltransferases FucT III, FucT IV, FucT VI and FucT IX catalyse the transfer of a fucose residue onto GlcNAc of type 1 or type 2 disaccharides, thus synthesizing  $\alpha$ 1,3-fucosylated and, in the case of FucT III,  $\alpha$ 1,4-fucosylated glycans [60-62]. FucT III is the only enzyme responsible for the synthesis of Le<sup>a</sup> and sLe<sup>a</sup> *in vivo* and is called the Lewis enzyme. The fucosyltransferases FucT I and FucT II are responsible for the Gal  $\alpha$ 1,2-fucosylation of Le<sup>a</sup> or Le<sup>x</sup>, generating Le<sup>b</sup> or Le<sup>y</sup> antigens, respectively (Figure 3). FucT I (H enzyme) preferentially fucosylates type 1, whereas FucT II (Se enzyme) determines the expression of H-type 2 (Fuca1-2Gal $\beta$ 1-4GlcNAc) and Le<sup>y</sup> antigens.

The biosynthesis of sialylated Lewis antigens (sLe<sup>a</sup> and sLe<sup>x</sup>) requires first the  $\alpha$ 2,3-sialylation of Gal prior to  $\alpha$ 1,3/4-fucosylation. ST3Gal III [63], preferentially acting on type 1 rather than on type 2 disaccharides, is involved in sLe<sup>a</sup> synthesis. ST3Gal IV and ST3Gal VI mainly act on type 2 disaccharide, leading to the biosynthesis of sLe<sup>x</sup> [64], but both also share acceptor specificity with ST3Gal III. For the subsequent fucosylation, FucT III is the main enzyme involved in the biosynthesis of sLe<sup>a</sup> [65], whereas FucT VII [62,66] shows a restricted substrate specificity since it can form only sLe<sup>x</sup> [67].

#### Overexpression of Lewis antigens in breast cancer

The expression of sialyl-Lewis antigens is usually altered in breast cancer. Normal breast commonly expresses Le<sup>x</sup>, mainly in the apical part of cell ducts, together with MUC1. sLe<sup>x</sup> and sLe<sup>y</sup> epitopes may also be sometimes detected in normal breast, but with a low intensity [16]. On the contrary, sLe<sup>a</sup> and sLe<sup>x</sup> expression is found to be increased in breast cancer tissues, including primary breast carcinoma lesions [68]. Furthermore, levels of expression of sLe<sup>x</sup> are higher in patients who had distant metastasis compared with patients presenting nonmetastatic cancers [69]. The expression of sialyl-Lewis



**Table 1. Fucosyltransferases and sialyltransferases potentially involved in biosynthesis of histoblood group Lewis antigens in breast tissues**

Gene name	Common name	Accession number	Main acceptor substrate
<i>α</i> 1,2-Fucosyltransferases			
<i>FUT1</i>	FucT I	NM_000148	<b>Galβ1-3</b> [Fucα1-4]0-1GlcNAc
<i>FUT2</i>	FucT II	NM_000511	<b>Galβ1-4</b> [Fucα1-3]0-1GlcNAc
<i>α</i> 1,3/4-Fucosyltransferases			
<i>FUT3</i>	FucT III	NM_000149	Galβ1-3/4 <b>GlcNAc</b>
<i>FUT6</i>	FucT VI	NM_000150	Galβ1-4 <b>GlcNAc</b>
<i>FUT7</i>	FucT VII	NM_004479	Neu5Acα2-3Galβ1-4 <b>GlcNAc</b>
<i>FUT9</i>	FucT IX	NM_006581	Galβ1-4 <b>GlcNAc</b>
<i>α</i> 2,3-Sialyltransferases			
<i>ST3GAL3</i>	ST3Gal III	NM_174972	<b>Galβ1-3</b> GlcNAc
<i>ST3GAL4</i>	ST3Gal IV	NM_006278	<b>Galβ1-4</b> GlcNAc
<i>ST3GAL6</i>	ST3Gal VI	NM_006100	<b>Galβ1-4</b> GlcNAc

The acceptor monosaccharide is indicated in bold.

antigens is weak in carcinoma *in situ*, is moderate in invasive carcinomas without metastases and is high in primary carcinoma with lymph node metastases, showing a good correlation with the metastatic risk in breast cancer patients [70]. Immunohistochemistry of breast cancer tissue suggests that sLe<sup>x</sup> expression is an independent prognostic indicator of survival regardless of the size of the primary tumour and lymph node involvement [71].

RNA analysis of tumours indicated that FucT VI expression correlates with the levels of sLe<sup>x</sup>. Furthermore, transfection of the *FUT6* gene into MCF-7 breast cancer cells showed that FucT VI is responsible for an

enhancement of sLe<sup>x</sup> expression [72]. The downregulation of ST3Gal IV increased MCF-7 cell adhesion and decreased MDA-MB-231 cell migration, suggesting that α2,3-sialylation played a crucial role in the adhesion and tumour metastases [73]. In parallel, the level of ST3Gal III expression is associated with axillary lymph node involvement and a reduced overall survival of patients [8,9].

A general reduced expression of type 1 carbohydrate antigens and increased expression of type 2 antigens in breast cancer tissues has also been reported [4,6], and the relative expression of type 2 carbohydrate antigens seems capable of serving as a prognostic factor. As much as 80%

of breast tumours have lost the expression of Le<sup>b</sup> antigen, which correlates with the degree of malignancy and invasiveness [74].

The expression level of sLe<sup>x</sup> is also increased in the serum of patients with advanced breast cancer [69,75] and is therefore a good candidate as a diagnosis marker. In combination with serum CA15-3, the most commonly used breast cancer marker, the measurement of serum sLe<sup>x</sup> improves monitoring – for 78.5% of detected metastatic breast cancer, compared with 61.5% when CA15-3 is used alone [76].

The Le<sup>y</sup> antigen is also overexpressed in 60 to 90% of cancer of epithelial origin, including breast cancer [77]. A clinical study showed that high expression of Le<sup>y</sup> correlated with high grade and poor prognosis, and significantly decreased patient survival in lymph node-negative breast carcinoma [78]. Showing restricted expression on epithelial surfaces in normal tissues, Le<sup>y</sup> also constitutes a good candidate for cancer immunotherapy [79].

#### Implication of sialyl-Lewis antigens in metastasis

The expression of sLe<sup>x</sup> on leukocytes contributes to their function in the inflammatory response via interaction with E-selectin expressed on endothelial cells. By exposing sLe<sup>x</sup> at the cell surface, malignant cells can use a similar strategy for the extravasation from the blood circulation and metastasis, targeting tissues where E-selectin is highly expressed.

Both P-selectin and E-selectin expression is significantly enhanced on endothelial cells of breast cancer patients [68], consistent with the proposal that sLe<sup>x</sup> binding to selectins aids cancer cell metastasis. A high tumour-associated ST3Gal III expression and a high E-selectin serum concentration were associated with a bad prognosis for relapse-free survival and overall survival in patients with node-negative breast cancer [8]. In both canine mammary carcinoma and the human inflammatory breast cancer model, the cellular expression of E-cadherin and sLe<sup>x</sup> are inversely correlated, suggesting a biologically coordinated mechanism of E-cadherin and sLe<sup>x</sup> expression essential for tumour establishment and metastatic progression [80].

### Gangliosides in breast cancer

#### Structure and biosynthesis of gangliosides

Gangliosides are expressed at the plasma membrane of cells, especially in developing tissues and the adult central nervous system. Gangliosides structurally and biosynthetically derive from lactosylceramide (LacCer), and additional sugar residues – including sialic acid, *N*-acetyl-galactosamine and galactose – are transferred in a step-wise manner by a series of specific GTs [81] (Figure 4). G<sub>M3</sub>, G<sub>D3</sub> and G<sub>T3</sub> are the starting points for the biosynthesis of a-series, b-series and c-series gangliosides,

respectively (Figure 4). The relative amounts of these glycosphingolipids determine the amount of complex gangliosides of each series. The amount of complex gangliosides is therefore dependent on the level of expression and activity of several GTs, including G<sub>M3</sub> synthase (ST3Gal V), G<sub>D3</sub> synthase (ST8Sia I), G<sub>T3</sub> synthase (ST8Sia V) and G<sub>M2/D2</sub> synthase (β1,4-GalNAc T1) (Table 2).


#### Overexpression or neo-expression of gangliosides in breast cancer

The neo-expression of gangliosides has been usually demonstrated in a variety of cancers – especially in neuroectoderm-derived tumours, in which they play a key role in invasion and metastasis. Breast tumour tissues were shown to be distinct from normal mammary tissues in terms of ganglioside composition. The gangliosides G<sub>M3</sub>, G<sub>D3</sub> and the derivatives 9-*O*-acetyl-G<sub>D3</sub> (CDw60 antigen) and 9-*O*-acetyl-G<sub>T3</sub>, which show a very restricted expression in normal breast tissues, are overexpressed in about 50% of IDC [82].

The *N*-glycolyl-G<sub>M3</sub> is also detected in IDC, with 100% efficiency in primary tumours for women diagnosed with stage II breast cancer [82,83]. Neu5Gc is normally absent in human due to the irreversible inactivation of the *CMAH* gene encoding the CMP-Neu5Ac hydroxylase. The lack of the enzyme responsible for the conversion of CMP-Neu5Ac in CMP-Neu5Gc results in the total absence of Neu5Gc in healthy human tissues. On the contrary, Neu5Gc was shown to be expressed on glycoproteins and gangliosides in melanoma, colon, retinoblastoma and breast cancers.

The expression of GTs implicated in ganglioside biosynthesis can be also altered in breast cancer tumours. Two clinical studies have shown that the G<sub>D3</sub> synthase (ST8Sia I) displayed higher expression among oestrogen receptor-negative breast cancer tumours [84]. *ST8SIA1* overexpression was associated with poor pathohistological grading in oestrogen receptor-negative tumours and lower survival of patients [85]. In contrast, a better prognosis for oestrogen receptor-positive samples with high expression of *ST8SIA1* was noticed [85].

It has been shown that silencing of G<sub>M3</sub> synthase in mammary murine 4T1 cells significantly inhibited cell migration, invasion and anchorage-independent growth *in vitro*, as well as lung metastasis *in vivo*. In parallel, overexpression of G<sub>M3</sub> synthase in nonmetastatic 67NR cells significantly restores the malignant phenotype [86]. We have recently demonstrated that overexpression of G<sub>D3</sub> synthase in MDA-MB-231 cells is associated with increased proliferation, especially in the absence of serum or exogenous growth factors [13]. G<sub>D3</sub> synthase-overexpressing cells bypass the need of serum for cell growth by the activation of Erk/MAPK, PI3K/Akt and SAPK/JNK pathways (unpublished results). These data



suggest that  $G_{M3}$  or  $G_{D3}$  synthase overexpression may contribute to increasing the malignant properties of breast cancer cells by mediating cell proliferation and migration. Recently, the role of ST6GalNAc V, a sialyltransferase involved in the biosynthesis of  $\alpha$ -series gangliosides, in breast cancer metastasis to the brain has been clearly demonstrated. Normally restricted to the brain, the expression of ST6GalNAc V enhances the adhesion of breast cancer cells to brain endothelium and their passage through the blood-brain barrier, highlighting the role of cell surface glycosylation in organ-specific metastatic interactions [12].

### Future perspectives for therapy

From the data summarized above, it appears that a limited number of TACAs are expressed in breast cancer and associated with restricted histopathological markers such as the oestrogen receptor status of tumours (Table 3). Breast cancer cells express truncated Core1-based

O-glycans, including T, Tn and sTn antigens. The expression of sialyl-Lewis antigens is also altered in breast carcinoma with a good correlation with metastatic risk, and complex gangliosides are overexpressed in IDC.

Although some TACAs have been shown as strong markers of a poor prognosis (for example, sLe<sup>x</sup> or sTn), they have not yet reached the clinic. Indeed, the strongest marker of prognosis in breast cancer remains the invasion of the lymph nodes determined during surgery. Unless another marker proves to be of superior quality or independent of lymph node involvement, it is very unlikely to be useful for clinic application. As TACAs are more or less associated with lymph node invasion and/or restricted to specific groups of patients, their use as prognosis markers appears rather limited. However, since TACAs seem to be functionally involved in cancer progression, one of their most promising uses in the future remains being targets for treatments designed for specific groups of patients.



**Table 2. Main glycosyltransferases involved in biosynthesis of gangliosides**

Gene name	Common name	Accession number	Main acceptor substrate
Glucosyltransferase			
<i>UGCG</i>	GlcCer synthase	NM_003358.1	Ceramide (Cer)
Galactosyltransferases			
<i>B4GALT6</i>	LacCer synthase	NM_004775	<b>Glc</b> α1-Cer
<i>B3GALT4</i>	β3-GalT4	NM_003782.3	<b>GalNAc</b> β1-4[NeuAc0-3]Galβ1-4Glcα1-Cer
N-Acetyl-galactosaminyltransferase			
<i>B4GALNACT1</i>	G <sub>M2'</sub> , G <sub>D2'</sub> , G <sub>T2</sub> synthase	NM_001478.2	[NeuAc0-3] <b>Gal</b> β1-4Glcα1-Cer
Sialyltransferases			
<i>ST3GAL1</i>	ST3Gal I	NM_003033.3	<b>Gal</b> β1-3GalNAcβ1-4[NeuAc0-3]Galβ1-4Glcα1-Cer
<i>ST3GAL2</i>	ST3Gal II	NM_006927.3	<b>Gal</b> β1-3GalNAcβ1-4[NeuAc0-3]Galβ1-4Glcα1-Cer
<i>ST3GAL5</i>	ST3Gal V	NM_003896.3	<b>Gal</b> β1-4Glcα1-Cer
<i>ST6GALNAC5</i>	ST6GalNAc V	NM_030965.1	NeuAcα2-3Galβ1-3 <b>GalNAc</b> β1-4Galβ1-4Glcα1-Cer
<i>ST8SIA1</i>	ST8Sia I	NM_003034.3	<b>NeuAc</b> α2-3Galβ1-4Glcα1-Cer
<i>ST8Sia5</i>	ST8Sia V	NM_013305.4	<b>NeuAc</b> α2-8NeuAcα2-3Galβ1-4Glcα1-Cer

The acceptor monosaccharide is indicated in bold.

**Table 3. Main tumour-associated carbohydrate antigens expressed in breast carcinomas**

TACA	DCIS	IDC	Altered pathway	Comments
Mucin-type antigens				
Thomsen-nouvelle	High	High	<i>COSMC</i> mutation, decrease in C1β3GalT activity	Expressed in almost 90% of breast cancers
Thomsen-Friedenreich	ND	High	Decrease in Core2 β6-GlcNAc-T expression, decrease in Core1 sialylation	
Sialyl-Thomsen-nouvelle	ND	High	Increase in ST6GalNAc I expression, decrease in Core2 β6-GlcNAc-T expression, alteration of <i>COSMC</i> function	25 to 30% of breast cancers, correlated with a decreased survival in node-positive patients
Lewis antigens				
Sialyl-Lewis <sup>x</sup>	Weak	High	Increased expression of FucT (FucT VI and FucT VII)	Higher in metastatic compared with nonmetastatic cancers
Sialyl-Lewis <sup>a</sup>	Weak	High	Increased expression of FucT (FucT III)	
Lewis <sup>y</sup>	Low	High	Increased expression of FucT	Overexpressed in 60 to 90% of breast cancer
Gangliosides				
G <sub>M3</sub>	ND	Increased	Increased expression of ST3Gal V	2.8-fold compared with normal tissue
G <sub>D3</sub>	ND	Increased	Increased expression of ST8Sia I	1.7-fold compared with normal tissue
9-O-acetyl-G <sub>D3</sub>	ND	Neo-expression	Increase of sialic acid acetylation	
9-O-acetyl-G <sub>T3</sub>	ND	Neo-expression	Increase of sialic acid acetylation	
N-glycolyl-G <sub>M3</sub>	ND	65 to 70%	Unknown	100% in stage II primary tumours

DCIS, ductal carcinoma *in situ*; IDC, invasive ductal carcinoma; ND, not determined; TACA, tumour-associated carbohydrate antigen.

Different methods can take advantage of TACA expression to treat breast cancer, such as blocking their biosynthesis or biological function, or using them as molecular targets in immunotherapy. Blocking the biosynthesis of glycans using chemical inhibitors has been shown to change the phenotype of breast cancer cells in culture, making them less aggressive [87,88]. Such an

approach, however, is difficult to adapt to human therapy because both healthy and diseased cells are using their glycans in a variety of functions. There are high risks that detrimental effects of a chemical inhibitor on healthy cells would counterbalance or avoid the beneficial effect of the molecule on cancer cells. In order to develop further, this field will have to await improvement in the

drug-targeting or drug delivery systems, such as the booming nanotechnologies [89].

Alternatively, one may consider specifically altering the expression of GTs involved in TACA biosynthesis. Little is still known about the molecular mechanisms leading to aberrant expression of GTs in cancer. Gain or loss of GT expression could be attributed to loss of chromosomal integrity often observed in cancers, leading to duplication, deletion or transposition of chromosomal segments involving glycosyltransferase genes. As an example, elevated ST6Gal I expression in murine W16 cells has been shown to be the result of DNA rearrangement in the *st6gal1* locus resulting from the transposition of a retroviral-like transposable element upstream of the *st6gal1* ORF [90]. Changes of gene expression do not always reflect genomic rearrangements, however, as comparison of the Comparative Genomic Hybridization Array and the RNA microarray do not always show absolute correlation. Expression of GT genes is often regulated at the transcriptional level [91], and several examples have shown that GT genes, including *ST6GAL1* [2] and *MGAT5* [3], can be regulated by oncogenes. Only a limited number of promoters of GT genes have been characterized [91], and this field is in need of further data in order to understand which factors of transcription, signalling pathways and, eventually, membrane receptors are involved in this cancer-specific deregulation. The knowledge of the intracellular molecular mechanisms regulating the expression of GTs might open new options to specifically impair TACA expression in cancer cells.

Functions of TACAs often involve interaction with glycan-specific receptors, such as selectins with sialyl-Lewis antigens. Attempts can be made to block this interaction, by masking either the glycans or its receptor. Several studies have shown that masking T antigens with specific peptides [92] or mAbs [93] can inhibit interaction of breast cancer cells with the endothelium and protect from metastasis. One advantage of this approach is that the lectin(s) recognizing the TACA does not need to be identified to block the interaction. Some limitations remain, however, such as the risk to induce unwanted physiological responses triggered by the same type of interactions (or loss of interaction), or the clinical viability of these compounds in terms of toxicity, body clearance or administration.

Active immunotherapy has been thoroughly investigated during the past 40 years as an alternative approach for cancer treatment. As glycans are naturally highly immunogenic, they have been extensively used for the design of immunotherapy strategies against breast cancer.

As early as in 1977, Springer and collaborators attempted to immunize patients with desialylated autologous red blood cells, in order to trigger a response to T

antigen. He reported a significantly improved 5-year survival (100%) and 10-year survival (44%) compared with the nonimmunized population [38]. Although of small scale and somewhat unorthodox, this study paved the way for further development of anti-TACA vaccines.

Ganglioside-based anti-idiotypic vaccines focusing on *N*-glycolyl- $G_{M3}$  have been developed for specific active immunotherapy against breast cancer [94]. The 1E10 mAb (Racotumomab) is an anti-idiotypic vaccine to P3 mAb, which recognized Neu5Gc- $G_{M3}$ . Vaccination with 1E10 of BALB/c mice challenged with the murine F3II mammary carcinoma leads to breast tumour growth suppression and a decrease of lung metastasis [95]. Moreover, the anti-tumour effect of 1E10 is enhanced by combination with a low-dose of cyclophosphamide [96]. Results of phase I clinical trials have shown that 1E10 vaccine was relatively well tolerated and able to produce a specific response against Neu5Gc-containing gangliosides in high-risk (stage III) or metastatic breast cancer [97,98].

Clustered sTn haptens conjugated to keyhole limpet hemocyanin (Theratope™; Biomira, Edmonton, AB, Canada) have also been used to immunize breast cancer patients [99]. Although the results of the first phase III clinical trial failed to show significant benefit, it has to be mentioned that patients were never screened for expression of sTn antigen in the tumours. The lack of significant benefit for the total population could therefore be explained by the low representation of patients presenting sTn-positive tumours (25 to 30%). Recently, we clearly demonstrated that Theratope™ induced tumour protection in a relevant murine model. The protection was dependent on both the presence of the sTn antigen in the tumour and the anti-sTn antibodies raised by immunization. Interestingly anti-sTn antibodies were able to recognize a wide range of sTn-positive proteins expressed by mammary carcinomas, suggesting that multi-targeting of the response was important to induce tumour protection [58].

Despite decades of research and evident proofs of concept in favour of anti-TACA cancer vaccines, one has to acknowledge that none of them have reached the clinic. One possible method of improvement is a better definition of patients who are most susceptible to benefit from the designed treatment. As TACAs are expressed in distinguishable groups of patients based on immunohistology or gene profiling expression, one may therefore consider personalizing the treatment according to expression of putative targets. Another way is to design multi-epitope targeting vaccines to cover a variety of targets found in different patient populations and to possibly trigger multi-antigenic response in each of the patients. Glycopeptide dendrimers have been developed to present simultaneously up to six different antigens (Tn, T, Globo H,  $G_{M2}$ , Le<sup>x</sup>, and MUC1-Tn antigens) and

have been assessed in immunization trials to compare multi-antigenic responses with mono-antigenic responses [100].

As the future of breast cancer treatment lies in a combination of patient-tailored and multi-targeting strategies, we believe that TACAs – because of their profile of expression and their proaggressive function – will be considered potent molecular targets.

#### Abbreviations

C1β3GalT, Core1 β1,3-galactosyltransferase (EC 2.4.1.122); GT, glycosyltransferase; HPA, *Helix pomatia* agglutinin; IDC, invasive ductal carcinoma; Le, Lewis; mAb, monoclonal antibody; MGL, macrophage galactose-type lectin; ORF, open reading frame; PNA, *Arachis hypogaea* agglutinin; sLe, sialyl-Lewis; sTn, sialyl-Thomsen-nouvelle; T, Thomsen-Friedenreich; TACA, tumour-associated carbohydrate antigen; Tn, Thomsen-nouvelle; VVA, *Vicia villosa* agglutinin isolectin B<sub>4</sub>.

#### Competing interests

The authors declare that they have no competing interests.

#### Acknowledgements

The present work was supported by the University of Sciences and Technologies of Lille, the Association pour la Recherche sur le Cancer (grant numbers 7936 and 5023), La Ligue régionale contre le Cancer and the PAI Hubert Curien France-Korea STAR.

#### Author details

<sup>1</sup>Structural and Functional Glycobiology Unit, UMR CNRS 8576, University of Sciences and Technologies of Lille, 59655 Villeneuve d'Ascq, France. <sup>2</sup>Université Lille Nord de France, F-59000 Lille, France. <sup>3</sup>Breast Cancer Biology Group, Research Oncology, King's College London, London SE1 9RT, UK.

Published: 8 June 2010

#### References

1. Varki A: **Selectin ligands.** *Proc Natl Acad Sci USA* 1994, **91**:7390-7397.
2. Le Marer N, Laudet V, Svensson EC, Cazlaris H, Van Hille B, Lagrou C, Stehelin D, Montreuil J, Verbert A, Delannoy P: **The c-Ha-ras oncogene induces increased expression of beta-galactoside alpha-2, 6-sialyltransferase in rat fibroblast (FR3T3) cells.** *Glycobiology* 1992, **2**:49-56.
3. Buckhaults P, Chen L, Fregien N, Pierce M: **Transcriptional regulation of N-acetylglucosaminyltransferase V by the src oncogene.** *J Biol Chem* 1997, **272**:19575-19581.
4. Narita T, Funahashi H, Satoh Y, Watanabe T, Sakamoto J, Takagi H: **Association of expression of blood group-related carbohydrate antigens with prognosis in breast cancer.** *Cancer* 1993, **71**:3044-3053.
5. Soares R, Marinho A, Schmitt F: **Expression of sialyl-Tn in breast cancer. Correlation with prognostic parameters.** *Pathol Res Pract* 1996, **192**:1181-1186.
6. Ura Y, Dion AS, Williams CJ, Olsen BD, Redfield ES, Ishida M, Herlyn M, Major PP: **Quantitative dot blot analyses of blood-group-related antigens in paired normal and malignant human breast tissues.** *Int J Cancer* 1992, **50**:57-63.
7. Miles DW, Happerfield LC, Smith P, Gillibrand R, Bobrow LG, Gregory WM, Rubens RD: **Expression of sialyl-Tn predicts the effect of adjuvant chemotherapy in node-positive breast cancer.** *Br J Cancer* 1994, **70**:1272-1275.
8. Hebbar M, Krzewinski-Recchi MA, Hornez L, Verdier A, Harduin-Lepers A, Bonnetterre J, Delannoy P, Peyrat JP: **Prognostic value of tumoral sialyltransferase expression and circulating E-selectin concentrations in node-negative breast cancer patients.** *Int J Biol Markers* 2003, **18**:116-122.
9. Recchi MA, Hebbar M, Hornez L, Harduin-Lepers A, Peyrat JP, Delannoy P: **Multiplex reverse transcription polymerase chain reaction assessment of sialyltransferase expression in human breast cancer.** *Cancer Res* 1998, **58**:4066-4070.
10. Julien S, Adriaenssens E, Ottenberg K, Furlan A, Courtand G, Vercoutter-Edouart AS, Hanisch FG, Delannoy P, Le Bourhis X: **ST6GalNAc I expression in MDA-MB-231 breast cancer cells greatly modifies their O-glycosylation**

pattern and enhances their tumourigenicity. *Glycobiology* 2006, **16**:54-64.

11. Julien S, Lagadec C, Krzewinski-Recchi MA, Courtand G, Le Bourhis X, Delannoy P: **Stable expression of sialyl-Tn antigen in T47-D cells induces a decrease of cell adhesion and an increase of cell migration.** *Breast Cancer Res Treat* 2005, **90**:77-84.
12. Bos PD, Zhang XH, Nadal C, Shu W, Gomis RR, Nguyen DX, Minn AJ, van de Vijver MJ, Gerald WL, Foekens JA, Massagué J: **Genes that mediate breast cancer metastasis to the brain.** *Nature* 2009, **459**:1005-1009.
13. Cazet A, Groux-Degroote S, Teylaert B, Kwon KM, Lehoux S, Slomianny C, Kim CH, Le Bourhis X, Delannoy P: **GD3 synthase overexpression enhances proliferation and migration of MDA-MB-231 breast cancer cells.** *Biol Chem* 2009, **390**:601-609.
14. Rakha EA, Boyce RW, Abd El-Rehim D, Kurien T, Green AR, Paish EC, Robertson JF, Ellis IO: **Expression of mucins (MUC1, MUC2, MUC3, MUC4, MUC5AC and MUC6) and their prognostic significance in human breast cancer.** *Mod Pathol* 2005, **18**:1295-1304.
15. Hanisch FG, Uhlenbruck G, Peter-Katalinic J, Egge H, Dabrowski J, Dabrowski U: **Structures of neutral O-linked polylactosaminoglycans on human skim milk mucins. A novel type of linearly extended poly-N-acetylglucosamine backbones with Gal beta(1-4)GlcNAc beta(1-6) repeating units.** *J Biol Chem* 1989, **264**:872-883.
16. Croce MV, Isla-Larrain M, Rabassa ME, Demichelis S, Colussi AG, Crespo M, Lacunza E, Segal-Eiras A: **Lewis x is highly expressed in normal tissues: a comparative immunohistochemical study and literature revision.** *Pathol Oncol Res* 2007, **13**:130-138.
17. Muller S, Hanisch FG: **Recombinant MUC1 probe authentically reflects cell-specific O-glycosylation profiles of endogenous breast cancer mucin. High density and prevalent core 2-based glycosylation.** *J Biol Chem* 2002, **277**:26103-26112.
18. Abecassis J, Collard R, Eber M, Pusel J, Fricker JP, Methlin G: **Proteinases and sialyltransferase in human breast tumors.** *Int J Cancer* 1984, **33**:821-824.
19. Springer GF: **T and Tn, general carcinoma autoantigens.** *Science* 1984, **224**:1198-1206.
20. Springer GF, Desai PR, Murthy MS, Scanlon EF: **Human carcinoma-associated precursor antigens of the NM blood group system.** *J Surg Oncol* 1979, **11**:95-106.
21. Schmitt FC, Figueiredo P, Lacerda M: **Simple mucin-type carbohydrate antigens (T, sialosyl-T, Tn and sialosyl-Tn) in breast carcinogenesis.** *Virchows Arch* 1995, **427**:251-258.
22. Tachibana K, Nakamura S, Wang H, Iwasaki H, Maebara K, Cheng L, Hirabayashi J, Narimatsu H: **Elucidation of binding specificity of Jacalin toward O-glycosylated peptides: quantitative analysis by frontal affinity chromatography.** *Glycobiology* 2006, **16**:46-53.
23. Wu AM, Wu JH, Yang Z, Singh T, Goldstein IJ, Sharon N: **Differential contributions of recognition factors of two plant lectins, *Amaranthus caudatus* lectin and *Arachis hypogaea* agglutinin, reacting with Thomsen-Friedenreich disaccharide (Galβ1-3GalNAcα1-Ser/Thr).** *Biochimie* 2008, **90**:1769-1780.
24. Lotan R, Skutelsky E, Danon D, Sharon N: **The purification, composition, and specificity of the anti-T lectin from peanut (*Arachis hypogaea*).** *J Biol Chem* 1975, **250**:8518-8523.
25. Pereira ME, Kabat EA, Lotan R, Sharon N: **Immunochemical studies on the specificity of the peanut (*Arachis hypogaea*) agglutinin.** *Carbohydr Res* 1976, **51**:107-118.
26. Hanisch FG, Baldus SE: **The Thomsen-Friedenreich (TF) antigen: a critical review on the structural, biosynthetic and histochemical aspects of a pancarcinoma-associated antigen.** *Histol Histopathol* 1997, **12**:263-281.
27. Cao Y, Stosiek P, Springer GF, Karsten U: **Thomsen-Friedenreich-related carbohydrate antigens in normal adult human tissues: a systematic and comparative study.** *Histochem Cell Biol* 1996, **106**:197-207.
28. Schindlbeck C, Jeschke U, Schulze S, Karsten U, Janni W, Rack B, Krajewski S, Sommer H, Friese K: **Prognostic impact of Thomsen-Friedenreich tumor antigen and disseminated tumor cells in the bone marrow of breast cancer patients.** *Breast Cancer Res Treat* 2007, **101**:17-25.
29. Schindlbeck C, Jeschke U, Schulze S, Karsten U, Janni W, Rack B, Sommer H, Friese K: **Characterisation of disseminated tumor cells in the bone marrow of breast cancer patients by the Thomsen-Friedenreich tumor antigen.** *Histochem Cell Biol* 2005, **123**:631-637.
30. Glinzky VV, Glinzky GV, Glinzky OV, Huxley VH, Turk JR, Mossine VV, Deutscher SL, Pienta KJ, Quinn TP: **Intravascular metastatic cancer cell homotypic aggregation at the sites of primary attachment to the endothelium.**

- Cancer Res* 2003, **63**:3805-3811.
31. Yu LG, Andrews N, Zhao Q, McKean D, Williams JF, Connor LJ, Gerasimenko OV, Hilkens J, Hirabayashi J, Kasai K, Rhodes JM: **Galectin-3 interaction with Thomsen-Friedenreich disaccharide on cancer-associated MUC1 causes increased cancer cell endothelial adhesion.** *J Biol Chem* 2007, **282**:773-781.
  32. Berger EG: **Tn-syndrome.** *Biochim Biophys Acta* 1999, **1455**:255-268.
  33. Ju T, Cummings RD: **Protein glycosylation: chaperone mutation in Tn syndrome.** *Nature* 2005, **437**:1252.
  34. Ju T, Lanneau GS, Gautam T, Wang Y, Xia B, Stowell SR, Willard MT, Wang W, Xia JY, Zuna RE, Laszik Z, Benbrook DM, Hanigan MH, Cummings RD: **Human tumor antigens Tn and sialyl Tn arise from mutations in Cosmc.** *Cancer Res* 2008, **68**:1636-1646.
  35. Yoo NJ, Kim MS, Lee SH: **Absence of COSMC gene mutations in breast and colorectal carcinomas.** *Apmis* 2008, **116**:154-155.
  36. Piller V, Piller F, Cartron JP: **Comparison of the carbohydrate-binding specificities of seven N-acetyl-D-galactosamine-recognizing lectins.** *Eur J Biochem* 1990, **191**:461-466.
  37. Brooks SA: **The involvement of Helix pomatia lectin (HPA) binding N-acetylgalactosamine glycans in cancer progression.** *Histol Histopathol* 2000, **15**:143-158.
  38. Springer GF: **Immunoreactive T and Tn epitopes in cancer diagnosis, prognosis, and immunotherapy.** *J Mol Med* 1997, **75**:594-602.
  39. Wang BL, Springer GF, Carlstedt SC: **Quantitative computerized image analysis of Tn and T (Thomsen-Friedenreich) epitopes in prognostication of human breast carcinoma.** *J Histochem Cytochem* 1997, **45**:1393-1400.
  40. Korourian S, Siegel E, Kieber-Emmons T, Monzavi-Karbassi B: **Expression analysis of carbohydrate antigens in ductal carcinoma in situ of the breast by lectin histochemistry.** *BMC Cancer* 2008, **8**:136.
  41. Kanska G, Guerry N, Caldefie-Chezet F, De Latour M, Guillot J: **Study of the expression of Tn antigen in different types of human breast cancer cells using VVA-B4 lectin.** *Oncol Rep* 2006, **15**:305-310.
  42. Kawaguchi T, Takazawa H, Imai S, Morimoto J, Watanabe T, Kanno M, Igarashi S: **Expression of Vicia villosa agglutinin (VVA)-binding glycoprotein in primary breast cancer cells in relation to lymphatic metastasis: is atypical MUC1 bearing Tn antigen a receptor of VVA?** *Breast Cancer Res Treat* 2006, **98**:31-43.
  43. Brooks SA, Leatham AJ: **Prediction of lymph node involvement in breast cancer by detection of altered glycosylation in the primary tumour.** *Lancet* 1991, **338**:71-74.
  44. Leatham AJ, Brooks SA: **Predictive value of lectin binding on breast-cancer recurrence and survival.** *Lancet* 1987, **1**:1054-1056.
  45. Danussi C, Coslovi A, Campa C, Mucignat MT, Spessotto P, Uggeri F, Paoletti S, Colombatti A: **A newly generated functional antibody identifies Tn antigen as a novel determinant in the cancer cell-lymphatic endothelium interaction.** *Glycobiology* 2009, **19**:1056-1067.
  46. van Vliet SJ, Saeland E, van Kooyk Y: **Sweet preferences of MGL: carbohydrate specificity and function.** *Trends Immunol* 2008, **29**:83-90.
  47. Saeland E, van Vliet SJ, Backstrom M, van den Berg VC, Geijtenbeek TB, Meijer GA, van Kooyk Y: **The C-type lectin MGL expressed by dendritic cells detects glycan changes on MUC1 in colon carcinoma.** *Cancer Immunol Immunother* 2007, **56**:1225-1236.
  48. Napolitano C, Rughetti A, Agervig Tarp MP, Coleman J, Bennett EP, Picco G, Sale P, Denda-Nagai K, Irimura T, Mandel U, Clausen H, Frati L, Taylor-Papadimitriou J, Burchell J, Nuti M: **Tumor-associated Tn-MUC1 glycoform is internalized through the macrophage galactose-type C-type lectin and delivered to the HLA class I and II compartments in dendritic cells.** *Cancer Res* 2007, **67**:8358-8367.
  49. Julien S, Delannoy P: **Sialyl-Tn antigen in cancer: from diagnosis to therapy.** In *Recent Research Developments in Cancer. Volume 5*. Edited by Pandalai SG. Kerala: Transworld Research Network; 2003:185-199.
  50. Julien S, Krzewinski-Recchi MA, Harduin-Lepers A, Gouyer V, Huet G, Le Bourhis X, Delannoy P: **Expression of sialyl-Tn antigen in breast cancer cells transfected with the human CMP-Neu5Ac: GalNAc alpha2,6-sialyltransferase (ST6GalNAc I) cDNA.** *Glycoconj J* 2001, **18**:883-893.
  51. Sewell R, Backstrom M, Dalziel M, Gschmeissner S, Karlsson H, Noll T, Gatgens J, Clausen H, Hansson GC, Burchell J, Nuti M: **The ST6GalNAc-I sialyltransferase localizes throughout the Golgi and is responsible for the synthesis of the tumor-associated sialyl-Tn O-glycan in human breast cancer.** *J Biol Chem* 2006, **281**:3586-3594.
  52. Gold DV, Mattes MJ: **Monoclonal antibody B72.3 reacts with a core region structure of O-linked carbohydrates.** *Tumour Biol* 1988, **9**:137-144.
  53. Reddish MA, Jackson L, Koganty RR, Qiu D, Hong W, Longenecker BM: **Specificities of anti-sialyl-Tn and anti-Tn monoclonal antibodies generated using novel clustered synthetic glycopeptide epitopes.** *Glycoconj J* 1997, **14**:549-560.
  54. Ogata S, Koganty R, Reddish M, Longenecker BM, Chen A, Perez C, Itzkowitz SH: **Different modes of sialyl-Tn expression during malignant transformation of human colonic mucosa.** *Glycoconj J* 1998, **15**:29-35.
  55. Kinney AY, Sahin A, Vernon SW, Frankowski RF, Annegers JF, Hortobagyi GN, Buzdar AU, Frye DK, Dhingra K: **The prognostic significance of sialyl-Tn antigen in women treated with breast carcinoma treated with adjuvant chemotherapy.** *Cancer* 1997, **80**:2240-2249.
  56. Leivonen M, Nordling S, Lundin J, von Boguslawski K, Haglund C: **STn and prognosis in breast cancer.** *Oncology* 2001, **61**:299-305.
  57. Clement M, Rocher J, Loirand G, Le Pendu J: **Expression of sialyl-Tn epitopes on beta1 integrin alters epithelial cell phenotype, proliferation and haptotaxis.** *J Cell Sci* 2004, **117**(Pt 21):5059-5069.
  58. Julien S, Picco G, Sewell R, Vercoutter-Edouart AS, Tarp M, Miles D, Clausen H, Taylor-Papadimitriou J, Burchell JM: **Sialyl-Tn vaccine induces antibody-mediated tumour protection in a relevant murine model.** *Br J Cancer* 2009, **100**:1746-1754.
  59. Ravn V, Dabelsteen E: **Tissue distribution of histo-blood group antigens.** *Apmis* 2000, **108**:1-28.
  60. Kaneo M, Kudo T, Iwasaki H, Ikehara Y, Nishihara S, Nakagawa S, Sasaki K, Shiina T, Inoko H, Saitou N, Narimatsu H: **Alpha1,3-fucosyltransferase IX (Fuc-TIX) is very highly conserved between human and mouse; molecular cloning, characterization and tissue distribution of human Fuc-TIX.** *FEBS Lett* 1999, **452**:237-242.
  61. Mollicone R, Gibaud A, Francois A, Ratcliffe M, Oriol R: **Acceptor specificity and tissue distribution of three human alpha-3-fucosyltransferases.** *Eur J Biochem* 1990, **191**:169-176.
  62. Natsuka S, Gersten KM, Zenita K, Kannagi R, Lowe JB: **Molecular cloning of a cDNA encoding a novel human leukocyte alpha-1,3-fucosyltransferase capable of synthesizing the sialyl Lewis x determinant.** *J Biol Chem* 1994, **269**:16789-16794.
  63. Kitagawa H, Paulson JC: **Cloning and expression of human Gal beta 1,3(4) GlcNAc alpha 2,3-sialyltransferase.** *Biochem Biophys Res Commun* 1993, **194**:375-382.
  64. Ellies LG, Sperandio M, Underhill GH, Yousif J, Smith M, Priatel JJ, Kansas GS, Ley K, Marth JD: **Sialyltransferase specificity in selectin ligand formation.** *Blood* 2002, **100**:3618-3625.
  65. Kukowska-Latallo JF, Larsen RD, Nair RP, Lowe JB: **A cloned human cDNA determines expression of a mouse stage-specific embryonic antigen and the Lewis blood group alpha(1,3/1,4)fucosyltransferase.** *Genes Dev* 1990, **4**:1288-1303.
  66. Sasaki K, Kurata K, Kojima N, Kurosawa N, Ohta S, Hanai N, Tsuji S, Nishi T: **Expression cloning of a GM3-specific alpha-2,8-sialyltransferase (GD3 synthase).** *J Biol Chem* 1994, **269**:15950-15956.
  67. Maly P, Thall A, Petryniak B, Rogers CE, Smith PL, Marks RM, Kelly RJ, Gersten KM, Cheng G, Saunders TL, Camper SA, Camphausen RT, Sullivan FX, Isogai Y, Hinds Gaul O, von Andrian UH, Lowe JB: **The alpha(1,3)fucosyltransferase Fuc-TVII controls leukocyte trafficking through an essential role in L-, E-, and P-selectin ligand biosynthesis.** *Cell* 1996, **86**:643-653.
  68. Renkonen J, Paavonen T, Renkonen R: **Endothelial and epithelial expression of sialyl Lewis(x) and sialyl Lewis(a) in lesions of breast carcinoma.** *Int J Cancer* 1997, **74**:296-300.
  69. Matsuura N, Narita T, Mitsuoka C, Kimura N, Kannagi R, Imai T, Funahashi H, Takagi H: **Increased level of circulating adhesion molecules in the sera of breast cancer patients with distant metastases.** *Jpn J Clin Oncol* 1997, **27**:135-139.
  70. Jeschke U, Mylonas I, Shabani N, Kunert-Keil C, Schindlbeck C, Gerber B, Friese K: **Expression of sialyl Lewis X, sialyl Lewis A, E-cadherin and cathepsin-D in human breast cancer: immunohistochemical analysis in mammary carcinoma in situ, invasive carcinomas and their lymph node metastasis.** *Anticancer Res* 2005, **25**:1615-1622.
  71. Nakagoe T, Fukushima K, Itoyanagi N, Ikuta Y, Oka T, Nagayasu T, Ayabe H, Hara S, Ishikawa H, Minami H: **Expression of ABH/Lewis-related antigens as prognostic factors in patients with breast cancer.** *J Cancer Res Clin Oncol* 2002, **128**:257-264.
  72. Matsuura N, Narita T, Hiraiwa N, Hiraiwa M, Murai H, Iwase T, Funahashi H, Imai T, Takagi H, Kannagi R: **Gene expression of fucosyl- and sialyl-transferases which synthesize sialyl Lewisx, the carbohydrate ligands for E-selectin, in**

- human breast cancer. *Int J Oncol* 1998, **12**:1157-1164.
73. Hsu CC, Lin TW, Chang WW, Wu CY, Lo WH, Wang PH, Tsai YC: **Soyasaponin-I-modified invasive behavior of cancer by changing cell surface sialic acids.** *Gynecol Oncol* 2005, **96**:415-422.
74. Idikio HA, Manickavel V: **Lewis blood group antigens (a and b) in human breast tissues. Loss of Lewis-b in breast cancer cells and correlation with tumor grade.** *Cancer* 1991, **68**:1303-1308.
75. Abd Hamid UM, Royle L, Saldova R, Radcliffe CM, Harvey DJ, Storr SJ, Pardo M, Antrobus R, Chapman CJ, Zitzmann N, Robertson JF, Dwek RA, Rudd PM: **A strategy to reveal potential glycan markers from serum glycoproteins associated with breast cancer progression.** *Glycobiology* 2008, **18**:1105-1118.
76. Kurebayashi J, Nomura T, Hirono M, Okubo S, Udagawa K, Shiiki S, Ikeda M, Nakashima K, Tanaka K, Sonoo H: **Combined measurement of serum sialyl Lewis X with serum CA15-3 in breast cancer patients.** *Jpn J Clin Oncol* 2006, **36**:150-153.
77. Hellstrom I, Garrigues HJ, Garrigues U, Hellstrom KE: **Highly tumor-reactive, internalizing, mouse monoclonal antibodies to Le(y)-related cell surface antigens.** *Cancer Res* 1990, **50**:2183-2190.
78. Madjd Z, Parsons T, Watson NF, Spendlove I, Ellis I, Durrant LG: **High expression of Lewis y/b antigens is associated with decreased survival in lymph node negative breast carcinomas.** *Breast Cancer Res* 2005, **7**:R780-R787.
79. Kelly MP, Lee FT, Smyth FE, Brechbiel MW, Scott AM: **Enhanced efficacy of 90Y-radiolabeled anti-Lewis Y humanized monoclonal antibody hu3S193 and paclitaxel combined-modality radioimmunotherapy in a breast cancer model.** *J Nucl Med* 2006, **47**:716-725.
80. Pinho SS, Reis CA, Gartner F, Alpaugh ML: **Molecular plasticity of E-cadherin and sialyl Lewis x expression, in two comparative models of mammary tumorigenesis.** *PLoS One* 2009, **4**:e6636.
81. Tettamanti G: **Ganglioside/glycosphingolipid turnover: new concepts.** *Glycoconj J* 2004, **20**:301-317.
82. Marquina G, Waki H, Fernandez LE, Kon K, Carr A, Valiente O, Perez R, Ando S: **Gangliosides expressed in human breast cancer.** *Cancer Res* 1996, **56**:5165-5171.
83. Oliva JP, Valdes Z, Casaco A, Pimentel G, Gonzalez J, Alvarez I, Osorio M, Velazco M, Figueroa M, Ortiz R, Escobar X, Orozco M, Cruz J, Franco S, Diaz M, Roque L, Carr A, Vazquez AM, Mateos C, Rubio MC, Pérez R, Fernández LE: **Clinical evidences of GM3 (NeuGc) ganglioside expression in human breast cancer using the 14F7 monoclonal antibody labelled with (99m)Tc.** *Breast Cancer Res Treat* 2006, **96**:115-121.
84. Ruckhaberle E, Rody A, Engels K, Gaetje R, von Minckwitz G, Schiffmann S, Grosch S, Geisslinger G, Holtrich U, Karn T, Kaufmann M: **Microarray analysis of altered sphingolipid metabolism reveals prognostic significance of sphingosine kinase 1 in breast cancer.** *Breast Cancer Res Treat* 2008, **112**:41-52.
85. Ruckhaberle E, Karn T, Rody A, Hanker L, Gatje R, Metzler D, Holtrich U, Kaufmann M: **Gene expression of ceramide kinase, galactosyl ceramide synthase and ganglioside GD3 synthase is associated with prognosis in breast cancer.** *J Cancer Res Clin Oncol* 2009, **135**:1005-1013.
86. Gu Y, Zhang J, Mi W, Yang J, Han F, Lu X, Yu W: **Silencing of GM3 synthase suppresses lung metastasis of murine breast cancer cells.** *Breast Cancer Res* 2008, **10**:R1.
87. Goss PE, Baker MA, Carver JP, Dennis JW: **Inhibitors of carbohydrate processing: a new class of anticancer agents.** *Clin Cancer Res* 1995, **1**:935-944.
88. Paszkiewicz-Gadek A, Porowska H, Lemancewicz D, Wolczynski S, Gindzienski A: **The influence of N- and O-glycosylation inhibitors on the glycosylation profile of cellular membrane proteins and adhesive properties of carcinoma cell lines.** *Int J Mol Med* 2006, **17**:669-674.
89. Tanaka T, Decuzzi P, Cristofanilli M, Sakamoto JH, Tasciotti E, Robertson FM, Ferrari M: **Nanotechnology for breast cancer therapy.** *Biomed Microdevices* 2009, **11**:49-63.
90. Lo NW, Dennis JW, Lau JT: **Overexpression of the alpha2,6-sialyltransferase, ST6Gal I, in a low metastatic variant of a murine lymphoblastoid cell line is associated with appearance of a unique ST6Gal I mRNA.** *Biochem Biophys Res Commun* 1999, **264**:619-621.
91. Taniguchi A: **Promoter structure and transcriptional regulation of human beta-galactoside alpha2, 3-sialyltransferase genes.** *Curr Drug Targets* 2008, **9**:310-316.
92. Glinisky VV, Huflejt ME, Glinisky GV, Deutscher SL, Quinn TP: **Effects of Thomsen-Friedenreich antigen-specific peptide P-30 on beta-galactoside-mediated homotypic aggregation and adhesion to the endothelium of MDA-MB-435 human breast carcinoma cells.** *Cancer Res* 2000, **60**:2584-2588.
93. Heimburg J, Yan J, Morey S, Gliniskii OV, Huxley VH, Wild L, Klick R, Roy R, Glinisky VV, Rittenhouse-Olson K: **Inhibition of spontaneous breast cancer metastasis by anti-Thomsen-Friedenreich antigen monoclonal antibody JAA-F11.** *Neoplasia* 2006, **8**:939-948.
94. Fernandez LE, Alonso DF, Gomez DE, Vazquez AM: **Ganglioside-based vaccines and anti-idiotype antibodies for active immunotherapy against cancer.** *Expert Rev Vaccines* 2003, **2**:817-823.
95. Vazquez AM, Gabri MR, Hernandez AM, Alonso DF, Beausoleil I, Gomez DE, Perez R: **Antitumor properties of an anti-idiotypic monoclonal antibody in relation to N-glycolyl-containing gangliosides.** *Oncol Rep* 2000, **7**:751-756.
96. Fuentes D, Avellanet J, Garcia A, Iglesias N, Gabri MR, Alonso DF, Vazquez AM, Perez R, Montero E: **Combined therapeutic effect of a monoclonal anti-idiotype tumor vaccine against NeuGc-containing gangliosides with chemotherapy in a breast carcinoma model.** *Breast Cancer Res Treat* 2010, **120**:379-389.
97. Diaz A, Alfonso M, Alonso R, Saurez G, Troche M, Catala M, Diaz RM, Perez R, Vazquez AM: **Immune responses in breast cancer patients immunized with an anti-idiotype antibody mimicking NeuGc-containing gangliosides.** *Clin Immunol* 2003, **107**:80-89.
98. Guthmann MD, Castro MA, Cinat G, Venier C, Koliren L, Bitton RJ, Vazquez AM, Fainboim L: **Cellular and humoral immune response to N-Glycolyl-GM3 elicited by prolonged immunotherapy with an anti-idiotypic vaccine in high-risk and metastatic breast cancer patients.** *J Immunother* 2006, **29**:215-223.
99. Miles D, Papazisis K: **Rationale for the clinical development of STn-KLH (Theratope) and anti-MUC-1 vaccines in breast cancer.** *Clin Breast Cancer* 2003, **3**(Suppl 4):S134-S138.
100. Slovin SF, Ragupathi G, Fernandez C, Diani M, Jefferson MP, Wilton A, Kelly WK, Morris M, Solit D, Clausen H, Livingston P, Scher HI: **A polyvalent vaccine for high-risk prostate patients: 'are more antigens better?'** *Cancer Immunol Immunother* 2007, **56**:1921-1930.

doi:10.1186/bcr2577

Cite this article as: Cazet A, et al.: Tumour-associated carbohydrate antigens in breast cancer. *Breast Cancer Research* 2010, **12**:204.