

Corrigendum to: Intravenous Injection of Coronavirus Disease 2019 (COVID-19) mRNA Vaccine Can Induce Acute Myopericarditis in Mouse Model

Can Li,¹ Yanxia Chen,¹ Yan Zhao,¹ David Christopher Lung,² Zhanhong Ye,¹ Wenchen Song,¹ Fei-Fei Liu,¹ Jian-Piao Cai,¹ Wan-Man Wong,¹ Cyril Chik-Yan Yip,¹ Jasper Fuk-Woo Chan,^{1,3,4} Kelvin Kai-Wang To,^{1,3} Siddharth Sridhar,^{1,3} Ivan Fan-Ngai Hung,^{3,5} Hin Chu,¹ Kin-Hang Kok,¹ Dong-Yan Jin,⁶ Anna Jinxia Zhang,¹ and Kwok-Yung Yuen^{1,3,4}

¹State Key Laboratory of Emerging Infectious Diseases, Carol Yu Centre for Infection, Department of Microbiology, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong Special Administrative Region, China; ²Department of Pathology, Queen Elizabeth Hospital and Hong Kong Children's Hospital, Hong Kong, Hong Kong Special Administrative Region, China; ³Department of Clinical Microbiology and Infection Control, The University of Hong Kong-Shenzhen Hospital, Shenzhen, Guangdong, China; ⁴Hainan Medical University-The University of Hong Kong Joint Laboratory of Tropical Infectious Diseases, The University of Hong Kong, Pokfulam, Hong Kong Special Administrative Region, China; ⁵Department of Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong Special Administrative Region, China; and ⁶School of Biomedical Sciences, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong Special Administrative Region, China


After the original publication of this manuscript [Li C, Chen Y, Zhao Y et al. Intravenous Injection of Coronavirus Disease 2019 (COVID-19) mRNA Vaccine Can Induce Acute Myopericarditis in Mouse Model. *Clin Infect Dis*, August 2021. <https://doi.org/10.1093/cid/ciab707>], it was discovered that Figure 5C on the TUNEL staining for apoptotic

changes was mistakenly taken from another group of mice. This does not affect the findings for this manuscript, and a revised figure is available in the version currently online. The statement 'Serum troponin level was significantly higher in IV group.' has also been inserted into the Results section of the abstract.

Correspondence: K.-Y. Yuen, State Key Laboratory of Emerging Infectious Diseases, Carol Yu Centre for Infection, Department of Microbiology, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong Special Administrative Region, China; and Department of Clinical Microbiology and Infection Control, The University of Hong Kong-Shenzhen Hospital, Shenzhen, China (kyyuen@hku.hk).

Clinical Infectious Diseases® 2021;73(12):2372–5

© The Author(s) 2021. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com.



<https://doi.org/10.1093/cid/ciab941>