

EPP1478

Postpartum depression screening scale-7: A valid and reliable short version both for Portugal and Brazil

A.T. Pereira^{1*}, M. Barros^{1,2}, M. Aguiar³, J. Azevedo¹, M. Marques¹, F. Carvalho¹, D. Pereira¹ and A. Macedo⁴

¹Institute Of Psychological Medicine, Faculty Of Medicine, University of Coimbra, Coimbra, Portugal; ²Departamento De Ciências Naturais, Universidade do Sudoeste da Bahia - UESB, Vitória da Conquista, Bahia, Brazil; ³Pos Graduação Em Psicologia Da Saúde, Universidade Federal da Bahia, Vitória da Conquista, Brazil and ⁴Institute Of Psychological Medicine, Faculty Of Medicine, University of Coimbra, Coimbra, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1618

Introduction: Screening programs for perinatal depression are systematically implemented in developed countries. To circumvent the most commonly pointed limitation by the primary healthcare professionals (the questionnaires length), we have developed shorter forms of the Beck and Gable Postpartum Depression Screening Scale-35. The shortest version consists of seven items, each one representing a dimension evaluated by the PDSS. This PDSS-7 demonstrated equal levels of reliability and validity as the 35-item PDSS with the advantage of being completed in as little as 1-2 minutes (Pereira et al. 2013).

Objectives: To analyze the construct validity of the PDSS-7 using Confirmatory Factor Analysis, to use both in Portugal and in Brazil.

Methods: The Portuguese sample was composed of 616 women (Mean age: 32.29±4.466; Mean gestation weeks=17.13±4.929). These participants were not the same who participated in the psychometric study that led to the selection of the seven items. The Brazilian sample was composed of 350 women (Mean age: 30.01±5.452; Mean gestation weeks=25.17±6.55). They all had uncomplicated pregnancies and completed the European/Brazilian Portuguese versions of PDSS-24 (Pereira et al. 2013/ Barros et al. 2021), which was composed of the same items and included the seven items that compose the PDSS-7.

Results: The unidimensional model of PDSS-7 presented a good fit in both samples (Portuguese/Brazilian: $\chi^2/d.f.=3.439/2.653$; RMSEA=.066/.069, CFI=.974/.981, TLI=.947/.957, GFI=.939/.957; $p<.001$). The PDSS-7 Cronbach's alphas were .82/.83 and all the items contribute to the internal consistency.

Conclusions: The PDSS-7 is a valid and precise, economic, fast and easy screening instrument for perinatal depression, a major public health problem, both in Portugal and in Brazil.

Keywords: PDSS-7; Perinatal depression; Postpartum depression

EPP1477

Can stress predict delivery date?: Role of chronic and acute stress to the threatened preterm labor as predictors of delivery date.

J. Buesa Lorenzo^{1*}, A. García-Blanco², M. Vento², A. Moreno-Giménez², L. Campos Berga¹, V. Diago³, D. Hervás⁴, C. Cháfer-Pericás² and P. Sáenz González²

¹Psychiatry, University and Polytechnic Hospital La Fe, Valencia, Spain; ²Neonatal Research Group, The Medical Research Institute

Hospital La Fe (IIS La Fe), Valencia, Spain; ³Gynaecology And Obstetrics, University and Polytechnic Hospital La Fe, Valencia, Spain and ⁴Data Science Unit, Biostatistics, And Bioinformatics, The Medical Research Institute Hospital La Fe (IIS La Fe), Valencia, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1619

Introduction: Threatened preterm labor (TPL) is a traumatic event during pregnancy that involves a threat to the physical integrity of the upcoming baby. Despite biomarkers would be the strongest delivery date predictors, an assessment of chronic and acute stress response to TPL diagnosis may improve this prediction.

Objectives: The objective is to predict delivery date in women with TPL based on their response to this diagnosis and chronic stressors, along with relevant obstetric variables.

Methods: A prospective cohort study was conducted with a sample was formed by 157 pregnant women with TPL diagnosis between 24 and 31 weeks. Determination of salivary cortisol, α -amylase levels, along with anxiety and depression symptoms were measured to estimate stress response to TPL. Cumulative life stressors as traumas, social and familiar functioning were also registered. To examine the effect of the possible predictor variables of delivery date, linear regression models were used.

Results: A correlation was found between the variables of response to chronic stress and between the variables of psychological response to stress. The main predictors of preterm delivery were low family adaptation, higher BMI, higher cortisol levels, and the week of diagnosis of TPL (<29 weeks of gestation).

Conclusions: The best predictor of delivery date was the combination of the stress response to the diagnosis of TPL measured by cortisol in saliva, cumulative life stressors (mainly family adaptation) and obstetric factors (week TPL and BMI). Through psychosocial therapeutic intervention programs, it is possible to influence this modifiable predictive factors of preterm birth in symptomatic women.

Keywords: stress; Threatened preterm labor; predictor

EPP1478

Together we stand, resilient we stay : The effect of minority stress and resilience on transgender mental health

A.B. Sahin^{1*} and D. Buyukgok²

¹Department Of Psychiatry, Basaksehir Cam and Sakura City Hospital, Istanbul, Turkey and ²Department Of Psychiatry, Istanbul University Istanbul Faculty of Medicine, Istanbul, Turkey

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1620

Introduction: Prejudice, stigmatization and discrimination behaviors causes social stress and lead vulnerability to mental and physical health problems in Transgender and Gender Nonconforming (TGNC) individuals. The prevalence of mental disorders that can be associated with "minority group stress", especially major depression and anxiety disorders, are known to be higher in the TGNC group in comparison to general population.