



POSTER PRESENTATION

Open Access

KIR-HLA footprints and NK cell-mediated recognition of HIV-1

CF Thobakgale^{1*}, J Carlson², J Wright¹, N van Teijlingen³, M Jaggernath¹, P Goulder⁴, BD Walker³, M Carrington⁵, D Heckerman², M Altfeld³, T Ndung'u¹

From AIDS Vaccine 2012

Boston, MA, USA. 9-12 September 2012

Background

Increasing data suggest an important role of KIR+ NK cells in the control of HIV-1, however the precise mechanisms on how NK cells recognize HIV-1-infected cells remain poorly understood. KIRs can bind to HLA class I molecules, but the binding affinity of these interactions is dependent on the HLA class I presented epitope. Recent reports have suggested that sequence variations within HIV-1 epitopes presented by HLA class I can affect the binding of inhibitory KIRs expressed on NK cells, potentially modulating NK cell responses to infected cells. Here we investigated whether HIV-1 might adapt to the combined KIR/HLA genotypes on a population level to identify areas within HIV-1 that might be targeted by KIR+ NK cells.

Methods

HIV-1 Gag was sequenced in 390 untreated chronically clade C infected individuals from KwaZulu-Natal, South Africa. All study subjects were HLA class I and KIR typed. Phylogenetically-corrected logistic regression analysis of KIR/HLA associated Gag sequence polymorphisms was performed and q-values were used for multiple test correction.

Results

A total of 93 sequence polymorphisms significantly associated with the combined HLA/KIR genotypes were identified ($p < 0.05$), 6 of them with a false-positive rate of less than 20% ($q < 0.2$). These significant associations were independent of previously identified KIR or HLA-linked polymorphisms.

Conclusion

This study identified several sequence polymorphisms within HIV-1 Gag that were significantly associated with the expression of combined KIR/HLA genotypes at the population level, indicating adaptation of HIV-1 to NK cell mediated immune pressure. KIR/HLA class I binding studies in the context of the sequence polymorphisms and studies for NK cell function are ongoing to determine the consequence of these sequence changes for NK cell-mediated recognition of HIV-1-infected cells.

Author details

¹University of KwaZulu-Natal, Durban, South Africa. ²Microsoft Research, Redmond, WA, USA. ³Ragon Institute of MIT, MGH and Harvard, Boston, MA, USA. ⁴University of Oxford, Oxford, UK. ⁵Cancer and Inflammation Program, Laboratory of Experimental Immunology, Frederick, MD, USA.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P170

Cite this article as: Thobakgale et al.: KIR-HLA footprints and NK cell-mediated recognition of HIV-1. *Retrovirology* 2012 **9**(Suppl 2):P170.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹University of KwaZulu-Natal, Durban, South Africa
Full list of author information is available at the end of the article