

CASE REPORT

Gut microbiota modulation and gold nanoparticle-mediated photothermal therapy for treatment of recalcitrant acne

Jimyung Seo  | Hyo Jin Roh | Jin Young Jung

Department of Dermatology, My Mirae Dermatology & Plastic Surgery Clinic, Seoul, Republic of Korea, Korea

Correspondence

Jin Young Jung, Department of Dermatology, My Mirae Dermatology & Plastic Surgery Clinic, 7, Gukjegeumyung-ro 2-gil, Yeongdeungpo-gu, Seoul, Republic of Korea 07327, Korea.
Email: yoracle@naver.com

Funding information

None

Abstract

Recent studies highlight that gut dysbiosis, an imbalanced state of intestinal microbiota, exacerbates skin inflammation. Here, we showed the presence of gut microbiota alterations in two patients with recalcitrant acne and investigated the impact of its therapeutic modulation together with gold nanoshell-mediated photothermal therapy (gold PTT).

KEYWORDS

acne, antibiotics, gold nanoshell, gold PTT, gut dysbiosis, gut microbiota alteration

1 | INTRODUCTION

Recent studies highlight that gut dysbiosis, an imbalanced state of intestinal microbiota, exacerbates skin inflammation.^{1,2} Here, we showed the presence of gut microbiota alterations in two patients with recalcitrant acne and investigated the impact of its therapeutic modulation together with gold nanoshell-mediated photothermal therapy (gold PTT).

2 | CASE PRESENTATION

2.1 | Case 1

A 30-year-old Korean male patient was suffered from adult acne that had worsened over the years. Oral isotretinoin was not tolerable due to his dry skin. Therefore, minocycline and doxycycline were used alternately for the acne treatment. Despite additional therapies involving fractional and needle-radiofrequency lasers, inflammatory papules recurred too often and caused pain. Gut microbiota

alteration was evaluated by microbial organic acid test (MOAT) identifying yeast and fungal, Clostridia bacterial, and other bacteria markers: Urine samples were sent to the Great Plains Laboratory, Inc. (Lenexa, KS, United States) for analysis.³ The result of his MAOT indicated an abnormal overgrowth of candida and harmful bacteria of the intestinal tract with an elevation of arabinose and hippuric acid of urine. We discontinued oral antibiotics and performed gold PTT as an alternative. Each treatment of gold PTT included facial washing and then applying an ample of gold nanoparticle suspension (inno.N Co. Ltd.) into the facial skin by sonophoresis after obtaining informed consent.⁴ A 40 khz low-frequency ultrasound (BellaSonic, Umbrella Inc.,) was used to transdermally deliver gold nanoparticles into sebaceous glands and follicles. The residual nanoparticle suspension of superficial skin was wiped off to prevent unexpected thermal injury to the epidermis. After that, laser purses from long-pulsed 1064 nm Nd:YAG laser (Cosjet SR, Won Technology,) was applied to the patient's facial skin for photobiomodulation (pulse duration 0.3 ms; fluence 10 J/cm²; 1500 laser shots) and photothermolysis (pulse duration 10/20/30 ms;

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.




FIGURE 1 Clinical improvement of acne lesions in patient #1 after gold nanoshell-mediated photothermal therapy (gold PTT) with gut microbiota modulation: (A) before treatment, (B) after nine sessions of gold PTT, and (C) 12 months after the last procedure




FIGURE 2 (A) Schematic diagram showing the timing of gold nanoshell-mediated photothermal therapy (gold PTT) and gut microbiota modulation. (B) Graph showing changes of the number of facial inflamed lesions in the two patients with recalcitrant acne. (C, D) Clinical photographs of patient # 2 before (left) and after six sessions of gold PTT (right)

fluence 12 J/cm^2 ; 500 shots, respectively, for each pulse duration) with an interval of 2–4 weeks. In addition, the patient received nystatin (1.5 million units/day) and rifaximin (1200 mg/day) for 2 weeks to restore yeast and bacterial dysbiosis.⁵ The MAOT result returned to normal and clinical improvement was well maintained at 1-year follow-up after the last procedure (Figure 1).

2.2 | Case 2


A 38-year-old Korean male patient presented with recurrent acne of the cheek that lasted for 15 years. He had been taking doxycycline repeatedly for a period whenever his acne got worse. The MAOT result revealed Clostridia

bacterial and other bacterial overgrowth of gut with increased urinary levels of hippuric acid and HPHA. Rifaximin (1200 mg/day) was prescribed for 2 weeks to reverse microbial dysbiosis, and the patient underwent gold PTT as described above without oral acne medications. The acne lesions were dramatically reduced after six treatment sessions of PTT (Figure 2). Moreover, no major recurrences were observed during a follow-up period of 14 months after the ninth session of PTT.

3 | DISCUSSION

The treatment of acne in adult patients may take several years; the chronic use of oral antibiotics can turn the gut

FIGURE 3 Summary of the function of gold nanoshell-mediated photothermal therapy (gold PTT) and gut microbiota modulation in acne treatment



microbiota into a dysbiotic state. It has been noted that doxycycline, but not isotretinoin, causes immediate and long-term changes of murine gut microbiota *in vivo*.¹ In order to discontinue oral acne antibiotics, we used gold PTT as an alternative. Gold PTT is recently highlighted as a safe and effective treatment for recurrent acne.⁴ About 150 nm-sized gold nanoparticles are designed to be applied topically and transported into sebaceous glands and follicles by the trans-appendageal route. When the gold nanoparticles are exposed to pulses of light, they exhibit plasmon resonance, strongly absorb light from 700 to 1200 nm, and cause thermal injury to surrounding tissues. Therefore, in gold PTT for acne treatment, light-absorbing gold nanoparticles can selectively destruct sebaceous follicles in accordance with the principles of selective photothermolysis.⁶ Gold PTT is practically painless with minimal systemic and local side effects; however, our clinical experience has found that acne improvement is maintained for about 3 months after gold PTT monotherapy in Korean patients. No acne recurrences over a year in these two patients thus suggest that therapeutic intervention for gut dysbiosis may allow a longer period of remission after this PTT (Figure 3). Although further studies are necessary to elucidate the mechanism by which gut microbiota imbalance affect acne exacerbation, recent studies highlight gut microbiota alterations in acne patients.^{1,2,7} Though not yet fully understood, the impaired integrity and increased permeability of intestinal epithelium caused by gut dysbiosis may result in systemic inflammatory effect, such as increased plasma interferon-gamma and interleukin-1 β , leading to skin inflammation.⁸ In addition, in a disturbed intestinal barrier, gut microbiota and their metabolites enter the bloodstream, accumulate in the skin, and influence skin homeostasis. Metabolites from the gut microbiota also promote acne by

overactivating the mTOR signaling pathway.¹ Therefore, therapeutic strategies for managing gut microbiota alterations can be beneficial in the treatment of refractory acne.

ACKNOWLEDGMENT

None.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

AUTHOR CONTRIBUTION

J.S. designed the study and wrote the manuscript. H.J.R. and J.Y.J. conceived and designed the study.

ETHICAL APPROVAL

This study conforms to the journal's recognized standards.

CONSENT

Written informed consent was obtained from the patient to publish this report in accordance with the journal's patient consent policy.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Jimyung Seo  <https://orcid.org/0000-0003-3416-9990>

REFERENCES

1. Lee YB, Byun EJ, Kim HS. Potential role of the microbiome in acne: a comprehensive review. *J Clin Med*. 2019;8(7):987.
2. Salem I, Ramser A, Isham N, et al. The gut microbiome as a major regulator of the gut-skin axis. *Front Microbiol*. 2018;9:1459.

3. Lord RS, Bralley JA. Clinical applications of urinary organic acids. Part 2. dysbiosis markers. *Altern Med Rev*. 2008;13(4):292-306.
4. Suh DH, Park TJ, Jeong JY, et al. Photothermal therapy using gold nanoparticles for acne in Asian patients: a preliminary study. *Dermatol Ther*. 2021;34(3):e14918.
5. Chang C. Short-course therapy for diarrhea-predominant irritable bowel syndrome: understanding the mechanism, impact on gut microbiota, and safety and tolerability of rifaximin. *Clin Exp Gastroenterol*. 2018;11:335-345.
6. Paithankar DY, Sakamoto FH, Farinelli WA, et al. Acne treatment based on selective photothermolysis of sebaceous follicles with topically delivered light-absorbing gold microparticles. *J Invest Dermatol*. 2015;135(7):1727-1734.
7. Yan HM, Zhao HJ, Guo DY, Zhu PQ, Zhang CL, Jiang W. Gut microbiota alterations in moderate to severe acne vulgaris patients. *J Dermatol*. 2018;45(10):1166-1171.
8. Brandsma E, Kloosterhuis NJ, Koster M, et al. A Proinflammatory gut microbiota increases systemic inflammation and accelerates atherosclerosis. *Circ Res*. 2019;124(1):94-100.

How to cite this article: Seo J, Roh HJ, Jung JY. Gut microbiota modulation and gold nanoparticle-mediated photothermal therapy for treatment of recalcitrant acne. *Clin Case Rep*. 2022;10:e05642. doi:[10.1002/ccr3.5642](https://doi.org/10.1002/ccr3.5642)