



De Novo Assembly and Annotation of the Complete Genome Sequence of *Myxococcus xanthus* DZ2

 Rodolfo Aramayo,^a  Beiyan Nan^a

^aDepartment of Biology, Texas A&M University, College Station, Texas, USA

ABSTRACT We report the assembly and annotation of a high-quality genome sequence for *Myxococcus xanthus* strain DZ2 (GenBank accession number [CP080538](#)), created using a combination of short reads generated using DNBSEQ technology (BGI Genomics) and long high-fidelity (HiFi) reads generated using Pacific Biosciences (PacBio) technology.

The *Myxococcus xanthus* isolate (GenBank accession number [CP080538](#)) reported here was originally acquired by David Zusman from the Roger Stanier collection at UC Berkeley and named DZ2 (1). While this work was in progress, Jain et al. (2) reported the assembly of a *Myxococcus xanthus* DZ2 isolate of similar origin ([CP070500](#)). While the assembly submitted under GenBank accession number [CP070500](#) represents a big improvement over the previously reported draft DZ2 assembly (3), the assembly reported here ([CP080538](#)) is both larger in size and different in gene content (Table 1).

Cells of *Myxococcus xanthus* strain DZ2, derived from frozen stock from the Roger Stanier collection at UC Berkeley, were grown in liquid CYE medium (10 g/L Casitone and 5 g/L yeast extract), 8 mM MgSO₄, and 10 mM 3-(*N*-morpholino)propanesulfonic acid (MOPS), pH 7.6, at 32°C, harvested by centrifugation, and frozen in liquid nitrogen. The pellet was ground into a fine powder, and DNA was extracted by lysing the cells with cetyltrimethylammonium bromide (CTAB) at 65°C. The DNA was purified using phenol/chloroform/isoamyl alcohol, followed by ethanol precipitation. DNA sequencing was performed by BGI Genomics using two different technologies: DNBSEQ and high-fidelity (HiFi) PacBio sequencing. Construction of the DNA libraries, DNA sequencing, and quality control (QC) of the long HiFi reads and the short reads derived from PCR-free rolling circle replication of the DNA nanoballs were all performed by BGI Genomics using their standard operating procedures (SOP) (4).

Sequencing generated a total of 436,127 HiFi PacBio subreads with an average length of 9,018 bp, totaling 3,933.4 Mbp (representing 420-fold genome coverage), and ~10.2 million 100-bp paired-end DNBSEQ reads, totaling 2,042.54 Mbp (representing 218-fold coverage). Small reads that passed our standard quality control (QC) protocol (5) and large reads with a quality value (QV) of >20 or >99% accuracy were used for assembly. Genome assembly was performed using HiCanu version 2.1.1-Java-1.8 (6), Unicycler version 0.4.8 (7), and Velvet version 1.2.10 (8, 9). We performed a total of 48 assemblies (36 HiCanu, 11 Unicycler, and 1 Velvet). The parameters used and the associated resulting data are described in a supplemental Zenodo repository (10). The largest circular contig of the HiCanu assemblies (contig [tig00000005](#) of Assembly 33), was ~23.8 kbp larger than the largest Unicycler assembly and was thus selected for analysis. We then used PSI-CD-HIT version 4.8.1/blastn version 2.12.0+ (11–14) to verify that the smaller homologous contigs generated by the other assemblies were contained within this largest HiCanu assembly (10). Given that all the smaller homologous contigs were indeed contained within the largest HiCanu contig ([tig00000005](#)), this contig was then declared as our genome assembly and submitted to the NCBI databases (10). Genome annotation was performed using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) version 5.2 (15). The [CP080538](#) assembly presented here differs from the [CP070500](#) assembly in many ways. The [CP080538](#) assembly is 6.4 kbp larger than the

Editor Frank J. Stewart, Montana State University

Copyright © 2022 Aramayo and Nan. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Rodolfo Aramayo, aramayo@tamu.edu.

The authors declare no conflict of interest.

Received 18 November 2021

Accepted 23 March 2022

Published 6 April 2022

TABLE 1 Comparative analysis of the highly related *Myxococcus xanthus* strain DZ2 assemblies

Characteristic ^a	Data for assembly (GenBank accession no.):	
	CP080538	CP070500
Genome size (bp)	9,365,783	9,359,382
Total no. of genes	7,581	7,576
Total no. of CDSs	7,499	7,494
No. of genes (coding)	7,402	7,408
No. of CDSs (with protein)	7,402	7,408
No. of RNA genes	82	82
No. of complete rRNAs (5S, 16S, 23S)	4, 4, 4	4, 4, 4
No. of tRNAs	66	66
No. of ncRNAs	4	4
No. of CDSs (without protein)	97	86
No. of pseudogenes:		
Total	97	86
With ambiguous residues	0	0
Frameshifted	49	37
Incomplete	52	56
With an internal stop	14	12
With multiple problems	15	16
No. of CRISPR arrays	4	4

^a CDS, coding DNA sequences; ncRNAs, noncoding RNAs; rRNAs, ribosomal RNAs; tRNAs, transfer RNAs.

CP070500 assembly, and while it has five more genes and 11 more pseudogenes, it also has six fewer coding sequences (CDSs). Both assemblies have the same number of rRNAs, tRNAs, ncRNAs, and CRISPR arrays (Table 1).

Data availability. The raw data associated with this publication have been deposited at NCBI under BioProject accession number [PRJNA748417](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA748417) and SRA accession numbers [SRX11508707](https://www.ncbi.nlm.nih.gov/sra/SRX11508707) (PacBio reads) and [SRX11508706](https://www.ncbi.nlm.nih.gov/sra/SRX11508706) (Illumina reads). The complete genome sequence has been deposited in GenBank under accession number [CP080538.1](https://www.ncbi.nlm.nih.gov/genbank/CP080538.1). The different genome assembly commands and resulting assembly files generated throughout this work have all been deposited at Zenodo (10).

ACKNOWLEDGMENTS

Funding was provided by the National Institute of General Medical Sciences (grant numbers R01GM58770 and R01GM123139 to R.A. and R01GM129000 to B.N.). This work was in part performed using the supercomputers Ada and Grace, managed and maintained by the Texas A&M University High Performance Research Computing Resources Center (<http://hprc.tamu.edu>).

We thank Michael Dickens for his invaluable help and support during the realization of this work and Tatiana Aramayo for her invaluable help editing the manuscript. We also thank Autumn McManis and Elias Topo for chromosome purification and PCR verification.

REFERENCES

- Blackhart BD, Zusman DR. 1985. "Frizzy" genes of *Myxococcus xanthus* are involved in control of frequency of reversal of gliding motility. *Proc Natl Acad Sci U S A* 82:8767–8770. <https://doi.org/10.1073/pnas.82.24.8767>.
- Jain R, Habermann BH, Mignot T. 2021. Complete genome assembly of *Myxococcus xanthus* strain DZ2 using long high-fidelity (HiFi) reads generated with PacBio technology. *Microbiol Resour Announc* 10:e0053021. <https://doi.org/10.1128/MRA.00530-21>.
- Müller S, Willett JW, Bahr SM, Darnell CL, Hummels KR, Dong CK, Vlamakis HC, Kirby JR. 2013. Draft genome sequence of *Myxococcus xanthus* wild-type strain DZ2, a model organism for predation and development. *Genome Announc* 1:e00217-13. <https://doi.org/10.1128/genomeA.00217-13>.
- BGI Genomics. DNBSQ de novo service overview. <https://www.flipsnack.com/BDEEDDF8D6/dnbsq-de-novo-service-overview-2021-10/full-view.html>.
- Maitra N, He C, Blank HM, Tsuchiya M, Schilling B, Kaeberlein M, Aramayo R, Kennedy BK, Polymenis M. 2020. Translational control of one-carbon metabolism underpins ribosomal protein phenotypes in cell division and longevity. *Elife* 9:e53127. <https://doi.org/10.7554/eLife.53127>.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res* 27:722–736. <https://doi.org/10.1101/gr.215087.116>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Zerbino DR, McEwen GK, Margulies EH, Birney E. 2009. Pebble and rock band: heuristic resolution of repeats and scaffolding in the Velvet short-read de novo assembler. *PLoS One* 4:e8407. <https://doi.org/10.1371/journal.pone.0008407>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <https://doi.org/10.1101/gr.074492.107>.
- Aramayo R. 2022. *Myxococcus xanthus* DZ2 genome assembly [data set]. Zenodo <https://doi.org/10.5281/zenodo.6359694>.

11. Li W, Godzik A. 2006. CD-HIT: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* 22:1658–1659. <https://doi.org/10.1093/bioinformatics/btl158>.
12. Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* 28:3150–3152. <https://doi.org/10.1093/bioinformatics/bts565>.
13. Li W, Fu L, Niu B, Wu S, Wooley J. 2012. Ultrafast clustering algorithms for metagenomic sequence analysis. *Brief Bioinform* 13:656–668. <https://doi.org/10.1093/bib/bbs035>.
14. Chen Q, Wan Y, Lei Y, Zobel J, Verspoor K. 2016. Evaluation of CD-HIT for constructing non-redundant databases, p 703–706. *In* 2016 IEEE International Conference on Bioinformatics and Biomedicine (BIBM). Institute of Electrical and Electronics Engineers, New York, NY. <https://doi.org/10.1109/bibm.2016.7822604>.
15. Tatusova T, DiCuccio M, Badretdin A, Chetvermin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.