

## Research Article

# Ex Vivo Study of Laban's Role in Decreasing Hemolysis Crisis in G6PD-Deficient Patients

Wissam Zam <sup>1</sup> and Loay Belal<sup>2</sup>

<sup>1</sup>Al-Andalus University for Medical Sciences, Department of Analytical and Food Chemistry, Tartous, Syria

<sup>2</sup>Tartous University, Department of Analytical and Food Chemistry, Tartous, Syria

Correspondence should be addressed to Wissam Zam; [ws.sarah2005@gmail.com](mailto:ws.sarah2005@gmail.com)

Received 20 October 2019; Accepted 8 January 2020; Published 21 January 2020

Academic Editor: C.S. Johnston

Copyright © 2020 Wissam Zam and Loay Belal. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In spite of the vast nutritional and environmental benefits provided by fava bean (*Vicia faba*), the ingestion of vicine/convicine provokes an acute hemolytic anemia called favism in individuals with a glucose-6-phosphate dehydrogenase (G6PD) deficiency. The elimination of these glycosides is a goal that could be accomplished using different processing methods including bacteriological treatment. Laban as a good source of lactic acid bacteria was tested in an *ex vivo* assay on human blood samples in order to determine its capacity in decreasing the hemolysis crisis induced by the ingestion of fava beans. Results indicate a significant decrease in human blood cell hemolysis after the treatment of fava beans by Laban. This decrease in hemolysis was also correlated with the G6PD deficiency categorization. The highest hemolysis level (mean:  $23.11 \pm 0.76\%$ ) was observed in samples with G6PD activity between 10 and 30%, while the lowest hemolysis level (mean:  $5.75 \pm 0.64\%$ ) was observed in samples with G6PD activity more than 60%. This decrease was correlated with a high antioxidant capacity of Laban ( $51.61 \pm 1.13\%$  expressed by the percentage inhibition of DPPH radical). The counts of isolates from MRS and M17 culture plates were  $6.75 \pm 0.095$  and  $7.91 \pm 0.061$  log cfu ml<sup>-1</sup>, respectively. In conclusion, the synergy between the antioxidant properties of Laban and the possible decrease of vicine and convicine concentrations by lactobacillus found in the fermented dairy products could explain the ability of Laban to reduce the hemolysis crisis *ex vivo*.

## 1. Introduction

Fava bean (*Vicia faba*) is a popular crop grown in different climates all around Asia, Europe, and Africa [1]. It is an important source of protein and various nutrients such as phytonutrients, dietary fiber, vitamins, and minerals [2]. In spite of these advantages, fava beans contain some anti-nutritional compounds including two glycosidic aminopyrimidine derivatives, vicine and convicine, which accumulate at maturity and are metabolized into isouramil and divicine upon hydrolysis [3]. These highly redox proteins rapidly oxidize NADPH and glutathione which are normally detoxified by catalase and glutathione peroxidase, in enzymatic reactions that depend on NADPH [4]. Hence, in favism, a disease that results from a deficiency of the enzyme glucose-6-phosphate dehydrogenase (G6PD),

erythrocytes are susceptible to oxidant stress due to poor antioxidant red cell ability and depletion of glutathione peroxidase and catalase [5].

Due to its high nutritional value and importance, several studies have been conducted to reduce the concentration of vicine and convicine from fava beans using different processing methods such as roasting and boiling [6]. Bacteriological treatment was also proved to be one of the most effective methods since  $\beta$ -glucosidase is widespread within lactic acid bacteria, but the level of expression is largely dependent on the strain [1, 7]. Laban is a popular food with a high nutritional value consumed in different areas. It is prepared by the fermentation of milk by lactic acid bacteria [8]. It is found to be useful in the treatment of multiple diseases such as obesity, cancer, and diabetes [9]. Bovine milk used in the preparation of Laban contains a wide variety

of compounds with antioxidant activity such as flavonoids and vitamin E [8, 9]. Furthermore, Laban contains lactic acid bacteria that were proved to have a  $\beta$ -glycosidase activity that hydrolyzes the glycoside bond [1, 10].

To our knowledge, this is the first study to investigate, through the *ex vivo* assays on human blood, the effect of Laban in decreasing the hemolysis crisis in patients with G6PD deficiency.

## 2. Materials and Methods

**2.1. Fava Bean Extract (FBE).** Fava beans grown in Tartous, Syria, were collected fresh after maturity and kept at  $-18^{\circ}\text{C}$  until use. The sample was extracted according to Sosulski and Pitz with slight modification [11]. Ground seed samples were extracted in duplicate with 0.1N NaOH for 20 minutes. The pH was adjusted to pH 4.2 by HCl 1N in order to precipitate most of the proteins in their isoelectric pH. Samples were centrifuged for 20 minutes at 10000 rpm, and supernatants were collected for further experiments. A blank was prepared for the spectrophotometric measurement.

**2.2. Fava Beans Treated by Laban (FBL).** Syrian fermented milk (Laban) was prepared using bovine milk one day before use. It was made by boiling the milk for 5 min, permitting it to cool to  $50^{\circ}\text{C}$ , and inoculating it for about 4 hr with 2.5 to 3.0% starter saved from a previous batch.

Ground seed samples of fava beans were incubated with Laban at a ratio of 1 : 2 for 30 minutes. Then, the sample was extracted as described above. A blank was also prepared for the spectrophotometric measurement.

**2.3. Human Blood Hemolysis.** This study was designed to evaluate the *in vitro* effect of Laban on the blood hemolysis caused by fava beans in children aged between 5 and 12. Before drawing the blood samples, parents of all participants were counseled and those who consented (written informed consent) to their children being part of the study were recruited.

Blood samples from children were drawn on EDTA tubes and used for the screening of the subjects for G6PD deficiency using fluorescent spot test (FST) [12]. Children with red blood cell G6PD value of  $<6.40\text{ U/gHb}$  were regarded as deficient and were recruited.

The determination of the human blood hemolysis was performed with appropriate local health regulations and ethical approval as described by Rizzello et al. [3]. Fully dissolved blood samples were used to determine maximum degeneration using purified water. 0.5 ml of each blood sample was incubated with 2.5 ml from each of FBE, FBL, and blanks for 5 min and then centrifuged to separate the cells from the plasma. An aliquot of plasma was diluted with Drabkin's reagent (leading to the conversion of hemoglobin to cyanmethemoglobin), and the OD540 was measured. Three samples for each extract, obtained by independent experiments, were twice analyzed, and the means of the data were statistically treated (as described below) to assess the significant differences against the control values.

**2.4. Determination of Antioxidant Activity (DPPH Assay).** The antioxidant activity was evaluated by the DPPH radical scavenging activity [13, 14]. DPPH radical solution (0.002%, w/v) in methanol was prepared, and a volume of 1800  $\mu\text{L}$  was added to 200  $\mu\text{L}$  of the sample diluted with phosphate buffer (0.1 M), well vortexed, and incubated for 60 min in dark room at room temperature. The absorbance of each sample at 520 nm was measured using Shimadzu UV-VIS double-beam spectrophotometer. Methanol was used as a blank, while DPPH solution in methanol served as control. The antioxidant activity was expressed as the percentage inhibition of DPPH radical. The experiments were conducted in triplicate, and the mean values were used.

**2.5. Microbial Enumeration and Isolation.** A  $10^{-1}$  dilution of Laban was made using physiological saline, and the microbial counts were determined according to the pour plate method of Houghtby et al. [15]. Total viable counts were determined using plate count agar (Scharlau Chemie S.A., Barcelona, Spain) incubated at  $32^{\circ}\text{C}$  for 48 h. Counts of lactic acid bacteria (LAB) were determined using de Man Rogosa Sharpe (MRS, Himedia) agar incubated anaerobically at  $35^{\circ}\text{C}$  for 48 h [16], while the count of *Streptococcus* strains was determined using M17 agar (Himedia) incubated anaerobically at  $37^{\circ}\text{C}$  for 48 h. Identification of bacterial colonies from the agar plates was performed by Gram staining, cell morphology, carbohydrate fermentation tests, sensitivity to different salt levels, and catalase reaction [17]. The counts were expressed as log 10 cfu/ml of the product.

## 3. Results and Discussion

The study included 57 children made up of 54 (94.73%) males and 3 (5.26%) females with a mean age of 10.07 years. Male children showed higher G6PD deficiency prevalence rates than females. These results are in accordance with different previous research studies that indicated that hemolysis induced by G6PD deficiency is most common in hemizygous males compared to homozygous females as the mutation would have to occur in both copies of the gene in females to cause the disorder, whereas in males only one abnormal copy of the gene is required for manifestation of the disease [18, 19].

Table 1 shows the distribution of G6PD deficiency using the categorization of the enzyme function into severe deficiency ( $<10\%$  G6PD activity), moderate deficiency (10–30%), mild deficiency (30–60%), and normal activity (60–100%) [20]. Of the 57 recruited children, twenty-one (36.84%) were moderately deficient, while eighteen (31.57%) were mildly deficient.

Red blood cells (RBCs) contain many antioxidant components, and the G6PD enzyme is essential for maintaining the NADPH supply, which is the key source of reducing equivalents such as glutathione reductase (GSH) and catalase [21]. Therefore, NADPH production by G6PD is critically important, and when it becomes limited, red cells are prone to being damaged [22]. Both vicine and convicine present in fava beans are converted in the gut into divicine

TABLE 1: Prevalence of G6PD deficiency levels and hemolysis of human blood cells treated with FBE and FBL.

G6PD deficiency categorization	Prevalence of G6PD deficiency levels Total samples tested 57 (100%)	Lysis (%) of human blood cells treated with FBE	Lysis (%) of human blood cells treated with FBL
<10% (<0.64 U/gHb)	0	—	—
10–30% (0.64–1.92 U/gHb)	21 (36.84%)	70.74 ± 1.26%–90.65 ± 2.09% Mean: 82.06 ± 1.63%	19.95 ± 0.95%–26.63 ± 0.68% Mean: 23.11 ± 0.76%
30–60% (1.92–3.84 U/gHb)	18 (31.57%)	67.13 ± 1.65%–76.09 ± 1.96% Mean: 70.61 ± 1.36%	9.68 ± 1.35%–21.48 ± 0.96% Mean: 14.13 ± 1.05%
>60% (>3.84 U/gHb)	18 (31.57%)	35.80 ± 2.06%–47.85 ± 1.05% Mean: 44.06 ± 1.49%	1.21 ± 0.23%–9.71 ± 0.68% Mean: 5.75 ± 0.64%

and isouramil which are highly redox proteins identified as the main factors of favism [23]. These molecules produce reactive oxygen species (ROS) including the superoxide anion and hydrogen peroxide, which rapidly oxidize NADPH and glutathione. Individuals affected by G6PD deficiency are unable to regenerate reduced glutathione and are undefended against oxidative stress [21]. Acute hemolysis caused by fava bean ingestion is described as being the best-studied natural model of oxidative damage [21]. As presented in Table 1, the incubation of blood samples with fava beans extract (FBE) caused an oxidative RBC damage and hemolysis. The percentage of this hemolysis is correlated with the G6PD deficiency levels as noticed in Table 1. Getachew et al. previously demonstrated that fava bean aglycones cause an irreversible rapid GSH depletion with a simultaneous increase in oxidized glutathione (GSSG) production in G6PD enzyme defective human RBCs [24]. This was correlated with progressive accumulation of denatured hemoglobin products into high molecular weight (HMW) proteins and the formation of both band 3 membrane proteins and hemichromes as HMW protein aggregates [23, 24].

In the most severe type of favism, therapeutic measurements include splenectomy, blood transfusion, food supplements such as folic acid, and antioxidants including vitamin E and selenium [25, 26]. However, favism incidence can be reduced by the decrease of both vicine and convicine from fava beans. Different processing methods such as heat treatment and continuous flow soaking have been studied and proved to achieve a reduction in the content of both glycosides [6, 27].  $\beta$ -Glycosidase from *Aspergillus oryzae*, *Fusarium graminearum*, and lactic acid bacteria was used in bioprocessing methods for a selective destruction of the pyrimidine glycosides [1].

In the condition of our study, a significant decrease of human blood cell hemolysis was observed after the treatment of fava beans by Laban (Table 1). This decrease in hemolysis was also correlated with the G6PD deficiency categorization. The highest hemolysis level (mean: 23.11 ± 0.76%) was observed in samples with G6PD activity between 10 and 30%, while the lowest hemolysis level (mean: 5.75 ± 0.64%) was observed in samples with G6PD activity more than 60%. In this sense and as Laban mechanism of action has yet to be determined, the antioxidant properties of Laban could partially explain its significant effect ( $P < 0.05$ ) in reducing the hemolysis induced by fava beans (Figure 1). DPPH free




FIGURE 1: Hemolysis of human blood cells treated with fava bean extract (FBE) and fava beans treated by Laban (FBL).

radical scavenging assay is widely common to investigate the antioxidant capacity of natural compounds [28]. Results proved that bovine fermented milk used in this study had a high antioxidant capacity of 51.61 ± 1.13% expressed by the percentage inhibition of DPPH radical. Our results are in accordance with Dina Tri et al. [29] and showed a relatively higher value than the value obtained by Fitrotin et al. [30] in the fermentation of sesame milk using *L. plantarum* Dad 13. This antioxidant capacity is mainly due to peptides released upon proteolysis during the fermentation period and the wide range of metabolic compounds formed by lactic acid bacteria which may act as electron donors reacting with free radicals to form more stable products [31, 32]. Additionally, reductones formed during fermentation could react with free radicals to stabilize and terminate radical chain reactions [33]. This could explain different results previously obtained for the antioxidant properties of fermented milk of different sources which greatly depend on several factors including the food matrix composition, overall peptidic profile, amino acid composition of the individual peptide, and the large differences between species or even strains used in fermentation [34].

It was previously shown that lactobacillus decreases the concentration of vicine and convicine proportionately with

the  $\beta$ -glycosidase enzyme concentration and the period of treatment [3]. The types of LAB usually found in the fermented dairy products are thermophilic and mesophilic strains of *Streptococcus*, *Lactococcus*, and *Lactobacillus* species [35]. In this study, colonies observed on MRS plates were large and irregular in shape. They had a light color with opaque centers. They were Gram-positive bacteria able to produce acid from lactose, glucose, and fructose. As a result, characteristics of these strains suggested that they belong to *Lactobacillus delbrueckii* subsp. *bulgaricus* species according to the criteria given by Teixeira [36]. Colonies observed on M17 plates were smaller and regular. They were Gram-positive and spherical bacteria resistant to NaCl (2%), but sensitive to higher salt concentrations. These results indicated that the isolates belonged to the *Streptococcus thermophilus* species [37]. The counts of isolates from MRS and M17 culture plates were  $6.75 \pm 0.095$  and  $7.91 \pm 0.061$  log cfu ml<sup>-1</sup>, respectively. Similar results of total bacteria count were reported for different traditional dairy products [16, 38–40].

#### 4. Conclusion and Further Prospective

In conclusion, the synergy between the antioxidant properties of Laban approved by the radical antiscavenging activity and the possible decrease of vicine and convicine concentrations by *Lactobacillus* found in the fermented dairy products could explain the ability of Laban to reduce the hemolysis crisis *ex vivo*. This decrease in hemolysis was correlated with the G6PD deficiency categorization.

Moreover, *Lactobacillus* was previously proved to increase the amount of antioxidant compounds through intestine which would have a more protecting effect against the hemolysis crisis. However, both vicine and convicine are converted in the gut into divicine and isouramil which are highly redox proteins identified as the main factors of favism. So, more research studies should be conducted in order to prove the potential synergistic effect between Laban and fava beans *in vivo*.

#### Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

#### Conflicts of Interest

The authors declare that they have no conflicts of interest.

#### Acknowledgments

This work was funded by Al-Andalus University for Medical Sciences and Tartous University.

#### References

[1] R. Coda, L. Melama, C. G. Rizzello et al., "Effect of air classification and fermentation by *Lactobacillus Plantarum* VTT E-133328 on faba bean (*Vicia faba* L.) flour nutritional properties," *International Journal of Food Microbiology*, vol. 193, pp. 34–42, 2015.

[2] D. Jezierny, R. Mosenthin, and E. Bauer, "The use of grain legumes as a protein source in pig nutrition: a review," *Animal Feed Science and Technology*, vol. 157, no. 3-4, pp. 111–128, 2010.

[3] C. G. Rizzello, I. Losito, L. Facchini et al., "Degradation of vicine, convicine and their aglycones during fermentation of faba bean flour," *Scientific Reports*, vol. 6, no. 1, p. 32452, 2016.

[4] O. M. Ighodaro and O. A. Akinloye, "First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): their fundamental role in the entire antioxidant defence grid," *Alexandria Journal of Medicine*, vol. 54, no. 4, pp. 287–293, 2018.

[5] G. Gaetani, M. Rolfo, S. Arena, R. Mangerini, G. Meloni, and A. Ferraris, "Active involvement of catalase during hemolytic crises of favism," *Blood*, vol. 88, no. 3, pp. 1084–1088, 1996.

[6] A. Cardador-Martínez, K. Maya-Ocaña, A. Ortiz-Moreno et al., "Effect of roasting and boiling on the content of vicine, convicine and L-3, 4-dihydroxyphenylalanine in *Vicia faba* L.," *Journal of Food Quality*, vol. 35, no. 6, pp. 419–428, 2012.

[7] C. G. Rizzello, T. Mueller, R. Coda et al., "Synthesis of 2-methoxy benzoquinone and 2, 6-dimethoxybenzoquinone by selected lactic acid bacteria during sourdough fermentation of wheat germ," *Microbial Cell Factories*, vol. 12, no. 1, p. 105, 2013.

[8] V. K. Shiby and H. N. Mishra, "Fermented milks and milk products as functional foods-a review," *Critical Reviews in Food Science and Nutrition*, vol. 53, no. 5, pp. 482–496, 2013.

[9] M. H. Tunick and D. L. Van Hekken, "Dairy products and health: recent insights," *Journal of Agricultural and Food Chemistry*, vol. 63, no. 43, pp. 9381–9388, 2015.

[10] I. T. Khan, M. Nadeem, M. Imran et al., "Antioxidant capacity and fatty acids characterization of heat treated cow and buffalo milk," *Lipids in Health and Disease*, vol. 16, no. 1, p. 163, 2017.

[11] F. W. Sosulski and W. J. Pitz, "Determination of vicine and convicine in faba bean cultivars by gas-liquid chromatography," *Canadian Institute of Food Science and Technology Journal*, vol. 12, pp. 93–97, 1979.

[12] N. LaRue, M. Kahn, M. Murray et al., "Comparison of quantitative and qualitative tests for glucose-6-phosphate dehydrogenase deficiency," *The American Journal of Tropical Medicine and Hygiene*, vol. 91, no. 4, pp. 854–861, 2014.

[13] W. Zam, "Effect of alginate and chitosan edible coating enriched with olive leaves extract on the shelf life of sweet cherries (*Prunus avium* L.)," *Journal of Food Quality*, vol. 2019, Article ID 8192964, 7 pages, 2019.

[14] H.-C. Wu, H.-M. Chen, and C.-Y. Shiau, "Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*)," *Food Research International*, vol. 36, no. 9-10, pp. 949–957, 2003.

[15] A. G. Houghtby, L. J. Maturin, and K. E. Koenig, "Microbiological count methods," in *Standard Methods for the Examination of Dairy Products*, R. T. Marshal, Ed., pp. 213–246, American Public Health Association, Washington, DC, USA, 16th edition, 1992.

[16] M. O. M. Abdalla and S. I. K. Hussain, "Enumeration and identification of microflora in rouib, a sudanese traditional fermented dairy product," *British Journal of Dairy Sciences*, vol. 1, no. 2, pp. 30–33, 2010.

[17] Y. Ren, W. Liu, and H. Zhang, "Identification of coccoidal bacteria in traditional fermented milk products from Mongolia, and the fermentation properties of the predominant species, *Streptococcus thermophilus*," *Korean Journal for Food Science of Animal Resources*, vol. 35, no. 5, pp. 683–691, 2015.

- [18] N. Cohan, M. Karimi, A. H. Khalili, M. H. Falahzadeh, B. Samadi, and R. M. Mahdavi, "The efficacy of a neonatal screening programme in decreasing the hospitalization rate of patients with G6PD deficiency in southern Iran," *Journal of Medical Screening*, vol. 17, no. 2, pp. 66-67, 2010.
- [19] A. Pinna, G. Solinas, C. Masia, A. Zinellu, C. Carru, and A. Carta, "Glucose-6-phosphate dehydrogenase (G6PD) deficiency in nonarteritic anterior ischemic optic neuropathy in a Sardinian population, Italy," *Investigative Ophthalmology & Visual Science*, vol. 49, no. 4, pp. 1328-1232, 2008.
- [20] M. De Niz, A. C. Eziefula, L. Othieno et al., "Tools for mass screening of G6PD deficiency: validation of the WST8/1-methoxy-PMS enzymatic assay in Uganda," *Malaria Journal*, vol. 12, no. 1, p. 210, 2013.
- [21] P. Arese, V. Gallo, A. Pantaleo, and F. Turrini, "Life and death of glucose-6-phosphate dehydrogenase (G6PD) deficient erythrocytes—role of redox stress and band 3 modifications," *Transfusion Medicine and Hemotherapy*, vol. 39, no. 5, pp. 328-334, 2012.
- [22] R. Zwieter, A. J. Verhoeven, and D. Roos, "Inborn defects in the antioxidant systems of human red blood cells," *Free Radical Biology and Medicine*, vol. 67, pp. 377-386, 2014.
- [23] D. C. McMillan, L. J. Bolchoz, and D. J. Jollow, "Favism: effect of divicine on rat erythrocyte sulfhydryl status, hexose monophosphate shunt activity, morphology, and membrane skeletal proteins," *Toxicological Sciences*, vol. 62, no. 2, pp. 353-359, 2001.
- [24] F. Getachew, A. Vandenberg, and J. Smits, "A practical toxicity bioassay for vicine and convicine levels in faba bean (*Vicia faba*)," *Journal of the Science of Food and Agriculture*, vol. 98, no. 13, pp. 5105-5111, 2018.
- [25] B. Darbandi, S. Zarezadeh, Z. A. Roshan, A. Hassanzadeh Rad, and A. Baghersalimi, "The efficacy of vitamin E and folic acid on the acute hemolysis caused by glucose-6 phosphate dehydrogenase," *Iranian Journal of Pediatric Hematology and Oncology*, vol. 7, no. 3, pp. 232-236, 2017.
- [26] I. Isaac, A. Mainasara, O. Erhabor et al., "Glucose-6-phosphate dehydrogenase deficiency among children attending the emergency paediatric unit of Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria," *International Journal of General Medicine*, vol. 6, no. 6, pp. 557-562, 2013.
- [27] J. Jamalain and M. Ghorbani, "Extraction of favism-inducing agents from whole seeds of faba bean (*Vicia faba* L varmajor)," *Journal of the Science of Food and Agriculture*, vol. 85, no. 6, pp. 1055-1060, 2005.
- [28] J.-M. Lü, P. H. Lin, Q. Yao, and C. Chen, "Chemical and molecular mechanisms of antioxidants: experimental approaches and model systems," *Journal of Cellular and Molecular Medicine*, vol. 14, no. 4, pp. 840-860, 2010.
- [29] D. T. Marya, Nurliyani, Widodo, and Sunarti, "Characterization and antioxidant activity of fermented milk produced with a starter combination," *Pakistan Journal of Nutrition*, vol. 16, no. 6, pp. 451-456, 2017.
- [30] U. Fitrotin, T. Utami, P. Hastuti, and U. Santoso, "Antioxidant properties of fermented sesame milk using *Lactobacillus plantarum* Dad 13," *International Research Journal of Biological Sciences*, vol. 4, no. 6, pp. 56-61, 2015.
- [31] C. Raveschot, B. Cudennec, F. Coutte et al., "Production of bioactive peptides by *Lactobacillus* species: from gene to application," *Frontiers in Microbiology*, vol. 9, p. 2354, 2018.
- [32] T. Kullisaar, M. Zilmer, M. Mikelsaar et al., "Two antioxidative lactobacilli strains as promising probiotics," *International Journal of Food Microbiology*, vol. 72, no. 3, pp. 215-224, 2002.
- [33] Y.-C. Wang, R.-C. Yu, and C.-C. Chou, "Antioxidative activities of soymilk fermented with lactic acid bacteria and bifidobacteria," *Food Microbiology*, vol. 23, no. 2, pp. 128-135, 2006.
- [34] K. Han, J. Cao, J. Wang et al., "Effects of lactobacillus helveticus fermentation on the Ca<sup>2+</sup> release and antioxidative properties of sheep bone hydrolysate," *Korean Journal for Food Science of Animal Resources*, vol. 38, no. 6, pp. 1144-1154, 2018.
- [35] O. N. Donkor, A. Henriksson, T. Vasiljevic, and N. P. Shah, "Proteolytic activity of dairy lactic acid bacteria and probiotics as determinant of viability and *in vitro* angiotensin-converting enzyme inhibitory activity in fermented milk," *Le Lait*, vol. 87, no. 6, pp. 21-38, 2007.
- [36] P. C. M. Teixeira, "Lactobacillus: *Lactobacillus bulgaricus*," in *Encyclopedia of Food Microbiology*, R. K. Robinson, C. A. Batt, and P. D. Patel, Eds., pp. 1136-1144, Academic Press, London, UK, 2000.
- [37] G. Zirnstein and R. Hutkins, "Streptococcus: *Streptococcus thermophilus*," in *Encyclopedia of Food Microbiology*, R. K. Robinson, C. A. Batt, and P. D. Patel, Eds., pp. 2127-2133, Academic Press, London, UK, 2000.
- [38] O. Samet-Bali, I. Felfoul, R. Lajnaf, H. Attia, and M. A. Ayadi, "Enumeration and identification of microflora in "Leben", a traditional Tunisian dairy beverage," *International Food Research Journal*, vol. 24, no. 3, pp. 927-932, 2017.
- [39] J. Ranasinghe and W. Perera, "Prevalence of Lactobacillus bulgaricus and *Streptococcus Thermophilus* stability in commercially available yogurts in Sri Lanka," *Asian Journal of Medical Sciences*, vol. 7, no. 5, pp. 97-101, 2016.
- [40] G. Chammas, R. Saliba, G. Corrieu, and C. Beal, "Characterisation of lactic acid bacteria isolated from fermented milk "Laban", " *International Journal of Food Microbiology*, vol. 110, no. 1, pp. 52-61, 2006.