

CASE REPORT

Open Access

Should we investigate mitochondrial disorders in progressive adult-onset undetermined ataxias?



José Luiz Pedroso^{1*}, Wladimir Bocca Vieira de Rezende Pinto², Orlando Graziani Povoas Barsottini¹ and Acary Souza Bulle Oliveira²

Abstract

Background: Despite the broad development of next-generation sequencing approaches recently, such as whole-exome sequencing, diagnostic workup of adult-onset progressive cerebellar ataxias without remarkable family history and with negative genetic panel testing for SCAs remains a complex and expensive clinical challenge.

Case presentation: In this article, we report a Brazilian man with adult-onset slowly progressive pure cerebellar ataxia, which developed neuropathy and hearing loss after fifteen years of ataxia onset, in which a primary mitochondrial DNA defect (MERRF syndrome - myoclonus epilepsy with ragged-red fibers) was confirmed through muscle biopsy evaluation and whole-exome sequencing.

Conclusions: Mitochondrial disorders are a clinically and genetically complex and heterogenous group of metabolic diseases, resulting from pathogenic variants in the mitochondrial DNA or nuclear DNA. In our case, a correlation with histopathological changes identified on muscle biopsy helped to clarify the definitive diagnosis. Moreover, in neurodegenerative and neurogenetic disorders, some symptoms may be evinced later during disease course. We suggest that late-onset and adult pure undetermined ataxias should be considered and investigated for mitochondrial disorders, particularly MERRF syndrome and other primary mitochondrial DNA defects, together with other more commonly known nuclear genes.

Keywords: Ataxia, MERRF syndrome, Mitochondrial diseases, Cerebellar ataxia

Dear Editor,

According to its etiological basis, hereditary ataxias are classified into six major groups: autosomal dominant spinocerebellar ataxias (SCA), autosomal recessive, congenital, mitochondrial, episodic and X-linked cerebellar ataxias [1, 2]. Despite the broad development of next-generation sequencing approaches recently, such as whole-exome sequencing (WES), diagnostic workup of adult-onset progressive cerebellar ataxias without remarkable family history and with

negative genetic panel testing for SCAs remains a complex and expensive clinical challenge [1–3]. In this article, we report a Brazilian man with adult-onset slowly progressive pure cerebellar ataxia, which developed neuropathy and hearing loss after fifteen years of ataxia onset, in which a primary mitochondrial DNA (mtDNA) defect was confirmed through muscle biopsy evaluation and WES.

A 66-year-old man presented with slow progressive ataxia that started 20 years before. When he was 46-year-old, mild loss of balance started. Parents were non-consanguineous. Family history was unremarkable. Examination disclosed moderate to severe cerebellar ataxia, dysmetria and dysarthria. An extensive investigation, during

* Correspondence: jpgedroso.neuro@gmail.com

¹Ataxia Unit, Department of Neurology and Neurosurgery, Universidade Federal de São Paulo (UNIFESP), Pedro de Toledo Street, 650. ZIP CODE: 04039-002. Vila Clementino, São Paulo, SP, Brazil

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

the first 15 years of disease onset, resulted negative. SCA genetic panel, Friedreich ataxia, autoimmune disorders (GAD, thyroid antibodies, celiac disease), paraneoplastic panel, vitamin levels, sensory and motor neuroconduction studies and needle electromyography (EMG) were normal. Basic metabolic work-up with plasma lactate, ammonia and lactate/pyruvate ratio was unremarkable. Brain magnetic resonance imaging (MRI) disclosed global cerebellar atrophy (Fig. 1). Gene panel testing for cerebellar ataxias including *SYNE1*, *SPG7* and *SACS* genes resulted negative. A first WES testing was inconclusive.

After fifteen years, the patient developed mild bilateral sensorineural hearing loss. Examination showed cerebellar ataxia, decreased deep tendon reflexes and distal weakness in lower limbs (Supplemental Video). Neuroconduction studies disclosed axonal sensorimotor polyneuropathy. The association of cerebellar ataxia with sensorineural deafness and axonal neuropathy leads to clinical suspicion of primary mtDNA defects. A second WES testing including coverage of the entire mtDNA was requested, and the A8344G pathogenic variant in the *MT-TK* gene was identified. Muscle biopsy disclosed increased subsarcolemmal mitochondrial proliferation with ragged-red fibers (Fig. 1). MERRF (myoclonus epilepsy with ragged-red fibers) genetic spectrum presenting with pure adult-onset cerebellar ataxia was diagnosed.

Mitochondrial disorders are a clinically and genetically complex and heterogenous group of metabolic diseases, resulting from pathogenic variants in the mtDNA or nuclear DNA [4, 5]. Furthermore, mitochondrial disorders are characterized by systemic involvement, and common

nervous system symptoms include seizures, cerebellar ataxia, neuropathy, encephalopathy, stroke-like episodes, visual loss and deafness [6]. Although most forms start in childhood, adult-onset is not uncommon, especially when associated with other multisystemic and neurological findings [4, 5]. It is quite unusual that mitochondrial disorders present with late-onset progressive ataxia as an isolated syndrome [7]. The most common syndromic mitochondrial diseases presenting with cerebellar ataxia as a cardinal sign include: (i) Kearns-Sayre syndrome; (ii) NARP (neuropathy, ataxia and retinitis pigmentosa) syndrome; (iii) *POLG* gene spectrum disorders, highlighting the mitochondrial recessive ataxia syndrome (MIRAS), including SANDO (sensory ataxic neuropathy, dysarthria and ophthalmoplegia) and Mitochondrial Spinocerebellar ataxia with epilepsy syndrome (MSCAE); (iv) MELAS syndrome (mitochondrial encephalopathy, lactic acidosis and stroke-like episodes); and (v) primary coenzyme Q10 deficiency [4–7].

Although MERRF may present with ataxia, typical clinical features include myoclonus, seizures, myopathy and variable degrees of cognitive impairment, visual loss, deafness and neuropathy [7, 8]. Hardly ever MERRF may present with pure ataxia, particularly in adult-onset phenotypes. In a large series of patients with progressive ataxia, 9% of the patients presented with muscle biopsy suggestive of mitochondrial disorder [7, 8]. In our case, the rise of hearing loss and neuropathy guided for a mitochondrial disorder. Next-generation sequencing associated with muscle biopsy confirmed the diagnosis of MERRF.




Fig. 1 (a) Axial T2-weighted, (b) axial FLAIR and (c) sagittal T2-weighted brain MR imaging showing global cerebellar atrophy (white arrows) with normal brainstem volume. Muscle biopsy disclosed increased subsarcolemmal mitochondrial proliferation with typical ragged-red fibers (RRF) in modified Gömöri Trichrome stain (d; white arrows) and ragged-blue fibers in modified succinate-dehydrogenase (SDH) reaction (E; black arrows)

This is a very instructive case of a patient with late-onset undetermined pure cerebellar ataxia that only lately developed hearing loss and neuropathy, and was diagnosed through genetics and muscle biopsy within the expanding clinical spectrum of MERRF syndrome. Despite the rare descriptions of mtDNA point mutations leading to pure cerebellar ataxia phenotypes, late-onset mtDNA defect cases most commonly remain undiagnosed [2, 8, 9]. With the greater availability and current use of WES in clinical practice, unusual genetic causes of ataxias may be identified and need to be properly correlated with clinical findings. However, on the other hand, for atypical late-onset pure presentations, interpretation of genetic findings on WES may become sometimes a problem rather than diagnostic solving [10–12].

In our case, a correlation with histopathological changes identified on muscle biopsy helped to clarify the definitive diagnosis. Moreover, in neurodegenerative and neurogenetic disorders, some symptoms may be evinced later during disease course. In conclusion, we suggest that late-onset and adult pure undetermined ataxias should be considered and investigated for mitochondrial disorders, particularly MERRF syndrome and other primary mtDNA defects, together with other more commonly known nuclear genes. Neuropathy and hearing loss during disease course may also aid in the proper evaluation of a suspected mitochondrial disorder.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s40673-020-00122-0>.

Additional file 1 Supplemental video. Patient with MERRF (myoclonus epilepsy with ragged-red fibers) presenting with gait ataxia, lower limb weakness related to neuropathy and dysmetria.

Abbreviations

SCA: Spinocerebellar ataxias; WES: Whole-exome sequencing; mtDNA: Mitochondrial DNA; EMG: Electromyography; MRI: Magnetic resonance imaging; MERRF: Myoclonus epilepsy with ragged-red fibers; NARP: Neuropathy, ataxia and retinitis pigmentosa; MIRAS: Mitochondrial recessive ataxia syndrome; SANDO: Sensory ataxic neuropathy, dysarthria and ophthalmoplegia; MELAS: Mitochondrial encephalopathy, lactic acidosis and stroke-like episodes

Submission declaration

This manuscript has not been published previously, is not under consideration for publication elsewhere, is approved in its final form by all authors and will not be published elsewhere in the same form.

Authors' contributions

All authors have participated in the research and article preparation. Conception and design of the study: A. Conception and Design, B. Acquisition of data, C. Analysis and interpretation of data; Drafting the article and revision: A. Writing of the first draft; B. Review and Critique. Final approval of the version. Pedroso JL: 1A, 1B, 1C, 2A, 2B, 3 (Nothing to disclose). Pinto WBVR: 1A, 1B, 1C, 2A, 2B, 3 (Nothing to disclose). Barsottini OGP: 1A, 1B, 2B, 3 (Nothing to disclose). Oliveira ASB: 1A, 1B, 1C, 2A, 2B, 3 (Nothing to disclose).

Funding

The authors have nothing to disclose.

Ethics approval and consent to participate

Full consent was obtained from the patient for the case report. All authors approved the final form of this manuscript and take full responsibility for the data.

Competing interests

Dr. Pedroso (jlpedroso.neuro@gmail.com) reports no conflicts of interest. Dr. Pinto (wladimirbvpinto@gmail.com) reports no conflicts of interest. Dr. Barsottini (orlandobarsottini@gmail.com) reports no conflicts of interest. Dr. Oliveira (acary.bulle@unifesp.br) reports no conflicts of interest.

Author details

¹Ataxia Unit, Department of Neurology and Neurosurgery, Universidade Federal de São Paulo (UNIFESP), Pedro de Toledo Street, 650. ZIP CODE: 04039-002. Vila Clementino, São Paulo, SP, Brazil. ²Division of Neuromuscular Diseases, Department of Neurology and Neurosurgery, Universidade Federal de São Paulo (UNIFESP), São Paulo, SP, Brazil.

Received: 11 August 2020 Accepted: 13 August 2020

Published online: 24 August 2020

References

1. Barsottini OG, Albuquerque MV, Braga-Neto P, Pedroso JL. Adult onset sporadic ataxias: a diagnostic challenge. *Arq Neuropsiquiatr*. 2014;72(3):232–40.
2. Sailer A, Houlden H. Recent advances in the genetics of cerebellar ataxias. *Curr Neurol Neurosci Rep*. 2012;12(3):227–36.
3. Klockgether T. Sporadic adult-onset ataxia. *Handb Clin Neurol*. 2018;155: 217–25.
4. Vernon HJ, Bindoff LA. Mitochondrial ataxias. *Handb Clin Neurol*. 2018;155: 129–41.
5. Molnar MJ, Kovacs GG. Mitochondrial diseases. *Handb Clin Neurol*. 2017;145: 147–55.
6. Sailer M, Matilla-Dueñas A, Soong BW, Pedroso JL, Barsottini OG, Mitoma H, et al. The classification of autosomal recessive cerebellar ataxias: a consensus statement from the society for research on the cerebellum and ataxias task force. *Cerebellum*. 2019;18(6):1098–125.
7. Bargiela D, Shanmugarajah P, Lo C, Blakely EL, Taylor RW, Horvath R, et al. Mitochondrial pathology in progressive cerebellar ataxia. *Cerebellum Ataxias*. 2015;2:16.
8. Chinnery PF. Mitochondrial disorders overview. 2000 Jun 8 [updated 2014 Aug 14]. In: Adam MP, Ardinger HH, Pagon RA, Wallace SE, LJM B, Stephens K, Amemiya A, editors. *GeneReviews*® [internet]. Seattle: University of Washington, Seattle; 1993–2020.
9. Rahman S, Copeland WC. POLG-related disorders and their neurological manifestations. *Nat Rev Neurol*. 2019;15(1):40–52.
10. Lieber DS, Calvo SE, Shanahan K, Slate NG, Liu S, Hershman SG, et al. Targeted exome sequencing of suspected mitochondrial disorders. *Neurology*. 2013;80(19):1762–70.
11. Németh AH, Kwasniewska AC, Lise S, Schnekenberg RP, Becker EBE, Bera KD, et al. Next generation sequencing for molecular diagnosis of neurological disorders using ataxias as a model. *Brain*. 2013;136:3106–18.
12. Cortese A, Simone R, Sullivan R, et al. Biallelic expansion of an intronic repeat in RFC1 is a common cause of late-onset ataxia. *Nat Genet*. 2019;51: 649–58.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.