

# Bcl-2 Inhibits Retinoic Acid–induced Apoptosis during the Neural Differentiation of Embryonal Stem Cells

Hitoshi Okazawa,\* Jun Shimizu,\* Masaki Kamei,\* Ichiro Imafuku,\* Hiroshi Hamada,‡ and Ichiro Kanazawa\*

\*Department of Neurology, Faculty of Medicine, University of Tokyo, Tokyo, Japan; and ‡Department of Developmental Biology and Cancer Prevention, Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan

**Abstract.** We report here that all *trans*-retinoic acid (RA), a classical morphogen, induces apoptosis during the neural differentiation of the embryonic stem cell line P19. The apoptotic cells showed, in addition to DNA cleavage, typical morphological changes including chromatin condensation, nuclear fragmentation, and cytoplasmic vacuolation. These apoptotic changes became obvious by 12 h after the addition of RA. The endogenous expression of *bcl-2* in surviving cells was down-regulated during this process, and the compelled expression of *bcl-2* by retroviral vectors reduced the number of apoptotic cells. Apoptosis was partially inhibited by adding antisense oligonucleotides against RA receptors (RARs) simultaneously or by transfecting a plasmid vector flanked with a RA-responsive element. Antisense oligonucleotides against retinoid X

receptors (RXRs), the receptors for 9 *cis*-RA, did not inhibit apoptosis induced by all *trans*-RA. Cycloheximide and actinomycin D, inhibitors of protein and RNA syntheses, respectively, suppressed apoptosis. No changes were seen in the expression of tumor necrosis factors, their receptors, Fas, FasL, p53, or *c-myc*, molecules which have been suggested to participate in the apoptotic process. Addition of neurotrophins to the culture medium did not affect apoptosis. These findings suggest that the signals mediated by RARs, the differentiation signals themselves, promote expression of molecules essential for apoptosis. Furthermore, we observed that RA induced apoptosis of cerebral neurons from murine embryos in primary culture, which suggests that RA might participate in cell death which occurs during neural development.

**A**POPTOSIS is a biological process which causes programmed cell death during development and may participate in various pathological processes. It is characterized by typical morphological changes including chromatin condensation, nuclear fragmentation, cytoplasmic vacuolation, membrane blebbing and cell shrinkage, as well as by the biochemical marker DNA cleavage. Apoptosis eliminates 20–80% of neurons during development (Hamburger and Oppenheim, 1982; Oppenheim, 1991) but leaves neurons which are indispensable for the function of the nervous system. This process seems essential because unnecessary neurons might disturb the development of neural networks. The apoptotic process is also observed under conditions stressful for neurons such as serum/neurotrophic factor deprivation (Greene and Tischer,

1976; Greene, 1978; Batistatou and Greene, 1991) or exposure to neurotoxins (Dispasquale et al., 1991; Edwards et al., 1991) including toxic levels of glutamate (Kure et al., 1991). Apoptosis has been postulated as a model for neuronal death in human neurodegenerative diseases because of morphological similarities at the cellular levels.

Apoptosis is regulated by the balance of various molecules including tumor suppressor genes, protooncogenes, and cell cycle regulators. Among the tumor suppressor genes, Rb and p53 are known to participate in apoptosis. Inhibition of Rb function by gene disruption (Morgensbesser et al., 1994) or by expression of human papilloma virus protein E7 (Pan and Griep, 1994) interferes with terminal differentiation of lens fiber cells, causes apoptosis of those cells, and results in microphthalmia. In this process, Rb acts as an inhibitor of DNA synthesis after withdrawal from the cell cycle and thereby prevent apoptosis. On the other hand, disruption of the p53 gene in mice reduces apoptotic cell death (Clarke et al., 1993; Lowe et al., 1993). Over-expression of p53 causes growth arrest and apoptosis (Yonish-Rouach et al., 1991), and p53 mutations are widely observed in various tumors (reviewed by Culotta and Koshland, 1993), suggesting that p53 mediates the apoptotic process to suppress the progression of tumors.

Address all correspondence to H. Okazawa, Department of Neurology, Faculty of Medicine, University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo, Japan. Tel.: (81) 3-3815-5411. Fax: (81) 3-3813-2129.

1. *Abbreviations used in this paper:* BMP, bone morphogenic protein; CRBP, cellular retinoid binding proteins; LTR, long terminal repeat; NEL, nick end labeling; pRARE-CAT, CAT reporter plasmid containing RA response elements; RA, retinoic acid; RAR, RA receptor; RXR, retinoid X receptor.

p53 acts as a transcription factor which regulates genes affecting apoptosis (Miyashita et al., 1994; Miyashita and Reed, 1995) such as *c-myc* (Askew et al., 1991; Evan et al., 1992; Shi et al., 1992; Bissonnette et al., 1992; Fanidi et al., 1992) and *c-jun* (Estus et al., 1994; Ham et al., 1995).

*bcl-2*, a protooncogene (Cleary and Sklar, 1985; Tsujimoto et al., 1985; Bakhshi et al., 1985), also affects the commitment to apoptosis (reviewed by White, 1994). The Bcl-2 protein forms heterodimers with other members of the *bcl-2* gene family including Bax (Oltvai et al., 1993), Mcl-1 (Kozopos et al., 1993), A1 (Lin et al., 1993), Bcl-x<sub>L</sub>, Bcl-x<sub>S</sub>, (Boise et al., 1993; Sato et al., 1994), Bad (Yang et al., 1995), and Bak (Chittenden et al., 1995; Kiefer et al., 1995). BAG-1, a protein nonhomologous to this family, also binds to Bcl-2 (Takayama et al., 1995). These partners determine whether the heterodimer acts as an inhibitor or promoter of apoptosis. Alterations in levels of reactive oxygen species (Hockenbery et al., 1993; Kane et al., 1993) and intracellular calcium (Bafy et al., 1993) by Bcl-2 have been suggested as possible anti-apoptosis mechanisms, while there is evidence suggesting other mechanisms by which Bcl-2 can inhibit apoptosis (Shimizu et al., 1995). Molecules that determine the cell fate for apoptosis have been discovered, whereas the upstream and downstream apoptotic pathways that link initial stimuli to DNA fragmentation have not yet been elucidated completely (reviewed by Reed, 1994).

To elucidate the specific features of neuronal apoptosis during development, an experimental system with which we can observe physiological stimuli inducing both neural differentiation and apoptosis is required. Deprivation of trophic factors is commonly used for the analysis of neuronal apoptosis (Greene and Tischer, 1976), but this system is not so suitable for the investigation of differentiation. Here, we present an embryonic cell line, P19, as a candidate system for the investigation of these processes. P19 is a widely used and well-documented system for the analysis of neural differentiation, because retinoic acid (RA) efficiently differentiates P19 cells to neurons and glias (Jones-Villeneuve et al., 1983; Rudnicki and McBurney, 1987; Staines et al., 1994). We report here that: (a) RA induces apoptosis of P19 cells during the neural differentiation; (b) *bcl-2* expression is down-regulated during this apoptotic process; (c) apoptosis is inhibited by dysregulated expression of Bcl-2 by retroviral infection; (d) the apoptotic signal is mediated by RA receptors (RARs); (e) de novo induction of some genes by the RAR-mediated signal is necessary for apoptosis; (f) apoptosis is not directly affected by the cytokine systems which have been suggested to participate in cell death; and (g) apoptosis is induced similarly in embryonic neurons by RA. These findings suggest a close relationship between apoptotic signals and morphogenic signals and the possibility that the RA-induced apoptosis of P19 can be used as a model for the analysis of apoptosis during neural differentiation.

## Materials and Methods

### Cell Culture and RA Treatment

P19 cells were maintained in  $\alpha$ -MEM (Sigma Chemical Co., St. Louis, MO) supplemented with 10% fetal bovine serum (ICN Radiochemicals,

Irvine, CA). To induce neural cell differentiation,  $1 \times 10^5$  P19 cells were cultured in 10 cm bacteriological dishes (Nipro) in 10 ml of medium containing 0.1  $\mu$ M of all *trans*-RA (Sigma Chemical Co., St. Louis, MO) for 4 d, transferred to tissue culture dishes (Corning, Corning, NY), and cultured in the absence of RA for 2 d. To induce mesodermal cell differentiation, they were cultured in tissue culture dishes for 2–4 d in the presence of RA. We used  $10^{-7}$  M RA in most experiments because this order of concentration is commonly used to differentiate P19 (Rudnicki and McBurney, 1987) and because it was optimal to observe apoptosis morphologically. However, RA concentrations were changed as described in the text and figure legends when necessary. RA stock solution ( $10^{-4}$  M all *trans*-RA in ethanol) was diluted with the culture medium just before use. Cycloheximide and/or actinomycin-D at concentrations of 1.0  $\mu$ g/ml and 0.1  $\mu$ g/ml, respectively, were added for the inhibition of protein/RNA synthesis simultaneously with RA.

### Ultrastructural Analysis of Apoptotic Cells

During the procedure for neural differentiation, dying cells were separated from cell aggregates. Culture medium was recovered to a centrifuge tube, and left to stand for 5 min. Cell aggregates dropped to the bottom during this time. The supernatants were carefully removed to new centrifuge tubes, and the floating cells were spun down at 200 g for 10 min. The collected cells were prefixed with 2.5% glutaraldehyde in 0.1 M phosphate buffer (pH 7.4), rinsed three times with 0.1 M phosphate buffer, postfixed in 1% OsO<sub>4</sub> in 0.1 M phosphate buffer, and embedded in 1% agarose, dehydrated, and re-embedded in Epon. Semi-thin sections were stained with 0.1% toluidine blue. Ultra-thin sections were contrasted with uranyl acetate and lead citrate.

### Estimation of Cell Death

To calculate the percentage cell death, we directly counted the dead cells in preparations stained by trypan blue. P19 cells ( $1 \times 10^5$ /bacteriological dish in 10 ml of  $\alpha$  medium - 10% FBS) or cultured neurons (the culture conditions were described later in Primary Culture of Neurons) were incubated with  $10^{-7}$  M of RA for 48 h. Thereafter they were scraped off from the bottom of the dishes and collected with the culture medium. An equal volume of 0.4% trypan blue solution (GIBCO BRL, Gaithersburg, MD) was added to the medium and left for 1 min. The numbers of stained and total cells were then counted within 5 min.

### Immunohistochemical Analysis of Differentiated Cells

Immunohistochemistry was performed to determine the characteristics of the differentiated P19 and P19-Bcl-2 stable transformant cells. Antibodies against neurofilament 160K, GFAP, vimentin, GalC, and fibronectin were used for this purpose (Neural cell typing kit; Boehringer-Mannheim Biochemicals, Indianapolis, IN). These cells were fixed with methanol precooled to  $-20^\circ\text{C}$  or 5% (vol/vol) acetic acid in ethanol in the case of staining with the anti-GalC antibody.

### DNA Preparation and Analysis

Genomic DNA was prepared as described previously (Okazawa et al., 1991). Briefly, dying cells were separated as described above and centrifuged at 200 g for 10 min. The pellets were resuspended in a 10-fold volume of RSB solution (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 1.5 mM MgCl<sub>2</sub>), left on ice for 10 min, homogenized in RSB with 0.5% NP-40, centrifuged at 800 g for 5 min. The resulting nuclear pellets were suspended in an equal volume of extraction buffer (50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 100 mM EDTA) containing SDS (final 0.5%) and proteinase K (final 200 ng/ml), incubated at  $37^\circ\text{C}$  for 6 h, extracted with phenol/chloroform (1:1) for more than 12 h, and centrifuged. The aqueous phase was recovered, and dialyzed with TE (10 mM Tris-HCl, pH 7.5, 1 mM Na<sub>2</sub>EDTA) for 1 d. The concentration of recovered DNA samples were checked, and 1- $\mu$ g aliquots of each DNA sample were separated on 3% nucieve/agarose (3:1) gels.

### DNA Fragmentation Assay

This assay was performed according to the reported method (Sellins and Cohen, 1987).  $1 \times 10^6$  cells were harvested by centrifugation at 200 g for 10 min. The pellet was lysed with 0.4 ml hypotonic lysing buffer (10 mM Tris, 1 mM EDTA, pH 7.5) containing 0.2% Triton-X 100, and lysates

were centrifuged at 13,000 g for separate intact from fragmented chromatin. The supernatant, containing fragmented DNA, was placed in a separate microfuge tube, and both pellet and supernatant were precipitated overnight at 4°C in 12.5% trichloroacetic acid. The precipitates were sedimented at 13,000 g for 10 min. The DNA in precipitates was hydrolyzed by heating 90°C for 10 min in 80 µl 5% trichloroacetic acid, and 0.16 ml of DPA reagent (0.15 g diphenylamine, 0.15 ml sulfuric acid, 0.05 ml acetaldehyde in 10 ml glacial acetic acid; Sigma Chemical Co.) was added to each tube. After overnight color development, optical densities of the samples were read at 600 nm. For the analysis of embryonic neurons in primary culture,  $1 \times 10^5$  cells were used for the reaction scaled down to 1/10× vol.

### RNA Preparation and Northern Blotting Analysis

RNA was prepared similarly to the step for separating nuclei. After centrifugation, Tris-HCl, pH 9.0 (final 100 mM), SDS (final 0.5%), and NaCl (final 100 mM), were added to the supernatant followed by extraction with phenol/chloroform, and precipitation with ethanol. Poly-A<sup>+</sup> RNA was separated by oligo-dT sepharose column chromatography (Pharmacia Fine Chemicals, Piscataway, NJ) according to the supplier's protocol. 20-µg aliquots of each RNA sample were separated on 1% agarose gels in 1× MOPS buffer (20 mM 3-[N-morpholino] propane-sulphonic acid, 5 mM sodium acetate, pH 7.0, 1 mM Na<sub>2</sub>EDTA) containing 4% formaldehyde, and blotted onto Hybond N (Amersham Corp., Arlington Heights, IL) with a vacuum blotter (Pharmacia Fine Chemicals). The filters were hybridized in 6× SSC, 5× Denhardt's solution, 1% SDS at 68°C and washed in 0.1× SSC, 0.1% SDS at 65°C. The same filters were then hybridized with *bcl-2*-specific cDNA (from nucleotide 1833 to 2120; Negrini et al., 1987), D<sub>1A</sub> dopamine receptor, and β-actin (Takara) probes, which were radiolabeled with a random primer labeling kit (Boehringer-Mannheim Biochemicals). The filters were then exposed for 7 d to Kodak XAR5 film.

### RT-PCR

For the specific amplification of *bcl-2* mRNA, we synthesized two primers based on the unique sequences of *bcl-2* with no homology to the genes of other family members. The primers used were 5'-AAGAAATTCGG-GAGAACAGGGTATG (from 1832 to 1848 of the mouse *bcl-2* gene, Negrini et al., 1987) and 5'-AAGGATCCGGTAGCGACGAGAGAAG (from 2137 to 2119). The expected size of the amplified product was 304 bp. The amplified products were digested and subcloned into pBluescriptSK<sup>+</sup> between EcoRI and BamHI site. The primers for *bax* were CAGCTCT-GAACAGATCATG and GCCCATCTTCTTCCAGAT which amplified 534 bp (from 43 to 576 of the mouse *bax* gene; Oltvai et al., 1993; GenBank/EMBL/DBJ accession number L22472). The primers for *bad* were CCAGGATCCAAATGGGAA and ATGTGGAGCAGAA-GATCA which amplified 651 bp (from 460 to 1110 of the mouse *bad* gene; Yang et al., 1995; GenBank/EMBL/DBJ accession number L37296). The primers for *bag-1* were CCGACAACATGGCCAAGA and TCCACTGCACTTCAATCA which amplified 679 bp (from 75 to 753 of the mouse *bag-1* gene; Takayama et al., 1995; GenBank/EMBL/DBJ accession number U17162). For the amplification of G3PDH, *c-myc*, p53, TNF-α, TNF-β, TNFR-I, TNFR-II, Fas, and FasL transcripts, commercial primers purchased from Clontech (Palo Alto, CA) were used. Briefly, 200 ng of total RNA, 0.1 µM of both directional primers, 1× RT-PCR buffer consisting of 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.1 mg/ml gelatin, 0.1% Triton X-100, 0.25 mM dNTPs, 5 U RNase inhibitor (Boehringer-Mannheim Biochemicals), 3.2 U AMV-reverse transcriptase XL (Life Science Inc., Gaithersburg, MD), and 2 U *Taq* polymerase (Takara), were mixed in a total volume of 25 µl. The solution was incubated at 41°C for 30 min, denatured at 94°C for 3 min, and then used for PCR. The thermal cycle conditions were as follows: 94°C for 1 min, 55°C for 1 min, and 72°C for 1 min. Preliminary experiments were performed to determine the optimum thermal cycle number within the linear amplification range; 35 and 30 thermal cycles were performed for *bcl-2* transcript and for the other gene transcripts, respectively.

### Retroviral Vector Construction and Production

The human *bcl-2* cDNA was subcloned into the pBC140 retroviral vector at the XhoI-cloning site (Shimizu et al., 1995). This vector expresses the cDNA insert driven by the CMV promoter and the selectable marker neomycin phosphotransferase driven by the leukemia virus long terminal re-

peat (LTR). Virus was generated from the recombinant retroviral vector plasmid pBC140B. pBC140B was transfected by calcium phosphate precipitation into the Ψ2 packaging cell line. The transfected cells were selected with 800 µM G418. A G418-resistant cell line, designated Ψ2/B4, was confirmed to produce viral particles sufficiently. P19 cells were transfected with supernatant from Ψ2/B4, selected with 300 µM of G418 for 2 wk. The expression of Bcl-2 in G418-resistant clones was tested by immunoblot analysis.

### Construction and Transfection of pRARE-CAT


Sense and antisense oligonucleotides corresponding to the RA response element (de-The et al., 1990) were synthesized, phosphorylated at their 5'-end, annealed and subcloned into pBLCAT2 (Luckow and Schutz, 1987) at HindIII sites. The sequences of the two oligonucleotides were 5'-AGCTGGGTAGGGTTCACCGAAAGTTCCTCG (from -59 to -33 of the RAR β gene; de-The et al., 1990) and 5'-AGCTCGAGTGAACTTTCGGTGAACCCTACCC. After the plasmid was checked by sequencing, it was amplified using a Qiagen Maxi kit (QIAGEN, Chatsworth, CA) and transfected into P19 cells by the calcium phosphate method (Okamoto et al., 1990) using a Eukaryotic Transfection kit (Stratagene Corp., La Jolla, CA). The P19 cells were treated with 15% glycerol in 1× BES (25 mM BES [N,N-bis[2-hydroxyethyl]-2-aminoethanesulfonic acid], 140 mM NaCl, 0.75 mM Na<sub>2</sub>HPO<sub>4</sub>·2H<sub>2</sub>O), pH 6.96, 12 h after the addition of plasmids. RA was added to the dishes 24 h after the addition of DNA. Apoptotic cells were collected according to a protocol similar to that described in Estimation of Cell Death and counted by photomicroscopy.

### Inhibition of Apoptosis by Antisense Oligonucleotides

Antisense and sense oligonucleotides were synthesized using an Applied Biosystems (Foster City, OR) nucleotide synthesizer, and purified by gel filtration. The sequences of antisense oligonucleotides were as follows, RARα1: 5'-ACTGCTATTGCTGGCCAT, RARα2: 5'-TTCCACACTCTC-GTACAT, RARβ1/3: 5'-GTGGCTGCTGGTCTCAT, RARβ2: 5'-ATCCATACAGTCAAACAT, RARγ: 5'-CTCCTTATTGGTGGCCAT, retinoid X receptor (RXR)α: 5'-GAAATGTTTGGTGTCCAT, RXRβ: 5'-AGTGGCGGGGGTGGCATT, RXRγ: 5'-GGAATAATTTCCAT-ACAT. These sequences were designed based on the previous reports (Zelent et al., 1989, 1991; Leroy et al., 1991; Mangelsdorf et al., 1992) to overlap the initiation codons of receptors. Sense oligonucleotides were made in the reverse orientations. Each oligonucleotide was added at 10 µM to the RA-containing culture medium and exchanged every 24 h. The optimal concentration was determined by preliminary experiments using various amounts of the oligonucleotides: 0.1, 1, 5, 10, and 20 µM. The effect on apoptosis reached a plateau at 20 µM. Stability of the oligonucleotides was checked before use. 10 µM of each oligonucleotide radiolabeled with <sup>32</sup>P-γ-ATP and polynucleotide kinase (Toyobo, Tokyo, Japan) was incubated in α-MEM medium with 10% FBS at 37°C for 24 h. Degradation was estimated by polyacrylamide gel electrophoresis as described (Akhtar et al., 1991). Percentage of the intact oligonucleotide band was calculated. The mean ± SD made in 4 independent experiments was 32 ± 2% in RARα1 sense, 31 ± 2% in RARα1 antisense, 33 ± 2% in RARα2 sense, 34 ± 2% in RARα2 antisense, 32 ± 3% in RARβ1/3 sense, 30 ± 3% in RARβ1/3 antisense, 34 ± 2% in RARβ2 sense, 30 ± 2% in RARβ2 antisense, 30 ± 2% in RARγ sense, 31 ± 2% in RARγ antisense, 31 ± 2% in RXRα sense, 33 ± 2% in RXRα antisense, 32 ± 2% in RXRβ sense, 33 ± 2% in RXRβ antisense, 30 ± 2% in RXRγ sense, and 34 ± 2% in RXRγ antisense. The stability was not statistically different among these oligonucleotides.

### Immunoblot Analysis

For immunoblot analysis, G418-selected cells were isolated, amplified to confluency in 10 cm tissue dishes, and scraped into PBS (pH 7.4). The cells were collected by centrifugation at 200 g for 10 min, and the resulting pellets were suspended in ice-cold lysis buffer (10 mM Tris-HCl, pH 7.4, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100) containing proteinase inhibitors. After incubation on ice for 10 min, samples were centrifuged at 15,000 g for 20 min. Aliquots of Triton-soluble fractions (each 100 µg) were subjected to SDS-PAGE, blotted onto Hybond ECL (Amersham Corp.), and incubated with anti-human Bcl-2 monoclonal antibody (Santa Cruz), and then detected using an ECL-kit (Amersham Corp.).



**Figure 1.** (a) DNA cleavage was induced by RA during the neural differentiation of P19 cells by RA. Genomic DNAs of dying cells were analyzed on 3% nusieve-agarose gels 48 h after RA treatment. P19 cells, cultured in bacteriological dishes in the presence of RA at various concentrations, form aggregates which differentiate into neural cells after being transferred to tissue dishes (Rudnicki and McBurney, 1987). During this process, cells which lose their viability detach from the cell aggregates. These dying cells were separated, and their genomic DNAs were analyzed. DNA cleavage was induced by treatment with  $10^{-9}$ – $10^{-5}$  M RA (lanes 4–8). Only a smear of RNA was observed at RA concentrations below  $10^{-9}$  M (lanes 1–3). (b) Observation of DNA cleavage with time.  $10^{-7}$  M RA was

added to the culture medium and partial DNA cleavage was detected after 6 h (lane 1). The typical DNA ladder formation at 180 bp was firstly observed 12 h after the addition of RA (lane 2). DNA cleavage was attenuated after P19 cells were transferred to tissue culture dishes and they ceased division and differentiated into neurons (lanes 6 and 7). The medium was changed at each analysis to collect the cells that had newly lost viability between time points. (c) DNA fragmentation assay showing the rapid increase of DNA leakage from nuclei. When P19 cells were treated with  $10^{-9}$  or  $10^{-7}$  M RA but not with  $10^{-11}$  M RA, the percentage of DNA outside of nuclei, which was calculated as described in Materials and Methods, increased promptly.

### Primary Culture of Neurons

At first, the meninges were removed from the brain surface of E16 murine embryos. Small pieces of cerebral cortex (3 mm × 3 mm × 1 mm depth) were dissected from the cerebrum, incubated in 0.25% trypsin (Immuno-Biological Laboratories) at 37°C for 5 min, and dissociated by gentle pipetting in Hank's  $Ca^{2+}$ ,  $Mg^{2+}$ -free BSS (Sigma Chemical Co.) containing 0.05% DNase I (Sigma Chemical Co.). Samples were then rinsed and resuspended in  $\alpha$ -medium (Sigma Chemical Co.) supplemented with 10% fetal bovine serum (ICN Radiochemicals) and 5% donor horse serum (CELLect; ICN Radiochemicals), and then seeded in poly-L-lysine hydrobromide (Sigma Chemical Co.)-coated dishes. The same lots of serum were used for analysis. For the estimation of neuronal survival, we seeded 1,000 neurons per single well in 16-chambers tissue slides (Corning). RA or oligonucleotides, at concentrations described in the figure legends, were added to the medium as appropriate.

### Nick End-Labeling Method

Nick end-labeling method was performed with a MEBSTAIN Apoptosis kit (MBL) basically according to the commercial protocol. Cultured neurons were fixed by incubation with precooled methanol at  $-20^{\circ}C$  for 10 min, washed with PBS three times, and incubated in 40  $\mu$ g/ml proteinase for 10 min at 37°C. After washing with TdT buffer, nicked DNA ends were labeled with biotin-dUTP by terminal deoxynucleotide transferase. The enzymes and solutions used above were contained in the kit. The tissue slides were washed with water twice, incubated with avidin-peroxidase (Sigma Chemical Co.), washed with PBS twice, and stained with DAB (3,3'-diaminobenzidine; Sigma Chemical Co.).

### Statistics

InStat 2.0 computer software (Macintosh) was used for statistical analyses.


Generally, multiple pair-wise comparisons were performed using analysis of variance and Bonferroni *t* test among groups of data.

## Results

### RA Induces Apoptosis during the Neural Differentiation of P19 Cells

P19 is one of the most commonly used embryonic cell lines for the analysis of differentiation as it possesses the characteristics of pluripotent stem cells (Jones-Villeneuve et al., 1983; Rudnicki and McBurney, 1987), and differentiates into neurons and glial cells, when treated with RA in the presence of cell-to-cell contact (Jones-Villeneuve et al., 1983; Rudnicki and McBurney, 1987). The differentiated P19 cells express various neuronal markers like GABA, GABA-transaminase, glutamic acid decarboxylase, neuropeptide Y, serotonin, DOPA decarboxylase, somatostatin, and ionotropic glutamate receptor (Ray and Gottlieb, 1994; Staines et al., 1994). RARs and RXRs have been proposed to mediate RA-induced signals based on their abundant expression (Pratt et al., 1990; Bain et al., 1994).

We have noticed in previous experiments (Okamoto et al., 1990; Okazawa et al., 1991) that a large number of cells tend to die during the neural differentiation of P19 cells by RA, and we suspected this phenomenon may be apoptosis.



**Figure 2.** Ultrastructure of dying P19 cells indicating apoptosis. P19 cells which lost their viability during RA-induced neural differentiation were collected and observed by electronmicroscopy. Various phases of apoptosis were found in different cells (*a*). At higher magnifications, chromatin condensation (*b*), cytoplasmic vacuolation, disruption of the nuclear membrane, nuclear fragmentation (*c*), and rupture of the cytoplasmic membrane (*d*) were observed.  $10^{-7}$  M RA was used for the morphological analysis presented here; however, we observed similar apoptotic changes on treatment with  $10^{-9}$ – $10^{-5}$  M RA. The percentage of apoptotic cells changed according to the concentration of RA used.

Firstly, to test this impression, genomic DNA of the dying cells was analyzed on agarose gels on the second day of RA treatment. As expected, DNA fragmentation at about 180 bp was observed on treatment with RA from  $10^{-9}$ – $10^{-5}$  M (Fig. 1 *a*). DNA fragmentation was confirmed by 12 h after exposure to  $10^{-7}$  M RA (Fig. 1 *b*) and was attenuated after day 5 when P19 cells ceased cell division and differentiated into neurons. This observation was quantitatively confirmed by DNA fragmentation assay (Burton, 1956; Sellins and Cohen, 1987). This assay detected the cellular change earlier than the agarose gel electrophoresis, because it observed the DNA leakage from nuclei (Fig. 1 *c*).


Secondly, the ultrastructural appearance of the dying cells was evaluated. Almost all the dying cells displayed typical apoptotic changes including chromatin condensation, nuclear fragmentation, and cytoplasmic vacuolation when treated with  $10^{-7}$  M RA (Fig. 2). The progression of apoptosis from the chromatin condensation to the rupture of the cytoplasmic membrane could be followed in different cells (Fig. 2). A subpopulation of cells showed degenerative changes when treated with a higher concentration of RA (data not shown).

These morphological changes and the DNA fragmentation were similarly induced by another form of RA treatment that evoked differentiation of P19 cells predominantly into endodermal cells. With this treatment, simply culturing P19 cells in tissue dishes for 2 d, apoptosis was observed within 12 h with RA at concentrations above  $10^{-10}$  M (data not shown). This observation suggests that RA-induced apoptosis is not restricted to a specific orientation of differentiation.


#### **Apoptotic Signaling Pathway of RA**

All *trans*-RA binds to RARs, while 9 *cis*-RA binds to RXRs and RARs (for review see Chambon, 1994). Both of these receptor families possess three members,  $\alpha$ ,  $\beta$ , and  $\gamma$ , which act as transcription factors when activated by their ligands (for review see Mangelsdorf, 1994). Therefore, it is possible that the RA-induced signal for the apoptosis is similarly transduced to the nucleus by these receptors and thereby elicits expression of genes which cause apoptosis.


To determine whether RARs mediate the apoptotic sig-



**Figure 3.** Effects of RARs antisense oligonucleotides on RA-induced apoptosis of P19 cells. P19 cells were cultured in  $\alpha$ -medium (Sigma Chemical Co.) -10% FBS in the presence of  $10^{-7}$  M of RA for 48 h. Percentage cell death was calculated by the formula (% number of trypan blue-positive cells from the experiment group - % number of trypan blue-positive cells background)/(100 - % number of trypan blue-positive cells background). Addition of a sense or an antisense oligonucleotide against a RAR family member did not affect the percentage cell death significantly. When all the antisense oligonucleotides against RARs were used in combination ( $\alpha/\beta/\gamma = 1:1:1$ , 10  $\mu$ M in total), the percentage cell death was repressed obviously, while combined use of the three antisense oligonucleotides (10  $\mu$ M in total) against RXRs did not reduce cell death efficiently. Each bar represents the mean  $\pm$  SD of determinations made in 12 fields from three separate cultures. \* $P < 0.01$ , \*\* $P < 0.05$  compared with culture treated by RA but not oligonucleotides.



**Figure 4.** Transfection with RA response element partially inhibited RA-induced apoptosis. Various amounts of the pRARE-CAT plasmid (from 0.1-20  $\mu$ g of DNA/10 cm dish) were transfected into P19 cells by the calcium phosphate method. P19 cells were transferred to bacteriological dishes 24 h after transfection.




**Figure 5.** Effects of cycloheximide and actinomycin-D on the apoptotic cell death evoked by RA.  $1 \times 10^5$  P19 cells were seeded in bacteriological dishes. In addition to  $10^{-7}$  M RA, 1  $\mu$ g/ml cycloheximide (CHX) and/or 0.1  $\mu$ g/ml actinomycin-D (AMD) were added to the media, and trypan blue-positive cells were counted after 48 h. The percentage cell death was determined by the formula (% number of trypan blue-positive cells from the experiment group - % number of trypan blue-positive cells background)/(100 - % number of trypan blue-positive cells background). Each bar represents the mean  $\pm$  SD of determinations made in 12 separate fields from three independent cultures. \* $P < 0.01$  compared to culture with RA alone.

nal, we performed three sets of experiments. Firstly, we synthesized antisense oligonucleotides against RARs (as well as RXRs) and observed whether they inhibited apoptosis. Oligonucleotides against specific members of the RAR family or against specific transcripts of an RAR family member reduced apoptotic cells mildly, while they obviously suppressed apoptosis with a statistical significance when used together (Fig. 3). As all the members of RAR and RXR families are expressed during the differentiation of P19 cells evoked by RA (Jonk et al., 1992; Bain and Gottlieb, 1994), these findings suggest that multiple members of the RAR family, but not of the RXR family for 9 *cis*-RA, participate in the apoptotic signal transduction from all *trans*-RA.

Secondly, we transfected P19 cells with a CAT reporter plasmid containing RA response elements (pRARE-

They were cultured in the presence of  $10^{-7}$  M of RA for additional 24 h and trypan blue-positive cells were counted microscopically. CAT activities were assayed simultaneously using another dish of P19 cells treated similarly. The percentage cell death was determined by the same formula as that shown in Fig. 3. The relative CAT activity was a ratio to the mean value of the transfections using 5  $\mu$ g of plasmid. The relationship between the percentage cell death and the relative CAT activity is shown in the figure. The negative relationship between these two parameters suggests that the RA response element absorbs the apoptotic signals mediated by some retinoic acid receptors.



**Figure 6.** (a) Expression of *bcl-2* was not detected by Northern blot analysis during RA-induced neural differentiation of P19 cells. PolyA<sup>+</sup> RNA was prepared from differentiated P19 cells cultured in the presence of RA and undifferentiated P19 cells cultured in the absence of RA. Expression of the dopamine receptor D<sub>1A</sub> gene was obviously up-regulated in differentiated P19 cells, while expression of *bcl-2* was not detected even when the x-ray film was exposed for 7 d. (b) Expression of *bcl-2* was detected by RT-PCR but suppressed further by RA. P19 cells were cultured in bacteriological dishes in the presence or absence of 10<sup>-7</sup> M RA for 12 or 24 h. 1 μg of total RNA was used for each RT-PCR. G3PDH was amplified as a reference (lanes 1, 2, 5, and 6). *bcl-2* (lane 4 or 8) was down-regulated obviously when compared to its reference (lane 3 or 7) and G3PDH (lane 2 or 6) amplified from the same RNA samples. The genes corresponding to these bands were confirmed by sequencing analysis. (c) Quantitative analysis of *bcl-2* expression. *bcl-2* and G3PDH bands were quantified using an image analyzer. Percentage *bcl-2* amount was calculated at each time point by the formula (*bcl-2* amount in the presence of RA × G3PDH amount in the absence of RA)/(*bcl-2* amount in the absence of RA × G3PDH amount in the presence of RA). Values represent the


mean ± SD determined in triplicate experiments. \*P < 0.01 compared to the value calculated with *bcl-2* before treatment. (d) Expression of the genes related to *bcl-2* did not change during RA-induced neural differentiation. RNA samples were prepared from non-differentiated P19 cells or 24 h treatment with 10<sup>-7</sup> M RA and aggregation. *bax*, *bad*, and *bag-1* gene expression was analyzed by RT-PCR. Specific bands for *bad* and *bag-1* stayed constant (arrowheads). *bax* gene was not amplified in both conditions, while a nonspecific band was observed (dot).

CAT) and observed its effect on apoptosis. We assumed that the transfected RA response element should compete for binding to RARs, with endogenous RA response elements which regulate the genes inducing apoptosis, and should thereby inhibit apoptosis. Since this experiment was dependent on the transfection efficiency, we used the CAT activity as a marker for the efficiency and corrected the effect of pRARE-CAT on the RA-induced apoptosis. As expected, we observed an inverse correlation between the percentage cell death and the CAT activity (Fig. 4).

This result is consistent with the idea that RARs mediate the apoptotic signal of RA. Although the effect was less prominent as that of antisense oligonucleotides, this can be explained by the difference in transfection efficiency between the two methods.

Finally, we observed the effects of protein/RNA synthesis inhibitors on RA-induced apoptosis. We added cycloheximide for the inhibition of protein synthesis and actinomycin-D for the inhibition of RNA synthesis to the culture media. Numbers of apoptotic cells were obviously de-





**Figure 7.** (a) G418-selected cell lines expressed Bcl-2 protein constitutively during neural differentiation evoked by RA. Protein samples were prepared from G418-selected cell lines and P19 before or after RA treatment. 100  $\mu$ g/lane of each sample was separated on 6% SDS-polyacrylamide gel and blotted onto a nylon membrane which was then incubated with anti-human Bcl-2 antibody. The results showed that G418-selected cell lines expressed Bcl-2 protein in both undifferentiated and differentiated states. In contrast to this stable expression, Bcl-2 protein was not detected in the parental P19 cells by immunoblot analysis. (b) Bcl-2 partially inhibited RA-induced apoptosis during neural differentiation. Trypan blue-positive cells were counted 48 h after addition of RA and

the percentage cell death was determined by the formula shown in Fig. 3. The percentage cell death was lower in the cell lines constitutively expressing Bcl-2 protein than in the parental P19 cells.

creased by both treatments (Fig. 5). These observations suggest that RA-induced apoptosis requires de novo synthesis of causative gene products through transcriptional activation by RARs.

### *bcl-2* Is Repressed Strictly during RA-induced Differentiation/Apoptosis

*bcl-2* expression was observed during the RA-induced differentiation for the following two reasons. Firstly, the importance of Bcl-2 function in neuronal death has been demonstrated by recent studies. Bcl-2 rescues neurons from the programmed cell death during development (Garcia et al., 1992), and inhibits neuronal apoptosis caused by neurotrophic factor deprivation (Allsopp et al., 1993), serum deprivation (Mah et al., 1993), or exposure to multiple neurotoxic chemicals including glutamate (Zhong et al., 1993; Reed, 1994). Secondly, the expression of *bcl-2* in neurons was confirmed at the cellular level (Castren et al., 1994; Merry et al., 1994).

We performed Northern blot analysis using polyA<sup>+</sup> RNA. However, neither differentiated nor undifferentiated P19 cells expressed amounts of *bcl-2* sufficient to be detected by this procedure, while the expression of dopamine receptor D<sub>1A</sub>, a reference gene, was up-regulated during differentiation (Fig. 6 a). Therefore, we checked the expression of *bcl-2* using RT-PCR. Interestingly, *bcl-2* transcript was further repressed by RA (Fig. 6 b, lanes 3, 4, 7, and 8), while the housekeeping gene expression of G3PDH estimated using the same RNA samples was not changed (lanes 1 and 2). The down-regulation

was obvious with a statistical analysis of repeated experiments (Fig. 6 c).

We additionally examined the expression of genes related to *bcl-2*: *bax*, *bad*, and *bag-1*. Bax has been shown to homodimerize as well as heterodimerize with Bcl-2 or Bcl-x<sub>L</sub>. When in excess, Bax counters the action of Bcl-2 (Oltvai et al., 1993), whereas Bax heterodimerizes with Bcl-2 to repress apoptosis (Yin et al., 1994). Bad replaces Bax from Bcl-2-Bax or Bcl-x<sub>L</sub>-Bax heterodimers, increases homodimers of Bax, and promotes cell death (Yang et al., 1995). BAG-1, although which is not homologous to the Bcl-2 family members, interacts with Bcl-2 and repress apoptosis (Takayama et al., 1995). RT-PCR analysis did not show obvious change in expression of these molecules (Fig. 6 d), indicating selective repression of *bcl-2*. These observations suggest that P19 cells are not sufficiently supported by the *bcl-2* homodimers or heterodimers at the critical period for their commitment to neuronal differentiation.




*bcl-2* expression in differentiated P19 cells was not equivalent to that of neurons in adult rat reported previously (Merry et al., 1994; Castren et al., 1994). As *bcl-2* expression is regulated developmentally (Merry et al., 1994; Castren et al., 1994; Ferrer et al., 1994), this discrepancy could have been due to the difference in maturity between differentiated P19 cells and neurons in vivo.

### Dysregulated *bcl-2* Expression Inhibits Apoptosis Induced by RA

To test whether dysregulated *bcl-2* expression can rescue P19 cells from RA-induced apoptosis, we established stable transformants of P19 cells which express *bcl-2* constitutively. P19 cells were transfected by a retroviral vector and selected with G418. The resultant clones (P19*bcl2*-7, -8, and -16) were confirmed to express the Bcl-2 protein in both undifferentiated and differentiated states by Western blot analysis (Fig. 7 a) and to possess a similar differentiation potential to P19 cells by morphological and immunohistochemical analyses (data not shown). We next observed the effects of dysregulated expression of *bcl-2* on RA-induced apoptosis using these clones. The percentage cell death, which was estimated 48 h after addition of RA, was clearly repressed in these clones when compared to the parental cell line (Fig. 7 b). This suggested that *bcl-2* expression can partially inhibit RA-induced apoptosis.

We addressed whether P19*bcl2*s and P19 cells could equally differentiate into neurons. When exposed to 10<sup>-7</sup> RA in tissue culture dishes, P19*bcl2*s cells develop a small number of neurons on the layer of nonneuronal cells (Fig. 8 a). When aggregated and exposed to 10<sup>-7</sup> RA in bacteriological dishes, they develop numerous neurons in subsequent culture using tissue culture dishes (Fig. 8 b). These features in differentiation were similar to those of P19 cells. The mean percentages of differentiated neurons were almost equal among the cell lines: 37% in P19, 36% in P19*bcl2*-7, 38% in P19*bcl2*-8, and 34% in P19*bcl2*-16, suggesting that the overexpression of *bcl-2* did not directly influence on the differentiation potential of P19 cells. As a result, the number of differentiated neurons from P19*bcl2*s increased correspondingly to the reduction of apoptosis (Fig. 8 c).






**Figure 8.** Neural differentiation of P19 cells constitutively expressing *bcl-2*.  $1 \times 10^5$  P19*bcl2*-7 cells were cultured with  $10^{-7}$  M RA in a 10 cm tissue grade culture dish for 4 d (a) or cultured with  $10^{-7}$  M RA in a 10 cm bacteriological grade dish for 2 d and additionally in a 10 cm tissue grade dish for 2 d in the absence of RA (b). Cells were incubated with a monoclonal anti-neurofilament 160K antibody (Boehringer-Mannheim) and detected by a biotinized secondary antibody and 3,3'-diaminobenzidine. In the former culture (a), small numbers of neurons extending neurites (arrowheads) were observed on the non-neuronal cell layer. Large numbers of neurons were differentiated in the latter culture (b). (c) Numbers of differentiated neurons increased in P19*bcl2*s. P19 and P19*bcl2*s were differentiated by RA/aggregation treatment as described in b. On day 4, the differentiated neurons were counted after immunostaining and the mean percentage of neurons was determined in eight fields. Relative neuronal number was calculated by the formula: (total cell number of each P19*bcl2* in a dish  $\times$  mean percentage of neurons in each P19*bcl2*)/(total cell number of P19 in a dish  $\times$  mean percentage of neurons in P19). Values represent the mean  $\pm$  SD made in four independent cultures.

### Apoptosis Is Not Associated with Changes in Cytokine Expression

It is possible to speculate that RA causes apoptosis either by repressing signals of neurotrophins or by enhancing

apoptotic signals of some cytokines. With regard to neurotrophic signals, the expression of *trkB*, the receptor for BDNF and NT-4/5, was shown to be increased during the neural differentiation of P19 cells by RT-PCR (Salvatore




**Figure 9.** RA-induced apoptosis of neurons from the murine embryonic brain. Embryonic neurons were prepared from the E16 cerebrum, incubated with trypsin, dispersed by gentle pipetting, and seeded in poly-L-lysine-coated dishes. They were then cultured in  $\alpha$ -MEM medium containing 10% FBS and 5% horse serum in either the presence (*b, d, f, and h*) or absence (*a, c, e, and g*) of  $10^{-7}$  M RA for 2 d. Surviving cells were observed by phase-contrast microscopy (*a-d*), immunostained with anti-160K neurofilament antibody (*e* and *f*), or labeled using a nick end-labeling (*NEL*) kit (*g* and *h*). Compared to non-treated neurons, the survival rate of RA-treated neurons was clearly lower (*d*). Their neurites were degenerated (*f*) and their nuclei were stained positively by NEL (*h*).


et al., 1995; and our unpublished observation). Therefore, supplementing the culture medium with neurotrophins should inhibit apoptosis, if neurotrophins play an important role in RA-induced apoptosis. However this possibility was excluded because addition of NGF, BDNF, NT-3, or NT-4 to the culture medium and the simultaneous use of all these neurotrophins had no effect on apoptosis (data not shown). We next analyzed the expression of molecules

that have been suggested to participate in cell death. RT-PCR analysis indicated that p53, *c-myc*, TNF-receptor-I, and FasL expression were stable during RA-induced neural differentiation. TNF- $\alpha$ , TNF- $\beta$ , TNF-receptor II, and Fas expression were hardly detected by this analysis (data not shown). These findings do not support the idea that apoptosis is a secondary phenomenon caused by a change in cytokine systems.





**Figure 10.** (a) DNA cleavage observed in embryonic neurons treated with RA for 48 h. DNA ladder formation was seen in neurons treated with  $10^{-9}$  or  $10^{-7}$  M RA (lane 3 or 4) but not with  $10^{-11}$  M RA (lane 2). DNA from neurons cultured in the absence of RA is shown as a reference (lane 1). (b) DNA fragmentation assay showing the rapid increase of DNA leakage from nuclei. When treated with  $10^{-9}$  or  $10^{-7}$  M RA but not with  $10^{-11}$  M RA, the percentage of DNA outside of nuclei increased promptly.



TNF-receptors I and II were reportedly up-regulated by RA in a neuroblastoma cell line (Chambaut-Guerin et al., 1995). However, their data showed the down-regulation of TNF-receptor I from days 4 to 6 and again the up-regulation from days 8 to 14, suggesting unstable expression of this receptor. The up-regulation of TNF-receptor II mRNA was observed only transiently on day 1. In our system using P19, the changes of TNF-receptors were not observed statistically. This discrepancy between the cell lines might reflect the difference in cellular maturity.

### RA Induces Apoptosis of Embryonic Neurons in Primary Culture

To test whether RA induces neuronal apoptosis *in vivo*, neurons were dissociated from the murine cerebrum at embryonic day 16 (E16). Survival of these neurons in primary culture was estimated 48 h after dissociation, by direct cell counting under phase-contrast microscopy, by immunostaining with anti-neurofilament 160K antibody, and by nick end labeling (NEL).


The survival of neurons from the cerebrum of E16 embryos was decreased obviously by RA treatment (Fig. 9). This neuronal death was judged to be apoptosis from the obvious nuclear staining shown by NEL experiments (Fig. 9) and the DNA fragmentation detected by agarose gel electrophoresis (Fig. 10 a). The quantitative DNA fragmentation assay confirmed that apoptosis occurred when exposed to  $10^{-9}$  or  $10^{-7}$  M RA (Fig. 10 b). A slight increase in the percentage DNA fragmentation of the reference

group was observed due to the inevitable death of primary neurons in culture (Fig. 10 b). This finding corresponded to the observations that a small portion of neurons morphologically died in the absence of RA (Fig. 9) and that the DNA of nontreated neurons showed a faint smear in agarose gel electrophoresis (Fig. 10 a). However, it was concluded undoubtedly that RA promoted apoptosis judging from the DNA ladder formation in agarose gel electrophoresis and the substantial increase of apoptosis observed by DNA fragmentation assay (Fig. 10 b).

Neurons in primary culture and P19 cells were almost similar in the range of effective concentration of RA to induce apoptosis. RA-induced apoptosis of embryonic neurons in culture was similarly inhibited by the use of antisense oligonucleotides against RARs (Fig. 11). Antisense oligonucleotides against RXRs were not capable to inhibit apoptosis in both cases. These findings suggest that P19 cells and embryonic neurons used common signals in the RA-induced apoptosis.

### Discussion

RA is one of the classical morphogens which induce cellular differentiation and participate in morphological pat-



**Figure 11.** Effects of antisense oligonucleotides against RARs on the survival of embryonic cerebral neurons. Neurons prepared from E16 murine cerebrum were cultured in conditioned medium containing 10% FBS and 5% horse serum. To induce apoptosis,  $10^{-7}$  M of RA was added to the medium. Either  $10 \mu\text{M}$  of an oligonucleotide or  $10 \mu\text{M}$  of mixed oligonucleotides ( $\alpha/\beta/\gamma = 1:1:1$ ) was added to the medium. Survival and death of neurons were estimated 48 h after cell dissociation by trypan blue staining. Percentage cell death was calculated as described in the legend for Fig. 3 to exclude natural death during primary culture. Values represent the mean  $\pm$  SD of determinations made in 12 separate fields from three independent cultures. There was a specific reduction of neuronal cell death when antisense oligonucleotides against RARs were used in combination. \* $P < 0.01$  compared to culture with RA alone.

tern formation. From the analysis of limb formation, RA is suspected to modulate or collaborate with the other morphogenic signals such as the sonic hedgehog gene (Shh) (Echelard et al., 1993; Riddle et al., 1993), homeobox genes (Morgan et al., 1992; Dolle et al., 1993; Charite et al., 1994), fibroblast growth factor 4 (FGF4) (Niswander et al., 1992, 1994), and bone morphogenic proteins (BMPs) (Niswander et al., 1993; Francis et al., 1994) to function as a morphogen (reviewed by Maden 1994; Tabin, 1995). It is well known that apoptosis is induced at the inter-digital areas during this limb formation. However, how RA contributes to this process has not been determined.

RA induces differentiation of various embryonic stem cell lines. P19 cells are differentiated by RA into neuronal cells which express various neuron-specific molecules and display typical morphological characteristics (Rudnicki and McBurney et al., 1986). We found that apoptosis occurs during this neural differentiation. Our experiments indicated that apoptosis was induced at least by  $10^{-9}$  M RA. It corresponds well to the minimum concentration of RA which efficiently activates RARs-mediated gene transcription (Zelent et al., 1989; Giguere et al., 1990). Furthermore, our approaches inhibiting the RARs function succeeded in partially suppressing apoptosis. These findings suggest that RARs-mediated signal transduction play a major role in this apoptosis. RARs should regulate expression of the genes affecting apoptosis as inhibitors of RNA/protein synthesis repressed the RA-induced apoptosis.

Among the various molecules affecting apoptosis, expression of *bcl-2* was selectively suppressed during RA-induced apoptosis of P19 cells. In addition, dysregulated expression of *bcl-2* by a retroviral vector inhibited apoptosis. However, the alteration of *bcl-2* expression would not be a direct cause of apoptosis because undifferentiated P19 cells which show low levels of *bcl-2* expression possess high viability. This suggests that the down-regulation of *bcl-2* provides conditions where the RA-induced apoptosis can progress without difficulty. Further investigations are necessary to elucidate the mechanisms which positively regulate RA-induced apoptosis.

RA-induced apoptosis of P19 cells possesses several unique characteristics: firstly, both neural differentiation and apoptosis progress simultaneously; secondly, apoptosis is induced by a morphogen which possibly acts *in vivo*; and thirdly, this apoptosis is Bcl-2-repressible like the programmed cell death of neurons *in vivo* (Garcia et al., 1992). In these features, our system mimicked the apoptosis which occurs during neural differentiation. This similarity was further confirmed because RA actually induces apoptosis of embryonic neurons. Meanwhile, it should be noted that the RA-induced apoptosis of P19 cells differs from the programmed neuronal death seen during development; the apoptotic process begins immediately after the RA treatment when P19 cells have not yet displayed neuronal characteristics, and is attenuated after the differentiated P19 cells cease cell division. This is in contrast to the programmed neuronal death which occurs just after the arrival of their axons in the target field and therefore after partial maturation of neurons (reviewed by Barde, 1989).

RA-induced apoptosis has been observed in other types of cells recently, and was first reported in HL-60 granulocytes (Park et al., 1994). Interestingly, *bcl-2* is repressed

during differentiation of HL-60 by RA. Furthermore, dysregulated expression of *bcl-2* inhibits apoptosis similarly in HL-60 (Park et al., 1994). The second case has been reported in another embryonic carcinoma cell line, F9, during differentiation to visceral endodermal cells (Atencia et al., 1994). These observations suggest that RA could act as a ubiquitous signal triggering both apoptosis and differentiation in various tissues.

The natures of the triggers of the programmed cell death which occurs *in vivo* during neural development are not known. We suppose that RA (or other retinoids) might act as an endogenous apoptotic signal during the development of the nervous system based on the following observations. Firstly, RA has been demonstrated to be present in the floor plate of the neural tube of embryos where apoptosis occurs during development (Hogan et al., 1992). Secondly, the receptors for RAs, RARs and RXRs, as well as cellular retinoid binding proteins (CRBPs) are expressed widely throughout the nervous system during development (Maden et al., 1990; Ruberte et al., 1992, 1993; Dolle et al., 1994). Thirdly, the ability of low levels of RA to trigger apoptosis during the neural development of stem cells was demonstrated in the present study.

We also expect that our system will be useful for the identification of specific molecules that are indispensable for the apoptosis induced by steroid receptors. It is well known that glucocorticoids induce Bcl-2-repressible apoptosis (Sentman et al., 1991; Strasser et al., 1991; Miyashita and Reed, 1992; Siegel et al., 1992). Androgen receptor, when its CAG repeat is expanded, causes death of motor neurons leading to a motor neuron disease in humans (LaSpada et al., 1991). These observations regarding steroid receptors and our present findings concerning RARs suggest that these receptors might share a common pathway of cell death especially in neurons. Analysis of the downstream signal transduction of the RA-induced apoptosis and comparison between these different systems might be useful for understanding the general mechanism of neuronal death.

We thank Drs. Yoshihide Tsujimoto and Yutaka Eguchi (Osaka University, Yamadaoka, Japan) for providing  $\psi$ 2/B4 cells which produce *bcl-2* expressing retroviral vectors.

Received for publication 15 September 1995 and in revised form 7 November 1995.

#### References

- Akhtar, S., R. Kohle, and R.L. Juliano. 1991. Stability of antisense DNA oligonucleotide analogs in cellular extracts and sera. *Life Sci.* 49:1793-1801.
- Allsopp, T.E., S. Wyatt, H.F. Patterson, and A.M. Davies. 1993. The proto-oncogene *bcl-2* can selectively rescue neurotrophic factor-dependent neurons from apoptosis. *Cell.* 73:295-307.
- Askew, D.S., R.A. Ashmun, B.C. Simmons, and J.L. Cleveland. 1991. Constitutive c-myc expression in an IL-3-dependent myeloid cell line suppresses cell cycle arrest and accelerates apoptosis. *Oncogene.* 6:1915-1922.
- Atencia, R., M. Garcia-Sanz, F. Unda, and J. Arechaga. 1994. Apoptosis during retinoic acid-induced F9 embryonal carcinoma cells. *Exp. Cell Res.* 214:663-667.
- Bafy, G., T. Miyashita, J.R. Williamson, and J.C. Reed. 1993. Apoptosis induced by withdrawal of Interleukin-3 [IL-3] from an IL-3 dependent hematopoietic cell line is associated with repartitioning of intracellular calcium and is blocked by enforced Bcl-2 oncoprotein production. *J. Biol. Chem.* 268: 6511-6519.
- Bain, G., and D.I. Gottlieb. 1994. Expression of retinoid X receptors in P19 embryonal carcinoma cells and embryonic stem cells. *Biochem. Biophys. Res. Commun.* 200:1252-1256.
- Bakshsi, A., J.P. Jensen, P. Goldman, J.J. Wright, O.W. McBride, A.L. Epstein, and S.J. Korsmeyer. 1985. Cloning the chromosomal breakpoint of t(14;18)

- human lymphomas: clustering around J<sub>H</sub> on chromosome 14 and near a transcription unit on 18. *Cell* 41:899-906.
- Barde, Y.-A. 1989. Trophic factors and neuronal survival. *Neuron* 2:1525-1534.
- Batistatou, A., and L.A. Greene. 1991. Aurintricarboxylic acid rescues PC12 cells and sympathetic neurons from death caused by nerve growth factor deprivation: correlation with suppression of endonuclease activity. *J. Cell Biol.* 115:461-471.
- Bissonnette, R.P., F. Echeverri, A. Mahboubi, and D.R. Green. 1992. Apoptotic cell death induced by c-Myc is inhibited by *bcl-2*. *Nature (Lond.)* 359:552-554.
- Boise, L.H., M. Gonzalez-Garcia, C.E. Posteme, L. Ding, T. Lindsten, L.A. Turka, X. Mao, G. Nunez, and C.B. Thompson. 1993. *bcl-x*, a *bcl-2*-related gene that functions as a dominant regulator of apoptotic cell death. *Cell* 74:597-608.
- Burton, K. 1956. A study of the conditions and mechanism of diphenylamine reaction for the colorimetric estimation of deoxyribonucleic acid. *Biochem. J.* 62:315-323.
- Castren, E., Y. Ohga, M.P. Berzaghi, G. Tzimagiorgis, H. Thoenen, and D. Lindholm. 1994. *bcl-2* messenger RNA is localized in neurons and adult rat brain. *Neuroscience* 61:165-177.
- Charite, J., W. de Graaff, S. Shen, and J. Deschamps. 1994. Ectopic expression of Hoxb-8 causes duplication of the ZPA in the forelimb and homeotic transformation of axial structures. *Cell* 78:589-601.
- Chambaut-Guerin, A.-M., M.-C. Martinez, C. Hamimi, X. Gauthereau, and J. Nunez. 1995. Tumor necrosis factor receptors in neuroblastoma SKNB cells and their regulation by retinoic acid. *J. Neurochem.* 65:537-544.
- Chambon, P. 1994. The retinoid pathway: molecular and genetic analyses. *Semin. Cell Biol.* 5:115-125.
- Chittenden, T., E.A. Harrington, R. O'Conner, C. Flemington, R.J. Lutz, G.I. Evan, and B.C. Guild. 1995. Induction of apoptosis by the Bcl-2 homologue Bak. *Nature (Lond.)* 374:733-736.
- Clarke, A.R., C.A. Purdie, D.J. Harrison, R.G. Morris, C.C. Bird, M.L. Hooper, and A.H. Wyllie. 1993. Thymocyte apoptosis induced by p53-dependent and independent pathways. *Nature (Lond.)* 362:849-852.
- Cleary, M.L., and J. Sklar. 1985. Nucleotide sequence of a (14;18) chromosomal breakpoint in follicular lymphoma and demonstration of a breakpoint-cluster region near a transcriptionally active locus on chromosome 18. *Proc. Natl. Acad. Sci. USA* 82:7439-7443.
- Culotta, E., and D.E. Koshland. 1993. p53 sweeps through cancer research. *Science (Wash. DC)* 262:1958-1961.
- de-The, H., M.M. Vivanco-Ruiz, P. Tiollais, H. Stunnenberg, and A. Dejean. 1990. Identification of a retinoic acid responsive element in the retinoic acid beta gene. *Nature (Lond.)* 343:177-180.
- Dispasquale, B., A.M. Marini, and R.J. Youle. 1991. Apoptosis and DNA degradation induced by 1-methyl-4-phenylpyridinium in neurons. *Biochem. Biophys. Res. Commun.* 181:1442-1448.
- Dolle, P., A. Dierich, M. LeMeur, T. Schimmang, B. Schubbaur, P. Chambon, and D. Double. 1993. Disruption of the Hox-13 gene induces localized heterochrony leading to mice with neotenic limbs. *Cell* 75:431-441.
- Dolle, P., V. Fraulob, P. Kastner, and P. Chambon. 1994. Developmental expression of murine retinoid X receptor (RXR) genes. *Mech. Dev.* 45:91-104.
- Echelard, Y., D.J. Epstein, B. St-Jacques, L. Shen, J. Mohler, J.A. McMahon, and A.P. McMahon. 1993. Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell* 75:1417-1430.
- Edwards, S.N., A.E. Buckmaster, and A.M. Tolkovsky. 1991. The death program in cultured sympathetic neurons can be suppressed at post-translational level by nerve growth factor, cyclic AMP and depolarization. *J. Neurochem.* 57:2140-2143.
- Estus, S., W.J. Zaks, R.S. Freeman, M. Gruda, R. Bravo, and E.M. Johnson, Jr. 1994. Altered gene expression in neurons during programmed cell death: identification of c-Jun as necessary for neuronal apoptosis. *J. Cell Biol.* 127:1717-1727.
- Evan, G.I., A.H. Wyllie, C.S. Gilbert, T.D. Littlewood, H. Land, M. Brooks, C.M. Waters, L.Z. Penn, and D.C. Hancock. 1992. Induction of apoptosis in fibroblasts by c-Myc protein. *Cell* 69:119-128.
- Fanidi, A., E.A. Harrington, and G.I. Evan. 1992. Cooperative interaction between *c-myc* and *bcl-2* proto-oncogenes. *Nature (Lond.)* 359:554-556.
- Ferrer, I., A. Tortosa, E. Condom, R. Blanco, A. Macaya, and A. Planas. 1994. Increased expression of *bcl-2* immunoreactivity in the developing cerebral cortex of the rat. *Neurosci. Lett.* 179:13-16.
- Francis, P.H., M.K. Richardson, P.M. Brickell, and C. Tickle. 1994. Bone morphogenetic proteins and a signalling pathway that controls patterning in the developing chick limb. *Development* 120:209-218.
- Garcia, I., I. Martinou, Y. Tsujimoto, and J.-C. Martinou. 1992. Prevention of programmed cell death of sympathetic neurons by the *bcl-2* protooncogene. *Science (Wash. DC)* 258:302-304.
- Giguere, V., M. Shago, R. Zirngibl, P. Tate, J. Rossant, and S. Varmuza. 1990. Identification of a novel isoform of the retinoic acid receptor  $\gamma$  expressed in mouse embryo. *Mol. Cell Biol.* 10:2335-2340.
- Greene, L.A., and A.S. Tischler. 1976. Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proc. Natl. Acad. Sci. USA* 73:2424-2428.
- Greene, L.A. 1978. Nerve growth factor prevents the death and stimulates neuronal differentiation of clonal PC12 pheochromocytoma cells in serum-free medium. *J. Cell Biol.* 78:747-755.
- Ham, J., C. Babij, J. Whitfield, C.M. Pfarr, M. Lallemand, and L.L. Rubin. 1995. A c-Jun dominant negative mutant protects sympathetic neurons against programmed cell death. *Neuron* 14:927-939.
- Hamburger, V., and R.W. Oppenheim. 1982. Naturally occurring neuronal death in vertebrates. *Neurosci. Commun.* 1:39-55.
- Hockenbery, D., G. Nunez, C. Millman, R.D. Schreiber, and S.J. Korsmeyer. 1990. Bcl-2 is an inner mitochondrial membrane protein that blocks programmed cell death. *Nature (Lond.)* 348:334-336.
- Hockenbery, D., Z. Oltvai, X.-M. Yin, C. Millman, and S.J. Korsmeyer. 1993. Bcl-2 functions in an antioxidant pathway apoptosis. *Cell* 75:241-251.
- Hogan, B.L.M., C. Thaller, and G. Eichele. 1992. Evidence that Hensen's node is a site of retinoic acid synthesis. *Nature (Lond.)* 359:237-241.
- Jones-Villeneuve, E.M., M.A. Rudnicki, J.F. Harris, and M.W. McBurney. 1983. Retinoic acid-induced neural differentiation of embryonal carcinoma cells. *Mol. Cell Biol.* 3:2271-2279.
- Jonk, L.J.C., M.E.J. de Jonge, F.A. E. Kruyt, C.L. Mummery, P.T. van der Saag, and W. Kruijer. 1992. Aggregation and cell cycle dependent retinoic acid receptor mRNA expression in P19 embryonal carcinoma cells. *Mech. Dev.* 36:165-172.
- Kane, D.J., T.A. Sarafin, S. Auton, H. Hahn, F.B. Gralla, J.C. Valentine, T. Ord, and D.E. Bredesen. 1993. Bcl-2 inhibition of neural cell death: decreased generation of reactive oxygen species. *Science (Wash. DC)* 262:1274-1276.
- Kiefer, M.C., M.J. Brauer, V.C. Powers, J.J. Wu, S.R. Umansky, L.D. Tomei, and P.J. Barr. 1995. Modulation of apoptosis by the widely distributed homologue Bak. *Nature (Lond.)* 374:736-739.
- Kozopas, K.M., T. Yang, H.L. Buchan, P. Zhou, and R.W. Craig. 1993. MCL-1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to BCL-2. *Proc. Natl. Acad. Sci. USA* 90:3516-3520.
- Kure, S., T. Tominaga, T. Yoshimoto, K. Tada, and K. Narisawa. 1991. Glutamate triggers internucleosomal DNA cleavage in neuronal cells. *Biochem. Biophys. Res. Commun.* 179:39-45.
- LaSpada, A.R., E.M. Wilson, D.B. Lubahn, A.E. Harding, and K.H. Fischbeck. 1991. Androgen receptor gene mutations in X-linked spinal and bulbar muscular atrophy. *Nature (Lond.)* 352:77-79.
- Leroy, P., A. Krust, A. Zelent, C. Mendelsohn, J.-M. Garnier, P. Kastner, A. Dierich, and P. Chambon. 1991. Multiple isoforms of the retinoic acid receptor  $\alpha$  are generated by alternative splicing and differential induction by retinoic acid. *EMBO J.* 10:59-69.
- Lin, E.Y., A. Orloffsky, M.S. Berger, and M.B. Prystowsky. 1993. Characterization of A1, a novel hemopoietic-specific early-response gene with sequence similarity to *bcl-2*. *J. Immunol.* 151:1979-1988.
- Lowe, S.W., E.M. Schmitt, S.W. Smith, B.A. Osborne, and T. Jacks. 1993. p53 is required for radiation-induced apoptosis in mouse thymocytes. *Nature (Lond.)* 362:847-849.
- Luckow, B., and G. Schutz. 1987. CAT constructions with multiple unique restriction sites for the functional analysis of eukaryotic promoters and regulatory elements. *Nucleic Acid Res.* 15:5490.
- Mah, S.P., L.T. Zhong, Y. Liu, A. Roghani, R.H. Edwards, and D.E. Bredesen. 1993. The protooncogene *bcl-2* inhibits apoptosis in PC12 cells. *J. Neurochem.* 60:1183-1186.
- Maden, M., D.E. Ong, and F. Chytil. 1990. Retinoid-binding protein distribution in the developing mammalian nervous system. *Development* 109:75-80.
- Maden, M. 1994. The limb bud - part two. *Nature (Lond.)* 371:560-561.
- Mangelsdorf, D.J., U. Borgmeyer, R.A. Heyman, J.Y. Zhou, E.S. Ong, A.E. Oro, A. Kakizuka, and R.M. Evans. 1992. Characterization of three RXR genes that mediate the action of 9-cis retinoic acid. *Genes & Dev.* 6:329-344.
- Mangelsdorf, D.J., K. Umehono, and R.M. Evans. 1994. The retinoid receptors. In *The retinoids*. 2nd edition. M.B. Sporn, A.B. Roberts, and D.S. Goodman, editors. Raven Press, New York. 319-349.
- Merry, D.E., D.J. Veis, W.F. Hickey, and S.J. Korsmeyer. 1994. *bcl-2* protein expression is widespread in the developing nervous system and retained in the adult PNS. *Development* 120:301-311.
- Miyashita, T., and J.C. Reed. 1993. Bcl-2 oncoprotein blocks chemotherapy-induced apoptosis in human leukemia cell line. *Blood* 81:151-157.
- Miyashita, T., S. Krajewski, M. Krajewska, H.G. Wang, H.K. Lin, D.A. Liebermann, B. Hoffman, and J.C. Reed. 1994. Tumor suppressor p53 is a regulator of *bcl-2* and *bax* gene expression in vitro and in vivo. *Oncogene* 9:1799-1805.
- Miyashita, T., and J.C. Reed. 1995. Tumor Suppressor p53 is a direct transcriptional activator of the human *bax* gene. *Cell* 80:293-299.
- Morgan, R.G., J.-C. Izpisua-Belmonte, D. Duboule, and C.J. Tabin. 1992. Targeted misexpression of Hox-4.6 in the avian limb bud causes apparent homeotic transformations. *Nature (Lond.)* 358:236-239.
- Morgenbesser, S.D., B.O. Williams, T. Jacks, and R.A. DePinho. 1994. p53-dependent apoptosis produced by Rb-deficiency in the developing mouse lens. *Nature (Lond.)* 371:72-74.
- Niswander, L., and G.R. Martin. 1992. *Fgf-4* expression during gastrulation, myogenesis, limb and tooth development in the mouse. *Development* 114:755-768.
- Niswander, L., and G.R. Martin. 1993. *Fgf4* and BMP-2 have opposite effects on limb growth. *Nature (Lond.)* 361:68-71.
- Niswander, L., S. Jeffrey, G.R. Martin, and C. Tickle. 1994. A positive feedback loop coordinates growth and patterning in the vertebrate limb. *Nature*

- (Lond.). 371:609–611.
- Negrini, M., E. Silini, C. Kozak, Y. Tsujimoto, and C.M. Croce. 1987. Molecular analysis of *bcl-2*: structure and expression of the murine gene homologous to the human gene involved in follicular lymphoma. *Cell*. 49:455–463.
- Okamoto, K., H. Okazawa, A. Okuda, M. Sakai, M. Muramatsu, and H. Hamada. 1990. A novel octamer binding transcription factor is differentially expressed in mouse embryonic cells. *Cell*. 60:461–472.
- Okazawa, H., K. Okamoto, T. Ishino-Kaneko, S. Takeda, Y. Toyoda, M. Muramatsu, and H. Hamada. 1991. The oct-3 gene, a gene for an embryonic transcription factor is controlled by retinoic acid repressible enhancer. *EMBO J*. 10:2997–3005.
- Oltvai, Z.N., C.L. Millman, and S.J. Korsmeyer. 1993. Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell*. 74:609–619.
- Oppenheim, R.W. 1991. Cell death during development of the nervous system. *Annu. Rev. Neurosci.* 14:453–501.
- Pan, H., and A.E. Griep. 1994. Altered cell cycle regulation in the lens of HPV-16 E6 or E7 transgenic mice: implications for tumor suppressor gene function in development. *Genes & Dev.* 8:1285–1299.
- Park, J.R., K. Robertson, D.D. Hickstein, S. Tsai, D.M. Hockenbery, and S.J. Collins. 1994. Dysregulated *bcl-2* expression inhibits apoptosis but not differentiation of retinoic acid-induced HL-60 granulocytes. *Blood*. 84:440–445.
- Pratt, M.A., J. Kralova, and M.W. McBurney. 1990. A dominant negative mutation of the alpha retinoic acid receptor gene in a retinoic acid-nonresponsive embryonal carcinoma cell. *Mol. Cell. Biol.* 10:6445–6453.
- Ray, W.J., and D.I. Gottlieb. 1994. Expression of ionotropic glutamate receptor genes by P19 embryonal carcinoma cells. *Biochem. Biophys. Res. Commun.* 197:1475–1482.
- Reed, J.C. 1994. Bcl-2 and the regulation of programmed cell death. *J. Cell Biol.* 124:1–6.
- Riddle, R.D., R.L. Johnson, E. Laufer, and C. Tabin. 1993. Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell*. 75:1401–1416.
- Ruberte, E., V. Friederich, G. Morris-Kay, and P. Chambon. 1992. Differential distribution patterns of CRABP I and CRABP II transcripts during mouse embryogenesis. *Development*. 115:973–987.
- Ruberte, E., V. Friederich, P. Chambon, and G. Morriss-Kay. 1993. Retinoic acid receptors and cellular retinoid binding proteins. III. Their differential transcript distribution during mouse nervous system development. *Development*. 118:267–282.
- Rudnicki, M.A., and M.W. McBurney. 1987. Cell culture methods and induction of differentiation of embryonal carcinoma cell lines. In *Teratocarcinomas and embryonic stem cells*. J. Robertson, editor. IRL Press, Oxford, UK. 19–49.
- Salvatore, A.M., M. Cozzolino, N. Gargano, S. Galanti, A. Levi, and S. Alema. 1995. Neuronal differentiation of P19 cells exhibits cell-specific regulation of neurotrophin receptors. *Neuroreport*. 6:873–877.
- Sato, T., M. Hanada, S. Bodrug, S. Irie, N. Iwama, L.H. Boise, C.B. Thompson, E. Golemis, L. Fong, H.-G. Wang, and J.C. Reed. 1994. Interactions among members of the *bcl-2* protein family analyzed with a two-hybrid system. *Proc. Natl. Acad. Sci. USA*. 91:9238–9242.
- Sellins, K.S., and J.J. Cohen. 1987. Gene induction by  $\gamma$ -irradiation leads to DNA fragmentation in lymphocytes. *J. Immunol.* 139:3199–3206.
- Sentman, C.L., J.R. Shutter, D. Hockenbery, O. Kanagawa, and S.J. Korsmeyer. 1991. Bcl-2 inhibits multiple forms of apoptosis but not negative selection in thymocytes. *Cell*. 67:879–888.
- Shi, Y., J.M. Glynn, L.J. Guilbert, T.G. Cotter, R.P. Bissonnette, and D.R. Green. 1994. Role for c-myc in activation-induced apoptotic cell death in T cell hybridomas. *Science (Wash. DC)*. 257:212–214.
- Shimizu, S., Y. Eguchi, H. Kosaka, W. Kamiike, H. Matsuda, and Y. Tsujimoto. 1995. Prevention of hypoxia-induced cell death by Bcl-2 and Bcl-xL. *Nature (Lond.)*. 374:811–813.
- Staines, W.A., D.J. Morassutti, K.R. Reul, A.I. Ally, and M.W. McBurney. 1994. Neurons derived from embryonal carcinoma cells have varied morphologies and neurotransmitters. *Neuroscience*. 58:735–751.
- Strasser, A., A.W. Harris, and S. Cory. 1991. Bcl-2 transgene inhibits T cell death and perturbs thymic self-censorship. *Cell*. 67:889–899.
- Tabin, C. 1995. The initiation of the limb bud: growth factors, Hox genes and retinoids. *Cell*. 80:671–674.
- Takayama, S., T. Sato, S. Krajewski, K. Kochei, S. Irie, J.A. Millan, and J.C. Reed. 1995. Cloning and functional analysis of BAG-1: a novel bcl-2-binding protein with anti-cell death activity. *Cell*. 80:279–284.
- Tsujimoto, Y., J. Cossman, E. Jaffe, and C.M. Croce. 1985. Involvement of the *bcl-2* gene in human follicular lymphoma. *Science (Wash. DC)*. 228:1440–1443.
- White, E. 1994. p53, guardian of Rb. *Nature (Lond.)*. 371:21–22.
- Yang, E., J. Zha, J. Jockei, L.H. Boise, C.B. Thompson, and S.J. Korsmeyer. 1995. Bad, a heterodimeric partner for Bcl-xL and Bcl-2, displaces Bax and promotes cell death. *Cell*. 80:285–291.
- Yin, X.-M., Z.N. Oltvai, and S.J. Korsmeyer. 1994. BH1 and BH2 domains are required for inhibition of apoptosis and heterodimerization with Bax. *Nature (Lond.)*. 369:321–323.
- Yonish-Rouach, E., D. Resnitzky, J. Lotem, L. Sachs, A. Kimchi, and M. Oren. 1991. Wild-type p53 induces apoptosis of myeloid leukemic cells that is inhibited by interleukin-6. *Nature (Lond.)*. 352:345–347.
- Zelent, A., A. Krust, M. Petkovich, P. Kastner, and P. Chambon. 1989. Cloning of murine  $\alpha$  and  $\beta$  retinoic acid receptors and novel receptor  $\gamma$  predominantly expressed in skin. *Nature (Lond.)*. 339:714–717.
- Zelent, A., C. Mendelssohn, P. Kastner, A. Krust, J.-M. Garnier, F. Ruffenach, and P. Chambon. 1991. Differentially expressed isoforms of the mouse retinoic acid receptor  $\beta$  are generated by usage of two promoters and alternative splicing. *EMBO J*. 10:71–81.
- Zhong, L.T., T. Sarafian, D.J. Kane, A.C. Charles, S.P. Mah, R.H. Edwards, and D. E. Bredesen. 1993. Bcl-2 inhibits death of central neural cells induced by multiple agents. *Proc. Natl. Acad. Sci. USA*. 90:4533–4537.