

SCIENTIFIC REPORTS



OPEN

Microcystis aeruginosa strengthens the advantage of *Daphnia similoides* in competition with *Moina micrura*

Hengxing Tang¹, Xinying Hou¹, Xiaofeng Xue¹, Rui Chen¹, Xuexia Zhu¹, Yuan Huang¹ & Yafen Chen²

Microcystis blooms are generally associated with zooplankton shifts by disturbing interspecific relationships. The influence of *Microcystis* on competitive dominance by different sized zooplanktons showed species-specific dependence. We evaluated the competitive responses of small *Moina micrura* and large *Daphnia similoides* to the presence of *Microcystis* using mixed diets comprising 0%, 20%, and 35% of toxic *M. aeruginosa*, and the rest of green alga *Chlorella pyrenoidosa*. No competitive exclusion occurred for the two species under the tested diet combinations. In the absence of *M. aeruginosa*, the biomasses of the two cladocerans were decreased by the competition between them. However, the *Daphnia* was less inhibited with the higher biomass, suggesting the competitive dominance of *Daphnia*. *M. aeruginosa* treatment suppressed the population growths of the two cladocerans, with the reduced carrying capacities. Nonetheless, the population inhibition of *Daphnia* by competition was alleviated by the increased *Microcystis* proportion in diet. As a result, the competitive advantage of *Daphnia* became more pronounced, as indicated by the higher *Daphnia*: *Moina* biomass ratio with increased *Microcystis* proportions. These results suggested that *M. aeruginosa* strengthens the advantage of *D. similoides* in competition with *M. micrura*, which contributes to the diversified zooplankton shifts observed in fields during cyanobacteria blooms.

Cyanobacteria blooms occur with increased frequency, persistence and wide water ranges due to the eutrophication associated with global warming¹. These aggravated blooms often lead to adverse changes in aquatic ecosystem properties, including toxin production, weakened trophic cascades, and deterioration of water quality^{2,3}.

Compared with other phytoplankton, cyanobacteria are generally accepted as poor food reducing zooplankton fitness. The production of toxic metabolites including microcystins usually causes sublethal or lethal effects for zooplankton survival⁴. The deficiency in nutrition like sterols and long-chained polyunsaturated fatty acids suppresses the carbon metabolism and thereby declines the zooplankton growth^{5,6}. In addition, the colonial or filamentous morphology in cyanobacteria inhibits the grazing activity by clogging the zooplankton filtering apparatus^{7,8}. Nonetheless, in the context of “arms-race” hypothesis, the zooplankton develop adaptations to alleviate the harmful effects by cyanobacteria^{9,10}. For example, some copepods can avoid the ingestion of toxic cells via detecting cyanobacterial metabolites based on the selective feeding^{11,12}. A short-time previous exposure to cyanobacteria improves the fitness of some cladocerans, which could be transferred to offspring via maternal effects^{13–15}. In addition, zooplankton can develop cyanobacteria-tolerant genotypes via rapid evolution¹⁶. These phenotypic and genotypic adaptations are thought to affect the species shifts and community structures of zooplankton during cyanobacteria blooms¹⁷.

Competition is one of the forces structuring zooplankton community. The competition between zooplankton in the presence of cyanobacteria has been widely studied. Most literatures stated that cyanobacteria support the competitive dominance from large sized species to small ones, e.g., from *Daphnia* to smaller cladocerans^{18–21}.

¹Jiangsu Key Laboratory for Biodiversity and Biotechnology, School of Biological Sciences, Nanjing Normal University, 1 Wenyuan Road, Nanjing, 210046, China. ²State Key Laboratory of Lake and Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, 73 East Beijing Road, Nanjing, 210008, China. Correspondence and requests for materials should be addressed to Y.H. (email: huangyuan@njnu.edu.cn) or Y.C. (email: chenyafen7109@163.com)




Figure 1. Population growth curves of *D. similoides* and *M. micrura* in monocultures with different *Microcystis* proportions in food. Both the two cladocerans have three replicates on each day. Some of the data points overlap because they have almost identical values. Lines represent non-linear regression (Logistic model).

Nonetheless, some investigations demonstrated that copepods or large cladocerans are superior competitors in cyanobacterial environment^{22–24}. Given these incompatible results in literatures and the zooplankton adaptations to cyanobacteria, the competition shift during blooms can be interpreted as the dominance by better adapted zooplankton species. As these adaptations are induced by exposure to cyanobacteria, it is hypothesized that the competitive advantage can be affected by varied cyanobacteria stress. To test the hypothesis, we co-cultivated the small-sized *Moina micrura* and large-sized *Daphnia similoides* by feeding diets comprising 0%, 20%, and 35% of toxic *M. aeruginosa*. The objective of the present study was to compare the competitions between the two cladocerans under different *Microcystis* stresses. As no competitive exclusion was observed during the cultivation, the species that has relatively higher biomass in competition was defined as the superior competitor.

Results

Population dynamics in monocultures. The biomasses of the two cladocerans generally increased with progressing culture time among all groups (Fig. 1). Nonetheless, the maximum biomass of both *Daphnia* and *Moina* decreased with increased *Microcystis* proportions in food. As the *Microcystis* proportion increased from 0% to 35%, the maximum biomass of cladocerans decreased from ~5.1 mg to ~1.4 mg per vessel for *Daphnia*, and decreased from ~2.2 mg to ~1.2 mg per vessel for *Moina*. *Microcystis* significantly affected the time reaching the maximum biomass (Fig. 1 and Table 1). Corresponding to the increased *Microcystis* proportion to 35% in food, the time reaching the maximum biomass was shortened from 16 days to 4 days for *Daphnia*, but was prolonged from 7 days to 16 days for *Moina* (Fig. 1).

Population dynamics in cocultures. In general, *Daphnia* had higher biomasses than *Moina* did in all cocultures (Fig. 2). When fed 100% *Chlorella*, the biomass of *Daphnia* rapidly increased to ~2.4 mg per vessel on day 7, after which the biomass increased slightly. By contrast, the biomass of *Moina* gradually decreased from day 7 when a peak biomass of ~0.9 mg per vessel was reached. With 20% *Microcystis* in food, *Daphnia* reached its peak biomass of 1.43 mg per vessel on day 4. The biomass of *Moina* increased slowly with a maximum value of 0.56 mg per vessel on day 13. The population dynamics of *Daphnia* at 35% *Microcystis* was comparable to that at 20% *Microcystis*. By contrast, there was only minor increase in the biomass of *Moina* before day 13, and *Moina* reached its peak biomass of ~0.6 mg per vessel on day 16 at 35% *Microcystis* (Fig. 2).

Traits	DF	SS	MS	F	P
Maximum biomass					
<i>D. similoides</i>					
Food combination (A)	2	17.793	8.896	181.100	<0.001
Presence of competitor (B)	1	8.046	8.046	163.787	<0.001
A × B	2	5.783	2.891	58.857	<0.001
<i>M. micrura</i>					
Food combination (A)	2	1.704	0.852	218.665	<0.001
Presence of competitor (B)	1	3.000	3.000	769.836	<0.001
A × B	2	0.443	0.221	56.788	<0.001
Time to maximum biomass					
<i>D. similoides</i>					
Food combination (A)	2	351.000	175.500	10.324	0.002
Presence of competitor (B)	1	72.000	72.000	4.235	0.062
A × B	2	147.000	73.500	4.324	0.039
<i>M. micrura</i>					
Food combination (A)	2	325.000	162.500	108.333	<0.001
Presence of competitor (B)	1	0.500	0.500	0.333	0.574
A × B	2	7.000	3.500	2.333	0.139
Carrying capacity (K)					
<i>D. similoides</i>					
Food combination (A)	2	15.301	7.651	248.312	<0.001
Presence of competitor (B)	1	5.680	5.680	184.364	<0.001
A × B	2	5.527	2.763	89.693	<0.001
<i>M. micrura</i>					
Food combination (A)	2	0.674	0.337	41.486	<0.001
Presence of competitor (B)	1	3.248	3.248	399.832	<0.001
A × B	2	0.246	0.123	15.159	<0.001
Population growth rate (r)					
<i>D. similoides</i>					
Food combination (A)	2	1.431	0.716	12.685	<0.001
Presence of competitor (B)	1	0.148	0.148	2.624	0.131
A × B	2	0.590	0.295	5.230	0.023
<i>M. micrura</i>					
Food combination (A)	2	1.750	0.875	152.820	<0.001
Presence of competitor (B)	1	0.0105	0.0105	1.839	0.200
A × B	2	0.0151	0.00757	1.319	0.304

Table 1. Results of two-way ANOVA on maximum biomass, time to maximum biomass, carrying capacity and population growth rate of *D. similoides* and *M. micrura* subjected to different food combinations and absence/presence of competitor (DF: degree of freedom; SS: sum of squares; MS: mean squares; F-F ratio).

Biomass inhibition of species and biomass ratio in competition. The biomass inhibition of the two species changed with time depending on the *Microcystis* proportion (Fig. 3a–c). Without *Microcystis* addition, the *Daphnia* biomass in competition was sharply inhibited by ~57% during the initial 3 days, whereas that of *Moina* was promoted by the presence of *Daphnia*, as indicated by the negative values of biomass inhibition rate. Nonetheless, the biomass inhibition of *Moina* dramatically increased to ~61%, and was higher than that of *Daphnia* after 7 days (Fig. 3a). With 20% *Microcystis* in food, although the biomass inhibition of *Daphnia* increased along with time, it reached a maximum value of ~59% at the end of experiment, which was rapidly achieved on day 3 in groups without *Microcystis*. The biomass inhibition rate of *Moina* varied around that of *Daphnia* (Fig. 3b). With 35% *Microcystis*, although negative values were observed for the biomass inhibition of *Moina* at the initial 7 days, it sharply increased to ~49% from day 9. Nonetheless, the biomass inhibition rate of *Daphnia* was negative and approached zero during the experiment period (Fig. 3c).

The *Daphnia*: *Moina* biomass ratio provided an intuitive understanding for the differences in population dynamics between the two cladocerans (Fig. 3d). In cocultures, the biomass of *Daphnia* was always higher than that of *Moina*, regardless of the *Microcystis* treatment, as indicated by the ratio values >1. There were remarkable increases in the biomass ratio with increased *Microcystis* proportions in food. Nonetheless, the peak value of the ratio at 35% *Microcystis* appeared later than that at 20% *Microcystis* did.




Figure 2. Population growth curves of *D. similoides* and *M. micrura* in cocultures with different *Microcystis* proportions in food. Both the two cladocerans have three replicates on each day. Some of the data points overlap because they have almost identical values. Lines represent non-line regression (Logistic model).

Environmental carrying capacity and population growth rate. The carrying capacities of both *Daphnia* and *Moina* were significantly affected by the *Microcystis* treatment, the presence of competitor and the interactions between them (Table 1). When fed 100% *Chlorella*, the carrying capacities of *Daphnia* and *Moina* were decreased by 51.7% and 62.4% by the presence of each other in cocultures. This phenomenon was also observed in populations fed 20% *Microcystis*, together with the overall decreased carrying capacities. At 35% *Microcystis*, the carrying capacity of *Daphnia* in cocultures was higher by 21.4% than that in monocultures, whereas the carrying capacity of *Moina* was decreased by 51.9% in competition (Fig. 4a). *Daphnia* had higher carrying capacities than the *Moina* did in all cultures except for the case in monocultures at 35% *Microcystis*. The *Daphnia*:*Moina* carrying capacity ratio in cocultures was always higher than that in monocultures, although decreasing trends were observed with increased *Microcystis* proportions in food (Fig. 4b).

The population growth rates (r) of the two cladocerans were significantly affected by the *Microcystis* treatment, but not the competition (Table 1). When fed 100% *Chlorella*, the r of *Moina* was higher by ~42% than that of *Daphnia*. *Microcystis* treatment increased the r of *Daphnia*, but decreased the r of *Moina*. The r of *Daphnia* was remarkable higher by 50.7% at 20% *Microcystis* and by 77.6% at 35% *Microcystis* than those of *Moina* in cocultures, leading to the increased *Daphnia*:*Moina* growth rate ratio with increased *Microcystis* proportions in food (Fig. 5).

Discussion

The present study revealed that the large-sized *D. similoides* was superior to the small-sized *M. micrura* in competition under favorable food condition. When fed 100% *Chlorella*, *D. similoides* had higher biomass than the *M. micrura* did in cocultures, although the biomasses of the two cladocerans were suppressed by the competition (Figs 1 and 2). This is in accordance with the previous conclusion that abundant edible food favours the larger species to be superior competitor^{25–27}. On condition that the carbon levels satisfy the food requirements of the animals, large species gathers food more efficiently, thereby decreasing the food availability of small species²⁸. The large animals also generally have the stronger ability to ingest the food. Other mechanisms, such as age at first reproduction and embryonic developmental time, also contribute to the competitive outcomes among zooplankton species²⁹.

Corresponding to the increased *Microcystis* proportion, different population responses to *Microcystis* were observed: faster to reach the carrying capacity with the subsequent increased population growth rate in *D. similoides*, but the opposite case in *M. micrura*. This result enriches the species-specific responses in zooplankton population growth to cyanobacteria³⁰. Exposure to *Microcystis* would promote the large zooplankton (e.g., *D.*




Figure 3. Biomass inhibition rates of species (a–c) and the biomass ratio between *D. similoides* (Ds) and *M. micrura* (Mm) in co-cultures (d) with different *Microcystis* proportions in food.




Figure 4. Carrying capacities (a) of *D. similoides* and *M. micrura* and their ratios in monocultures or cocultures (b) with different *Microcystis* proportions in food.

magna) to reach its maturity earlier, with the shortened reproduction age²³. At a population level, these changes in life history traits facilitate the large species reaching its carrying capacity faster. Nonetheless, the biomasses and carrying capacities of the two cladocerans were finally decreased by *Microcystis* treatment (Figs 1, 2 and 4). This is highly related to the nutritional deficiencies and the toxicity of microcystins of the cyanobacteria for the zooplankton^{7,31,32}. Given the biomass inhibition by *Microcystis* treatment, the biomass difference between the two cladocerans in competition was enlarged with increased *Microcystis* proportion, as indicated by the increasing




Figure 5. Population growth rates (a) of *D. similoides* and *M. micrura* and their ratios in monocultures or cocultures (b) with different *Microcystis* proportions in food.

Daphnia: *Moina* biomass ratio (Fig. 3). It is concluded that the competitive advantage of *D. similoides* over *M. micrura* was strengthened by *Microcystis* treatment.

Because of the relatively larger gape size with higher filtration on the filamentous cyanobacteria, large *Daphnia* is generally assumed to be more vulnerable to *Microcystis* than small species^{33,34}. Nonetheless, the present strain of *M. aeruginosa* grows as unicell in laboratory, and the *D. similoides* and *M. micrura* are supposed to graze both the toxic and non-toxic cells equally due to their non-selective filtration. *M. micrura* assimilated little *Microcystis* when fed only *Microcystis* or even a mixture of *Microcystis* and *Chlorella*³⁵. Given the present low proportions (<35%) of *Microcystis* in diet, large *Daphnia* can minimise the negative influence from cyanobacteria via microcystins detoxification³⁶. Zhang *et al.*³⁷ studied that the large *D. similoides* assimilates low abundance of *Microcystis* with improved reproduction. Using a combined stable-isotope and fatty-acid approach, de Kluijver *et al.*³⁸ found that *D. similis* consumes live *Microcystis* cells. This supplies additional material and energy for the *D. similoides* growth in comparison with *M. micrura*. Repeated toxic cyanobacteria- exposure can also increase the tolerance of large *Daphnia* population to toxic *Microcystis* via improving antioxidant systems^{39,40}. In addition, the low nutritional value of cyanobacteria for *Daphnia* promotes the offspring tolerance to toxic *Microcystis*⁴¹. Gustafsson *et al.*¹³ studied that previous-exposure to toxic *Microcystis* increased the offspring fitness in *Microcystis* environment. This adaptation may result from improved survival, enhanced reproduction or faster development of offspring^{42,43}. Although some small cladocerans are also studied to develop adaptation³⁴, the severer biomass inhibition by *Microcystis* in *M. micrura* indicated its weaker adaptation compared with that in *D. similoides* in the current study. A comparative study on the phenotypic adaptations between the two species based on individual performances will be performed in the next work. In the presence of *M. micrura*, we surprisingly observed that the biomass of *D. similoides* at 35% *Microcystis* was slightly higher than that at 20% *Microcystis*. As the *Moina* was severer inhibited by competition with increased *Microcystis*, it is presumed that *Daphnia* in the 35% *Microcystis*-treated cocultures consumed more good food, thereby leading to the relatively higher biomass.

There is a great variation in the influence of cyanobacteria on zooplankton competition. The present observation is not consistent with the general recognition that *Microcystis* promotes the dominance of small-sized cladocerans, but instead supports several investigations demonstrating that large-sized cladocerans are superior competitors in cyanobacteria environment^{22,24,44}. Besides the concentration tested in present study, many bloom-related variables, such as morphology and toxic property, affect the zooplankton shifts^{7,45-47}. In natural systems, the zooplankton composition during bloom is also regulated by other factors such as temperature and planktivorous fish^{48,49}. Under the background of global warming, increasing temperature would enhance the effects of cyanobacteria on zooplankton with expansive blooms⁵⁰⁻⁵². For example, the cladoceran offspring tolerance to toxic *Microcystis* can be promoted by maternal warming⁵³. Fish predation also drives the zooplankton fluctuation³⁴. The planktivorous fishes affect zooplankton via directly predation or by reducing edible phytoplankton abundance to zooplankton⁵⁵. Recent study showed that planktivorous fishes associated with cyanobacteria promote the zooplankton community shift towards species with good escape ability and r -strategy in survival^{56,57}. Therefore, besides the exploration on bloom-relevant factors influencing zooplanktons alone or in combinations, the trophic interactions between planktivorous fish, zooplankton and cyanobacteria require deeper studies to assess the zooplankton community structures during the expansive blooms under warming climate.

Materials and Methods

Cladocerans and algal food. Both the *Daphnia similoides* and *Moina micrura* were collected from Taihu Lake in China. The animals were then cultivated in laboratory by feeding 100% *Chlorella pyrenoidosa* at 25 °C for about three years. *C. pyrenoidosa* was pre-cultured in liquid BG-11 medium at 25 °C and illuminated at 45 $\mu\text{mol m}^{-2} \text{s}^{-1}$ provided by fluorescent lamps in a light-dark period of 14:10 h. Log-phase *C. pyrenoidosa* were harvested by centrifugation at 6300 $\times g$ for 10 min and used as food. *M. aeruginosa* PCC7806 was obtained from the Freshwater Algae Culture Collection of the Institute of Hydrobiology (Wuhan, China). The cyanobacteria produce at least two types of microcystin (MC-LR and MC-RR) with a total content of 3.6 pg per cell via the

high-performance liquid chromatography detection⁵⁸. The cyanobacteria were axenically cultured under the same above conditions.

Experimental protocol. Three food compositions were tested as 100% *C. pyrenoidosa* + 0% *M. aeruginosa*, 80% *C. pyrenoidosa* + 20% *M. aeruginosa*, and 65% *C. pyrenoidosa* + 35% *M. aeruginosa*, with the total carbon content of 1 mg L⁻¹. The two cladocerans all died in several days when fed on 50% *Microcystis* in diet based on our pre-experiment. The present-used proportions of *M. aeruginosa* were not highly toxic to cause elimination of either *M. micrura* or *D. similoides*, and thus suitable for competition experiments. Within each food composition, three cladoceran-treatments were set up: (1) 5 *D. similoides* (Ds) cultivated alone; (2) 5 *M. micrura* (Mm) cultivated alone; (3) 5 Ds and 5 Mm cultivated together. The initial sizes for *D. similoides* and *M. micrura* were averaged 0.65 mm and 1.2 mm in length on account of a Nis-elements image analyzer coupled with a Nikon light microscope. The experiment was carried out in 1-L beakers containing 500 mL culture media with one specific food composition. The experiment was performed in triplicate, resulting in 3 (food composition) × 3 (cultivation pattern) × 3 (replicate) = 27 beakers. All beakers were maintained at 25°C in a temperature-controlled chamber and illuminated by 45 μmol m⁻² s⁻¹ fluorescent light with a light–dark period of 14:10 h. To maintain constant food concentrations, we replaced 50% of the medium in each beaker daily with fresh medium with appropriate food abundance. The dry biomass of the animals was estimated via measuring the body length on the basis of the regression curves by Culver *et al.*⁵⁹. The experiment was not terminated until no remarkable increase was detected in the population abundance. When the experiment was finished, the cladoceran populations were composed of individuals in different ages with a length range of 0.7–1.6 mm for *D. similoides* and 0.4–0.9 mm for *M. micrura*.

The cladoceran biomass versus time was fitted by using the Logistic model:

$$B_t = \frac{K}{1 + \frac{K - B_0}{B_0} e^{-rt}},$$

where B_0 and B_t represent the cladoceran biomasses at initial time (t_0) and time t , r represents the population growth rate, and K represents the environmental carrying capacity. The biomass inhibition of species in competition was calculated as: biomass inhibition rate (%) = [(biomass in monocultures) – (biomass in cocultures)]/biomass in monocultures × 100%. The biomass ratio of the two cladocerans in cocultures was defined as the biomass of *Daphnia* relative to that of *Moina* (*Daphnia*: *Moina*).

Statistical analysis. All data are presented as mean ± 1 SE. Two-way ANOVA was used to compare differences between groups in terms of population growth rate, maximum biomass, time to maximum biomass, and the carrying capacity with food combination and absence/presence of competitor as the fixed factors. Significant analyses were followed by Tukey's post-hoc tests to locate meaningful differences. Statistical analysis was performed using Sigmaplot 11.0 software.

References

- O'neil, J. M., Davis, T. W., Burford, M. A. & Gobler, C. J. The rise of harmful cyanobacteria blooms: the potential roles of eutrophication and climate change. *Harmful Algae* **14**, 313–334 (2012).
- Paerl, H. W. & Paul, V. J. Climate change: links to global expansion of harmful cyanobacteria. *Water Res.* **46**, 1349–1363 (2012).
- Rondel, C. *et al.* A cyanobacterial bloom prevents fish trophic cascades. *Freshwater Biol.* **53**, 637–651 (2008).
- Leflaive, J. & Ten-Hage, L. Algal and cyanobacterial secondary metabolites in freshwaters: a comparison of allelopathic compounds and toxins. *Freshwater Biol.* **52**, 199–214 (2007).
- Martin-Creuzburg, D., Von Elert, E. & Hoffmann, K. H. Nutritional constraints at the cyanobacteria–*Daphnia magna* interface: the role of sterols. *Limnol. Oceanogr.* **53**, 456–468 (2008).
- Muller-Navarra, D. C., Brett, M. T., Liston, A. M. & Goldman, C. R. A highly unsaturated fatty acid predicts carbon transfer between primary producers and consumers. *Nature* **403**, 74 (2000).
- Wilson, A. E., Sarnelle, O. & Tillmanns, A. R. Effects of cyanobacterial toxicity and morphology on the population growth of freshwater zooplankton: Meta-analyses of laboratory experiments. *Limnol. Oceanogr.* **51**, 1915–1924 (2006).
- DeMott, W. R., Gulati, R. D. & Van Donk, E. *Daphnia* food limitation in three hypereutrophic Dutch lakes: Evidence for exclusion of large-bodied species by interfering filaments of cyanobacteria. *Limnol. Oceanogr.* **46**, 2054–2060 (2001).
- Ger, K. A., Hansson, L. A. & Lüring, M. Understanding cyanobacteria–zooplankton interactions in a more eutrophic world. *Freshwater Biol.* **59**, 1783–1798 (2014).
- Ger, K. A. *et al.* The interaction between cyanobacteria and zooplankton in a more eutrophic world. *Harmful Algae* **54**, 128–144 (2016).
- Ger, K. A., Panosso, R. & Lüring, M. Consequences of acclimation to *Microcystis* on the selective feeding behavior of the calanoid copepod *Eudiaptomus gracilis*. *Limnol. Oceanogr.* **56**, 2103–2114 (2011).
- Engström, J. *et al.* Feeding interactions of the copepods *Eurytemora affinis* and *Acartia bifilosa* with the cyanobacteria *Nodularia* sp. *J. Plankton Res.* **22**, 1403–1409 (2000).
- Gustafsson, S., Rengefors, K. & Hansson, L. A. Increased consumer fitness following transfer of toxin tolerance to offspring via maternal effects. *Ecology* **86**, 2561–2567 (2005).
- Jiang, X. D. *et al.* Fitness benefits and costs of induced defenses in *Daphnia carinata* (Cladocera: Daphnidae) exposed to cyanobacteria. *Hydrobiologia* **702**, 105–113 (2013).
- Schwarzenberger, A. & Von Elert, E. Cyanobacterial protease inhibitors lead to maternal transfer of increased protease gene expression in *Daphnia*. *Oecologia* **172**, 11–20 (2013).
- Sarnelle, O. & Wilson, A. E. Local adaptation of *Daphnia pulicaria* to toxic cyanobacteria. *Limnol. Oceanogr.* **50**, 1565–1570 (2005).
- Chislock, M. F., Sarnelle, O., Olsen, B. K., Doster, E. & Wilson, A. E. Large effects of consumer offense on ecosystem structure and function. *Ecology* **94**, 2375–2380 (2013).
- Infante, A. & Riehl, W. The effect of cyanophyta upon zooplankton in a eutrophic tropical lake (Lake Valencia, Venezuela). *Hydrobiologia* **113**, 293–298 (1984).
- Hansson, L. A., Gustafsson, S., Rengefors, K. & Bomark, L. Cyanobacterial chemical warfare affects zooplankton community composition. *Freshwater Biol.* **52**, 1290–1301 (2007).

20. Chen, F. Z., Ye, J. L., Shu, T. T., Sun, Y. & Li, J. Zooplankton response to the lake restoration in the drinking-water source in Meiliang Bay of subtropical eutrophic Lake Taihu, China. *Limnol. Ecol. Manage. Inland Waters* **42**, 189–196 (2012).
21. Jiang, X. D. *et al.* Increasing dominance of small zooplankton with toxic cyanobacteria. *Freshwater Biol.* **62**, 429–443 (2017).
22. Kim, H. W., Joo, G. J., Chang, K. H. & Hwang, S. J. Zooplankton community dynamics during the summer *Microcystis* bloom in the lower part of the Nakdong River, South Korea. *P. Internat. Assoc. Theor. Appl. Limnol.* **27**, 1044–1049 (2001).
23. Bouvy, M., Pagano, M. & Troussellier, M. Effects of a cyanobacterial bloom (*Cylindrospermopsis raciborskii*) on bacteria and zooplankton communities in Ingazeira reservoir (northeast Brazil). *Aquat. Microb. Ecol.* **25**, 215–227 (2001).
24. Sikora, A. & Dawidowicz, P. Do the presence of filamentous cyanobacteria and an elevated temperature favor small-bodied *Daphnia* in interspecific competitive interactions? *Fund. Appl. Limnol.* **185**, 307–314 (2014).
25. Romanovsky, Y. E. & Feniova, I. Y. Competition among Cladocera: effect of different levels of food supply. *Oikos* **44**, 243–252 (1985).
26. Chen, R. *et al.* Food availability and initial relative abundance determine the outcome of interspecific competition between two different-sized cladocerans. *Int. Rev. Hydrobiol.* **101**, 105–112 (2016).
27. Divya, S. P., Kathiresan, K., Asha, P., Sekar, V. & Rajasekaran, R. Experimental study of the interspecific competition between two sibling marine herbivorous rotifers in relation to food availability and initial population density. *Acta Oceanol. Sin.* **31**, 113–126 (2012).
28. Lampert, W. & Sommer, U. *Limnology: the Ecology of Lakes and Streams* (Oxford University Press, 2007).
29. DeMott, W. R. The role of competition in zooplankton succession in *Plankton Ecology: Succession in Plankton Communities* (ed. Sommer, U.) 195–252 (Springer, 1989).
30. Tillmanns, A. R., Wilson, A. E., Pick, F. R. & Sarnelle, O. Meta-analysis of cyanobacterial effects on zooplankton population growth rate: species-specific responses. *Fund. Appl. Limnol.* **171**, 285–295 (2008).
31. Von Elert, E. & Wolffrom, T. Supplementation of cyanobacterial food with polyunsaturated fatty acids does not improve growth of *Daphnia*. *Limnol. Oceanogr.* **46**, 1552–1558 (2001).
32. Bednarska, A., Pietrzak, B. & Pijanowska, J. Effect of poor manageability and low nutritional value of cyanobacteria on *Daphnia magna* life history performance. *J. Plankton Res.* **36**, 838–847 (2014).
33. Ferrão-Filho, Ad. S. & Azevedo, S. M. F. Effects of unicellular and colonial forms of toxic *Microcystis aeruginosa* from laboratory cultures and natural populations on tropical cladocerans. *Aquat. Ecol.* **37**, 23–35 (2003).
34. Guo, N. C. & Xie, P. Development of tolerance against toxic *Microcystis aeruginosa* in three cladocerans and the ecological implications. *Environ. Pollut.* **143**, 513–518 (2006).
35. Hanazato, T. & Yasuno, M. Evaluation of *Microcystis* as food for zooplankton in a eutrophic lake. *Hydrobiologia* **144**, 251–259 (1987).
36. Pflugmacher, S. *et al.* Identification of an enzymatically formed glutathione conjugate of the cyanobacterial hepatotoxin microcystin-LR: the first step of detoxication. *BBA-Gen. Subjects* **1425**, 527–533 (1998).
37. Zhang, X. L., Deng, D. G., Zhang, K., Wang, W. P. & Ji, L. Combined effects of colonial size and concentration of *Microcystis aeruginosa* on the life history traits of *Daphnia similoides*. *Acta Biol. Hung.* **66**, 282–292 (2015).
38. de Kluijver, A., Yu, J. L., Houtekamer, M., Middelburg, J. J. & Liu, Z. W. Cyanobacteria as a carbon source for zooplankton in eutrophic Lake Taihu, China, measured by ¹³C labeling and fatty acid biomarkers. *Limnol. Oceanogr.* **57**, 1245–1254 (2012).
39. Gustafsson, S. & Hansson, L. A. Development of tolerance against toxic cyanobacteria in *Daphnia*. *Aquat. Ecol.* **38**, 37–44 (2004).
40. Wojtal-Frankiewicz, A. *et al.* Microcystin assimilation and detoxification by *Daphnia* spp. in two ecosystems of different cyanotoxin concentrations. *J. Limnol.* **72**, 13 (2013).
41. Lyu, K. *et al.* Maternal consumption of non-toxic *Microcystis* by *Daphnia magna* induces tolerance to toxic *Microcystis* in offspring. *Freshwater Biol.* **61**, 219–228 (2016).
42. Jiang, X. D. *et al.* Maternal effects of inducible tolerance against the toxic cyanobacterium *Microcystis aeruginosa* in the grazer *Daphnia carinata*. *Environ. Pollut.* **178**, 142–146 (2013).
43. Li, Q. M. & Jiang, X. D. Offspring tolerance to toxic *Microcystis aeruginosa* in *Daphnia pulex* shaped by maternal food availability and age. *Fund. Appl. Limnol.* **185**, 315–319 (2014).
44. George, D. G. & Edwards, R. W. Population dynamics and production of *Daphnia hyalina* in a eutrophic reservoir. *Freshwater Biol.* **4**, 445–465 (1974).
45. Pérez-Morales, A., Sarma, S. S. S. & Nandini, S. Feeding and filtration rates of zooplankton (rotifers and cladocerans) fed toxic cyanobacterium (*Microcystis aeruginosa*). *J. Environ. Biol.* **35**, 1013 (2014).
46. Chen, F. Z. & Xie, P. The toxicities of single-celled *Microcystis aeruginosa* PCC7820 and liberated *M. aeruginosa* to *Daphnia carinata* in the absence and presence of the green alga *Scenedesmus obliquus*. *J. Freshwater Ecol.* **19**, 539–545 (2004).
47. Chen, F. Z. & Xie, P. The effects of fresh and decomposed *Microcystis aeruginosa* on cladocerans from a subtropical Chinese lake. *J. Freshwater Ecol.* **18**, 97–104 (2003).
48. Kaczkowski, Z. *et al.* Relationships among cyanobacteria, zooplankton and fish in sub-bloom conditions in the Sulejow Reservoir. *J. Limnol.*, doi:10.4081/jlimnol.2017.1521 (2017).
49. Havens, K. E., Fulton, R. S. III, Beaver, J. R., Samples, E. E. & Colee, J. Effects of climate variability on cladoceran zooplankton and cyanobacteria in a shallow subtropical lake. *J. Plankton Res.* **38**, 418–430 (2016).
50. Gilbert, J. J. Effect of temperature on the response of planktonic rotifers to a toxic cyanobacterium. *Ecology* **77**, 1174–1180 (1996).
51. Hochmuth, J. D., Janssen, C. R. & De Schampelaere, K. A. Temperature and food concentration have limited influence on the mixture toxicity of copper and *Microcystis aeruginosa* to *Daphnia magna*. *Environ. Toxicol. Chem.* **35**, 742–749 (2016).
52. Jiang, X. D. *et al.* Cyanobacteria alter competitive outcomes between *Daphnia* and *Bosmina* in dependence on environmental conditions. *Fund. Appl. Limnol.* **184**, 11–22 (2014).
53. Lyu, K. *et al.* Cladoceran offspring tolerance to toxic *Microcystis* is promoted by maternal warming. *Environ. Pollut.* **227**, 451–459 (2017).
54. Havens, K. E., Beaver, J. R., Manis, E. E. & East, T. L. Inter-lake comparisons indicate that fish predation, rather than high temperature, is the major driver of summer decline in *Daphnia* and other changes among cladoceran zooplankton in subtropical Florida lakes. *Hydrobiologia* **750**, 57–67 (2015).
55. DeVries, D. R. & Stein, R. A. Complex interactions between fish and zooplankton: quantifying the role of an open-water planktivore. *Can. J. Fish. Aquat. Sci.* **49**, 1216–1227 (1992).
56. Li, Y., Xie, P., Zhang, J., Tao, M. & Deng, X. W. Effects of filter-feeding planktivorous fish and cyanobacteria on structuring the zooplankton community in the eastern plain lakes of China. *Ecol. Eng.* **99**, 238–245 (2017).
57. Chen, Y., Xu, B., Yang, W. & Jiang, X. D. Combined effects of calcium deficiency and increasing cyanobacteria on competitive dominance of zooplankton. *Fund. Appl. Limnol.* **187**, 11–20 (2015).
58. Gan, N. Q., Huang, Q., Zheng, L. L. & Song, L. R. Quantitative assessment of toxic and nontoxic *Microcystis* colonies in natural environments using fluorescence *in situ* hybridization and flow cytometry. *Sci. China Life Sci.* **53**, 973–980 (2010).
59. Culver, D. A., Boucherle, M. M., Bean, D. J. & Fletcher, J. W. Biomass of freshwater crustacean zooplankton from length–weight regressions. *Can. J. Fish. Aquat. Sci.* **42**, 1380–1390 (1985).

Acknowledgements

We thank the two anonymous reviewers for their helpful comments and suggestions which significantly improved this manuscript. This study was supported by the National Natural Science Foundation of China (31500373), Natural Science Youth Foundation of Jiangsu Province (BK20150972), “333 High Level Talent Project” in Jiangsu Province (BRA2017452), and the Priority Academic Program Development of Jiangsu Higher Education Institutions.

Author Contributions

Y.H. conceived this study and designed the experiments; H.X.T., X.Y.H. and R.C. performed the experiments; X.F.X., X.X.Z. and Y.H. conducted the data analysis and prepared figures and tables. Y.H. and Y.F.C. wrote the manuscript. All of the authors reviewed and approved the manuscript.

Additional Information

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017