

Published in final edited form as:

Nat Genet. 2020 December 01; 52(12): 1314–1332. doi:10.1038/s41588-020-00713-x.

Discovery of rare variants associated with blood pressure regulation through meta-analysis of 1.3 million individuals

A full list of authors and affiliations appears at the end of the article.

These authors contributed equally to this work.

Abstract

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence to: Patricia B. Munroe; Joanna M. M. Howson.

Corresponding authors: Joanna M. M. Howson: JMMHowson@gmail.com, Patricia B. Munroe: P.B.Munroe@qmul.ac.uk.

*Current address (if different to the affiliations), Mark McCarthy: Genentech, South San Francisco, CA, USA.

²⁶/⁷These authors jointly supervised the work.

Author Contributions

These authors contributed to the drafting of the manuscript: P. Surendran, E.V.F., N.L., I.N., A.C.M., B.M.P., E.B., D.I.C., D.L., P.B.M., and J.M.M.H. The following authors were involved in the central analyses: P. Surendran, E.V.F., N.L., I.N., S. Karthikeyan, J. Cook, D.J.L., F.D., C.N.-C., P.B.M., and J.M.M.H. All authors critically reviewed and approved the final version of the manuscript. The following authors performed bioinformatics analyses: S. Karthikeyan, L.C., B.M., C.Y., A.T.K., J.H.C., B.P.P., I.D.S., C.P.C., J.M.E., A.A., E.B.F., C.N.F., L.A.L., D.S.P., J.R.S., S. Burgess, M.K., J.P., E.Y., M.R.B., M.T., P.B.M., and J.M.M.H. Study Analysts: J.N.H., A.G., V.T., G. Thorleifsson, B.P.P., J.M.E., A.A., P.L.A., L.F.B., J.C.B., V.S.B., J.A.B., E.W.D., F.D., S.F.N., J.D.F., C.F., T.F., H.G., O.G., F.G., D.F.G., X.G., S.E.H., A.S.H., A.H., J.E.H., S.H., S. Kanoni, J.K., M.G.L., R.L., J. Lindström, Y.L., J. Luan, A.M., G.M., N.G.M., H.M., C.M., D.M., M. Müller-Nurasyid, G.P., M.P., A.P., R. Rainer, M. Richard, T.G.R., N. Sepúlveda, X.S., A.V.S., J.A.S., A.S., P. Sulem, S. Thériault, U.T., S. Trompet, T.V.V., D.R.V.E., G.V., S.W., S.M.W., J.Y., R.Y., B.Y., W. Zhang, J.Z., W. Zhao (UPenn), W. Zhao (UMich), E.E., L.L.B., K.C., T.L., I.L., M.N., N.W.R., M. Reedik, T.S., I.T., H.R.W., H.Z., S.F., B.J.H., M.I.M., T.D.S., A.S.B., H.H., C. Liu, A.K.M., A.P.M., A.C.M., D.I.C., and J.M.M.H. Study Principal Investigators (PI) or Co-PIs: N.F., D.O.M., I.B., P.S.B., C.C., J. Connell, A.F.D., J.M.G., N.G., T.H., F. Karpe, H.A.K., K.K., M. Moitry, C.N.A.P., O. Pedersen, N.P., A.R., F.R., P. J. Sever, E. Tönu, C.J.W., P. Almgren, P. Amouyel, F.W.A., A.I.B., M.B., M.L.B., E.P.B., J.E.B., J.C.C., Y.I.C., R.C., D.C., A.C., G.D.S., R.A.d.B., I.J.D., G.D., P.D., E.D.A., P.E., S.B.F., J.F., I.F., M.F., P.W.F., P.F., G.G., T.R.G., L.G., V.G., T.B.H., C.H., B.J.H., K.H., E.L., T.J., M.J., J.W.J., S.L.K., F. Kee, J.S.K., C.K., L.J.L., L. Lind, R.J.F.L., A.a.S.M., L.M., M.I.M., O.M., K.L.M., A.D.M., B.G.N., M.O., C.J.P., S.P., W.P., O. Polasek, D.J.P., A.M.P., M.A.P., C.L.R., K.R., P.M.R., O.R., F.R.R., J.I.R., I.R., V.S., N.J.S., N. Sattar, W.H.S., B.H.S., N. Soranzo, T.D.S., J.M.S., S.S., K.D.T., L.A.T., N.J.T., M.D.T., P.v.d.H., P.v.d.M., V.S.R., N.V., J.V., U.V., D.R. Weir, E.Z., F.J.C., N.J.W., C. Langenberg, M.T., A.S.B., M.J.C., J.D., T.L.E., A.M.H., C.M.L., A.P.M., C.O., B.M.P., D.S., K.S., E.B., D.I.C., D.L., P.B.M., and J.M.M.H. Study phenotyping: A.T.K., J.D.F., M.G.L., Y.L., S.A., E.A., S. Blankenberg, R.d.M., M.D., G.E., A.F., M.L.G.-G., G. Hallmans, G. Heiss, P.J., E.K., A.K., K.K., T.L., L. Lannfelt, W.L., L.W.M., M.N., G.J.P., K.L.R., M. Reedik, F.R., R. Rettig, J.R., P.J. Schreiner, E.L.S., J.S., G. Thorgeirsson, E. Trabetti, T.T., S.T.T., I.T., I.V., A.V., P. Amouyel, J.E.B., J.C.C., Y.I.C., R.A.d.B., J.F., G.G., V.G., B.J.H., F. Kee, J.S.K., L. Lind, R.J.F.L., O.M., W.P., O. Polasek, P.M.R., I.R., N. Sattar, W.H.S., T.D.S., J.M.S., P.v.d.H., P.v.d.M., N.V., J.V., D.R. Weir, B.M.P., D.I.C., and D.L.

Competing Interests

The following authors affiliated with deCODE genetics/Amgen Inc. are employed by the company: Vinicius Tragante, Gudmar Thorleifsson, Anna Helgadóttir, Patrick Sulem, Gudmundur Thorgeirsson, Hilma Holm, Daniel F. Gudbjartsson, Unnur Thorsteinsdóttir, Kari Stefansson. Bruce Psaty serves on the Steering Committee of the Yale Open Data Access Project funded by Johnson & Johnson. John Danesh reports grants, personal fees and non-financial support from Merck Sharp & Dohme (MSD), grants, personal fees and non-financial support from Novartis, grants from Pfizer, and grants from AstraZeneca outside the submitted work. Adam Butterworth reports grants outside of this work from AstraZeneca, Biogen, Merck, Novartis, and Pfizer and personal fees from Novartis. Veikko Salomaa has participated in a conference trip sponsored by Novo Nordisk and received an honorarium for participating in an advisor board meeting, outside the present study. He also has ongoing research collaboration with Bayer Ltd, outside the present study. Dennis Mook-Kanamori is a part-time clinical research consultant for Metabolon, Inc. Mark I. McCarthy has served on advisory panels for Pfizer, Novo Nordisk, Zoe Global, has received honoraria from Merck, Pfizer, Novo Nordisk and Eli Lilly, and research funding from Abbvie, Astra Zeneca, Boehringer Ingelheim, Eli Lilly, Janssen, Merck, NovoNordisk, Pfizer, Roche, Sanofi Aventis, Servier, and Takeda. As of June 2019, he is an employee of Genentech, and a holder of Roche stock. Eric B. Fauman is an employee of and owns stock in Pfizer, Inc. Mark J. Caulfield is Chief Scientist for Genomics England, a UK Government company. Joanna M. M. Howson became a full-time employee of Novo Nordisk, and I.N. became a full-time employee of Gilead during revision of the manuscript.

Genetic studies of blood pressure (BP) to date have mainly analyzed common variants (minor allele frequency, $MAF > 0.05$). In a meta-analysis of up to >1.3 million participants, we discovered 106 new BP-associated genomic regions and 87 rare ($MAF < 0.01$) variant BP associations ($P < 5 \times 10^{-8}$), of which 32 were in new BP-associated loci and 55 were independent BP-associated SNVs within known BP-associated regions. Average effects of rare variants (44% coding) were ~8 times larger than common variant effects and indicate potential candidate causal genes at new and known loci (e.g. *GATA5*, *PLCB3*). BP-associated variants (including rare and common) were enriched in regions of active chromatin in fetal tissues, potentially linking fetal development with BP regulation in later life. Multivariable Mendelian randomization suggested possible inverse effects of elevated systolic and diastolic BP on large artery stroke. Our study demonstrates the utility of rare variant analyses for identifying candidate genes and the results highlight potential therapeutic targets.

Increased blood pressure (BP) is a major risk factor for cardiovascular disease (CVD) and related disability worldwide¹. Its complications are estimated to account for ~10.7 million premature deaths annually¹. Genome-wide association studies (GWAS) and exome array-wide association studies (EAWAS) have identified over 1,000 BP-associated single nucleotide variants (SNVs)^{2–19} for this complex, heritable, polygenic trait. The majority of these are common SNVs ($MAF > 0.05$) with small effects on BP. Most reported associations involve non-coding SNVs, and due to linkage disequilibrium (LD) between common variants, these studies provide limited insights into the specific causal genes through which their effects are mediated. The exome array was designed to facilitate analyses of rare coding variants ($MAF < 0.01$) with potential functional consequences. Over 80% of SNVs on the array are rare, ~6% are low frequency ($0.01 < MAF < 0.05$), and ~80% are missense, *i.e.* the variants implicate a candidate causal gene through changes to the amino acid sequence. Previously, using the exome array, we identified four BP loci with rare variant associations (*RBM47*, *COL21A1*, *RRAS*, *DBH*)^{13,14} and a rare nonsense BP variant in *ENPEP*, encoding an aminopeptidase with a known role in BP regulation¹³. These findings confirmed the utility of rare variant studies for identifying potential causal genes. These rare variant associations had larger effects on BP (typically ~1.5 mmHg per minor allele) than common variants identified by previous studies (typically ~0.5 mmHg per minor allele), many of which had power to detect common variants with large effects. Here, we combine the studies from our previous two exome array reports with additional studies, including the UK Biobank (UKBB) study, to analyze up to ~1.319 million participants and investigate the role of rare SNVs in BP regulation.

Results

We performed an EAWAS and a rare variant GWAS (RV-GWAS) of imputed and genotyped SNVs to identify variants associated with BP traits, hypertension (HTN), and inverse normal transformed systolic BP (SBP), diastolic BP (DBP), and pulse pressure (PP) using (i) single variant analysis and (ii) a gene-based test approach. An overview of our study design for both the EAWAS and for the RV-GWAS is provided in Figure 1.

Blood pressure associations in the EAWAS

We performed a discovery meta-analysis to identify genetic variants associated with BP in up to ~1.32 million individuals. To achieve this, we first performed a meta-analysis of 247,315 exome array variants in up to 92 studies (870,217 participants, including UKBB) for association with BP, Stage 1 (Fig. 1, Methods, and Supplementary Information). There were 362 BP loci known at the time of the analysis (Supplementary Table 1), 240 of which were covered on the exome array. To improve statistical power for discovery for a subset of variants significant in Stage 1 at $P < 5 \times 10^{-8}$ outside of the known BP regions (Supplementary Table 1a), we requested summary association statistics from three additional studies (Million Veteran Program (MVP), deCODE, and GENOA). We then performed meta-analyses of the three data request studies and Stage 1 results to discover novel variants associated with BP. In total, 343 SNVs (200 genomic regions; Methods) were associated ($P < 5 \times 10^{-8}$) with one or more BP traits in the Stage 2 single variant European (EUR) EAWAS meta-analyses involving up to ~1.168 million individuals (Table 1, Fig. 2, Supplementary Table 2, and Supplementary Information). A further seven SNVs (seven genomic regions) were only associated ($P < 5 \times 10^{-8}$) in the pan-ancestry (PA) meta-analyses of ~1.319 million individuals (Supplementary Table 2). All 350 SNV-BP associations were novel at the time of analysis (204 loci), 220 have subsequently been reported^{20,21}, and 130 SNVs (99 loci) remain novel, including nine rare and 13 low-frequency SNVs (Fig. 2, Supplementary Table 2, Supplementary Fig. 1).

All nine novel rare BP-associated SNVs identified in the EAWAS were conditionally independent of common variant associations within the respective regions (Supplementary Table 3) using the multi-SNP-based conditional and joint association analysis (GCTA v1.91.4)²² with the Stage 1 EUR EAWAS results (Methods and Supplementary Table 4). In addition to the rare variants, there were 147 additional distinct ($P < 1 \times 10^{-6}$) common SNV-BP associations (46% were missense variants), and 18 distinct low-frequency SNVs (89% were missense). Approximately 59% of the distinct BP-associated SNVs were coding or in strong LD ($r^2 > 0.8$) with coding SNVs. In total, 42 of the 99 novel loci had two or more distinct BP-associated SNVs in the conditional analyses. Of the 50 loci that were previously identified using UKBB^{16,17} and were on the exome array, 43 replicated at $P < 0.001$ (Bonferroni correction for 50 known variants) in samples independent of the original discovery (Supplementary Table 5).

Blood pressure associations from EUR RV-GWAS

We tested a further 29,454,346 (29,404,959 imputed and 49,387 genotyped) rare SNVs for association with BP in 445,360 UKBB participants²³ using BOLT-LMM²⁴ (Fig. 1 and Methods). The SNVs analyzed as part of the EAWAS were not included in the RV-GWAS. Similar to EAWAS, within RV-GWAS we performed a single discovery meta-analyses to identify rare SNVs associated with BP. In Stage 1 (UKBB), 84 rare SNVs outside of the known BP loci (at the time of our analyses) were associated with one or more BP traits at $P < 1 \times 10^{-7}$ (Supplementary Table 6). Additional data were requested from MVP for the 84 BP-associated SNVs in up to 225,112 EUR from the MVP, and 66 were available. Meta-analyses of Stage 1 (UKBB) and results obtained from MVP were performed for novel rare variant discovery. We identified 23 unique rare SNVs associated with one or more BP traits

($P < 5 \times 10^{-8}$) with consistent direction of effects in a meta-analysis of UKBB and MVP (min $P_{\text{heterogeneity}} = 0.02$) (Table 1, Fig. 2, Supplementary Table 7, and Supplementary Fig. 1). Two of the SNVs, rs55833332 (p.Arg35Gly) in *NEK7* and rs200383755 (p.Ser19Trp) in *GATA5*, were missense. Eleven rare SNVs were genome-wide significant in UKBB alone but were not available in MVP and await further support in independent studies (Supplementary Table 7).

Rare and low frequency variant associations at established BP loci

It is difficult to prioritize candidate genes at common variant loci for functional follow up. We believe analysis of rare (MAF < 0.01) and very low frequency coding variants (MAF 0.02) in known loci may provide further support for or identify a candidate causal gene at a locus. Twelve of the 240 BP-associated regions had one or more conditionally independent rare variant associations ($P < 10^{-6}$ in the GCTA joint model of the EUR Stage 1 EAWAS; Methods, Table 2, and Supplementary Table 3). A further nine loci had one or more conditionally independent BP-associated SNVs with MAF ≥ 0.02 (Table 2 and Supplementary Table 8). In total, 183 SNVs (rare and common) across 110 known loci were not identified previously.

We used FINEMAP²⁵ to fine-map 315 loci known at the time of our analysis and available in UKBB GWAS, which provides dense coverage of genomic variation not available on the exome array. Of these, 36 loci had one or more conditionally independent rare variant associations (Supplementary Table 8), and 251 loci had multiple common variants associations. We also replicated rare variant associations that we reported previously^{13,14} at *RBM47*, *COL21A1*, *RRAS*, and *DBH* ($P < 5 \times 10^{-5}$) in UKBB (independent of prior studies). Overall, from both FINEMAP and GCTA, we identified 40 loci with one or more rare SNV associations, independent of previously reported common variant associations (Table 3, Fig. 2, Supplementary Table 8, and Supplementary Information).

We note that, of 256 known variants identified without UKBB participants (Supplementary Table 1a), 229 replicated at $P < 1.95 \times 10^{-4}$ (Bonferroni adjusted for 256 variants) in UKBB.

Gene-based tests to identify BP-associated genes

To test whether rare variants in aggregate affect BP regulation, we performed gene-based tests for SBP, DBP, and PP using SKAT²⁶ (<https://genome.sph.umich.edu/wiki/RareMETALS>), including SNVs with MAF ≥ 0.01 that were predicted by VEP²⁷ to have high or moderate impact (Methods). We performed separate analyses within the Stage 1 EAWAS and the UKBB RV-GWAS. Six genes in the EAWAS (*FASTKD2*, *CPXM2*, *CENPJ*, *CDC42EP4*, *OTOP2*, *SCARF2*) and two in the RV-GWAS (*FRY*, *CENPJ*) were associated with BP ($P < 2.5 \times 10^{-6}$, Bonferroni adjusted for ~20,000 genes) and were outside known and new BP loci (Supplementary Tables 1 and 9). To ensure these associations were not attributable to a single (sub-genome-wide significant) rare variant, we also performed SKAT tests conditioning on the variant with the smallest P -value in the gene (Methods and Supplementary Table 9). *FRY* had the smallest conditional P -value ($P = 0.0004$), but did not pass our pre-determined conditional significance threshold (conditional SKAT $P < 0.0001$;

Methods), suggesting that all gene associations are due to single (sub-genome-wide significant) rare variants and not due to the aggregation of multiple rare variants.

Amongst the known loci, five genes (*NPR1*, *DBH*, *COL21A1*, *NOX4*, *GEM*) were associated with BP due to multiple rare SNVs independent of the known common variant associations (conditional $P = 1 \times 10^{-5}$; Methods, Supplementary Information, and Supplementary Table 9) confirming the findings in the single variant conditional analyses above (Supplementary Table 8).

We also performed gene-based tests using a MAF = 0.05 threshold to assess sensitivity to the MAF = 0.01 threshold. The results were concordant with the MAF = 0.01 threshold findings, and two new genes (*PLCB3* and *CEP120*) were associated with BP due to multiple SNVs and were robust to conditioning on the top SNV in each gene (Supplementary Information and Supplementary Table 9).

Rare variant BP associations

In total, across the EAWAS and the RV-GWAS, there were 32 new BP-associated rare variants spanning 18 new loci (Table 1 and Fig. 2). Of these 32, five (representing five loci) were genome-wide significant for HTN, 22 (ten loci) for SBP, 14 (six loci) for DBP, and 15 (ten loci) for PP (Supplementary Tables 1, 2, 3, 6, and 7). Ten of the new rare variants were missense. Within previously reported loci, there were 55 independent rare-variant associations (representing 40 loci) from either the EAWAS or RV-GWAS, making a total of 87 independent rare BP-associated SNVs. We identified 45 BP-associated genes, eight of which were due to multiple rare variants and independent of common variant associations ($P < 1 \times 10^{-4}$, Methods). Twenty-one rare variants were located within regulatory elements (e.g. enhancers), highlighting genetic influence on BP levels through gene expression (Fig. 2). The rare variants contributed to BP variance explained (Supplementary Information).

Power calculations are provided in the Supplementary Information and show that our study had 80% power to detect an effect of 0.039 SD for a MAF = 0.01 (Extended Data Fig. 1). As anticipated, given statistical power, some rare variants displayed larger effects on BP regulation than common variants (Fig. 2 and Supplementary Tables 3, 7, and 8); mean effects of rare SNVs for SBP and DBP were ~7.5 times larger than common variants (mean effect ~0.12 SD/minor allele for rare SNVs, ~0.035 SD/minor allele for low-frequency and ~0.016 SD/minor allele for common SNVs) and for PP were 8.5 times larger for rare variants compared to common (mean effect ~0.135 SD/minor allele for rare SNVs, ~0.04 SD/minor allele for low-frequency and ~0.016 SD/minor allele for common SNVs). Our study was exceptionally well-powered to detect common variants (MAF > 0.05) with similarly large effects but found none, consistent with earlier BP GWAS and genetic studies of some other common complex traits^{28,29,36}.

Overlap of rare BP associations with monogenic BP genes

Twenty-four genes are reported in ClinVar to cause monogenic conditions with hypertension or hypotension as a primary phenotype. Of these, three (*NR3C2*, *AGT*, *PDE3A*) were associated with BP in SKAT tests in the EAWAS ($P < 0.002$, Bonferroni adjusted for 24

tests; Supplementary Table 10). These genes also had genome-wide significant SNV-BP associations in the EAWAS and/or RV-GWAS (Supplementary Table 10).

Functional annotation of rare BP-associated SNVs

None of the BP-associated rare SNVs (from known or novel loci) had been previously reported as expression quantitative trait loci (eQTL) in any tissue ($P > 5 \times 10^{-8}$; Supplementary Table 11 and Methods). We used GTEx v7 data to examine in which tissues the genes closest to the rare BP-SNVs were expressed (Extended Data Fig. 2 and Supplementary Table 4). Many of the eQTL gene transcripts were expressed in BP-relevant tissues (e.g. kidney, heart, and arteries). We observed significant enrichment (Bonferroni adjusted $P < 0.05$) in liver, kidney, heart left ventricle, pancreas, and brain tissues, where the BP genes were down-regulated. In contrast, the BP genes were up-regulated in tibial artery, coronary artery, and aorta (Extended Data Fig. 3). There were 33 genes at 30 known loci with novel BP rare variants (from Supplementary Table 12); distinct known common BP variants at these known loci were eQTLs for 52% of these genes, providing additional evidence that the rare variants implicate plausible candidate genes (Supplementary Table 12).

We tested whether genes near rare BP-associated SNVs were enriched in gene sets from Gene Ontology (GO), KEGG, Mouse Genome Informatics (MGI), and Orphanet (Methods and Supplementary Table 4). These (rare variant) genes from both known and novel loci were enriched in BP-related pathways (Bonferroni adjusted $P < 0.05$; Methods and Supplementary Table 13), including “regulation of blood vessel size” (GO) and “renin secretion” (KEGG). Genes implicated by rare SNVs at known loci were enriched in “tissue remodeling” and “artery aorta” (GO). Genes implicated by rare SNVs at new BP-loci were enriched in rare circulatory system diseases (that include hypertension and rare renal diseases) in Orphanet.

Potential therapeutic insights from the rare BP-associated SNVs

Twenty-three of the genes near rare or low-frequency BP-associated variants in novel and known loci were potentially druggable as suggested by the “druggable genome”³⁰ (Supplementary Information and Supplementary Tables 4 and 14). Six genes (four with rare variants) are already drug targets for CVD conditions, while 15 others are in development or used for other conditions. As an example, the renin-angiotensin-aldosterone system (RAAS) is one of the principal homeostatic mechanisms for BP control, and aldosterone is the main mineralocorticoid (secreted by adrenal glands) and binds receptors, including *NR3C2*, resulting in sodium retention by the kidney and increased potassium excretion. Spironolactone is an aldosterone antagonist widely used in heart failure and as a potassium-sparing anti-hypertensive medication that targets *NR3C2* (Open targets: <https://www.opentargets.org>).

Overlap of new BP-associations with metabolites

To identify novel BP variants that are metabolite QTLs, we performed *in silico* lookups of new sentinel and conditionally independent BP variants for association with 913 plasma metabolites measured using the Metabolon HD4 platform in ~14,000 individuals (Methods

and Supplementary Table 4). Nine BP-associated variants were associated with 25 metabolites ($P < 5 \times 10^{-8}$) involved in carbohydrate, lipids, cofactors and vitamins, nucleotide (cysteine), and amino acid metabolism (Supplementary Table 15), while 11 were unknown.

We performed MR analyses to assess the influence of the 14 known metabolites (Supplementary Table 15) on BP. Lower levels of 3-methylglutaryl carnitine(2) (acyl carnitines involved in long-chain fatty acid metabolism in mitochondria and in leucine metabolism) were significantly associated with increased DBP ($P < 0.003$, 0.05/14 metabolites; Supplementary Table 16). There was no suggestion of reverse causation, i.e. BP did not affect 3-methylglutaryl carnitine(2) ($P > 0.04$; Supplementary Table 16). We further tested whether the association with 3-methylglutaryl carnitine(2) was due to pleiotropic effects of other metabolites in a multivariable MR framework, but found it was still causally associated with DBP (Supplementary Information and Supplementary Table 16).

New BP-associated SNVs are gene eQTLs across tissues

Sentinel variants from 66 new BP loci were associated ($P < 5 \times 10^{-8}$) with gene expression (or had $r^2 > 0.8$ in 1000G EUR with eQTLs) in publicly available databases (Methods and Supplementary Tables 4 and 11). We performed colocalization for 49 of the 66 BP loci (169 genes) with significant eQTLs available in GTEx v7, jointly across all 48 tissues and the BP traits using HyPrColoc³¹ (Methods), to verify that the eQTL and BP-SNV associations were due to the same SNVs and not due to LD or spurious pleiotropy³². The BP associations and eQTL colocalized at 17 BP loci with a single variant (posterior probability, $PPa > 0.6$), i.e. the expression and BP associations were due to the same underlying causal SNV (Fig. 3 and Supplementary Table 17). A further 10 loci had $PPa > 0.6$ for colocalization of BP associations and eQTL for multiple nearby genes (Fig. 3). Colocalization analyses were also performed for the 35 eQTLs in whole blood from the Framingham Heart Study, and five additional loci were consistent with a shared SNV between BP and gene expression (Supplementary Table 17).

Given the central role of the kidney in BP regulation, we investigated if BP-associated SNVs from the EAWAS were kidney eQTLs using TRANScriptome of renaL humAn TissuE study and The Cancer Genome Atlas study ($n = 285$; Methods^{33,34}). We observed significant eQTL associations ($P < 5 \times 10^{-8}$) at three newly identified BP loci (*MFAP2*, *NFU1*, and *AAMDC*, which were also identified in GTEx) and six at previously published loci (*ERAPI1*, *ERAP2*, *KIAA0141*, *NUDT13*, *RP11-582E3.6*, and *ZNF100*; Supplementary Table 18).

New BP-associated SNVs are pQTLs

Eighteen BP loci had sentinel variants (or were in LD with BP SNVs, $r^2 > 0.8$ in 1000G EUR) that were also protein QTL (pQTL) in plasma. Across the 18 loci, BP-SNVs were pQTLs for 318 proteins (Supplementary Table 19). Low-frequency SNVs in *MCL1* and *LAMA5* were cis-pQTL for MCL1 and LAMA5, respectively. The BP-associated SNV, rs4660253, is a cis-pQTL and cis-eQTL for *TIE1* across eight tissues in GTEx including heart (Fig. 3 and Supplementary Table 17). The DBP-associated SNV, rs7776054, is in

strong LD with rs9373124, which is a trans-pQTL for erythropoietin, a hormone mainly synthesized by the kidneys, which has links to hypertension.

Pathway and enrichment analyses

The over-representation of rare and common BP SNVs in DNaseI-hypersensitive sites (DHS), which mark open chromatin, was tested using GARFIELD (Methods and Supplementary Table 4). The most significant enrichment in DHS hotspots for SBP-associated SNVs was in fetal heart tissues, with an ~3-fold enrichment compared to ~2-fold in adult heart (Fig. 3 and Supplementary Information). This difference in enrichment was also reflected in fetal muscle compared to adult muscle for SBP-associated SNVs. The most significant enrichment for DBP- and PP-associated SNVs (~3-fold) was in blood vessels (Fig. 3 and Supplementary Information). There was also enrichment across SBP, DBP and PP in fetal and adult kidney and fetal adrenal gland. In support, complementary enrichment analyses with FORGE (Methods) showed similar enrichments including in fetal kidney and fetal lung tissues (Z -score = 300; Supplementary Table 13 and Supplementary Information).

Mendelian randomization with CVD

Twenty-six new BP loci were also associated with cardiometabolic diseases and risk factors in PhenoScanner³⁵ (<http://www.phenoscanter.medschl.cam.ac.uk>) (Methods, Fig. 3, Supplementary Information, and Supplementary Tables 4, 20, and 21). Given that BP is a key risk factor for CVD, we performed Mendelian randomization (MR) analyses to assess the causal relationship of BP with any stroke (AS), ischemic stroke (IS), large artery stroke (LAS), cardio-embolic stroke (CE), small vessel stroke (SVS), and coronary artery disease (CAD) using all the distinct BP-associated SNVs from our study (both known and new; Supplementary Table 4 and Methods). BP was a predictor of all stroke types analyzed and CAD (Fig. 4 and Supplementary Fig. 4). Notably, SBP had the strongest effect on all CVD phenotypes, with the most profound effect on LAS, increasing risk by >2-fold per SD (Supplementary Table 22). BP had weakest effect on CE, which may reflect the greater role of atrial fibrillation versus BP in CE risk. Multi-variable MR analyses, including both SBP and DBP, showed that the effect of DBP attenuated to zero once SBP was accounted for (consistent with observational studies³⁷), except for LAS (Fig. 4, Supplementary Table 22, and Methods), where SBP/DBP had a suggestive inverse relationship, perhaps reflecting arterial stiffening. An inverse relationship between DBP and stroke above age 50 years has also been reported³⁷.

Discussion

Unlike most previous BP studies that focused primarily on common variant associations, the novelty of this investigation is the extensive analysis of rare variants, both individually and in aggregate within a gene. Many of the new rare variants are located in genes that potentially have a role in BP regulation, as evidenced by support from existing mouse models (21 genes) and/or have previously been implicated in monogenic disorders (11 genes) whose symptoms include hyper-/hypotension or impaired cardiac function/development (Supplementary Table 12). For example, rs139600783 (p.Pro274Ser) was associated with increased DBP and is located in the *ARHGAP31* gene that causes Adams-

Oliver syndrome, which can be accompanied by pulmonary hypertension and heart defects. A further three (of the six) genes that cause Adams-Oliver syndrome are located in BP-associated loci (*DLL4*¹⁶, *DOCK6*^{13,15}, and *NOTCH1*, a new BP locus). A missense variant rs200383755 (p.Ser19Trp, predicted deleterious by SIFT), located in the *GATA5*, encoding a transcription factor, is associated with increased SBP and DBP. *GATA5* mutations cause congenital heart defects, including bicuspid aortic valve and atrial fibrillation, while a *Gata5*-null mouse model had increased SBP and DBP at 90 days³⁸.

Within the known loci, we detected new rare variant associations at several candidate genes, e.g. a rare missense SNV rs1805090 (MAF = 0.0023) in the angiotensinogen (*AGT*) gene was associated with increased BP independently of the known common variant association. *AGT* is known to have an important role in BP regulation, and the variant is predicted to be among the top 1% of most deleterious substitutions³⁹. The established common variant at *FOXS1* was not associated with BP in the conditional analysis, but new rare variants in *FOXS1* (rs45499294, p.Glu74Lys; MAF = 0.0037) and *MYLK2* (rs149972827; MAF = 0.0036; Supplementary Information) were associated with BP. Two BP-associated SNVs (rs145502455, p.Ile806Val; rs117874826, p.Glu564Ala) highlight *PLCB3* as a candidate gene. Phospholipase C is a key enzyme in phosphoinositide metabolism, with *PLCB3* as the major isoform in macrophages⁴⁰, and a negative regulator of VEGF-mediated vascular permeability, a key process in ischemic disease and cancer⁴¹. *PLCβ3* deficiency is associated with decreased atherogenesis, increased macrophage apoptosis in atherosclerotic lesions, and increased sensitivity to apoptotic induction *in vitro*⁴⁰. Variants in *SOS2* have previously been linked to kidney development/function⁴² and also cause Noonan syndromes 1 and 9, which are rare inherited conditions characterized by craniofacial dysmorphic features and congenital heart defects, including hypertrophic cardiomyopathy⁴³. Here we report the rare variant rs72681869 (p.Arg191Pro) in *SOS2* as associated with SBP, DBP, PP, and HTN, highlighting *SOS2* as a candidate gene. Previously, we identified a rare missense BP-associated variant in *RRAS*, a gene causing Noonan syndrome¹³. Our discoveries of rare missense variants at known BP loci provide additional support for candidate genes at these loci.

We report new low-frequency variant associations, such as the missense variant rs45573936 (T>C, Ile216Thr) in *SLC29A1*. The minor allele is associated with both decreased SBP and DBP (Table 1), and the SNV has been shown to affect the function of the encoded protein, equilibrative nucleoside transporter (ENT1)⁴⁴. Best et al.⁴⁵ showed that loss of function of ENT1 caused an (~2.75-fold) increase in plasma adenosine and (~15%) lower BP in mice. Drugs, including dipyridamole and S-(4-Nitrobenzyl)-6-thioinosine (NBTI, NBMPR), are currently used as ENT1 inhibitors for their anti-cancer, anti-cardio, and neuro-protective properties, and our results provide the genetic evidence to indicate that ENT1 inhibition might lower BP in humans.

We found greater enrichment of SBP-associated SNVs in DHS hotspots in fetal vs. adult heart muscle tissue. These results suggest that BP-associated SNVs may influence the expression of genes that are critical for fetal development of the heart. This is consistent with our finding that some BP-associated genes also cause congenital heart defects (see above). Furthermore, *de novo* mutations in genes with high expression in the developing

heart, as well as in genes that encode chromatin marks that regulate key developmental genes, have previously been shown to be enriched in congenital heart disease patients^{46,47}. A recent study of atrial fibrillation genetics, for which BP is a risk factor, described enrichment in DHS in fetal heart⁴⁸. The authors hypothesized that the corresponding genes acting during fetal development increase risk of atrial fibrillation⁴⁸. Together, these data suggest that early development and/or remodeling of cardiac tissues may be an important driver of BP regulation later in life.

The BP measures we have investigated here are correlated; amongst the 107 new genetic BP loci, only two are genome-wide significant across all four BP traits (*RP11-284M14.1* and *VTN*; Fig. 2). None of the new loci were unique to HTN (Fig. 2), perhaps as HTN is derived from SBP and DBP, or perhaps due to reduced statistical power for a binary trait. The results from our study indicate rare BP-associated variants contribute to BP variability in the general population, and their identification has provided information on new candidate genes and potential causal pathways. We have primarily focused on the exome array, which is limited. Future studies using both exome and whole genome sequencing in population cohorts (e.g. UKBB and TOPMed) will lead to identification of further rare variant associations and may advance the identification of causal BP genes across the ~1,000 reported BP loci.

Online Methods

The statistical methods used and analytical packages used are further detailed in the Life Sciences Reporting Summary.

Participants

The cohorts contributing to Stage 1 of the EAWAS comprised 92 studies from four consortia (CHARGE, CHD Exome+, GoT2D:T2DGenes, ExomeBP), and UK Biobank (UKBB) totalling 870,217 individuals of European (EUR, $n = 810,865$), African Ancestry (AA, $n = 21,077$), South Asian (SAS, $n = 33,689$), and Hispanic (HIS, $n = 4,586$) ancestries. Study-specific characteristics, sample quality control and descriptive statistics for the new studies are provided in Supplementary Tables 23 and 24 (and in Supplementary Table 1 and 2 of Surendran *et al.*¹³ (<https://media.nature.com/original/nature-assets/ng/journal/v48/n10/extref/ng.3654-S2.xlsx>) and Supplementary Table 20 of Liu *et al.*¹⁴ (<https://media.nature.com/original/nature-assets/ng/journal/v48/n10/extref/ng.3660-S1.pdf>) for the previously published studies).

For EAWAS, summary association statistics were requested (for the SNVs with $P < 5 \times 10^{-8}$, outside of known BP loci) from the following cohorts: 127,478 Icelanders from deCODE; 225,113 EUR, 63,490 AA, 22,802 HIS, 2,695 NAm (Native Americans), and 4,792 EAS (East Asians) from the Million Veterans Program (MVP); and 1,505 EUR and 792 AA individuals from the Genetic Epidemiology Network of Arteriopathy (GENOA). In total, following the data request, 448,667 individuals of EUR ($n = 354,096$), AA ($n = 63,282$), HIS ($n = 22,802$), NAm ($n = 2,695$), and EAS ($n = 4,792$) ancestries were available for meta-analyses with Stage 1. Study specific characteristics are provided in Supplementary Tables 23 and 24.

Stage 1 of the RV-GWAS used data from 445,360 EUR individuals from UKBB (Supplementary Tables 23 and 24, Supplementary Information), and rare variants were followed up in a data request involving 225,112 EUR individuals from MVP.

All participants provided written informed consent, and the studies were approved by their local research ethics committees and/or institutional review boards. The BioVU biorepository performed DNA extraction on discarded blood collected during routine clinical testing, and linked to de-identified medical records.

Phenotypes

SBP, DBP, PP and HTN were analyzed. Details of the phenotype measures for the previously published studies can be found in the Supplementary Information of the Surendran *et al.* and Liu *et al.* papers (<https://media.nature.com/original/nature-assets/ng/journal/v48/n10/extref/ng.3654-S2.xlsx>; <https://media.nature.com/original/nature-assets/ng/journal/v48/n10/extref/ng.3660-S1.pdf>), and further details of the additional studies are provided in Supplementary Table 24 and Supplementary Information. Typically, the average of two baseline measurements of SBP and DBP were used. For individuals known to be taking BP-lowering medication, 15 and 10 mmHg were added to the raw SBP and DBP values, respectively, to obtain medication-adjusted values⁴⁹. PP was defined as SBP minus DBP after medication adjustment. For HTN, individuals were classified as hypertensive cases if they satisfied at least one of the following criteria: (i) SBP \geq 140 mmHg, (ii) DBP \geq 90 mmHg, or (iii) use of antihypertensive or BP-lowering medication. All other individuals were considered controls. Further information on study-specific BP measurements is provided in Supplementary Table 24. Residuals from the null model obtained after regressing the medication-adjusted trait on the covariates (age, age², sex, BMI, principal components (PCs) to adjust for population stratification, in addition to any study-specific covariates) within a linear regression model were ranked and inverse normalized (Supplementary Information).

Genotyping

The majority of the studies were genotyped using one of the Illumina HumanExome BeadChip arrays (Supplementary Table 24). An exome chip quality control standard operating procedure (SOP: <https://ruder02.u.hpc.mssm.edu/Exome-chip-QC.pdf>) developed by A. Mahajan, N.R.R. and N.W.R. at the Wellcome Trust Centre for Human Genetics, University of Oxford was used by some studies for genotype calling and quality control, while the CHARGE implemented an alternative approach⁵⁰ (Supplementary Table 24 and Supplementary Tables 3 and 21, respectively, of Surendran *et al.*¹³ and Liu *et al.*¹⁴). All genotypes were aligned to the plus strand of the human genome reference sequence (build 37) before any analyses and any unresolved mappings were removed. UKBB, MVP, and deCODE were genotyped using GWAS arrays (Supplementary Table 24).

Exome array meta-analyses

Study-specific analyses were performed to test for the association of 247,315 SNVs with SBP, DBP, PP and HTN in 810,865 individuals of European ancestry (75 EUR studies) and additionally in 59,352 individuals of non-European ancestry comprising of SAS (5 studies),

AA (10 studies), and HIS (2 studies) individuals (Supplementary Information). Study-specific association summaries were meta-analyzed in Stage 1 using an inverse-variance-weighted fixed-effect meta-analyses implemented in METAL⁵². Fixed effect and random effects meta-analyses showed concordant results (Supplementary Table 2). For the binary trait (HTN), we performed sample-size-weighted meta-analysis.

Minimal inflation in the association test statistic, λ , was observed ($\lambda = 1.18$ for SBP, 1.20 for DBP, 1.18 for PP, and 1.18 for HTN in the EUR meta-analyses; and $\lambda = 1.19$ for SBP, 1.20 for DBP, 1.18 for PP, and 1.16 for HTN in the PA meta-analyses). The meta-analyses were performed independently at three centres, and results were found to be concordant across the centres. Following Stage 1, SNVs outside of known BP-associated regions with $P < 5 \times 10^{-8}$ were looked up in individuals from the MVP, deCODE, and GENOA studies (data request). Two meta-analyses of the three additional studies for each trait were performed by two independent analysts, one involving EUR individuals (354,096 participants) only and one PA (448,667 participants). Likewise, two Stage 2 meta-analyses for each trait were performed by two independent analysts, one EUR (1,167,961 participants) and one PA (1,318,884 participants). SNVs with (a conservative) $P < 5 \times 10^{-8}$ in the Stage 2 meta-analysis, with consistent directions of effect in Stage 1 and data request studies and no evidence of heterogeneity ($P > 0.0001$), were considered potentially novel⁵³.

RV-GWAS

Rare SNVs with $P < 5 \times 10^{-8}$ (a widely accepted significance threshold^{54,55}) in the inverse variance-weighted meta-analysis of UKBB and MVP, with consistent directions of effect in Stage 1 and MVP and no evidence of heterogeneity ($P > 0.0001$), were considered potentially novel.

Quality control

As part of the sample QC, plots comparing inverse of the standard error as a function of the square root of study sample size for all studies were manually reviewed for each trait, and phenotype-specific study outliers were excluded. In addition, inflation of test static was manually reviewed for each study and for each phenotype and confirmed minimal or no inflation prior to Stage 1 meta-analyses. For EAWAS and RV-GWAS, we performed our own QC for genotyped variants as we were specifically interested in rare variants and knew that these were most vulnerable to clustering errors. Full details of UKBB QC are provided in the Supplementary Note. To ensure that the variants we reported are not influenced by technical artefacts and not specific to a certain ancestry, we ensured that there was no heterogeneity and also that the variants had consistent direction of effects between Stage 1 and the data request studies (MVP+deCODE+GENOA). In addition, we ensured that the association was not driven by a single study. For variants reported in RV-GWAS and EAWAS, we reviewed the cluster plots for clustering artefacts and removed poorly clustered variants. Lastly, for RV-GWAS, if the variant was available in UKBB whole exome data (~50K individuals), we ensured that the minor allele frequencies were consistent with the imputed MAF despite restricting the reporting of only variant with a good imputation quality (INFO > 0.8).

Definition of known loci

For each known variant, pairwise LD was calculated between the known variant and all variants within the 4-Mb region in the 1000 Genomes phase 3 data restricted to samples of European (EUR) ancestry. Variants with $r^2 > 0.1$ were used to define a window around the known variant. The region start and end were defined as the minimum position and maximum position of variants in LD within the window ($r^2 > 0.1$), respectively. Twelve variants were not in 1000 Genomes, and for these variants, a 500-kb window around the known variant was used. The window was extended by a further 50 kb and overlapping regions were merged (Supplementary Table 1).

Conditional analyses

Within the new BP loci, we defined a region based on LD (Supplementary Table 1) within which conditional analysis was performed (five variants were not in the 1000G panel, and for these we established a 500-kb window definition). Conditional and joint association analysis as implemented in Genome-wide Complex Trait Analysis (GCTA v1.91.4)²² was performed using the EAWAS results to identify independent genetic variants associated with BP traits within newly identified and known regions available in the exome array. We restricted this analysis to the summary data from Stage 1 EUR EAWAS meta-analyses ($n = 810,865$) as LD patterns were modelled using individual level genotype data from 57,718 EUR individuals from the CHD Exome+ consortium. Variants with $P_{\text{joint}} < 1 \times 10^{-6}$ were considered conditionally independent.

We used the UKBB GWAS results and FINEMAP²⁵ v1.1 to fine-map the known BP-associated regions in order to identify rare variants that are associated with BP independently of the known common variants (Supplementary Note; due to lack of statistical power, we did not use UKBB GWAS data alone to perform conditional analyses within the new EAWAS loci). For each known region, we calculated pairwise Pearson correlation for all SNVs within a 5-Mb window of the known SNVs using LDstore v1.1. Z -scores calculated in the UKBB single-variant association analyses were provided as input to FINEMAP along with the correlation matrix for the region. We selected the configuration with the largest Bayes Factor (BF) and largest posterior probability as the most likely causal SNVs. We considered causal SNVs to be significant if the configuration cleared a threshold of $\log_{10}\text{BF} > 5$ and if the variants in the configuration had an unconditional association of $P < 1 \times 10^{-6}$. We examined the validity of the SNVs identified for the most likely configuration by checking marginal association P -values and LD (r^2) within UKBB between the selected variants. For loci that included rare variants identified by FINEMAP, we validated the selected configuration using a linear regression model in R.

Gene-based tests

Gene-based tests were performed using the sequence kernel association test (SKAT)²⁶ as implemented in the rareMETALS package version 7.1 (<https://genome.sph.umich.edu/wiki/RareMETALS>) (which allows for the variants to have different directions and magnitudes of effect) to test whether rare variants in aggregate within a gene are associated with BP traits. For the EAWAS, two gene-based meta-analyses were performed for inverse-normal transformed DBP, SBP, and PP, one of EUR and a second PA including all studies with

single-variant association results and genotype covariance matrices (up to 691,476 and 749,563 individuals from 71 and 88 studies were included in the EUR and PA gene-based meta-analyses, respectively).

In UKBB, we considered summary association results from 364,510 unrelated individuals only. We annotated all SNVs on the exome array using VEP²⁷. A total of 15,884 (EUR) and 15,997 genes (PA) with two or more variants with MAF ≥ 0.01 annotated with VEP as high or moderate effects were tested. The significance threshold was set at $P < 2.5 \times 10^{-6}$ (Bonferroni adjusted for $\sim 20,000$ genes).

A series of conditional gene-based tests were performed for each significant gene. To verify the gene association was due to more than one variant (and not due to a single sub-genome-wide significance threshold variant), gene tests were conditioned on the variant with the smallest P -value in the gene (top variant). Genes with $P_{\text{conditional}} < 1 \times 10^{-4}$ were considered significant, which is in line with locus-specific conditional analyses used in other studies⁵⁶. In order to ensure that gene associations located in known or newly identified BP regions (Supplementary Note and Supplementary Table 1) were not attributable to common BP-associated variants, analyses were conditioned on the conditionally independent known/novel common variants identified using GCTA within the known or novel regions, respectively, for the EAWAS (or identified using FINEMAP for the GWAS). Genes mapping to either known or novel loci with $P_{\text{conditional}} < 1 \times 10^{-5}$, were considered significant. The P -value to identify gene-based association not driven by a single variant was set in advance of performing gene-based tests and was based on an estimation of the potential number of genes that could be associated with BP.

Mendelian randomization with CVDs

We used two-sample MR to test for causal associations between BP traits and any stroke (AS), any ischemic stroke (IS), large artery stroke (LAS), cardioembolic stroke (CE), small vessel stroke (SVS), and coronary artery disease (CAD). All the new and known BP-associated SNVs (including conditionally independent SNVs) listed in Supplementary Tables 2, 3, 5, 7 and 8, were used as instrumental variables (IVs). In addition to trait specific analyses, we performed an analysis of “generic” BP, in which we used the SNVs associated with any of the traits. Where variants were associated with multiple BP traits, we extracted the association statistics for the trait with the smallest P -value (or the largest posterior probability for the known loci). To exclude potentially invalid (pleiotropic) genetic instruments, we used PhenoScanner³⁵ to identify SNVs associated with CVD risk factors, cholesterol (LDL/HDL/triglycerides (TG)), smoking, type 2 diabetes (T2D) and atrial fibrillation (AF) (Supplementary Table 22) and removed these from the list of IVs. We extracted estimates for the associations of the selected instruments with each of the stroke subtypes from the MEGASTROKE PA GWAS results (67,162 cases; 454,450 controls)⁶³ and from a recent GWAS for CAD⁶⁴. We applied a Bonferroni correction ($P < 0.05/6 = 0.0083$) to account for the number of CVD traits. We used the inverse-variance weighting method with a multiplicative random-effects because we had hundreds of IVs for BP⁶⁵. We performed MR-Egger regression, which generates valid estimates even if not all the genetic instruments are valid, as long as the Instrument Strength Independent of Direct Effect

assumption holds⁶⁶. We note that MR-Egger has been shown to be conservative⁶⁶, but has the useful property that the MR-Egger-intercept can give an indication of (unbalanced) pleiotropy, which allowed us to test for pleiotropy amongst the IVs. We used MR-PRESSO to detect outlier IVs⁶⁷. To assess instrument strength, we computed the F-statistic⁶⁸ for the association of genetic variants with SBP, DBP and PP, respectively (Supplementary Information and Supplementary Table 22). We also assessed heterogeneity using the Q-statistic. Although these methods may have different statistical power, the rationale is that, if these methods give a similar conclusion regarding the association of BP and CVD, then we are more confident in inferring that the positive results are unlikely to be driven by violation of the MR assumptions⁶⁹.

Moreover, we used multivariable MR (mvMR) to estimate the effect of multiple variables on the outcome^{65,70}. This is useful when two or more correlated risk factors are of interest, e.g. SBP and DBP, and may help to understand whether both risk factors exert a causal effect on the outcome, or whether one exerts a leading effect on the outcome. Thus, we used multiple genetic variants associated with SBP and DBP to simultaneously estimate the causal effect of SBP and DBP on CVDs.

All analyses were performed using R version 3.4.2 with R packages ‘TwoSampleMR’ and ‘MendelianRandomization’ and “MRPRESSO”.

Metabolite quantitative trait loci and Mendelian randomization analyses

Plasma metabolites were measured in up to 8,455 EUR individuals from the INTERVAL study^{71,72} and up to 5,841 EUR individuals from EPIC-Norfolk⁷³ using the Metabolon HD4 platform. In both studies, 913 metabolites passed QC and were analyzed for association with ~17 million rare and common genetic variants. Genetic variants were genotyped using the Affymetrix Axiom UK Biobank array and imputed using the UK10K+1000Genomes or the HRC reference panel. Variants with INFO > 0.3 and MAC > 10 were analyzed. Phenotypes were log-transformed within each study, and standardized residuals from a linear model adjusted for study-specific covariates were calculated prior to the genetic analysis. Study-level genetic analysis was performed using linear mixed models implemented in BOLT-LMM to account for relatedness within each study, and the study-level association summaries were meta-analyzed using METAL prior to the lookup of novel BP variants for association with metabolite levels.

The same methodology for MR analyses as implemented for CVDs was also adopted to test the effects of metabolites on BP. Causal analyses were restricted to the list of 14 metabolites that overlapped our BP-associations and were known. We used a Bonferroni significance threshold ($P < 0.05/14 = 0.0036$), adjusting for the number of metabolites being tested. We also tested for a reverse causal effect of BP on metabolite levels. The IVs for the BP traits were the same as those used for MR with CVDs. For the mvMR analysis of metabolites with BP, we included 3-methylglutarylcarnitine(2) and the three metabolites that shared at least one IV with 3-methylglutarylcarnitine(2) in the mvMR model. A union set of genetic IVs for all the metabolites were used in the mvMR model to simultaneously estimate the effect size of each metabolite on DBP.

Colocalization of BP associations with eQTLs

Details of kidney-specific eQTL are provided in Supplementary Information. Using the phenoscanner lookups to prioritize BP regions with eQTLs in GTEx version 7, we performed joint colocalization analysis with the HyPrColoc package in R³¹ (<https://github.com/jrs95/hyprcoloc>; regional colocalization plots, <https://github.com/jrs95/gassocplot>). HyPrColoc approximates the COLOC method developed by Giambartolomei et al.⁶² and extends it to allow colocalization analyses to be performed jointly across many traits simultaneously and pinpoint candidate shared SNV(s). Analyses were restricted to SNVs present in all the datasets used (for GTEx data this was 1 Mb upstream and downstream of the center of the gene probe), data were aligned to the same human genome build 37 and strand, and a similar prior structure as the colocalization analysis with cardiometabolic traits was used ($P = 0.0001$ and $\gamma = 0.99$).

Gene set enrichment analyses

In total, 4,993 GO biological process, 952 GO molecular function, 678 GO cellular component, 53 GTEx, 301 KEGG, 9537 MGI, and 2645 Orphanet gene sets were used for enrichment analyses (Supplementary Information).

We restricted these analyses to the rare BP-associated SNVs (Supplementary Table 4). For each set of gene sets, the significance of the enrichment of the genetically identified BP genes was assessed as the Fisher's exact test for the over-abundance of BP genes in the designated gene set based on a background of all human protein coding genes or, in the case of the MGI gene sets, a background of all human protein-coding genes with an available knock-out phenotype in the MGI database.


Results were deemed significant if after multiple testing correction for the number of gene sets in the specific set of gene sets the adjusted P -value < 0.05 . Results were deemed suggestive if the adjusted P -value was between 0.05 and 0.1.

Functional enrichment using BP-associated variants

To assess enrichment of GWAS variants associated with the BP traits in regulatory and functional regions in a wide range of cell and tissue types, we used GWAS Analysis of Regulatory or Functional Information Enrichment with LD Correction (GARFIELD). The GARFIELD method has been described extensively elsewhere^{76,77}. In brief, GARFIELD takes a non-parametric approach that requires GWAS summary statistics as input. It performs the following steps: (i) LD-pruning of input variants; (ii) calculation of the fold enrichment of various regulatory/functional elements; and (iii) testing these for statistical significance by permutation testing at various GWAS significance levels, accounting for MAF, the distance to the nearest transcription start site, and the number of LD proxies of the GWAS variants. We used the SNVs from the full UKBB GWAS of BP traits as input to GARFIELD (Supplementary Table 4).


Extended Data

Power Estimation




Extended Data Fig. 1. Power estimation for stage 2 meta-analyses

Power calculations were performed assuming that, for any given variant, there were 1,318,884 individuals for EAWAS PA analyses, 1,164,961 participants for EAWAS EA analyses, and 670,472 participants for RV-GWAS analyses. Calculations were performed in R (https://genome.sph.umich.edu/wiki/Power_Calculations:_Quantitative_Traits).



Extended Data Fig. 2. Expression of genes implicated by the rare SNVs in GTEx v7 tissues
 We used FUMA GWAS to perform these analyses. We included genes closest to the identified rare variants from the EAWAS and the RV-GWAS.



Extended Data Fig. 3. Tissue enrichment of rare variant gene expression levels in GTEx v7
 We used FUMA GWAS to perform these analyses. We included genes closest to the identified rare variants from the EAWAS and the RV-GWAS.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Authors

Praveen Surenran^{#1,2,3,4}, Elena V. Feofanova^{#5}, Najim Lahrouchi^{#6,7,8}, Ioanna Ntalla^{#9}, Savita Karthikeyan^{#1}, James Cook¹⁰, Lingyan Chen¹, Borbala Mifsud^{9,11}, Chen Yao^{12,13}, Aldi T. Kraja¹⁴, James H. Cartwright⁹, Jacklyn N. Hellwege¹⁵, Ayush Giri^{15,16}, Vinicius Tragante^{17,18}, Gudmar Thorleifsson¹⁸, Dajiang J. Liu¹⁹, Bram P. Prins¹, Isobel D. Stewart²⁰, Claudia P. Cabrera^{9,21}, James M. Eales²², Artur Akbarov²², Paul L. Auer²³, Lawrence F. Bielak²⁴, Joshua C. Bis²⁵, Vickie S. Braithwaite^{20,26,27}, Jennifer A. Brody²⁵, E. Warwick Daw¹⁴, Helen R. Warren^{9,21},

Fotios Drenos^{28,29}, Sune Fallgaard Nielsen³⁰, Jessica D. Faul³¹, Eric B. Fauman³², Cristiano Fava^{33,34}, Teresa Ferreira³⁵, Christopher N. Foley^{1,36}, Nora Franceschini³⁷, He Gao^{38,39}, Olga Giannakopoulou^{9,40}, Franco Giulianini⁴¹, Daniel F. Gudbjartsson^{18,42}, Xiuqing Guo⁴³, Sarah E. Harris^{44,45}, Aki S. Havulinna^{45,46}, Anna Helgadottir¹⁸, Jennifer E. Huffman⁴⁷, Shih-Jen Hwang^{48,49}, Stavroula Kanoni^{9,50}, Jukka Kontto⁴⁶, Martin G. Larson^{51,52}, Ruifang Li-Gao⁵³, Jaana Lindström⁴⁶, Luca A. Lotta²⁰, Yingchang Lu^{54,55}, Jian'an Luan²⁰, Anubha Mahajan^{56,57}, Giovanni Malerba⁵⁸, Nicholas G. D. Masca^{59,60}, Hao Mei⁶¹, Cristina Menni⁶², Dennis O. Mook-Kanamori^{53,63}, David Mosen-Ansorena³⁸, Martina Müller-Nurasyid^{64,65,66}, Guillaume Paré⁶⁷, Dirk S. Paul^{1,2,68}, Markus Perola^{46,69}, Alaitz Poveda⁷⁰, Rainer Rauramaa^{71,72}, Melissa Richard⁷³, Tom G. Richardson⁷⁴, Nuno Sepúlveda^{75,76}, Xueling Sim^{77,78}, Albert V. Smith^{79,80,81}, Jennifer A. Smith^{24,31}, James R. Staley^{1,74}, Alena Stanáková⁸², Patrick Sulem¹⁸, Sébastien Thériault^{83,84}, Unnur Thorsteinsdottir^{18,80}, Stella Trompet^{85,86}, Tibor V. Varga⁷⁰, Digna R. Velez Edwards⁸⁷, Giovanni Veronesi⁸⁸, Stefan Weiss^{89,90}, Sara M. Willems²⁰, Jie Yao⁹¹, Robin Young^{1,92}, Bing Yu⁵, Weihua Zhang^{38,39,93}, Jing-Hua Zhao^{1,20,68}, Wei Zhao⁹⁴, Wei Zhao²⁴, Evangelos Evangelou^{38,95}, Stefanie Aeschbacher⁹⁶, Eralda Asllanaj^{97,98}, Stefan Blankenberg^{90,99,100,101}, Lori L. Bonnycastle¹⁰², Jette Bork-Jensen¹⁰³, Ivan Brandslund^{104,105}, Peter S. Braund^{59,60}, Stephen Burgess^{1,36,68}, Kelly Cho^{106,107,108}, Cramer Christensen¹⁰⁹, John Connell¹¹⁰, Renée de Mutsert⁵³, Anna F. Dominiczak¹¹¹, Marcus Dörr^{90,112}, Gudny Eiríksdóttir⁷⁹, Alikei-Eleni Farmaki^{113,114}, J. Michael Gaziano^{106,107,108}, Niels Grarup¹⁰³, Megan L. Grove-Gaona⁵, Göran Hallmans¹¹⁵, Torben Hansen¹⁰³, Christian T. Have¹⁰³, Gerardo Heiss³⁷, Marit E. Jørgensen¹¹⁶, Pekka Jousilahti⁴⁶, Eero Kajantie^{46,117,118,119}, Mihir Kamat^{1,68}, AnneMari Käräjämäki^{120,121}, Fredrik Karpe^{57,122}, Heikki A. Koistinen^{46,123,124}, Csaba P. Kovesdy¹²⁵, Kari Kuulasmaa⁴⁶, Tiina Laatikainen^{46,126}, Lars Lannfelt¹²⁷, I-Te Lee^{128,129,130,131}, Wen-Jane Lee^{132,133}, LifeLines Cohort Study

Rudolf A. de Boer¹⁸², Pim van der Harst^{240,241,242}, Peter van der Meer²⁴², Niek Verweij²⁴⁴

, Allan Linneberg^{134,135}, Lisa W. Martin¹³⁶, Marie Moitry¹³⁷, Girish Nadkarni⁵⁴, Matt J. Neville^{57,122}, Colin N. A. Palmer¹³⁸, George J. Papanicolaou¹³⁹, Oluf Pedersen¹⁰³, James Peters^{1,3,140}, Neil Poulter¹⁴¹, Asif Rasheed¹⁴², Katrine L. Rasmussen³⁰, N. William Rayner^{56,57}, Reedik Mägi¹⁴³, Frida Renström^{70,115}, Rainer Rettig^{90,144}, Jacques Rossouw¹⁴⁵, Pamela J. Schreiner¹⁴⁶, Peter J. Sever¹⁴¹, Emil L. Sigurdsson^{147,148}, Tea Skaaby¹⁴⁹, Yan V. Sun¹⁵⁰, Johan Sundstrom¹⁵¹, Gudmundur Thorgeirsson^{18,80,152}, Tõnu Esko^{143,153}, Elisabetta Trabetti⁵⁸, Philip S. Tsao¹⁵⁴, Tiinamaija Tuomi^{155,156,157}, Stephen T. Turner¹⁵⁸, Ioanna Tzoulaki^{38,95,265}, Ilonca Vaartjes^{159,160}, Anne-Claire Vergnaud³⁸, Cristen J. Willer^{161,162,163}, Peter W.F. Wilson¹⁶⁴, Daniel R. Witte^{165,166,167}, Ekaterina Yonova-Doing¹, He Zhang¹⁶¹, Naheed Aliya¹⁶⁸, Peter Almgren¹⁶⁹, Philippe Amouyel^{170,171,172,173}, Folkert W. Asselbergs^{17,174,175}, Michael R. Barnes^{9,21}, Alexandra I. Blakemore^{28,176}, Michael Boehnke⁷⁷, Michiel L. Bots^{159,160}, Erwin P. Bottinger⁵⁴, Julie E. Buring^{41,177}, John C. Chambers^{38,39,93,178,179}, Yii-Der Ida Chen⁹¹, Rajiv Chowdhury¹, David Conen^{83,180}, Adolfo Correa¹⁸¹, George Davey

Smith⁷⁴, Rudolf A. de Boer¹⁸², Ian J. Deary^{44,183}, George Dedoussis¹¹³, Panos Deloukas^{9,21,50,184}, Emanuele Di Angelantonio^{1,2,3,68,185,186}, Paul Elliott^{38,39,187,188}, EPIC-CVD

Adam S. Butterworth^{1,2,3,68,185}, **John Danesh**^{1,2,3,68,185,186}

, EPIC-InterAct

Claudia Langenberg²⁰, **Panos Deloukas**^{9,21,50,184}, **Mark I. McCarthy**^{56,57,122},

Paul W. Franks^{70,190,191,192}, **Olov Rolandsson**¹⁹¹, **Nicholas J. Wareham**²⁰

, Stephan B. Felix^{90,112}, Jean Ferrières¹⁸⁹, Ian Ford⁹², Myriam Fornage^{5,73}, Paul W.

Franks^{70,190,191,192}, Stephen Franks¹⁹³, Philippe Frossard¹⁴², Giovanni

Gambaro¹⁹⁴, Tom R. Gaunt⁷⁴, Leif Groop^{195,196}, Vilmondur Gudnason^{79,80}, Tamara

B. Harris¹⁹⁷, Caroline Hayward⁴⁷, Branwen J. Hennig^{27,198}, Karl-Heinz

Herzig^{199,200}, Erik Ingelsson^{201,202,203,204}, Jaakko Tuomilehto^{46,205,206,207}, Marjo-

Riitta Jarvelin^{28,38,39,208}, J. Wouter Jukema^{86,209}, Sharon L. R. Kardia²⁴, Frank

Kee²¹⁰, Jaspal S. Kooner^{39,93,179,211}, Charles Kooperberg²¹², Lenore J. Launer¹⁹⁷,

Lars Lind¹⁵¹, Ruth J. F. Loos^{54,213}, Abdulla al Shafi Majumder²¹⁴, Markku

Laakso¹²⁶, Mark I. McCarthy^{56,57,122}, Olle Melander³⁴, Karen L. Mohlke²¹⁵, Alison

D. Murray²¹⁶, Børge Grønne Nordestgaard³⁰, Marju Orho-Melander³⁴, Chris J.

Packard²¹⁷, Sandosh Padmanabhan²¹⁸, Walter Palmas²¹⁹, Ozren Polasek²²⁰,

David J. Porteous^{221,222}, Andrew M. Prentice^{27,223}, Michael A. Province¹⁴, Caroline

L. Relton⁷⁴, Kenneth Rice²²⁴, Paul M. Ridker^{41,177}, Olov Rolandsson¹⁹¹, Frits R.

Rosendaal⁵³, Jerome I. Rotter²²⁵, Igor Rudan²²⁶, Veikko Salomaa⁴⁶, Nilesh J.

Samani^{59,60}, Naveed Sattar¹¹¹, Wayne H.-H. Sheu^{128,129,227,228}, Blair H. Smith²²⁹,

Nicole Soranzo^{230,231,232}, Timothy D. Spector⁶², John M. Starr^{44,233}, Sebert

Sylvain^{234,235,236}, Kent D. Taylor²³⁷, Timo A. Lakka^{71,72,238}, Nicholas J. Timpson⁷⁴,

Martin D. Tobin^{60,239}, Understanding Society Scientific Group

Bram P. Prins¹, **Eleftheria Zeggini**^{245,246}

, Pim van der Harst^{240,241,242}, Peter van der Meer²⁴², Ramachandran S.

Vasan^{51,243}, Niek Verweij²⁴⁴, Jarmo Virtamo⁴⁶, Uwe Völker^{89,90}, David R. Weir³¹,

Eleftheria Zeggini^{245,246}, Fadi J. Charchar^{59,247,248}, Million Veteran Program

Jacklyn N. Hellwege¹⁵, **Ayush Giri**^{15,16}, **Digna R. Velez Edwards**⁸⁷, **Kelly**

Cho^{106,107,108}, **J. Michael Gaziano**^{106,107,108}, **Csaba P. Kovcsy**¹²⁵, **Yan V.**

Sun¹⁵⁰, **Philip S. Tsao**¹⁵⁴, **Peter W. F. Wilson**¹⁶⁴, **Todd L. Edwards**²⁵⁰, **Adriana**

M. Hung²⁵¹, **Christopher J. O'Donnell**²⁵⁷

, Nicholas J. Wareham²⁰, Claudia Langenberg²⁰, Maciej Tomaszewski^{22,249}, Adam

S. Butterworth^{1,2,3,68,185}, Mark J. Caulfield^{9,21}, John Danesh^{1,2,3,68,185,186}, Todd L.

Edwards²⁵⁰, Hilma Holm¹⁸, Adriana M. Hung²⁵¹, Cecilia M. Lindgren^{35,252,253},

Chunyu Liu²⁵⁴, Alisa K. Manning^{108,255}, Andrew P. Morris^{10,252,256}, Alanna C.

Morrison⁵, Christopher J. O'Donnell²⁵⁷, Bruce M. Psaty^{25,258,259,260}, Danish

Saleheen^{1,261,262}, Kari Stefansson^{18,80}, Eric Boerwinkle^{5,263,267}, Daniel I.

Chasman^{41,177,267}, Daniel Levy^{51,264,267}, Christopher Newton-Cheh^{6,7,267}, Patricia

B. Munroe^{9,21,267}, Joanna M. M. Howson^{1,68,265,267}

Rudolf A. de Boer¹⁸², **Pim van der Harst**^{240,241,242}, **Peter van der Meer**²⁴², **Niek Verweij**²⁴⁴

Adam S. Butterworth^{1,2,3,68,185}, **John Danesh**^{1,2,3,68,185,186}

Claudia Langenberg²⁰, **Panos Deloukas**^{9,21,50,184}, **Mark I. McCarthy**^{56,57,122},
Paul W. Franks^{70,190,191,192}, **Olov Rolandsson**¹⁹¹, **Nicholas J. Wareham**²⁰
Bram P. Prins¹, **Eleftheria Zeggini**^{245,246}
Jacklyn N. Hellwege¹⁵, **Ayush Giri**^{15,16}, **Digna R. Velez Edwards**⁸⁷, **Kelly
 Cho**^{106,107,108}, **J. Michael Gaziano**^{106,107,108}, **Csaba P. Kovesdy**¹²⁵, **Yan V.
 Sun**¹⁵⁰, **Philip S. Tsao**¹⁵⁴, **Peter W. F. Wilson**¹⁶⁴, **Todd L. Edwards**²⁵⁰, **Adriana
 M. Hung**²⁵¹, **Christopher J. O'Donnell**²⁵⁷

Affiliations

¹British Heart Foundation Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK ²British Heart Foundation Centre of Research Excellence, University of Cambridge, Cambridge, UK ³Health Data Research UK Cambridge, Wellcome Genome Campus and University of Cambridge, Cambridge, UK ⁴Rutherford Fund Fellow, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK ⁵Human Genetics Center, The University of Texas Health Science Center at Houston, Houston, TX, USA ⁶Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA, USA ⁷Cardiovascular Research Center, Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA ⁸Amsterdam UMC, University of Amsterdam, Heart Center, Department of Clinical and Experimental Cardiology, Amsterdam Cardiovascular Sciences, Amsterdam, The Netherlands ⁹William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK ¹⁰Department of Biostatistics, University of Liverpool, Liverpool, UK ¹¹College of Health and Life Sciences, Hamad Bin Khalifa University, Doha, Qatar ¹²Framingham Heart Study, Framingham, MA, USA ¹³Population Sciences Branch, Division of Intramural Research, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, USA ¹⁴Division of Statistical Genomics, Department of Genetics and Center for Genome Sciences and Systems Biology, Washington University School of Medicine, St. Louis, MO, USA ¹⁵Division of Epidemiology, Department of Medicine, Institute for Medicine and Public Health, Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Tennessee Valley Healthcare System (626)/Vanderbilt University, Nashville, TN, USA ¹⁶Division of Quantitative Sciences, Department of Obstetrics & Gynecology, Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Tennessee Valley Healthcare System (626)/Vanderbilt University, Nashville, TN, USA ¹⁷Department of Cardiology, Division Heart & Lungs, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands ¹⁸deCODE genetics/Amgen, Inc., Reykjavik, Iceland ¹⁹Institute of Personalized Medicine, Penn State College of Medicine, Hershey, PA, USA ²⁰MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge, UK ²¹National Institute for Health Research Barts Cardiovascular Biomedical Research Centre, Queen Mary University of London, London, UK ²²Division of Cardiovascular Sciences, Faculty of Medicine, Biology and Health, University of Manchester, Manchester, UK ²³Joseph J. Zilber School of Public Health, University of Wisconsin, Milwaukee, WI, USA ²⁴Department of

Epidemiology, University of Michigan, Ann Arbor, MI, USA ²⁵Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle, WA, USA ²⁶MRC Nutrition and Bone Health Group, University of Cambridge, Cambridge, UK. Formerly, MRC Human Nutrition Research, Cambridge, UK ²⁷MRC Unit The Gambia at London School of Hygiene & Tropical Medicine, Banjul, The Gambia ²⁸Department of Life Sciences, College of Health and Life Sciences, Brunel University London, London, UK ²⁹Institute of Cardiovascular Science, Faculty of Population Health Sciences, University College London, London, UK ³⁰Department of Clinical Biochemistry, Herlev and Gentofte Hospital, Copenhagen University Hospital, Herlev, Denmark ³¹Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, MI, USA ³²Internal Medicine Research Unit, Pfizer, Cambridge MA, USA ³³Department of Medicine, University of Verona, Verona, Italy ³⁴Department of Clinical Sciences Malmö, Lund University, Malmö, Sweden ³⁵The Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK ³⁶MRC Biostatistics Unit, University of Cambridge, Cambridge, UK ³⁷Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA ³⁸Department of Epidemiology and Biostatistics, MRC Centre for Environment and Health, School of Public Health, Imperial College London, London, UK ³⁹National Institute for Health Research (NIHR) Imperial Biomedical Research Centre, Imperial College London, London, UK ⁴⁰Centre for Genomic Health, Queen Mary University of London, London, UK ⁴¹Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA, USA ⁴²School of Engineering and Natural Sciences, University of Iceland, Reykjavik, Iceland ⁴³The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, LABioMed at Harbor-UCLA Medical Center, Torrance, CA, USA ⁴⁴Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK ⁴⁵Centre for Genomic and Experimental Medicine, University of Edinburgh, Edinburgh, UK ⁴⁶Department of Public Health Solutions, Finnish Institute for Health and Welfare, Helsinki, Finland ⁴⁷MRC Human Genetics Unit, IGMM, University of Edinburgh, Western General Hospital, Edinburgh, UK ⁴⁸Population Sciences, Branch, National Heart, Lung, and Blood Institute, National Institute of Health, Bethesda, MD, USA ⁴⁹Boston University's and Boston University's and National Heart, Lung and Blood Institute's Framingham Heart Study, Framingham, MA, USA ⁵⁰Centre for Genomic Health, Life Sciences, Queen Mary University of London, London, UK ⁵¹Boston University's and National Heart, Lung and Blood Institute's Framingham Heart Study, Framingham, MA, USA ⁵²Biostatistics Department, Boston University School of Public Health, Boston, MA, USA ⁵³Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, The Netherlands ⁵⁴The Charles Bronfman Institute for Personalized Medicine at Mount Sinai, Icahn School of Medicine at Mount Sinai, New York, NY, USA ⁵⁵Division of Epidemiology, Department of Medicine, Vanderbilt-Ingram Cancer Center, Vanderbilt Epidemiology Center, Vanderbilt University School of Medicine, Nashville, TN, USA ⁵⁶Wellcome Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, Oxford, UK ⁵⁷Oxford Centre for Diabetes,

Endocrinology and Metabolism, Radcliffe Department of Medicine, University of Oxford, Oxford, UK ⁵⁸Department of Neurosciences, Biomedicine and Movement Sciences, University of Verona, Verona, Italy ⁵⁹Department of Cardiovascular Sciences, University of Leicester, Leicester, UK ⁶⁰National Institute for Health Research Leicester Biomedical Research Centre, Leicester, UK ⁶¹Department of Data Science, School of Population Health, University of Mississippi Medical Center, Jackson, MS, USA ⁶²Department of Twin Research and Genetic Epidemiology, King's College London, London, UK ⁶³Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, The Netherlands ⁶⁴Institute of Genetic Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany ⁶⁵Department of Medicine I, Ludwig-Maximilians-University Munich, Munich, Germany ⁶⁶Chair of Genetic Epidemiology, IBE, Faculty of Medicine, LMU Munich, Germany ⁶⁷Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Ontario, Canada ⁶⁸National Institute for Health Research Cambridge Biomedical Research Centre, University of Cambridge and Cambridge University Hospitals, Cambridge, UK ⁶⁹Clinical and Molecular Metabolism Research Program (CAMM), Faculty of Medicine, University of Helsinki, Helsinki, Finland ⁷⁰Department of Clinical Sciences, Genetic and Molecular Epidemiology Unit, Lund University, Skåne University Hospital Malmö, Malmö, Sweden ⁷¹Kuopio Research Institute of Exercise Medicine, Kuopio, Finland ⁷²Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital, Kuopio, Finland ⁷³Institute of Molecular Medicine, McGovern Medical School, The University of Texas Health Science Center at Houston, Houston, TX, USA ⁷⁴MRC Integrative Epidemiology Unit (IEU), Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK ⁷⁵Department of Infection Biology, Faculty of Tropical and Infectious Diseases, London School of Hygiene & Tropical Medicine, London, UK ⁷⁶Centre of Statistics and Applications of University of Lisbon, Lisbon, Portugal, Lisbon ⁷⁷Department of Biostatistics and Center for Statistical Genetics, University of Michigan, Ann Arbor, MI, USA ⁷⁸Saw Swee Hock School of Public Health, National University of Singapore, Singapore ⁷⁹Icelandic Heart Association, Kopavogur, Iceland ⁸⁰Faculty of Medicine, University of Iceland, Reykjavik, Iceland ⁸¹Department of Biostatistics, University of Michigan, Ann Arbor, MI, USA ⁸²University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland ⁸³Population Health Research Institute, McMaster University, Hamilton, Ontario, Canada ⁸⁴Department of Molecular Biology, Medical Biochemistry and Pathology, Laval University, Quebec City, Quebec, Canada ⁸⁵Department of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, The Netherlands ⁸⁶Department of Cardiology, Leiden University Medical Center, Leiden, The Netherlands ⁸⁷Vanderbilt Genetics Institute, Vanderbilt Epidemiology Center, Department of Obstetrics and Gynecology, Vanderbilt University Medical Center; Tennessee Valley Health Systems VA, Nashville, TN, USA ⁸⁸Research Center in Epidemiology and Preventive Medicine, Department of Medicine and Surgery, University of Insubria, Varese, Italy ⁸⁹Interfaculty Institute for Genetics and Functional Genomics, University Medicine and University of Greifswald, Greifswald,

Germany ⁹⁰DZHK (German Centre for Cardiovascular Research), partner site Greifswald, Greifswald, Germany ⁹¹The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, LABioMed at Harbor-UCLA Medical Center, Torrance, CA, USA ⁹²Robertson Centre for Biostatistics, University of Glasgow, Glasgow, UK ⁹³Department of Cardiology, Ealing Hospital, Middlesex, UK ⁹⁴Department of Biostatistics and Epidemiology, University of Pennsylvania, Philadelphia, PA, USA ⁹⁵Department of Hygiene and Epidemiology, University of Ioannina Medical School, Ioannina, Greece ⁹⁶Division of Cardiology, University Hospital, Basel, Switzerland ⁹⁷Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany ⁹⁸Department of Epidemiology, Erasmus MC, University Medical Centre, Rotterdam, The Netherlands ⁹⁹Department of General and Interventional Cardiology, University Heart Center Hamburg, Hamburg, Germany ¹⁰⁰University Medical Center Hamburg Eppendorf, Hamburg, Germany ¹⁰¹German Centre for Cardiovascular Research (DZHK), partner site Hamburg/Kiel/Lübeck, Hamburg, Germany ¹⁰²Medical Genomics and Metabolic Genetics Branch, National Human Genome Research Institute, NIH, Bethesda, MD, USA ¹⁰³Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark ¹⁰⁴Department of Clinical Biochemistry, Lillebaelt Hospital, Vejle, Denmark ¹⁰⁵Institute of Regional Health Research, University of Southern Denmark, Odense, Denmark ¹⁰⁶Massachusetts Veterans Epidemiology Research and Information Center (MAVERIC), VA Boston Healthcare System, Boston, MA, USA ¹⁰⁷Division of Aging, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA ¹⁰⁸Department of Medicine, Harvard Medical School, Boston, MA, USA ¹⁰⁹Medical department, Lillebaelt Hospital, Vejle, Denmark ¹¹⁰University of Dundee, Ninewells Hospital & Medical School, Dundee, UK ¹¹¹Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK ¹¹²Department of Internal Medicine B, University Medicine Greifswald, Greifswald, Germany ¹¹³Department of Nutrition and Dietetics, School of Health Science and Education, Harokopio University, Athens, Greece ¹¹⁴Department of Population Science and Experimental Medicine, Institute of Cardiovascular Science, University College London, London, UK ¹¹⁵Department of Public Health and Clinical Medicine, Umeå University, Umeå, Sweden ¹¹⁶Steno Diabetes Center, Copenhagen, Gentofte, Denmark ¹¹⁷PEDEGO Research Unit, MRC Oulu, Oulu University Hospital and University of Oulu, Finland ¹¹⁸Department of Clinical and Molecular Medicine, Norwegian University of Science and Technology, Trondheim, Norway ¹¹⁹Hospital for Children and Adolescents, Helsinki University Central Hospital and University of Helsinki, Helsinki, Finland ¹²⁰Department of Primary Health Care, Vaasa Central Hospital, Vaasa, Finland ¹²¹Diabetes Center, Vaasa Health Care Center, Vaasa, Finland ¹²²Oxford NIHR Biomedical Research Centre, Oxford University Hospitals Trust, Oxford, UK ¹²³Department of Medicine, University of Helsinki and Helsinki University Central Hospital, Helsinki, Finland ¹²⁴Minerva Foundation Institute for Medical Research, Helsinki, Finland ¹²⁵Nephrology Section, Memphis VA Medical Center, Memphis, TN, USA ¹²⁶Department of Medicine,

University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland
¹²⁷Department of Public Health and Caring Sciences, Uppsala University, Uppsala, Sweden
¹²⁸Division of Endocrinology and Metabolism, Department of Internal Medicine, Taichung Veterans General Hospital, Taichung, Taiwan
¹²⁹School of Medicine, National Yang-Ming University, Taipei, Taiwan
¹³⁰School of Medicine, Chung Shan Medical University, Taichung, Taiwan
¹³¹College of Science, Tunghai University, Taichung, Taiwan
¹³²Department of Medical Research, Taichung Veterans General Hospital, Taichung, Taiwan
¹³³Department of Social Work, Tunghai University, Taichung, Taiwan
¹³⁴Center for Clinical Research and Prevention, Bispebjerg and Frederiksberg Hospital, The Capital Region, Copenhagen, Denmark
¹³⁵Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark
¹³⁶George Washington University School of Medicine and Health Sciences, Washington DC, USA
¹³⁷Department of Public health, Strasbourg University hospital, University of Strasbourg, Strasbourg, France
¹³⁸Medical Research Institute, University of Dundee, Ninewells Hospital and Medical School, Dundee, UK
¹³⁹Epidemiology Branch, NHLBI, Bethesda, MD, USA
¹⁴⁰Department of Immunology and Inflammation, Imperial College London, London, UK
¹⁴¹International Centre for Circulatory Health, Imperial College London, London, UK
¹⁴²Centre for Non-Communicable Diseases, Karachi, Pakistan
¹⁴³Institute of Genomics, University of Tartu, Tartu, Estonia
¹⁴⁴Institute of Physiology, University Medicine Greifswald, Karlsburg, Germany
¹⁴⁵Division of Cardiovascular Sciences, NHLBI, Bethesda, MD, USA
¹⁴⁶Division of Epidemiology and Community Health, University of Minnesota, Minneapolis, MN, USA
¹⁴⁷Department of Family Medicine, University of Iceland, Reykjavik, Iceland
¹⁴⁸Development Centre for Primary Health Care in Iceland, Iceland
¹⁴⁹Center for Clinical Research and Disease Prevention, Bispebjerg and Frederiksberg Hospital, The Capital Region, Copenhagen, Denmark
¹⁵⁰Department of Epidemiology, Emory University Rollins School of Public Health; Department of Biomedical Informatics, Emory University School of Medicine, Atlanta, GA, USA
¹⁵¹Department of Medical Sciences, Uppsala University, Uppsala, Sweden
¹⁵²Department of Internal Medicine, Division of Cardiology, Landspítali - The National University Hospital of Iceland, Reykjavik, Iceland
¹⁵³Medical and Population Genetics, Broad Institute, Cambridge, MA, USA
¹⁵⁴VA Palo Alto Health Care System, Division of Cardiovascular Medicine, Stanford University School of Medicine, Stanford, CA, USA
¹⁵⁵Folkhälsan Research Centre, Helsinki, Finland
¹⁵⁶Department of Endocrinology, Helsinki University Central Hospital, Helsinki, Finland
¹⁵⁷Department of Clinical Sciences, Diabetes and Endocrinology, Lund University Diabetes Centre, Malmö, Sweden. Institute for Molecular Medicine Helsinki (FIMM), Helsinki University, Helsinki, Finland
¹⁵⁸Division of Nephrology and Hypertension, Mayo Clinic, Rochester, MN, USA
¹⁵⁹Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, University of Utrecht, University of Utrecht, The Netherlands
¹⁶⁰Center for Circulatory Health, University Medical Center Utrecht, University of Utrecht, The Netherlands
¹⁶¹Department of Internal Medicine, Division of Cardiovascular Medicine, University

of Michigan, Ann Arbor, MI, USA ¹⁶²Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA ¹⁶³Department of Human Genetics, University of Michigan, Ann Arbor, MI, USA ¹⁶⁴Atlanta VAMC and Emory Clinical Cardiovascular Research Institute, Atlanta, GA, USA ¹⁶⁵Department of Public Health, Aarhus University, Aarhus, Denmark ¹⁶⁶Danish Diabetes Academy, Odense, Denmark ¹⁶⁷Steno Diabetes Center Aarhus, Aarhus, Denmark ¹⁶⁸International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Mohakhali, Dhaka, Bangladesh ¹⁶⁹Dep of Medicine, Lund University, Malmö, Sweden ¹⁷⁰Univ. Lille, U1167 - RID-AGE - Facteurs de risque et déterminants moléculaires des maladies liées au vieillissement, Lille, France ¹⁷¹Inserm, U1167, Lille, France ¹⁷²CHU Lille, U1167, Lille, France ¹⁷³Institut Pasteur de Lille, U1167, Lille, France ¹⁷⁴Health Data Research UK, Institute of Health Informatics, University College London, London, UK ¹⁷⁵Institute of Cardiovascular Science, Faculty of Population Health Sciences, University College London, London, UK ¹⁷⁶Section of Investigative Medicine, Imperial College London, Hammersmith Hospital Campus, London, UK ¹⁷⁷Harvard Medical School, Boston, MA, USA ¹⁷⁸Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore, Singapore ¹⁷⁹Imperial College Healthcare NHS Trust, London, UK ¹⁸⁰Cardiovascular Research Institute Basel, Basel, Switzerland ¹⁸¹Jackson Heart Study, Department of Medicine, University of Mississippi Medical Center, Jackson, MS, USA ¹⁸²University of Groningen, University Medical Center Groningen, Department of Cardiology, Groningen, The Netherlands ¹⁸³Department of Psychology, University of Edinburgh, Edinburgh, UK ¹⁸⁴Princess Al-Jawhara Al-Brahim Centre of Excellence in Research of Hereditary Disorders (PACER-HD), King Abdulaziz University, Jeddah, Saudi Arabia ¹⁸⁵National Institute for Health Research Blood and Transplant Research Unit in Donor Health and Genomics, University of Cambridge, Cambridge, UK ¹⁸⁶Department of Human Genetics, Wellcome Sanger Institute, Hinxton, UK ¹⁸⁷Health Data Research UK – London at Imperial College London, London, UK ¹⁸⁸UKDRI, Dementia Research Institute at Imperial College London, London, UK ¹⁸⁹Department of Cardiology and Department of Epidemiology, INSERM UMR 1027, Toulouse University Hospital, Toulouse, France ¹⁹⁰Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA ¹⁹¹Department of Public Health & Clinical Medicine, Umeå University, Umeå, Sweden ¹⁹²Oxford Center for Diabetes, Endocrinology & Metabolism, Radcliff Department of Medicine, University of Oxford, Oxford, UK ¹⁹³Institute of Reproductive & Developmental Biology, Imperial College London, London, UK ¹⁹⁴Division of Nephrology, Department of Medicine, University of Verona, Verona, Italy ¹⁹⁵Department of Clinical Sciences, Diabetes and Endocrinology, Lund University Diabetes Centre, Malmö, Sweden ¹⁹⁶Institute for Molecular Medicine Helsinki (FIMM), Helsinki University, Helsinki, Finland ¹⁹⁷Laboratory of Epidemiology and Population Sciences, National Institute of Aging, Bethesda, MD, USA ¹⁹⁸Wellcome Trust, London, UK ¹⁹⁹Institute of Biomedicine, Medical Research Center (MRC), University of Oulu, and University Hospital Oulu, Oulu, Finland ²⁰⁰Department of Gastroenterology and Metabolism, Poznan University of Medical Sciences, Poznan, Poland ²⁰¹Department of Medicine, Division

of Cardiovascular Medicine, Stanford University School of Medicine, Stanford, CA, USA ²⁰²Department of Medical Sciences, Molecular Epidemiology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden ²⁰³Stanford Cardiovascular Institute, Stanford University, Stanford, CA, USA ²⁰⁴Stanford Diabetes Research Center, Stanford University, Stanford, CA, USA ²⁰⁵Department of Public Health, University of Helsinki, Helsinki, Finland ²⁰⁶Saudi Diabetes Research Group, King Abdulaziz University, Jeddah, Saudi Arabia ²⁰⁷National Institute of Public Health, Madrid, Spain ²⁰⁸Unit of Primary Care, Oulu University Hospital, Oulu, Finland ²⁰⁹Netherlands Heart Institute, Utrecht, The Netherlands, Utrecht, The Netherlands ²¹⁰Centre for Public Health, Queens University Belfast, Belfast, UK ²¹¹National Heart and Lung Institute, Imperial College London, London, UK ²¹²Fred Hutchinson Cancer Research Center, Division of Public Health Sciences, Seattle, WA, USA ²¹³The Mindich Child Health and Development Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA ²¹⁴National Institute of Cardiovascular Diseases, Sher-e-Bangla Nagar, Dhaka, Bangladesh ²¹⁵Department of Genetics, University of North Carolina, Chapel Hill, NC, USA ²¹⁶The Institute of Medical Sciences, Aberdeen Biomedical Imaging Centre, University of Aberdeen, Aberdeen, UK ²¹⁷University of Glasgow, Glasgow, UK ²¹⁸Institute of Cardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK ²¹⁹Department of Medicine, Columbia University Medical Center, New York, NY, USA ²²⁰Department of Public Health, University of Split School of Medicine, Split, Croatia ²²¹Centre for Genomic and Experimental Medicine, Institute of Genetics & Molecular Medicine, University of Edinburgh, Western General Hospital, Edinburgh, UK ²²²Centre for Cognitive Ageing and Cognitive Epidemiology, Department of Psychology, The University of Edinburgh, Edinburgh, UK ²²³MRC International Nutrition Group at London School of Hygiene & Tropical Medicine, London, UK ²²⁴Department of Biostatistics, University of Washington, Seattle, WA, USA ²²⁵Institute for Translational Genomics and Population Sciences, Departments of Pediatrics and Medicine, LABioMed at Harbor-UCLA Medical Center, Torrance, CA, USA ²²⁶Centre for Global Health Research, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, UK ²²⁷School of Medicine, National Defense Medical Center, Taipei, Taiwan ²²⁸Institute of Medical Technology, National Chung-Hsing University, Taichung, Taiwan ²²⁹Division of Population Health and Genomics, Ninewells Hospital and Medical School, University of Dundee, Dundee, UK ²³⁰Department of Human Genetics, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, UK ²³¹Department of Haematology, University of Cambridge, Cambridge, UK ²³²The National Institute for Health Research Blood and Transplant Unit (NIHR BTRU) in Donor Health and Genomics at the University of Cambridge, Cambridge, UK ²³³Alzheimer Scotland Research Centre, University of Edinburgh, Edinburgh, UK ²³⁴Center for Life Course Health Research, Faculty of Medicine, University of Oulu, Oulu, Finland ²³⁵Biocenter Oulu, University of Oulu, Oulu, Finland ²³⁶Department of Genomics of Complex Diseases, School of Public Health, Imperial College London, London, UK ²³⁷Institute for Translational Genomics and Population Sciences, Los Angeles Biomedical Research Institute at Harbor/

UCLA Medical Center, Torrance, CA, USA ²³⁸Institute of Biomedicine/Physiology, University of Eastern Finland, Kuopio Campus, Kuopio, Finland ²³⁹Department of Health Sciences, University of Leicester, Leicester, UK ²⁴⁰University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen, The Netherlands ²⁴¹Durrer Center for Cardiogenetic Research, ICIN-Netherlands Heart Institute, Utrecht, The Netherlands ²⁴²University of Groningen, University Medical Center Groningen, Department of Cardiology, Groningen, The Netherlands ²⁴³Boston University Schools of Medicine and Public Health, Boston, MA, USA ²⁴⁴University Medical Center Groningen, Groningen, The Netherlands ²⁴⁵Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridge, UK ²⁴⁶Institute of Translational Genomics, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany ²⁴⁷School of Health and Life Sciences, Federation University Australia, Ballarat, Victoria, Australia ²⁴⁸Department of Physiology, University of Melbourne, Melbourne, Victoria, Australia ²⁴⁹Division of Medicine, Manchester University NHS Foundation Trust, Manchester Academic Health Science Centre, Manchester, UK ²⁵⁰Division of Epidemiology, Department of Medicine, Institute for Medicine and Public Health, Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Tennessee Valley Healthcare System (626)/Vanderbilt University, Nashville, TN, USA ²⁵¹VA Tennessee Valley Healthcare System, Division of Nephrology & Hypertension, Department of Medicine, Vanderbilt Center for Kidney Disease, Vanderbilt University Medical Center, Nashville, TN, USA ²⁵²Wellcome Centre for Human Genetics, University of Oxford, Oxford, UK ²⁵³Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, MA, USA ²⁵⁴Boston University School of Public Health, Boston, MA, USA ²⁵⁵Center for Human Genetics Research, Massachusetts General Hospital, Boston, MA, USA ²⁵⁶Division of Musculoskeletal and Dermatological Sciences, The University of Manchester, Manchester, UK ²⁵⁷VA Boston Healthcare, Section of Cardiology and Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA ²⁵⁸Department of Epidemiology, University of Washington, Seattle, WA, USA ²⁵⁹Department of Health Services, University of Washington, Seattle, WA, USA ²⁶⁰Kaiser Permanente Washington Health Research Institute, Seattle, WA, USA ²⁶¹Department of Biostatistics and Epidemiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA ²⁶²Center for Non-Communicable Diseases, Karachi, Pakistan ²⁶³Human Genome Sequencing Center, Baylor College of Medicine, Houston, TX, USA ²⁶⁴Population Sciences Branch, National Heart, Lung, and Blood Institute, National Institute of Health, Bethesda, MD, USA ²⁶⁵Department of Genetics, Novo Nordisk Research Centre Oxford, Oxford, UK

Acknowledgements

P. Surendran is supported by a Rutherford Fund Fellowship from the Medical Research Council grant MR/S003746/1. N. Lahrouchi is supported by the Foundation “De Drie Lichten” in The Netherlands and the Netherlands Cardiovascular Research Initiative, an initiative supported by the Dutch Heart Foundation (CVON2012-10 PREDICT and CVON2018-30 PREDICT2). J. N. Hellwege was supported by the Vanderbilt Molecular and Genetic Epidemiology of Cancer (MAGEC) Training Program (T32CA160056, PI X.-O. Shu). N.

Franceschini is supported by the National Institute of Health awards HL140385, MD012765 and DK117445. F. W. Asselbergs is supported by UCL Hospitals NIHR Biomedical Research Centre. P. Deloukas's work was supported by the British Heart Foundation (BHF) grant RG/14/5/30893. R. J. F. Loos is funded by R01DK110113, U01HG007417, R01DK101855, R01DK107786. C. Hayward is supported by an MRC University Unit Programme Grant MC_UU_00007/10 (QTL in Health and Disease) and MRC University Unit Programme Grant MC_PC_U127592696. M. I. McCarthy* is a Wellcome Senior Investigator (098381; 212259) and an NIHR Senior Investigator (NF-SI-0617-10090). The research was supported by the National Institute for Health Research (NIHR) Oxford Biomedical Research Centre (BRC), and by the Wellcome (090532, 106130, 098381, 203141, 212259). T. Ferreira* is supported by the NIHR Biomedical Research Centre, Oxford. M. Tomaszewski is supported by British Heart Foundation (PG/17/35/33001 and PG/19/16/34270) and Kidney Research UK (RP_017_20180302). J. Danesh* is funded by the National Institute for Health Research (Senior Investigator Award). C. M. Lindgren* is supported by the Li Ka Shing Foundation, WT-SSI/John Fell funds and by the NIHR Biomedical Research Centre, Oxford, by Widenlife and NIH (5P50HD028138-27). J. M. M. Howson* was funded by the National Institute for Health Research (Cambridge Biomedical Research Centre at the Cambridge University Hospitals NHS Foundation Trust).

*The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Full acknowledgements and full lists of consortia members are provided in the Supplementary Note.

Data availability

Summary association results for all the traits are available for download from: <https://app.box.com/s/1ev9iakptips70k8t4cm8j347iff0ef2u> and from the CHARGE dbGaP Summary site, (<https://www.ncbi.nlm.nih.gov/gap/>) accession number phs000930.

References

1. Forouzanfar MH, et al. Global burden of hypertension and systolic blood pressure of at least 110 to 115 mm Hg, 1990-2015. *JAMA*. 2017; 317:165–182. [PubMed: 28097354]
2. Newton-Cheh C, et al. Genome-wide association study identifies eight loci associated with blood pressure. *Nat Genet*. 2009; 41:666–676. [PubMed: 19430483]
3. Cho YS, et al. A large-scale genome-wide association study of Asian populations uncovers genetic factors influencing eight quantitative traits. *Nat Genet*. 2009; 41:527–534. [PubMed: 19396169]
4. Levy D, et al. Genome-wide association study of blood pressure and hypertension. *Nat Genet*. 2009; 41:677–687. [PubMed: 19430479]
5. Kato N, et al. Meta-analysis of genome-wide association studies identifies common variants associated with blood pressure variation in east Asians. *Nat Genet*. 2011; 43:531–538. [PubMed: 21572416]
6. Wain LV, et al. Genome-wide association study identifies six new loci influencing pulse pressure and mean arterial pressure. *Nat Genet*. 2011; 43:1005–1011. [PubMed: 21909110]
7. International Consortium for Blood Pressure Genome-Wide Association Studies. et al. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature*. 2011; 478:103–109. [PubMed: 21909115]
8. Johnson AD, et al. Association of hypertension drug target genes with blood pressure and hypertension in 86,588 individuals. *Hypertension*. 2011; 57:903–910. [PubMed: 21444836]
9. Johnson T, et al. Blood pressure loci identified with a gene-centric array. *Am J Hum Genet*. 2011; 89:688–700. [PubMed: 22100073]
10. Tragante V, et al. Gene-centric meta-analysis in 87,736 individuals of European ancestry identifies multiple blood-pressure-related loci. *Am J Hum Genet*. 2014; 94:349–360. [PubMed: 24560520]
11. Simino J, et al. Gene-age interactions in blood pressure regulation: a large-scale investigation with the CHARGE, Global BPgen, and ICBP Consortia. *Am J Hum Genet*. 2014; 95:24–38. [PubMed: 24954895]
12. Kato N, et al. Trans-ancestry genome-wide association study identifies 12 genetic loci influencing blood pressure and implicates a role for DNA methylation. *Nat Genet*. 2015; 47:1282–1293. [PubMed: 26390057]

13. Surendran P, et al. Trans-ancestry meta-analyses identify rare and common variants associated with blood pressure and hypertension. *Nat Genet.* 2016; 48:1151–1161. [PubMed: 27618447]
14. Liu C, et al. Meta-analysis identifies common and rare variants influencing blood pressure and overlapping with metabolic trait loci. *Nat Genet.* 2016; 48:1162–1170. [PubMed: 27618448]
15. Ehret GB, et al. The genetics of blood pressure regulation and its target organs from association studies in 342,415 individuals. *Nat Genet.* 2016; 48:1171–1184. [PubMed: 27618452]
16. Hoffmann TJ, et al. Genome-wide association analyses using electronic health records identify new loci influencing blood pressure variation. *Nat Genet.* 2017; 49:54–64. [PubMed: 27841878]
17. Warren HR, et al. Genome-wide association analysis identifies novel blood pressure loci and offers biological insights into cardiovascular risk. *Nat Genet.* 2017; 49:403–415. [PubMed: 28135244]
18. Kraja AT, et al. New blood pressure-associated loci identified in meta-analyses of 475 000 individuals. *Circ Cardiovasc Genet.* 2017; 10:e001778. [PubMed: 29030403]
19. Wain LV, et al. Novel blood pressure locus and gene discovery using genome-wide association study and expression data sets from blood and the kidney. *Hypertension.* 2017
20. Evangeou E, et al. Genetic analysis of over 1 million people identifies 535 new loci associated with blood pressure traits. *Nat Genet.* 2018; 50:1412–1425. [PubMed: 30224653]
21. Giri A, et al. Trans-ethnic association study of blood pressure determinants in over 750,000 individuals. *Nat Genet.* 2019; 51:51–62. [PubMed: 30578418]
22. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet.* 2011; 88:76–82. [PubMed: 21167468]
23. Bycroft C, et al. The UK Biobank resource with deep phenotyping and genomic data. *Nature.* 2018; 562:203–209. [PubMed: 30305743]
24. Loh PR, et al. Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nat Genet.* 2015; 47:284–290. [PubMed: 25642633]
25. Benner C, et al. FINEMAP: efficient variable selection using summary data from genome-wide association studies. *Bioinformatics.* 2016; 32:1493–1501. [PubMed: 26773131]
26. Wu MC, et al. Rare-variant association testing for sequencing data with the sequence kernel association test. *Am J Hum Genet.* 2011; 89:82–93. [PubMed: 21737059]
27. McLaren W, et al. The Ensembl Variant Effect Predictor. *Genome Biol.* 2016; 17:122. [PubMed: 27268795]
28. Marouli E, et al. Rare and low-frequency coding variants alter human adult height. *Nature.* 2017; 542:186–190. [PubMed: 28146470]
29. Liu DJ, et al. Exome-wide association study of plasma lipids in 300,000 individuals. *Nat Genet.* 2017; 49:1758–1766. [PubMed: 29083408]
30. Finan C, et al. The druggable genome and support for target identification and validation in drug development. *Sci Transl Med.* 2017; 9
31. Foley CN, et al. A fast and efficient colocalization algorithm for identifying shared genetic risk factors across multiple traits. *bioRxiv.* 2019
32. Solovieff N, Cotsapas C, Lee PH, Purcell SM, Smoller JW. Pleiotropy in complex traits: challenges and strategies. *Nat Rev Genet.* 2013; 14:483–495. [PubMed: 23752797]
33. Xu X, et al. Molecular insights into genome-wide association studies of chronic kidney disease-defining traits. *Nat Commun.* 2018; 9:4800. [PubMed: 30467309]
34. Rowland J, et al. Uncovering genetic mechanisms of kidney aging through transcriptomics, genomics, and epigenomics. *Kidney Int.* 2019; 95:624–635. [PubMed: 30784661]
35. Staley JR, et al. PhenoScanner: a database of human genotype-phenotype associations. *Bioinformatics.* 2016; 32:3207–3209. [PubMed: 27318201]
36. Turcot V, et al. Protein-altering variants associated with body mass index implicate pathways that control energy intake and expenditure in obesity. *Nat Genet.* 2018; 50:26–41. [PubMed: 29273807]
37. Vishram JK, et al. Impact of age on the importance of systolic and diastolic blood pressures for stroke risk: the MOnica, Risk, Genetics, Archiving, and Monograph (MORGAM) Project. *Hypertension.* 2012; 60:1117–1123. [PubMed: 23006731]

38. Messaoudi S, et al. Endothelial Gata5 transcription factor regulates blood pressure. *Nat Commun.* 2015; 6:8835. [PubMed: 26617239]
39. Kircher M, et al. A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet.* 2014; 46:310–315. [PubMed: 24487276]
40. Wang Z, et al. Phospholipase C beta3 deficiency leads to macrophage hypersensitivity to apoptotic induction and reduction of atherosclerosis in mice. *J Clin Invest.* 2008; 118:195–204. [PubMed: 18079968]
41. Hoepfner LH, et al. Revealing the role of phospholipase Cbeta3 in the regulation of VEGF-induced vascular permeability. *Blood.* 2012; 120:2167–2173. [PubMed: 22674805]
42. Li M, et al. SOS2 and ACP1 loci identified through large-scale exome chip analysis regulate kidney development and function. *J Am Soc Nephrol.* 2017; 28:981–994. [PubMed: 27920155]
43. Tidyman WE, Rauen KA. Pathogenetics of the RASopathies. *Hum Mol Genet.* 2016; 25:R123–R132. [PubMed: 27412009]
44. Kim JH, et al. Functional role of the polymorphic 647 T/C variant of ENT1 (SLC29A1) and its association with alcohol withdrawal seizures. *PLoS One.* 2011; 6:e16331. [PubMed: 21283641]
45. Best KA, Bone DB, Vilas G, Gros R, Hammond JR. Changes in aortic reactivity associated with the loss of equilibrative nucleoside transporter 1 (ENT1) in mice. *PLoS One.* 2018; 13:e0207198. [PubMed: 30408123]
46. Zaidi S, et al. De novo mutations in histone-modifying genes in congenital heart disease. *Nature.* 2013; 498:220–223. [PubMed: 23665959]
47. Jin SC, et al. Contribution of rare inherited and de novo variants in 2,871 congenital heart disease probands. *Nat Genet.* 2017; 49:1593–1601. [PubMed: 28991257]
48. Nielsen JB, et al. Genome-wide study of atrial fibrillation identifies seven risk loci and highlights biological pathways and regulatory elements involved in cardiac development. *Am J Hum Genet.* 2018; 102:103–115. [PubMed: 29290336]
49. Tobin MD, Sheehan NA, Scurrah KJ, Burton PR. Adjusting for treatment effects in studies of quantitative traits: antihypertensive therapy and systolic blood pressure. *Stat Med.* 2005; 24:2911–2935. [PubMed: 16152135]
50. Grove ML, et al. Best practices and joint calling of the HumanExome BeadChip: the CHARGE Consortium. *PLoS One.* 2013; 8:e68095. [PubMed: 23874508]
51. Liu DJ, et al. Meta-analysis of gene-level tests for rare variant association. *Nat Genet.* 2014; 46:200–204. [PubMed: 24336170]
52. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics.* 2010; 26:2190–2191. [PubMed: 20616382]
53. Fadista J, Manning AK, Florez JC, Groop L. The (in)famous GWAS P-value threshold revisited and updated for low-frequency variants. *Eur J Hum Genet.* 2016; 24:1202–1205. [PubMed: 26733288]
54. Flannick J, et al. Exome sequencing of 20,791 cases of type 2 diabetes and 24,440 controls. *Nature.* 2019; 570:71–76. [PubMed: 31118516]
55. Mahajan A, et al. Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. *Nat Genet.* 2018; 50:1505–1513. [PubMed: 30297969]
56. Mahajan A, et al. Refining the accuracy of validated target identification through coding variant fine-mapping in type 2 diabetes. *Nat Genet.* 2018; 50:559–571. [PubMed: 29632382]
57. Yengo L, et al. Meta-analysis of genome-wide association studies for height and body mass index in ~700000 individuals of European ancestry. *Hum. Mol Genet.* 2018; 27:3641–3649.
58. Willer CJ, et al. Discovery and refinement of loci associated with lipid levels. *Nat Genet.* 2013; 45:1274–1283. [PubMed: 24097068]
59. Dupuis J, et al. New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet.* 2010; 42:105–116. [PubMed: 20081858]
60. Scott RA, et al. An expanded genome-wide association study of type 2 diabetes in Europeans. *Diabetes.* 2017; 66:2888–2902. [PubMed: 28566273]

61. Nikpay M, et al. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet.* 2015; 47:1121–1130. [PubMed: 26343387]
62. Giambartolomei C, et al. Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* 2014; 10:e1004383. [PubMed: 24830394]
63. Malik R, et al. Multiancestry genome-wide association study of 520,000 subjects identifies 32 loci associated with stroke and stroke subtypes. *Nat Genet.* 2018; 50:524–537. [PubMed: 29531354]
64. van der Harst P, Verweij N. Identification of 64 novel genetic loci provides an expanded view on the genetic architecture of coronary artery disease. *Circ Res.* 2018; 122:433–443. [PubMed: 29212778]
65. Burgess S, et al. Using published data in Mendelian randomization: a blueprint for efficient identification of causal risk factors. *Eur J Epidemiol.* 2015; 30:543–552. [PubMed: 25773750]
66. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol.* 2015; 44:512–525. [PubMed: 26050253]
67. Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet.* 2018; 50:693–698. [PubMed: 29686387]
68. Pierce BL, Ahsan H, Vanderweele TJ. Power and instrument strength requirements for Mendelian randomization studies using multiple genetic variants. *Int J Epidemiol.* 2011; 40:740–752. [PubMed: 20813862]
69. Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology. *Int J Epidemiol.* 2016; 45:1866–1886. [PubMed: 28108528]
70. Sanderson E, Davey Smith G, Windmeijer F, Bowden J. An examination of multivariable Mendelian randomization in the single-sample and two-sample summary data settings. *Int J Epidemiol.* 2019; 48:713–727. [PubMed: 30535378]
71. Di Angelantonio E, et al. Efficiency and safety of varying the frequency of whole blood donation (INTERVAL): a randomised trial of 45 000 donors. *Lancet.* 2017; 390:2360–2371. [PubMed: 28941948]
72. Astle WJ, et al. The allelic landscape of human blood cell trait variation and links to common complex disease. *Cell.* 2016; 167:1415–1429 e19. [PubMed: 27863252]
73. Day N, et al. EPIC-Norfolk: study design and characteristics of the cohort European Prospective Investigation of Cancer. *Br J Cancer.* 1999; 80(1):95–103. [PubMed: 10466767]
74. Cancer Genome Atlas Research Network. et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet.* 2013; 45:1113–1120. [PubMed: 24071849]
75. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol.* 2016; 34:525–527. [PubMed: 27043002]
76. Iotchkova V, et al. Discovery and refinement of genetic loci associated with cardiometabolic risk using dense imputation maps. *Nat Genet.* 2016; 48:1303–1312. [PubMed: 27668658]
77. Iotchkova V, et al. GARFIELD classifies disease-relevant genomic features through integration of functional annotations with association signals. *Nat Genet.* 2019; 51:343–353. [PubMed: 30692680]
78. Zhu X, et al. Meta-analysis of correlated traits via summary statistics from GWASs with an application in hypertension. *Am J Hum Genet.* 2015; 96:21–36. [PubMed: 25500260]
79. Newton-Cheh C, et al. Association of common variants in NPPA and NPPB with circulating natriuretic peptides and blood pressure. *Nat Genet.* 2009; 41:348–53. [PubMed: 19219041]




Figure 1. Study design for single variant discovery.

a, Exome array-wide association study (EAWAS) of SBP, DBP, PP and HTN. In Stage 1, we performed two fixed effect meta-analyses for each of the blood pressure (BP) phenotypes SBP, DBP, PP and HTN: one meta-analysis including 810,865 individuals of European (EUR) ancestry and a second pan-ancestry (PA) meta-analysis including 870,217 individuals of EUR, South Asians (SAS), East Asians (EAS), African Ancestry (AA), Hispanics (HIS) and Native Americans (NA_m) (Supplementary Tables 23 and 24; Methods). Summary association statistics for SNVs with $P < 5 \times 10^{-8}$ in Stage 1 that were outside of previously

reported BP loci (Methods, Supplementary Tables 1 and 25) were requested in independent studies (up to 448,667 participants; Supplementary Table 24). In Stage 2, we performed both a EUR and a PA meta-analyses for each trait of Stage 1 results and summary statistics from the additional studies. Only SNVs that were associated with a BP trait at $P < 5 \times 10^{-8}$ in the combined Stage 2 EUR or PA meta-analyses and had concordant directions of effect across studies ($P_{\text{heterogeneity}} > 1 \times 10^{-4}$; Methods) were considered significant. Further details are provided in the Methods and Supplementary Information. **b**, Rare variant GWAS (RV-GWAS) of SBP, DBP and PP. For SNVs outside of the previously reported BP loci (Methods, Supplementary Tables 1 and 6) with $P < 1 \times 10^{-7}$ in Stage 1, summary association statistics were requested from MVP (up to 225,112 participants; Supplementary Table 24). In Stage 2, we performed meta-analyses of Stage 1 and MVP for SBP, DBP and PP in EUR. SNVs that were associated with a BP trait at $P < 5 \times 10^{-8}$ in the combined Stage 2 EUR with concordant directions of effect across UKBB and MVP ($P_{\text{heterogeneity}} > 1 \times 10^{-4}$; Methods) were considered significant. Justification of the significance thresholds used and further information on the statistical methods are detailed in the Methods and Supplementary Information. *Total number of participants analyzed within each study that provided single variant association summaries following the data request—EAWAS EUR: Million Veterans Program (MVP: 225,113), deCODE (127,478) and GENOA (1,505); EAWAS PA: Million Veterans Program (MVP: 225,113 EUR; 63,490 AA; 22,802 HIS; 2,695 Nam; 4,792 EAS), deCODE (127,478 participants from Iceland) and GENOA (1,505 EUR; 792 AA); RV-GWAS EUR: Million Veterans Program (MVP: 225,112 EUR).




Figure 2. New BP associations.

a. Fuji plot of the genome-wide significant BP-associated SNVs from the Stage 2 EAWAS and Stage 2 rare variant GWAS. The first four circles (from inside-out) and the last circle (locus annotation) summarize pleiotropic effects, while circles 5 to 8 summarize the genome-wide significant associations. Every dot or square represents a BP-associated locus, and large dots represent novel BP-associated loci, while small dots represent loci containing novel variants identified in this study, which are in linkage disequilibrium with a variant reported by Evangelou et al.²⁰ and/or Giri et al.²¹. All loci are independent of each other, but

due to the scale of the plot, dots for loci in close proximity overlap. *Loci with rare variant associations. **b**, Venn diagram showing the overlap of the 107 new BP loci across the analyzed BP traits. **c**, Functional annotation from VEP of all the identified rare variants in known and novel regions. **d**, Plots of minor allele frequency against effect estimate on the transformed scale for the BP-associated SNVs. Blue squares are new BP-associated SNVs, black dots represent SNVs at known loci, and red dots are newly identified distinct BP-associated SNVs at known loci. Effect estimates and SEs for the novel loci are taken from the Stage 2 EUR analyses (up to 1,164,961 participants), while for the known are from the Stage 1 analyses (up to 810,865 participants). Results are from the EAWAS where available and the GWAS (up to 670,472 participants) if the known variants were not on the exome array (data from Supplementary Tables 1, 3, 7, 8, and 25 were used).




Figure 3. Annotation of BP loci.

a. BP associations shared with eQTL from GTEx through multi-trait colocalization analyses. Expressed gene and the colocalized SNV are provided on the y-axis. BP trait and eQTL tissues are provided on the x-axis. The color indicates whether the candidate SNV increases BP and gene expression (brown), decreases BP and gene expression (orange), or has the inverse effects on BP and gene expression (blue). **b.** Enrichment of BP-associated SNVs in DNase I hypersensitivity hot spots (active chromatin). The top plot is for SBP, middle is for DBP, and bottom represents PP. Height of the bar indicates the fold enrichment in the listed

tissues, with error bars representing the 95% confidence intervals. The colors represent the enrichment P -value.




Figure 4. Phenome-wide associations of the new BP loci.

a. Modified Fuji plot of the genome-wide significant associated SNVs from the Stage 2 EAWAS and Stage 2 rare variant GWAS (novel loci only). Each dot represents a novel locus where a conditionally independent variant or a variant in LD with the conditionally independent variant has been previously associated with one or more traits unrelated to blood pressure, and each circle represents different trait category (Supplementary Table 20). Locus annotation is plotted in the outer circle, and * sign denotes loci where the conditionally independent signal maps to a gene which is different to the one closest to the

sentinel variant. **b**, Bar chart showing the distribution of traits (x -axis) and number of distinct BP-associated variants per trait (y -axis) that the SNVs in **a** are associated with. **c**, Bar chart of the number of traits included in **b** (y -axis) by trait category (x -axis). The color coding for **a** and **b** is relative to **c**.




Figure 5. Causal association of BP with stroke and coronary artery disease.

Mendelian randomization analyses of the effect of blood pressure on stroke and coronary artery disease. **a**, Univariable analyses. **b**, Multivariable analyses (Methods). Analyses were performed using summary association statistics (Methods). The causal estimates are on the odds ratio (OR) scale (the square in the plot). The whiskers on the plots are the 95% confidence intervals for these ORs. Results on the standard deviation scale are provided in Supplementary Table 22. The genetic variants for the estimation of the causal effects in this plot are sets of SNVs after removing the confounding SNVs and invalid instrumental

variant. OR, odds ratio (P -value from the inverse variance weighted two sample Mendelian randomization method). n , number of disease cases.

Table 1
Rare and low-frequency SNV-blood pressure associations in participants of European ancestry from the (Stage 2) EAWAS and (Stage 2) RV-GWAS that map to new BP loci

Locus	rsID	Chr:Pos	Gene	EA/OA	Amino acids	Consequence	Trait	EAF	β	P	Het P	n
Exome array-wide association study (EAWAS)												
10	rs11580946	1:150,551,327	<i>MCL1</i>	A/G	p.Val1227Ala	missense	PP	0.016	-0.37	2.74×10^{-9}	0.24	1,159,900
11	rs61747728 [†]	1:179,526,214	<i>NPHS2</i>	T/C	p.Gln229Arg	missense	DBP	0.040	0.26	8.74×10^{-13}	0.22	1,160,530
16	rs4149909	1:242,023,898	<i>EXO1</i>	G/A	p.Ser279Asn	missense	SBP	0.033	0.36	2.46×10^{-8}	0.09	1,158,190
32	rs3821033 [†]	2:219,507,302	<i>ZNF142</i>	T/C	p.Thr1313Ala	missense	DBP	0.033	-0.29	1.42×10^{-13}	0.75	1,160,530
	rs16859180 [†]	2:219,553,468	<i>STK36</i>	T/C	p.Trp477Arg	missense	DBP	0.049	-0.26	1.11×10^{-16}	0.34	1,160,530
44	rs145072852	3:101,476,645	<i>CEP97</i>	T/C	p.Phe399Leu	missense	PP	0.004	1.05	1.42×10^{-13}	0.01	1,158,820
46	rs139600783	3:119,109,769	<i>ARHGAP31</i>	T/C	p.Ser274Pro	missense	HTN	0.008	5.85	5.05×10^{-9}	0.19	975,381
50	rs73181210	3:169,831,268	<i>PHC3</i>	C/T	p.Glu692Lys	missense	DBP	0.009	-0.66	9.14×10^{-15}	0.04	1,159,580
52	rs11937432 [†]	4:2,233,709	<i>HAUS3</i>	G/A	p.Thr586Ile	missense	DBP	0.046	0.21	9.56×10^{-10}	0.26	1,160,520
58	rs1229984	4:100,239,319	<i>ADH1B</i>	T/C	p.His48Arg	missense	PP	0.026	-0.75	2.97×10^{-25}	0.54	686,104
63	rs143057152	4:149,075,755	<i>NR3C2</i>	T/C	p.His771Arg	missense	SBP	0.003	1.75	4.14×10^{-14}	0.22	1,128,880
71	rs61755724	5:132,408,967	<i>HSPA4</i>	A/G	p.Thr159Ala	missense	DBP	0.024	0.26	9.75×10^{-9}	0.36	1,160,530
72	rs33956817	5:137,278,682	<i>FAM13B</i>	C/T	p.Met802Val	missense	SBP	0.044	0.31	1.76×10^{-8}	0.27	1,158,190
77	rs34471628 [†]	5:172,196,752	<i>DUSP1</i>	G/A	p.His187Tyr	missense	DBP	0.039	-0.23	3.00×10^{-10}	0.42	1,153,300
85	rs45573936	6:44,198,362	<i>SLC29A1</i>	C/T	p.Ile295Thr	missense	DBP	0.027	-0.38	3.70×10^{-19}	0.59	1,160,530
100	rs144867634	7:111,580,166	<i>DOCK4</i>	C/T	p.Val326Met	missense/splice region	DBP	0.025	-0.26	2.62×10^{-8}	0.04	1,160,530
109	rs5635308 [†]	8:17,419,461	<i>SLC7A2</i>	A/G	p.Met545Val	missense	DBP	0.025	0.31	1.40×10^{-10}	0.26	1,160,530
114	rs76767219	8:81,426,196	<i>ZBTB10</i>	A/C	p.Glu346Ala	missense	SBP	0.034	-0.44	4.41×10^{-13}	0.18	1,160,830
119	rs61752533 [†]	8:145,108,151	<i>OPLAH</i>	A/G	-	synonymous	DBP	0.049	-0.21	2.05×10^{-10}	0.86	1,085,170
	rs34674752 [†]	8:145,154,222	<i>SHARPIN</i>	A/G	p.Ser294Pro	missense	DBP	0.049	-0.19	5.89×10^{-10}	0.91	1,132,350
146	rs117874826	11:64,027,666	<i>PLCB3</i>	C/A	p.Ala564Glu	missense	SBP	0.014	0.71	4.67×10^{-12}	0.42	1,153,360
	rs145502455	11:64,031,030	<i>PLCB3</i>	A/G	p.Ile806Val	missense	SBP	0.005	0.90	5.01×10^{-9}	0.04	1,156,310
154	rs141325069	12:20,769,270	<i>PDE3A</i>	A/G	p.Gln459Arg	missense	SBP	0.003	1.45	6.25×10^{-11}	0.82	1,134,260
158	rs77357563	12:114,837,349	<i>TBX5</i>	A/C	p.Tyr111Asp	missense	PP	0.005	-1.01	7.72×10^{-22}	0.22	1,152,080

Locus	rsID	Chr:Pos	Gene	EA/OA	Amino acids	Consequence	Trait	EAF	β	P	Het P	n
Exome array-wide association study (EAWAS)												
159	rs13141	12:121,756,084	ANAPC5	A/G	p.Val630Ala	missense	DBP	0.011	0.52	1.98x10 ⁻¹²	0.63	1,156,950
168	rs17880989 [†]	14:23,313,633	MMP14	A/G	p.Ile355Met	missense	DBP	0.027	0.32	2.02x10 ⁻¹⁴	0.95	1,160,530
169	rs61754158	14:31,774,324	HEATR5A	T/C	p.Arg1670Gly	missense	SBP	0.009	-0.70	6.28x10 ⁻⁹	0.04	1,119,230
170	rs72681869	14:50,655,357	SOS2	C/G	p.Arg191Pro	missense	SBP	0.010	-1.22	2.25x10 ⁻²²	0.25	1,144,040
177	rs150843673	15:81,624,929	TMC3	T/G	p.Ser1045Ter	stop/lost	DBP	0.021	0.36	1.43x10 ⁻¹²	0.14	1,154,000
181	rs61739285	16:27,480,797	GTF3C1	T/C	p.His1630Arg	missense	DBP	0.035	0.24	4.71x10 ⁻¹⁰	0.04	1,155,020
186	rs62051555	16:72,830,539	ZFX3	G/C	p.His2014Gln	missense	PP	0.048	0.47	1.19x10 ⁻²⁵	0.43	797,332
206	rs11699758	20:60,901,762	LAMA5	T/C	p.Ile1757Val	missense	PP	0.034	-0.26	6.68x10 ⁻¹¹	0.54	1,154,410
	rs13039398	20:60,902,402	LAMA5	A/G	p.Trp1667Arg	missense	PP	0.033	-0.26	1.89x10 ⁻¹⁰	0.44	1,133,830
Rare variant – genome-wide association study (RV-GWAS)												
215	rs55833332	1:198,222,215	NEK7	G/C	p.Gly35Arg	missense	PP	0.008	0.62	4.58x10 ⁻⁸	0.08	670,129
	rs143554274	1:198,455,391	ATP6V1G3	T/C	-	intergenic	PP	0.008	0.71	1.26x10 ⁻⁹	0.14	670,128
216	rs12135454	1:219,310,461	LYPLAL1-AS1	T/C	-	intron	PP	0.010	-0.62	1.61x10 ⁻⁸	0.22	665,523
	rs12128471	1:219,534,485	RPI1-392017.1	A/G	-	intergenic	PP	0.010	-0.68	2.99x10 ⁻⁹	0.19	670,130
217	rs114026228	4:99,567,918	TSPAN5	C/T	-	intron	PP	0.008	-0.65	5.20x10 ⁻⁹	0.03	670,128
	rs145441283	4:99,751,794	EIF4E	G/A	-	intergenic	PP	0.010	-0.71	2.01x10 ⁻¹¹	0.08	670,128
219	rs187207161	6:122,339,304	HMGB3P18	C/T	-	intergenic	PP	0.009	-0.63	2.16x10 ⁻¹⁰	0.02	670,130
221	rs149165710	8:121,002,676	DEPTOR	A/G	-	intron	PP	0.003	1.32	2.78x10 ⁻¹²	0.03	665,523
222	rs184289122	10:106,191,229	CFAP58	G/A	-	intron	SBP	0.008	1.31	1.66x10 ⁻¹³	0.53	670,472
	rs7076147	10:106,250,394	RPI1-12704.3	G/A	-	intergenic	SBP	0.010	1.11	1.71x10 ⁻¹⁴	0.75	670,472
	rs75337836	10:106,272,188	RPI1-12704.3	T/G	-	intergenic	SBP	0.010	1.12	2.67x10 ⁻¹⁵	0.54	670,472
	rs142760284	10:106,272,601	RPI1-12704.3	A/C	-	intergenic	SBP	0.009	1.22	2.19x10 ⁻¹⁵	0.92	670,472
	rs576629818	10:106,291,923	RPI1-12704.3	T/C	-	intergenic	SBP	0.009	1.24	1.02x10 ⁻¹⁵	0.71	670,472
	rs556058784	10:106,322,283	RPI1-12704.2	G/A	-	intergenic	SBP	0.009	1.26	4.54x10 ⁻¹⁶	0.57	665,861
	rs535313355 [‡]	10:106,399,140	SORCS3	C/T	-	upstream gene	SBP	0.009	1.36	1.04x10 ⁻¹⁷	0.22	670,472
	rs181200083 [‡]	10:106,520,975	SORCS3	C/A	-	intron	SBP	0.009	1.60	1.08x10 ⁻²¹	0.58	665,861
	rs540369678 [‡]	10:106,805,351	SORCS3	T/A	-	intron	SBP	0.010	1.18	2.29x10 ⁻¹⁴	0.16	670,472
	rs117627418	10:107,370,555	RPI1-45P22.2	T/C	-	intergenic	SBP	0.009	1.11	1.98x10 ⁻¹¹	0.1	665,861

Locus	rsID	Chr:Pos	Gene	EA/OA	Amino acids	Consequence	Trait	EAF	β	<i>P</i>	Het <i>P</i>	<i>n</i>
Exome array-wide association study (EAWAS)												
224	rs138656258	14: 31,541,910	<i>AP4SI</i>	G/T	-	intron	SBP	0.007	-0.93	1.15x10 ⁻⁸	0.13	665,861
228	rs6061911	20: 60,508,289	<i>CDH4</i>	C/T	-	intron	SBP	0.010	-0.85	4.67x10 ⁻⁸	0.09	665,861
	rs114580352	20: 60,529,963	<i>TAF4</i>	A/G	-	intron	SBP	0.009	-0.84	1.99x10 ⁻⁸	0.04	665,860
	rs11907239	20: 60,531,853	<i>TAF4</i>	A/G	-	intron	SBP	0.009	-0.82	4.99x10 ⁻⁸	0.05	670,472
	rs200383755	20: 61,050,522	GATA5	C/G	p.Trp19Ser	missense	DBP	0.006	1.00	1.01x10⁻¹³	0.49	670,172

Newly identified rare and low-frequency SNV-inverse normal transformed blood pressure associations are reported from Stage 2 of the exome array study and genome-wide association study. The reported associations are for the trait with the smallest *P*-value in the Stage 1 meta-analysis; the full results are provided in Supplementary Tables 2 and 7. SNVs are ordered by trait, chromosome, and position. Gene, gene containing the SNV or the nearest gene; rsID, dbSNP rsID; Chr:Pos, Chromosome:NCBI Build 37 position; EA/OA, effect allele (also the minor allele) and other allele; EAF, effect allele frequency based on Stage 1; Consequence, consequence of the SNV to the transcript as annotated by VEP; Amino acids, reference and variant amino acids from VEP; Trait, blood pressure trait for which association is reported; β , effect estimate, in mmHg, from the Stage 2 meta-analysis of the *untransformed*BP trait or the *Z*-score from the HTN analyses in Stage 2; *P*, *P*-value for association with the listed inverse normal transformed blood pressure trait from the Stage 2 meta-analyses; Het_*P*, *P*-value for heterogeneity; *n*, sample size. Bold type indicates rare missense variants.

[†] Novel variants identified in this study that are in linkage disequilibrium (LD: $r^2 > 0.6$ rare SNVs and $r^2 > 0.1$ common SNVs) with a variant that has been reported by Evangelou et al.²⁰ and/or Giri et al.²¹ within +/- 500 kb of the novel variant.

Table 2
Conditionally independent rare and very low-frequency SNV (MAF < 0.02) associations from exome array at known loci in Stage 1 EUR studies

Locus ID	rsID	Chr:bp	Gene	EA/OA	AA	Consequence	Trait	EAF	β_{joint}	P_{joint}	n	Ref
18	rs116245325	1: 153665650	<i>NPR1</i> ⁺	T/C	p.Phe1034Leu	Missense	SBP	0.001	0.1660	7.49x10 ⁻⁹	758,252	14
	rs61757359	1: 153658297		A/G	p.Ser541Gly	Missense		0.003	-0.0812	6.10x10 ⁻⁹	794,698	
	rs35479618	1: 153662423		A/G	p.Lys967Glu	Missense		0.017	0.0694	1.19x10 ⁻²⁸	774,862	
28	rs1805090	1: 230840034	<i>AGT</i> ⁺	T/G	p.Met392Leu	Missense	DBP	0.002	0.1070	6.00x10 ⁻¹⁰	759,349	8
	rs699	1: 230845794		G/A	p.Thr268Met	Missense	DBP	0.408	0.0225	2.12x10 ⁻⁴⁵	806,731	
94	rs111620813	4: 8293193	<i>HTRA3</i> ⁺	A/G	p.Met269Val	Missense	PP	0.011	-0.0432	1.38x10 ⁻⁸	798,063	18
	rs7437940	4: 7887500	<i>AFAPI</i>	T/C	-	Intron	PP	0.406	-0.0131	1.62x10 ⁻¹⁶	806,708	
102	rs112519623	4: 103184239	<i>SLC39A8</i> ⁺	A/G	p.Phe449Leu	Missense	DBP	0.016	-0.0391	3.02x10 ⁻¹⁰	803,151	6
	rs13107325	4: 103188709		T/C	p.Thr391Ala	Missense	DBP	0.072	-0.0615	9.69x10 ⁻⁸⁸	806,731	
	rs4699052	4: 104137790	<i>CENPE</i>	T/C	-	Intergenic	DBP	0.388	-0.0121	7.31x10 ⁻¹⁴	806,731	
105	rs6825911	4: 111381638	<i>ENPEP</i>	T/C	-	Intron	DBP	0.205	-0.0215	1.47x10 ⁻²⁸	801,965	
	rs33966350	4: 111431444		A/G	p.T _{er} -4.1.3.T _{rp}	Stop/lost	DBP	0.013	0.0735	2.40x10 ⁻²⁵	798,385	
144	rs4712056	6: 53989526	<i>MLIP</i>	G/A	p.Val159Ile	Missense	PP	0.360	0.0091	1.86x10 ⁻⁸	806,708	14,16,13
	rs115079907	6: 55924005	<i>COL21A1</i> ⁺	T/C	p.Arg882Gly	Missense	PP	0.003	0.2060	8.33x10 ⁻¹⁷	783,546	
	rs12209452	6: 55924962		G/A	p.Pro821Leu	Missense	PP	0.049	0.0411	5.49x10 ⁻²⁶	743,036	
	rs200999181	6: 55935568		A/C	p.Val665Gly	Missense	PP	0.001	0.3350	4.74x10 ⁻⁴³	764,864	
	rs35471617	6: 56033094		A/G	p.Met343Thr	Missense/splice region	PP	0.073	0.0249	1.03x10 ⁻¹⁵	806,708	
	rs2764043	6: 56035643		G/A	p.Pro277Leu	Missense	PP	0.002	0.1530	5.11x10 ⁻¹⁴	785,643	
	rs1925153	6: 56102780		T/C	-	Intron	PP	0.448	-0.0096	1.03x10 ⁻⁸	786,734	
	rs4294007	6: 57512510	<i>PRM2</i>	T/G	-	Splice acceptor	PP	0.379	0.0096	1.13x10 ⁻⁷	632,625	
	rs507666	9: 136149399	<i>ABO</i>	A/G	-	Intron	DBP	0.189	-0.0293	7.53x10 ⁻⁴⁷	796,103	13,15
208	rs3025343	9: 136478355	<i>LLONNC01-254D11.1</i>	A/G	-	Exon (noncoding transcript)	DBP	0.112	-0.0126	4.91x10 ⁻⁷	806,731	
	rs77273740	9: 136501728	<i>DBH</i>	T/C	p.Trp65Arg	Missense	DBP	0.027	-0.0846	3.85x10 ⁻¹¹	790,500	

Locus ID	rsID	Chr:bp	Gene	EA/OA	AA	Consequence	Trait	EAF	β _joint	P _joint	n	Ref
	rs3025380	9:136501756	<i>DBH</i>	C/G	p-Ala74Gly	Missense	DBP	0.005	-0.1030	5.37×10^{-18}	795,263	
	rs74853476	9:136501834	<i>DBH</i>	T/C	-	Splice donor	DBP	0.002	0.1000	3.69×10^{-8}	775,793	
223	rs201422605	10:95993887	<i>PLCE1</i>	G/A	p.Val678Met	Missense	SBP	0.003	-0.0837	1.41×10^{-7}	795,009	7,14
	rs11187837	10:96035980		C/T	-	Intron	SBP	0.110	-0.0198	4.23×10^{-14}	801,969	
	rs17417407	10:95931087		T/G	p.Leu548Arg	Missense	SBP	0.167	-0.0122	9.97×10^{-9}	806,735	
	rs9419788	10:96013705		G/A	-	Intron	SBP	0.387	0.0137	9.63×10^{-16}	806,735	
229	rs60889456	11:723311	<i>EPSS1L2</i> ⁺	T/C	p.Leu471Pro	Missense	PP	0.017	0.0303	6.37×10^{-7}	799,021	17
	rs7126805	11:828916	<i>CRACR2B</i>	G/A	p.Gln77Arg	Missense	PP	0.271	-0.0134	1.43×10^{-13}	752,026	
246*	rs56061986	11:89182686	<i>NOX4</i> ⁺	C/T	p.Gly67Ser	Missense	PP	0.003	-0.1080	2.25×10^{-11}	798,273	17,16
	rs139341533	11:89182666		A/C	p.Phe97Leu	Missense	PP	0.004	-0.0947	6.82×10^{-14}	785,947	
	rs10765211	11:89228425		A/G	-	Intron	PP	0.342	-0.0176	8.77×10^{-27}	806,708	
250	rs117249984	11:107375422	<i>ALKBH8</i>	A/C	p.Tyr653Asp	Missense	SBP	0.019	-0.0304	2.90×10^{-7}	805,695	16
	rs3758911	11:107197640	<i>CWFI9L2</i>	C/T	p.Cys894Tyr	Missense	SBP	0.341	0.0113	1.54×10^{-11}	806,735	
304	rs61738491	16:30958481	<i>FBXL19</i> ⁺	A/G	p.Gln652Arg	Missense	PP	0.010	-0.0460	1.25×10^{-8}	796,459	17,16
	rs35675346	16:30936081		A/G	p.Lys10Glu	Missense	PP	0.241	-0.0125	1.06×10^{-11}	802,932	
130*	rs114280473	5:122714092	<i>CEP120</i> ⁺	A/G	p.Phe712Leu	Missense	PP	0.006	-0.0584	9.98×10^{-8}	805,632	13, 12, 14, 15
	rs2303720	5:122682334		T/C	p.His947Arg	Missense	PP	0.029	-0.0419	3.44×10^{-18}	806,708	
	rs1644318	5:122471989	<i>PRDM6</i>	C/T	-	Intron	PP	0.387	0.0192	2.43×10^{-32}	790,025	
179*	rs3735080	7:150217309	<i>GIMAP7</i>	T/C	p.Cys83Arg	Missense	DBP	0.237	-0.0092	6.56×10^{-7}	806,731	9, 14, 10
	rs3807375	7:150667210	<i>KCNH2</i>	T/C	-	Intron	DBP	0.364	-0.0084	3.94×10^{-7}	806,731	
	rs3918234	7:150708035	<i>NOS3</i> ⁺	T/A	p.Leu982Gln	Missense	DBP	0.004	-0.0727	1.33×10^{-7}	786,541	
	rs891511	7:150704843		A/G	-	Intron	DBP	0.331	-0.0231	1.56×10^{-40}	778,271	
	rs10224002	7:151415041	<i>PRKAG2</i>	G/A	-	Intron	DBP	0.286	0.0186	7.41×10^{-27}	806,731	
190*	rs138582164	8:95264265	<i>GEM</i> ⁺	A/G	p.Ter199Arg	Stop lost	PP	0.001	0.2810	1.90×10^{-17}	735,507	16, 78
195*	rs112892337	8:135614553	<i>ZEAT</i> ⁺	C/G	p.Cys470Ser	Missense	SBP	0.005	-0.0831	4.39×10^{-12}	792,203	17
	rs12680655	8:135637337		G/C	-	Intron	SBP	0.398	0.0118	1.81×10^{-13}	797,982	
259*	rs145878042	12:48143315	<i>RAPGEF3</i> ⁺	G/A	p.Pro258Leu	Missense	SBP	0.012	-0.0453	9.28×10^{-10}	805,791	16, 13

Locus ID	rsID	Chr:bp	Gene	EA/OA	AA	Consequence	Trait	EAF	β _joint	P _joint	n	Ref
	rs148755202	12: 48191247	HDAC7	T/C	p.His166Arg	Missense	SBP	0.016	0.0310	9.07x10⁻⁷	806,735	
	rs1471997	12: 48723595	HIFNT	A/G	p.Gln174Arg	Missense	SBP	0.216	0.0130	1.15x10 ⁻¹¹	806,735	
	rs1126930**	12: 49399132	PRKAG1	C/G	p.Ser98Thr	Missense	SBP	0.035	0.0408	1.45x10 ⁻²¹	793,216	
	rs52824916***	12: 49993678	FAM186B	T/C	p.Gln582Arg	Missense	SBP	0.088	-0.0155	1.70x10 ⁻⁸	806,735	
	rs7302981**	12: 50537815	CERS5	A/G	p.Cys75Arg	Missense	SBP	0.375	0.0219	1.52x10 ⁻⁴¹	806,735	
312 *	rs61753655	17: 1372839	MYO1C⁺	T/C	p.Lys866Glu	Missense	SBP	0.011	0.0653	6.48x10⁻¹⁸	806,735	17, 16
	rs1885987	17: 2203025	SMG6	G/T	p.Thr341Asn	Missense	SBP	0.371	-0.0127	3.94x10 ⁻¹⁵	806,735	
339 *	rs34093919	19: 41117300	LTBP4⁺	A/G	p.Asn715Asp	Missense/splice region	PP	0.014	-0.0631	4.18x10⁻²⁰	805,764	19
	rs814501	19: 41038574	SPTBN4	G/A	p.Gly1331Ser	Missense	PP	0.482	-0.0115	2.40x10 ⁻¹³	806,708	
346	rs45499294	20: 30433126	FOXS1⁺	T/C	p.Lys74Glu	Missense	SBP	0.004	-0.0732	2.36x10⁻⁸	801,284	16

GCTA was used to perform conditional analyses of the meta-analysis results from the exome array study from the Stage 1 meta-analysis of EUR studies in known blood pressure regions (defined in Supplementary Table 1). All SNVs had $P < 0.0001$ for heterogeneity. The trait selected in this table is the trait for which the rare variant had the smallest P -value. We provide all conditionally independent variants at these loci, i.e. rare, very low frequency (MAF < 0.02), low frequency, and common. The full detailed listing of results is provided in Supplementary Table 8. Bold font highlights variants with MAF < 0.02. Locus ID, the known locus identifier used in Supplementary Table 1; Chr:Position, chromosome and NCBI Build 37 physical position; EA/OA, Effect allele/other allele; AA, amino acid change; Effect, predicted consequence of the SNV from VEP; EAF, effect allele frequency; β _joint, effect estimate for the SNV in the joint analysis from GCTA; P _joint, the P -value for association of the rare variant from the joint analysis in GCTA; Gene, nearest gene; Trait, blood pressure trait analyzed; Ref, reference of the first reports of association in the listed region.

* Indicates that one or more of the previously reported variants in the locus were not on exome array.

** Indicates that the listed variant is the known variant or its proxy ($r^2 > 0.8$ in 1000G EUR).

† Indicates that the listed gene had an unconditional SKAT P -value < 2×10^{-6} (see Supplementary Table 9).

Table 3
Newly identified independent BP-associated rare SNVs (MAF < 0.01) at known loci in UK Biobank only

Locus ID	rsID	Chr:Position	Gene	Info	EA/OA	Consequence	Trait	Unconditional SNV analysis			FINEMAP output		Ref
								EAF	β	P-value	Common SNVs in top configuration	PP of n SNVs	
5	rs41300100	1:11908146	<i>NPPA</i>	0.82	G/C	5' UTR	SBP	0.010	-0.10	4.70×10^{-21}	0.55	122.50	9,2,79
18	rs756799918	1:153464738	<i>RN7SL44P</i>	0.89	T/C	intergenic	SBP	0.0004	0.26	4.30×10^{-7}	0.36	27.49	14
28	rs1805090	1:230840034	<i>AGT</i>	NA	T/G	missense	SBP	0.0025	0.11	6.80×10^{-8}	0.79	26.23	8
28	rs539645495	1:230860071	<i>RPI1-99116_A.2</i>	0.97	G/A	intron, non-coding transcript	DBP	0.0024	0.13	3.20×10^{-9}	0.83	30.97	8
33	rs56152193	2:20925891	<i>LDAH</i>	0.76	C/G	intron	PP	0.0006	-0.23	8.10×10^{-7}	0.36	17.95	17,16
55	rs759606582	2:178325956	<i>AGPS</i>	0.96	G/A	intron	PP	0.0003	0.29	1.90×10^{-7}	0.57	7.48	16
72	rs559344473	3:48899332	<i>SLC25A20</i>	0.74	T/G	intron	DBP	0.0012	-0.17	2.50×10^{-6}	0.25	35.71	17,16,6,11
73	rs76920163	3:53857055	<i>CHDH</i>	0.96	G/T	intron	SBP	0.0059	0.10	3.80×10^{-13}	0.58	29.45	18,16
85	rs144980716	3:53776904	<i>CACNA1D</i>	0.91	A/G	intron	PP	0.0065	0.07	2.60×10^{-8}	0.57	18.42	13
86	rs45513277	3:143113550	<i>ATP1B3</i>	0.75	G/A	intron	PP	0.0008	0.20	6.00×10^{-6}	0.54	7.040	16
92	rs186525102	3:185539249	<i>IGF2BP2</i>	0.85	A/G	intron	SBP	0.0086	-0.06	6.70×10^{-7}	0.56	11.97	17
94	rs111620813	4:8293193	<i>HTRA3</i>	NA	A/G	missense	PP	0.0100	-0.05	2.00×10^{-6}	0.53	12.54	18
132	rs181585444	5:129963509	<i>AC005741.2</i>	0.83	C/T	intergenic	DBP	0.0003	-0.30	3.80×10^{-6}	0.55	10.70	14,13
137	rs46907130	6:8156072	<i>EEF1E1</i>	0.90	T/C	intergenic	SBP	0.0017	-0.14	1.90×10^{-7}	0.70	8.57	16
141	rs72854120	6:39248533	<i>KCNK17</i>	0.91	C/T	intergenic	SBP	0.0073	-0.08	3.10×10^{-9}	0.76	10.49	16
141	rs72854118	6:39248092	<i>KCNK17</i>	0.91	G/A	intergenic	DBP	0.0072	-0.07	2.70×10^{-7}	0.85	11.12	16
164	rs138890991	7:40804309	<i>SUGCT</i>	0.94	C/T	intron	PP	0.0100	0.06	1.60×10^{-7}	0.77	19.08	17
179	rs561912039	7:150682950	<i>NOS3</i>	0.74	T/C	intergenic	DBP	0.0017	-0.13	6.40×10^{-6}	0.34	81.75	9,14,10

Locus ID	rsID	Chr:Position	Gene	Info	EA/OA	Consequence	Trait	Unconditional SNV analysis			FINEMAP output			
								EAF	β	P-value	Common SNVs in top configuration	PP of n SNVs	$\log_{10}BF$	Ref
183	rs70342886	8:23380012	<i>SLC25A37</i>	0.85	C/G	intergenic	DBP	0.0001	-0.48	9.80×10^{-7}	rs7788497, rs891511	0.58	15.74	16
190	rs201196388	8:95265263	<i>GEM</i>	NA	T/C	splice donor	PP	0.0005	0.26	2.40×10^{-9}	rs7842120	0.34	31.80	16,78
193	rs32252660	8:120587297	<i>ENPP2</i>	0.79	T/C	intron	DBP	0.0025	-0.11	4.10×10^{-7}	rs2170363 rs7017173	0.81	26.53	6
193	rs181416549	8:120678125	<i>ENPP2</i>	0.84	A/G	intron	PP	0.0026	0.20	5.10×10^{-21}	rs35362581, rs80309268	0.95	113.21	6
212	rs138765972	10:20554597	<i>PLXDC2</i>	0.94	C/T	intron	DBP	0.0075	-0.07	4.40×10^{-8}	rs61841505	0.49	9.06	16
219	rs192036851	10:64085523	<i>RPI1-120C12.3</i>	0.92	C/T	intergenic	SBP	0.0062	0.06	6.40×10^{-6}	rs10995311	0.28	19.55	16,13
234	rs150090666	11:14865399	<i>PDE3B</i>	NA	T/C	stop gained	DBP	0.0010	-0.16	5.20×10^{-7}	rs11023147, rs2597194	0.55	12.93	16
242	rs139620213	11:61444612	<i>DAGLA</i>	0.89	T/C	upstream gene	PP	0.0019	0.11	5.90×10^{-6}	rs2524299	0.48	6.64	15
246	rs40659338	11:89183302	<i>NOX4</i>	0.85	C/T	intron	PP	0.0027	-0.14	2.60×10^{-10}	rs2289125, rs494144	0.62	58.09	17,16
260	rs186600986	12:53769106	<i>SPI</i>	0.91	A/G	upstream gene	PP	0.0030	-0.09	1.10×10^{-6}	rs73099903	0.48	12.91	19
266	rs137937061	12:111001886	<i>PPTC7</i>	0.74	A/G	intron	SBP	0.0048	-0.09	1.30×10^{-6}	rs9739637, rs35160901, rs10849937, rs3184504	0.34	55.74	16,4,5
268	rs190870203	12:123997554	<i>RILPL1</i>	0.85	T/G	intron	PP	0.0020	0.12	1.70×10^{-7}	rs4759375	0.72	9.50	13
270	rs541261920	13:30571753	<i>RPI1-629E24.2</i>	0.79	G/C	intergenic	SBP	0.0005	0.24	9.20×10^{-6}	rs7338758	0.54	10.09	16
281	rs149250178	14:100143685	<i>HHPL1</i>	0.75	A/G	3' UTR	DBP	0.0004	-0.29	2.30×10^{-6}	rs7151887	0.51	7.93	16
299	rs139491786	16:2086421	<i>SLC9A3r2</i>	NA	T/C	missense	DBP	0.0068	-0.12	1.60×10^{-20}	rs28590346, rs34165865, rs62036942, rs8061324	0.57	50.80	16
304	rs2234710	16:30907835	<i>BCL7C</i>	0.79	T/G	upstream gene	SBP	0.0075	-0.08	2.30×10^{-9}	-	0.52	6.29	17,16
304*	rs148753960	16:31047822	<i>STX4</i>	0.89	T/C	intron	PP	0.0099	-0.07	1.80×10^{-9}	rs7500719	0.42	12.21	17,16
317	rs756906294	17:42323081	<i>SLC4A1</i>	0.73	T/C	downstream gene	PP	0.0030	0.01	8.30×10^{-6}	rs66838809	0.27	18.94	17
322	rs16946721	17:61106371	<i>TANC2</i>	0.91	G/A	intron	DBP	0.0100	-0.07	1.40×10^{-11}	rs1867624, rs4291	0.51	20.91	17,16
333	rs55670943	19:11441374	<i>RAB3D</i>	0.87	C/T	intron	SBP	0.0085	-0.10	2.10×10^{-17}	rs12976810, rs4804157,	0.78	85.45	13-15

Locus ID	rsID	Chr:Position	Gene	Info	EA/OA	Consequence	Trait	Unconditional SNV analysis			FINEMAP output			
								EAF	β	P-value	Common SNVs in top configuration	PP of n SNVs	$\log_{10}BF$	Ref
346*	rs149972827	20:30413439	<i>MYLK2</i>	0.98	A/G	intron	SBP	0.0036	-0.10	6.20×10^{-9}	rs160838, rs167479	0.85	9.86	16
362	rs115089782	22:42329632	<i>CENPM</i>	0.93	T/C	intergenic	SBP	0.0001	0.53	4.20×10^{-6}	rs139919	0.44	14.12	17,13

FINEMAP²⁵ was used to identify the most likely causal variants within the known loci (defined in Supplementary Table 1) using the BOLT-LMM results in UKBB, the full detailed listing of results is provided in Supplementary Table 8. Locus ID, the known locus identifier provided in Supplementary Table 1; Chr:Position, chromosome and physical position in Build 37; Info, imputation information score, NA indicates that the SNV was genotyped and not imputed; EA/OA, Effect allele and other allele, respectively; AA, amino acid change; Effect, predicted effect of the listed SNV; EAF, effect allele frequency; β , single variant effect estimate for the rare variant in the BOLT-LMM analysis; P-value, the single variant P-value from the mixed model in the BOLT-LMM analysis; PP of n SNVs, the posterior probability of the number of causal variants; Log₁₀BF, log₁₀ Bayes factor for the top configuration; Gene, nearest gene; Trait, blood pressure trait analyzed; Ref, reference of the first reports of association in the listed region.

rs540659338 identified in UK Biobank in *NOX4* has $r^2 = 1$ in 1000G EUR with rs56061986 identified in the GCTA analysis in Table 4.

* Variants at these loci are in LD with GCTA variants (Table 2): at locus 304, $r^2 = 0.876$ between rs148753960 and rs61738491; at locus 346, $r^2 = 0.952$ between rs149972827 and rs45499294.