


Neuronal Excitability

# Synaptotagmin-7 Enhances Facilitation of $\text{Ca}_v2.1$ Calcium Channels

Alaeddine Djillani, Jeremy Bazinet, and  William A. Catterall<https://doi.org/10.1523/ENEURO.0081-22.2022>

Department of Pharmacology, University of Washington, Seattle, WA 98195-7280

## Visual Abstract



Synaptotagmin-7 interacts with the synprint site on  $\text{Ca}_v2.1$  channels and enhances calcium-dependent facilitation in an isoform-specific manner

Voltage-gated calcium channel  $\text{Ca}_v2.1$  undergoes  $\text{Ca}^{2+}$ -dependent facilitation and inactivation, which are important in short-term synaptic plasticity. In presynaptic terminals,  $\text{Ca}_v2.1$  forms large protein complexes that include synaptotagmins. Synaptotagmin-7 (Syt-7) is essential to mediate short-term synaptic plasticity in many

### Significance Statement

Short-term synaptic plasticity mediated by regulation of  $\text{Ca}_v2.1$  channels plays a crucial role in information processing, learning, and memory. Our results reveal a novel mode of regulation of  $\text{Ca}_v2.1$  channels by the high-affinity  $\text{Ca}^{2+}$  sensor synaptotagmin-7 (Syt-7) through direct interaction with the synprint site.  $\text{Ca}_v2.1$ /Syt-7 interaction enhances short-term facilitation of the P/Q-type  $\text{Ca}^{2+}$  current that triggers neurotransmitter release. This unexpected intersection of Syt-7 and  $\text{Ca}_v2.1$  may regulate short-term,  $\text{Ca}^{2+}$ -dependent synaptic plasticity, along with SNARE proteins and other calcium-binding proteins in presynaptic terminals. Understanding the mechanism by which Syt-7 enhances facilitation of  $\text{Ca}_v2.1$  channels is an important step toward deciphering the molecular mechanisms of short-term synaptic plasticity in the brain.

synapses. Here, based on evidence that  $\text{Ca}_v2.1$  and  $\text{Syt-7}$  are both required for short-term synaptic facilitation, we investigated the direct interaction of  $\text{Syt-7}$  with  $\text{Ca}_v2.1$  and probed its regulation of  $\text{Ca}_v2.1$  function. We found that  $\text{Syt-7}$  binds specifically to the  $\alpha_{1A}$  subunit of  $\text{Ca}_v2.1$  through interaction with the synaptic-protein interaction (synprint) site. Surprisingly, this interaction enhances facilitation in paired-pulse protocols and accelerates the onset of facilitation.  $\text{Syt-7}\alpha$  induces a depolarizing shift in the voltage dependence of activation of  $\text{Ca}_v2.1$  and slows  $\text{Ca}^{2+}$ -dependent inactivation, whereas  $\text{Syt-7}\beta$  and  $\text{Syt-7}\gamma$  have smaller effects. Our results identify an unexpected, isoform-specific interaction between  $\text{Ca}_v2.1$  and  $\text{Syt-7}$  through the synprint site, which enhances  $\text{Ca}_v2.1$  facilitation and modulates its inactivation.

**Key words:** calcium channels; P/Q-type calcium current; protein interactions; synaptic facilitation; synaptotagmin-7; synprint site

## Introduction

Inward  $\text{Ca}^{2+}$  currents conducted by voltage-gated  $\text{Ca}^{2+}$  ( $\text{Ca}_v$ ) channels couple action potentials and other depolarizing stimuli to many  $\text{Ca}^{2+}$ -dependent intracellular processes, including neurotransmission, hormone secretion, and muscle contraction (Zamponi et al., 2015). In presynaptic nerve terminals,  $\text{Ca}_v2.1$ ,  $\text{Ca}_v2.2$ , and  $\text{Ca}_v2.3$  channels conduct P/Q-type, N-type, and R-type  $\text{Ca}^{2+}$  currents that trigger rapid neurotransmission (for review, see Olivera et al., 1994; Zamponi et al., 2015; Nanou and Catterall, 2018). However, only P/Q-type  $\text{Ca}^{2+}$  currents conducted by  $\text{Ca}_v2.1$  channels can mediate short-term synaptic facilitation at the calyx of Held in mice (Inchauspe et al., 2004), pointing to a unique role of these  $\text{Ca}^{2+}$  channels in short-term synaptic plasticity.

In transfected nonneuronal cells,  $\text{Ca}^{2+}$  entry mediated by  $\text{Ca}_v2.1$  channels causes calcium-dependent facilitation (CDF) and inactivation (CDI) during single depolarizations and in trains of repetitive depolarizing pulses (Lee et al., 1999, 2000; DeMaria et al., 2001; Catterall and Few, 2008; Christel and Lee, 2012; Ben-Johny and Yue, 2014). Both CDF and CDI of  $\text{Ca}_v2.1$  channels are dependent on calmodulin (CaM; Lee et al., 1999, 2000; DeMaria et al., 2001). CaM preassociates with the C-terminal domain of the pore-forming  $\alpha_1$  subunit of  $\text{Ca}_v2.1$  channels (Erickson et al., 2001). Following  $\text{Ca}^{2+}$  binding, CaM initially interacts with the nearby IQ-like motif (IM) and causes CDF, whereas further binding of  $\text{Ca}^{2+}$ /CaM to the more distal CaM-binding domain (CBD) induces CDI of  $\text{Ca}_v2.1$

channels (DeMaria et al., 2001; Lee et al., 2003). Introducing the IM-AA mutation into the IQ-like motif of  $\text{Ca}_v2.1$  impairs CDF and CDI, providing a tool to assess the significance of these processes in synaptic transmission and short-term synaptic plasticity (Zühlke et al., 1999; DeMaria et al., 2001; Lee et al., 2003).

CDF of  $\text{Ca}_v2.1$  channels contributes significantly to short-term synaptic facilitation. Expression of  $\text{Ca}_v2.1$  in cultured superior cervical ganglion neurons, whose endogenous  $\text{Ca}_v2.2$  channels were specifically blocked by  $\omega$ -conotoxin GVIA, was sufficient to restore synaptic transmission and induce  $\text{Ca}^{2+}$ -dependent synaptic facilitation, which was impaired by introducing the IM-AA mutation in  $\text{Ca}_v2.1$  channels (Mochida et al., 2003a, 2008). In mice in which  $\text{Ca}_v2.1$  channels contained the IM-AA mutation, synaptic facilitation was substantially decreased at the neuromuscular junction as well as in hippocampal CA3-to-CA1 synapses and CA3-to-parvalbumin-expressing basket cell synapses (Nanou et al., 2016a,b, 2018). These results support an important role for facilitation of  $\text{Ca}_v2.1$  channels in short-term synaptic facilitation.

In addition to  $\text{Ca}_v2.1$  channels, the high-sensitivity  $\text{Ca}^{2+}$  sensor synaptotagmin-7 ( $\text{Syt-7}$ ) has been proposed to support short-term synaptic facilitation by binding residual  $\text{Ca}^{2+}$  in the nerve terminal following the action potential, thereby increasing interaction with the SNARE complex and enhancing  $\text{Ca}^{2+}$ -dependent synaptic vesicle exocytosis (Jackman et al., 2016). Previous studies using  $\text{Syt-7}$  KO mice have shown that  $\text{Syt-7}$  is required for short-term plasticity in several types of synapses in the hippocampus, cerebral cortex, and cerebellum (Jackman et al., 2016; Turecek and Regehr, 2018). Because  $\text{Ca}_v2.1$  channels and  $\text{Syt-7}$  are located near each other in the active zones of nerve terminals (Müller et al., 2010), and  $\text{Ca}_v2.1$  and  $\text{Syt-7}$  are both implicated in synaptic facilitation, we have tested the hypothesis that these two proteins interact directly with each other and regulate  $\text{Ca}^{2+}$  entry through  $\text{Ca}_v2.1$  channels. Our results reveal direct interactions of  $\text{Syt-7}$  with  $\text{Ca}_v2.1$  that enhance facilitation of the  $\text{Ca}_v2.1$   $\text{Ca}^{2+}$  current. These data suggest that interaction of  $\text{Syt-7}$  with  $\text{Ca}_v2.1$  channels may contribute to short-term synaptic facilitation.

## Materials and Methods

### Cell lines and transfection

Cells from tsA-201 cell line were maintained in DMEM (Invitrogen by Life Technologies) supplemented with 10%

Received February 18, 2022; accepted April 22, 2022; First published April 27, 2022.

The authors declare no competing financial interests.

Author contributions: A.D., J.B., and W.A.C. designed research; A.D. and J.B. performed research; A.D. analyzed data; A.D. and W.A.C. wrote the paper.

This work was supported by the National Institutes of Health Research Grant R35 NS111573.


Acknowledgment: We thank our colleague Dr. Jin Li (Department of Pharmacology, University of Washington) for advice and technical assistance with the project. We also thank Dr. Edwin Chapman (University of Wisconsin) and Dr. Katsuhiko Mikoshiba (University of Tokyo and Brain Science Institute, Riken) for generously providing  $\text{Syt-7}$  cDNAs.

Correspondence should be addressed to William A. Catterall at [wcatt@uw.edu](mailto:wcatt@uw.edu).

<https://doi.org/10.1523/ENEURO.0081-22.2022>

Copyright © 2022 Djillani et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license, which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.



**Figure 1.** Direct binding of Syt-7 $\alpha$  to the synprint site of P/Q-type Ca<sub>v</sub>2.1 channels. **A, B**, In brain lysates (**A**) and transfected tsA-201 cells (**B**), co-immunoprecipitation experiments show the binding of Syt-7 $\alpha$  and  $\alpha_{1A}$  subunit of Ca<sub>v</sub>2.1 channels. **C**, Binding of Syt-7 $\alpha$  (GST-Syt-7 $\alpha$ ) to the His-tagged synprint site (724–981) of  $\alpha_{1A}$  subunit of P/Q-type Ca<sub>v</sub>2.1 channel. GST-Syt-7 $\alpha$  were immobilized on glutathione-Sepharose beads and incubated with His-Ca<sub>v</sub>2.1 or His-Ca<sub>v</sub>1.2 using different Ca<sup>2+</sup> concentrations (10  $\mu$ M to 1 mM). The binding experiments were performed in a Ca<sup>2+</sup> buffering system containing: 5 mM N-hydroxyethyl ethylenediaminetriacetic, 50 mM HEPES (pH 7.2), and 150 mM NaCl. The free Ca<sup>2+</sup> concentrations were estimated using the MAX CHELATOR software. The Ca<sup>2+</sup> concentrations used were 10  $\mu$ M, 50  $\mu$ M, 100  $\mu$ M, 300  $\mu$ M, 500  $\mu$ M, 700  $\mu$ M, and 1 mM. After extensive washing, GST-Syt-7 $\alpha$  bound to the beads were eluted with 15 mM reduced glutathione (GSH) in 50 mM Tris-HCl (pH 8) and proceed to electrophoresis and immunoblotting. The bound His-Ca<sub>v</sub>2.1 synprint (724–981) proteins were detected with anti-His antibody. Because different Ca<sup>2+</sup> concentrations were used in co-immunoprecipitation experiments, segments from different immunoblots were spliced together to show comparisons clearly. Those protein bands are delineated for clarification. The immunoblots presented here are representative of at least three experiments for each co-immunoprecipitation or immunoblot. A related co-immunoprecipitation experiment conducted with different experimental conditions is presented in Extended Data Figure 1-1.

fetal bovine serum (Fisher Scientific), 1% glutamine (Sigma-Aldrich), 1% penicillin and streptomycin (Sigma-Aldrich). The cells were maintained at 37°C under 5% CO<sub>2</sub>. Cells were plated in 35-mm tissue culture dishes to achieve 70% confluency and then transfected using TransIT-LT1 transfection reagent (Mirus) with a total of 5- $\mu$ g plasmid including: 2, 1.5, 1  $\mu$ g of  $\alpha_{1A}$ ,  $\beta_{2A}$ , and  $\alpha_{2\delta}$  subunits composing the Ca<sub>v</sub>2.1 channel and a ratio 3  $\mu$ l of transfection reagent to 1  $\mu$ g of cDNA plasmid. 0.22  $\mu$ g eGFP was added to the plasmid mix to identify the transfected cells.


### Construction and expression of fusion proteins

Recombinant glutathione S-transferase (GST)-Syt-7 $\alpha$  fusion proteins were synthesized from the expression plasmid in the vector pGEX-2T. His-fusion proteins containing the synprint site region from the intracellular loop between domain II and III of the P/Q-type Ca<sub>v</sub>2.1 (synprint 724–981) or the equivalent synprint site from the L-type Ca<sub>v</sub>1.2 (680–800) used as a control, were expressed using the expression plasmid pET-28b. GST and His recombinant proteins were expressed in *Escherichia coli* BL26 cells, a protease-deficient strain (NEB). Fusion proteins were extracted by mild sonication (10 times 10 s with 1-min break) in lysis buffer containing: Tris 50 mM

(pH 7.4), NaCl 150 mM, Na-deoxycholate 1%, NaF 10 mM, EDTA 1 mM, Triton X-100 1%, and glycerol 5%, supplemented with protease inhibitors Calpain I, Calpain II, and cComplete protease inhibitor cocktail (Sigma-Aldrich). GST-Syt-7 $\alpha$  proteins were purified using glutathione Sepharose beads (Millipore Sigma) and eluted with 15 mM reduced glutathione (GSH) in 50 mM Tris (pH). His-synprints from Ca<sub>v</sub>2.1 and Ca<sub>v</sub>1.2 were purified by binding to Ni<sup>2+</sup>-charged HisPur Ni-NTA Resin (ThermoFisher) and eluted with 250 and 500 mM imidazole in PBS. The amount of proteins used was standardized based on Coomassie Blue-stained SDS gels or estimated with a standard curve relating the intensity of the immunoblotting signal to the amount of a standard fusion protein applied.

### Co-immunoprecipitation experiments

Immunoprecipitation experiments were performed using Dynabeads Protein G (Invitrogen) in TBS buffer with a Ca<sup>2+</sup>-buffering system containing 50 mM Tris/HCl, 140 mM NaCl, 50 mM HEPES (pH 7.2), 5 mM N-(2-hydroxyethyl)ethylenediamine-N,N',N'-triacetic acid (HydroxyEDTA), 0.3% Triton X-100 and different Ca<sup>2+</sup> concentrations varying from 10  $\mu$ M to 5 mM. The Ca<sup>2+</sup>-buffering system



**Figure 2.** Syt-7 $\alpha$  accelerates the onset of facilitation of Ca<sub>v</sub>2.1 channels. Inset top, Pulse protocol. Currents recorded with 10 mM extracellular Ca<sup>2+</sup> and 0.5 mM EGTA in the intracellular recording solution were elicited by test pulses to +10 mV before (P1) and 5 ms after (P2) 10-mV preconditioning prepulses of the indicated durations. Inset, Example traces from control and Syt-7 $\alpha$  transfected tsA cells following P1 and P2 pulses. **A**, Effect of Syt-7 $\alpha$  on facilitation as a function of prepulse duration. Facilitation was obtained by normalizing the peak current from P2 to that from P1. Single-exponential fits of the data are shown. **B**, in tsA-201 cells co-expressing Ca<sub>v</sub>2.1 channel with Syt-7 $\alpha$ , the slope is significantly increased compared with control cells. Data are represented as mean  $\pm$  SEM.

was used to produce free Ca<sup>2+</sup> concentrations calculated using MAX CHELATOR software (UC Davis). Dynabeads were incubated with antibodies directed against Ca<sub>v</sub>2.1 channels or Syt-7 $\alpha$  for 1 h at 4°C. Then, whole brain lysates or transfected tsA cell lysates were added to the beads and incubated at 4°C under rotation overnight. Nonspecific proteins were washed three times with a washing buffer. Proteins attached to the beads were eluted using an elution buffer. Proteins were blotted with antibodies against Syt-7 (mouse monoclonal antibody N275/14, Product Number MABN665, Millipore Sigma) or Ca<sub>v</sub>2.1 (rabbit polyclonal antibody catalog #ACC-001, Alomone Labs). The antibodies used for immunoblotting were titrated to assure that the concentration used was in the linear response range. The co-immunoprecipitation experiments and western blots have been repeated at least three times showing reproducible results.

### Study of Syt-7 binding to the synprint site by affinity chromatography


GST-Syt-7 $\alpha$  proteins were bound to glutathione-Sepharose beads (Millipore Sigma) in TBS-Ca<sup>2+</sup> buffer incubated at 4°C for 1 h under constant rotation. To remove unbound proteins, the mixture was washed two times with a washing buffer. Glutathione-Sepharose

beads coupled with GST-fusion proteins were added to similar amount of purified His-Ca<sub>v</sub>2.1 synprint (724–981) or His-Ca<sub>v</sub>1.2 synprint (680–800). The mixture was incubated under constant rotation for 1 h at 4°C. The binding experiments were conducted in presence of TBS-Ca<sup>2+</sup> buffering system with 0.1% Triton X-100. The beads were washed three times with washing buffer and bound complexes were eluted with 15 mM of reduced glutathione and 50 mM Tris-HCl (pH 8). Eluates were separated from beads by centrifugation at 10,000  $\times$  g for 1 min and processed for 10–20% SDS/tricine gradient gel electrophoresis and immunoblotted with anti-His antibody.

### Electrophysiological recording

Calcium current (*I*<sub>Ca</sub>) or Barium current (*I*<sub>Ba</sub>) were recorded at least 48 h after tsA-201 cell transfection using whole-cell configuration of the patch-clamp technique. Data acquisition was conducted using patch-clamp amplifier (HEKA Elektronik GmbH). Voltage-clamp protocols and facilitation protocols were applied, and data were acquired using Pulse (HEKA Elektronik GmbH). Currents were filtered at 5 kHz. Leak and capacitance transient currents were subtracted using a P/4 protocol.

Recording pipettes were pulled from borosilicate glass to achieve initial bath resistances of 1.5–3.0 M $\Omega$  and filled



**Figure 3.** Effect of Syt-7 $\alpha$  on prepulse facilitation of Ca<sub>v</sub>2.1 at physiological Ca<sup>2+</sup> levels. Inset top, Pulse protocol. Currents recorded with 2 mM extracellular Ca<sup>2+</sup> and 0.5 mM EGTA in the intracellular recording solution were elicited by test pulses to +10 mV before (P1) and 5 ms after (P2) 10-mV conditioning prepulses of the indicated durations. Inset, Example traces from control and Syt-7 $\alpha$  transfected tsA cells following P1 and P2 pulses. Main panel, Graph shows the effect of Syt-7 $\alpha$  on facilitation as a function of prepulse duration. Facilitation was obtained by normalizing the peak current from P2 to that from P1. Single-exponential fits of the data are shown. Data are represented as mean  $\pm$  SEM.


with an intrapipette solution containing (in mM): 120 N-methyl-D-glucamine (NMDG), 60 HEPES, 1 MgCl<sub>2</sub>, 2 Mg-ATP, and 0.5 EGTA. The extracellular patch-clamp solution contained (in mM): 150 Tris, 1 MgCl<sub>2</sub>, and 10 CaCl<sub>2</sub> or BaCl<sub>2</sub> depending on the experimental protocols. The pH of both intrapipette and extracellular solutions was adjusted to 7.3 using methanesulfonic acid. tsA-201 cell membrane capacitance (C<sub>m</sub>) varied from 15–25 pF and access resistance (R<sub>s</sub>) varied from 8 to 20 M $\Omega$ . All averaged data represent the mean  $\pm$  SEM of at least 10 cells. For peak current measurement using current-voltage (I/V) curves, Ca<sub>v</sub>2.1 P/Q-type current was generated using steps of depolarization from -80 to +60 mV every 10-mV step with a holding potential at -80 mV. A total of 10 mM CaCl<sub>2</sub> or BaCl<sub>2</sub> was used in the external patch-clamp solution. For facilitation protocol experiments, 10 mM of CaCl<sub>2</sub> were used in the external solution. Several facilitation protocols have been used to study the role of the three isoforms of Syt-7 (Syt-7 $\alpha$ , Syt-7 $\beta$ , and Syt-7 $\gamma$ ) in Ca<sup>2+</sup> current facilitation. Paired-pulse facilitation protocols were evoked by applying two 1-s-spaced depolarizing pulses P1 and P2 from -80 to +10 mV. A preconditioning 50-ms

depolarizing step from -80 to +10 mV was applied only 5 ms before P2. In order to study the voltage dependence of Ca<sub>v</sub>2.1 channel activation, P1 and P2 were applied using variable voltages from -120 to +40 mV. P2 over P1 ratios were calculated and compared between transfected tsA-201 cells with and without Syt-7 $\alpha$ , Syt-7 $\beta$ , or Syt-7 $\gamma$ . The second protocol of paired-pulse facilitation was used to study the effect of changing voltages in the preconditioning pulse on P2. P1 and P2 pulses were maintained from -80 to +10 mV; however, the preconditioning pulse was applied with variable voltages from -120 to +40 mV. Finally, onset of facilitation was studied by increasing preconditioning pulse duration to 10 ms and measuring the ratios of P2 over P1.

#### Statistical analysis

Statistical analyses were performed using GraphPad Prism 5 (GraphPad Software) and Origin Pro (OriginLab Inc.). All data are shown as the mean  $\pm$  SEM. The statistical details of the experiments can be found the results section and figure legends. A Student's *t* test was used to





**Figure 4.** Effect of Syt-7 $\alpha$  on the decay from facilitation of Ca<sub>v</sub>2.1 channels. Inset top, Pulse protocol for measuring decay of facilitation. Ca<sup>2+</sup> currents are elicited by test pulses to +10 mV before (P1) and after (P2) a conditioning prepulse to +10 mV for 5 ms. Inset bottom left, Decay from facilitation measured by comparing  $\tau$  between control and Syt-7 $\alpha$  transfected cells. Inset bottom right, Comparison of P2/P1 facilitation ratio at  $\Delta t = 0$  between control and Syt-7 $\alpha$ -expressing cells. Main panel, Effect of Syt-7 $\alpha$  on the decay from facilitation. The facilitation ratio was obtained by normalizing the peak current from P2 to that from P1 and was plotted against the interval between the conditioning prepulse and P2. Shown are results obtained with 50-ms conditioning prepulse. Graph shows the effect of Syt-7 $\alpha$  on decay of facilitation as a function of interpulse duration. Data are represented as mean  $\pm$  SEM.

compare two sets of data. The significance was defined using a threshold of  $p = 0.05$  throughout the study. Error bars indicate SEM. Sample sizes are described in the figure legends.

## Results

### Syt-7 binds to the Ca<sub>v</sub>2.1 channel in mouse brain


To determine whether the slow, high-affinity Ca<sup>2+</sup> sensor Syt-7 binds to the presynaptic Ca<sub>v</sub>2.1 channels *in vivo*, co-immunoprecipitation studies were performed on membrane preparations from mouse brain lysates. Ca<sub>v</sub>2.1 channels extracted from these neuronal membranes were immunoprecipitated with specific anti-Ca<sub>v</sub>2.1 antibodies, and the resulting complexes were probed with anti-Syt-7 antibody using the Dynabeads/Protein G co-immunoprecipitation protocol. The resulting immunoblots revealed Ca<sub>v</sub>2.1/Syt-7 $\alpha$  interaction with anti-Syt-7 when anti-Ca<sub>v</sub>2.1 was used as the precipitating antibody (Fig. 1A, top). In a complementary experiment, Ca<sub>v</sub>2.1 channels were immunoprecipitated with anti-Syt-7 antibodies and detected in immunoblots with anti-Ca<sub>v</sub>2.1 antibodies (Fig. 1A, bottom). Co-immunoprecipitation experiments using a

modified experimental protocol yielded comparable results (Extended Data Fig. 1-1). Together, these results show that Ca<sub>v</sub>2.1 channels and Syt-7 are associated with each other in mouse brain membranes.

### Syt-7 binds to the synprint site of the Ca<sub>v</sub>2.1 channel

Human embryonic kidney tsA-201 cells were transfected with the  $\alpha_{1A}$ ,  $\beta_{2A}$ , and  $\alpha_{2\delta_1}$  subunits of Ca<sub>v</sub>2.1 channels together with Syt-7 $\alpha$ , the most abundant isoform of Syt-7 (Fukuda et al., 2002). A specific complex of Ca<sub>v</sub>2.1 and Syt-7 was co-immunoprecipitated from lysates of tsA-201 cells transfected with Ca<sub>v</sub>2.1 $\alpha_{1A}$  subunit and Syt-7 $\alpha$ , using either anti-Syt-7 or anti-Ca<sub>v</sub>2.1 as the precipitating antibody (Fig. 1B). These results demonstrate a physical interaction between Syt-7 $\alpha$  and Ca<sub>v</sub>2.1 in intact cells expressing these proteins *in vitro*, and suggest that other neuron-specific proteins are not required for this protein-protein interaction.

To investigate which domain of the pore-forming  $\alpha_1$  subunit of Ca<sub>v</sub>2.1 channels binds Syt-7 $\alpha$ , *in vitro* binding experiments were performed using recombinant fusion proteins (Materials and Methods). Full-length Syt-7 $\alpha$  protein




**Figure 5.** Syt-7 $\alpha$  potentiates Ca<sub>v</sub>2.1 facilitation in a paired-pulse protocol following change in prepulse voltage. Inset top, Pulse protocol shown represents paired pulse protocol. Ca<sup>2+</sup> current was recorded using 10 mM Ca<sup>2+</sup> and 0.5 mM EGTA in the external and internal solutions, respectively. Pulse 1 (P1; depolarization from -80 to +10 mV) elicits the first Ca<sup>2+</sup> current. A second 5-ms pulse (P2) generating a second I<sub>Ca</sub> is applied 2 ms after a 50-ms conditioning prepulse with variable voltages (-40 to 60 mV). Inset bottom, Example traces from control and Syt-7 $\alpha$  transfected tsA cells following P1 and P2 pulses. Main panel, Graph shows the effects of Syt-7 $\alpha$  isoform on facilitation as a function of prepulse voltage. The facilitation ratio was obtained by normalizing the peak current from P2 to that from P1. Data are represented as mean  $\pm$  SEM.

was expressed as a GST-fusion protein, and the synprint site in Ca<sub>v</sub>2.1 (724–981) was expressed as a His-fusion protein. As a control, the equivalent synprint site from the cardiac Ca<sup>2+</sup> channel, Ca<sub>v</sub>1.2 (680–800), was expressed as a His-fusion protein. GST-Syt-7 $\alpha$  proteins were immobilized by binding to glutathione-Sepharose beads and incubated with a constant concentration of His-Ca<sub>v</sub>2.1 synprint peptide (724–981) or His-Ca<sub>v</sub>1.2 synprint peptide (680–800) using different free Ca<sup>2+</sup> concentrations varying from 10  $\mu$ M to 1 mM. After extensive washing, binding of His-Ca<sub>v</sub>2.1 synprint (724–981) to GST-Syt-7 $\alpha$  was revealed by immunoblot analysis using an anti-His antibody. As shown in Figure 1C, GST-Syt-7 $\alpha$  bound to His-Ca<sub>v</sub>2.1 (724–981) synprint in a Ca<sup>2+</sup>-dependent manner *in vitro*, with binding first detected at 50  $\mu$ M free Ca<sup>2+</sup> concentration and increasing to a maximum at 500  $\mu$ M Ca<sup>2+</sup>. In contrast, the negative control peptide His-Ca<sub>v</sub>1.2 (680–800) from the corresponding segment of cardiac Ca<sub>v</sub>1.2 channels did not bind to GST-Syt-7 $\alpha$ . These results demonstrate specific binding of Syt-7 to the synprint site from Ca<sub>v</sub>2.1 channels in preference to the corresponding segment of the cardiac Ca<sub>v</sub>1.2 channel.

### Syt-7 $\alpha$ increases the rate and extent of Ca<sup>2+</sup>-dependent facilitation

Previous studies have shown the key role of Ca<sub>v</sub>2.1 channels (Lee et al., 2000; Mochida et al., 2003a, 2008; Inchauspe et al., 2004) and Syt-7 (Jackman et al., 2016; Turecek and Regehr, 2018) in synaptic facilitation, but it is not known whether functional interactions between these two proteins modulate paired-pulse facilitation of P/Q-type Ca<sup>2+</sup> currents using pulse protocols similar to those in studies of short-term synaptic facilitation. In order to characterize the mechanism by which Syt-7 $\alpha$  increases Ca<sup>2+</sup>-dependent facilitation, the effects of Syt-7 $\alpha$  on the onset and decay of facilitation were measured with 10 mM Ca<sup>2+</sup> in the external solution to mimic the high local Ca<sup>2+</sup> concentration near the intracellular mouth of Ca<sub>v</sub>2.1 channels in nerve terminals during synaptic transmission. In a paired-pulse protocol, the rate of onset of facilitation was determined by plotting facilitation of I<sub>Ca</sub> as a function of prepulse duration ( $\Delta t$ ; Fig. 2, inset). In cells expressing only Ca<sub>v</sub>2.1 channels, the facilitation ratio increased with prepulse duration according to a single-exponential time course (Fig. 2A, black). Facilitation ratio reached a plateau




**Figure 6.** Effect of Syt-7 $\alpha$  on prepulse facilitation of Ca<sub>v</sub>2.1 channel. Facilitation of voltage-dependent activation of Ca<sup>2+</sup> currents. Inset, Pulse protocol to study the voltage dependence of activation before (open circle or squares; P1) and after (closed circle or squares; P2) a depolarizing prepulse from -80 to +10 mV. Tail currents were measured by holding potential at -40 mV for 5 ms after test pulses (P1, P2) to variable voltages (-40 to +80 mV). Peak tail currents were normalized to the largest tail current measured during the nonfacilitated prepulses (P1) and plotted against the test pulse voltage. **A**, In control tsA cells, the protocol shows an increase in facilitation P2 normalized to P1. **B**, Syt-7 $\alpha$  potentiated facilitation amplitude of Ca<sub>v</sub>2.1 and induced a right shift in prepulse facilitation curve. **C**, Overlaying the two graphs in **A**, **B** shows the increase in amplitude of facilitation and the right shift in voltage dependency of activation. **D**, Difference in voltage shift in P1 and P2 between cells co-expressing Ca<sub>v</sub>2.1 and Syt-7 $\alpha$  and control cells. Data are represented as mean  $\pm$  SEM.

at a prepulse duration of 20 ms and declined during prepulses of 50 ms or longer (P2/P1 ratio =  $1.29 \pm 0.06$ ,  $n = 20$ ). In tsA-201 cells co-expressing Ca<sub>v</sub>2.1 channels with Syt-7 $\alpha$ , facilitation increased more rapidly, reached a higher plateau at prepulse durations of 20–30 ms, and declined slowly during prepulses with  $\Delta t > 30$  ms (Fig. 2A,B, red; P2/P1 ratio =  $3.03 \pm 1.13$ ,  $n = 9$ ). The increase in facilitation ratio was significant at all prepulse durations compared with control cells ( $p < 0.0001$ ; Fig. 2A,B). The rate of increase in facilitation was significantly steeper in cells co-expressing Ca<sub>v</sub>2.1 with Syt-7 $\alpha$  (slope =  $0.03 \pm 0.02$  ms<sup>-1</sup>;  $n = 9$ ,  $p = 0.01$ ) compared with control cells (slope =  $0.005 \pm 0.001$  ms<sup>-1</sup>;  $n = 20$ ; Fig. 2A,B, red), and the half-time was shorter ( $\tau = 6.37 \pm 1.22$  ms,  $n = 7$ ,  $p = 0.03$ ) compared with control cells ( $\tau = 10.75 \pm 1.41$  ms,  $n = 17$ ;

Fig. 2A,B, red). Taken together, these data show that Syt-7 $\alpha$  increases the rate of the onset of facilitation of Ca<sub>v</sub>2.1 channels and increases the facilitation ratio at all prepulse durations tested.

Intracellular Ca<sup>2+</sup> concentrations near presynaptic Ca<sup>2+</sup> channels rise to nearly 100  $\mu$ M during rapid stimulation (Berridge et al., 2000). To mimic that condition, we have used 10 mM extracellular Ca<sup>2+</sup> in our standard experimental protocol to generate high Ca<sup>2+</sup> influx. However, at the physiological level of extracellular Ca<sup>2+</sup>, with 2 mM CaCl<sub>2</sub> the external recording solution, applying paired-pulse protocols revealed a significant acceleration of the onset and increase of the extent of Ca<sup>2+</sup>-dependent facilitation in cells co-expressing Ca<sub>v</sub>2.1 combined with Syt-7 $\alpha$  compared with control





**Figure 7.** Effect of Syt-7 $\alpha$  prepulse facilitation of Ca $_v$ 2.1 channel at physiological levels. Inset, Voltage protocol. Currents recorded with 2 mM extracellular Ca $^{2+}$  and 0.5 mM EGTA in the intracellular recording solution were elicited by test pulses to +10 mV before (P1) and 5 ms after (P2) 10-mV conditioning prepulses of the indicated durations. **A–C**, Effect of Syt-7 $\alpha$  on facilitation as a function of prepulse voltage. Facilitation was obtained by normalizing the peak current from P2 to that from P1. Single-exponential fits of the data are shown. Data are represented as mean  $\pm$  SEM.

(Fig. 3), as we observed with 10 mM external Ca $^{2+}$  concentration.


### Syt-7 $\alpha$ induces a rapidly decaying form of Ca $^{2+}$ -dependent facilitation

Syt-7 $\alpha$  accelerates the onset of  $I_{Ca}$  facilitation and increases facilitation amplitude at all potentials. However, the increased facilitation caused by Syt-7 $\alpha$  decayed rapidly ( $\tau = 4.29 \pm 1.68$  ms,  $p = 0.02$ ,  $n = 7$ ), compared with facilitation of Ca $_v$ 2.1 observed for tsA-201 cells in the absence of Syt-7 $\alpha$  ( $\tau = 12.87 \pm 3.77$  ms,  $n = 5$ ; Fig. 4, inset, left). The facilitation ratio P2/P1 at the first interpulse duration point was significantly greater for Ca $_v$ 2.1-Syt-7 $\alpha$  cells (facilitation ratio =  $2.73 \pm 0.71$ ,  $p = 0.004$ ,  $n = 7$ ) than for control cells (facilitation ratio =  $1.36 \pm 0.08$ ,  $n = 5$ ; Fig. 4, inset, right). Although the facilitation ratio in the presence of Syt-7 $\alpha$  decays rapidly, the integral of calcium current during the first 10 ms following stimulation is substantially increased (Fig. 4), illustrating the potential physiological significance of this increase in Ca $_v$ 2.1 channel activity.

### Syt-7 $\alpha$ increases voltage-dependent facilitation in paired-pulse protocols

In order to study the effect of Syt-7 on the voltage dependence of Ca $_v$ 2.1 activation and its consequences on facilitation, we measured facilitation using a paired-pulse protocol with variable stimulus potentials. In this protocol, facilitation induced by a 50-ms-long prepulse to a variable voltage (–40 to +60 mV) was measured by comparing  $I_{Ca}$  elicited by a test pulse before (P1) and after (P2) the conditioning prepulse (Fig. 5, inset). In cells expressing Ca $_v$ 2.1 alone, paired-pulse facilitation increased to a maximum at a prepulse voltage of +20 mV and remained at a plateau until +60 mV (facilitation ratio at 20 mV =  $1.09 \pm 0.04$ ,  $n = 8$ ; Fig. 5). Co-expression of Ca $_v$ 2.1 with Syt-7 $\alpha$  increased the maximum paired-pulse ratio to  $1.2 \pm 0.05$  ( $n = 8$  at +40 mV,  $p < 0.01$ ), approximately doubling the increase in Ca $^{2+}$  current induced by paired-pulse facilitation in the absence of Syt-7.

We also expressed Ca $_v$ 2.1 channels without or with Syt-7 and measured the voltage dependence of activation



**Figure 8.** Syt-7 isoforms  $\alpha$ ,  $\beta$ , and  $\gamma$  co-immunoprecipitate with  $\text{Ca}_v2.1$  channels. **A**, Left, In transfected tsA-201 cells, co-immunoprecipitation experiments show the binding of Syt-7 $\beta$  and  $\alpha_{1A}$  subunit of  $\text{Ca}_v2.1$ .  $\text{Ca}_v2.1$  channels were immunoprecipitated with an anti- $\text{Ca}_v2.1$  antibody. The immunoprecipitates were resolved by SDS/PAGE and immunoblotted with antibody against Syt-7. Right, The reverse experiment where Syt-7 $\beta$  were immunoprecipitated with an anti-Syt antibody and blotted with anti- $\text{Ca}_v2.1$  antibody. **B**, Left, In tsA-201 cells co-transfected with the  $\alpha_{1A}$  subunit of  $\text{Ca}_v2.1$  channels together with Syt-7 $\gamma$ ,  $\text{Ca}_v2.1$  channels were immunoprecipitated with an anti- $\text{Ca}_v2.1$  antibody and blotted with anti-Syt antibody. Right, Reverse experiment where Syt-7 $\gamma$  were immunoprecipitated with an anti-Syt antibody and blotted with anti- $\text{Ca}_v2.1$  antibody. **C**, Left, In tsA-201 cells co-transfected with the  $\alpha_{1A}$  subunit of  $\text{Ca}_v2.1$  channels along with Syt-7 $\alpha$ ,  $\beta$ , and  $\gamma$ ,  $\text{Ca}_v2.1$  channels were immunoprecipitated with an anti- $\text{Ca}_v2.1$  antibody and blotted with anti-Syt antibody. Right, Reverse experiment where the three isoforms of Syt-7 were immunoprecipitated with an anti-Syt antibody and blotted with anti- $\text{Ca}_v2.1$  antibody. Because different  $\text{Ca}^{2+}$  concentrations were used in co-immunoprecipitation experiments, segments from different immunoblots were spliced together to show specific comparisons. Those protein bands are delineated for clarification. The immunoblots presented here are representative of at least three experiments for each co-immunoprecipitation or immunoblot. Whole-cell voltage clamp experiments show that  $\text{Ca}_v2.1$  expressed alone gives similar peak  $\text{Ba}^{2+}$  and  $\text{Ca}^{2+}$  currents as  $\text{Ca}_v2.1 + \text{Syt7-}\alpha\beta\gamma$  (Extended Data Fig. 8-1).


of the resulting  $\text{Ca}^{2+}$  currents (Fig. 6). In this paired-pulse protocol described in Figure 6, inset, voltages were varied from  $-40$  to  $+80$  mV in both pulses P1 and P2 following a constant prepulse voltage of  $-80$  to  $+10$  mV before P2. Previous studies (Lee et al., 1999, 2000) showed that this protocol induced facilitation of  $\text{Ca}_v2.1$  channels. As shown in Figure 6A–C, Syt-7 $\alpha$  significantly increased  $I_{\text{Ca}}$  across the positive voltage range and increased maximum facilitation at potentials of  $+40$  mV and higher in the presence of  $10$  mM  $\text{Ca}^{2+}$ . Syt-7 $\alpha$  induced a significant  $\sim 5$ - to  $15$ -mV positive shift in the voltage dependence of  $\text{Ca}_v2.1$  activation, as observed by comparing the half-activation voltage ( $V_{50}$ ) at P1 ( $V_{50} = 14.28 \pm 3.28$ ,  $n = 7$ ,  $p = 0.002$ ) versus control cells ( $V_{50} = 3.96 \pm 1.72$ ,  $n = 19$ ). During pulse P2, Syt-7 $\alpha$  induced a significant  $\sim 3.6$ - to  $11$ -mV positive shift in the voltage dependence of  $\text{Ca}_v2.1$  activation ( $V_{50} = 7.96 \pm 2.3$ ,  $n = 7$ ,  $p = 0.02$ ) versus control cells ( $V_{50} = 0.7 \pm 1.4$ ,  $n = 19$ ; Fig. 6D). This Syt-7 $\alpha$  effect was also observed at physiological  $\text{Ca}^{2+}$  levels, where both facilitation amplitude and the positive shift in voltage dependence of  $\text{Ca}_v2.1$  activation were evident (Fig. 7). Together, these results show that Syt-7 $\alpha$  induces strong facilitation of  $\text{Ca}^{2+}$  currents at membrane potentials in the range of the peak of action potentials ( $\sim 0$  to  $+40$  mV).

#### Differential modulation of $\text{Ca}_v2.1$ by isoforms of Syt-7

Among the three  $\text{Ca}_v2$  subfamily members, only the  $\text{Ca}_v2.1$  channel supports short-term synaptic facilitation (Inchauspe et al., 2004); however, Syt-7 isoforms may

have subtype-specific modulatory effects on  $\text{Ca}_v2.1$ . Three splice variants of Syt-7 exist in mouse and human: the major form Syt-7 $\alpha$  and two minor forms, Syt-7 $\beta$  and Syt-7 $\gamma$  (Fukuda et al., 2002). Syt-7 $\beta$  and Syt-7 $\gamma$  contain additional 44 and 116 amino acids, respectively, in the connecting segment between their transmembrane domain and the cytoplasmic C2  $\text{Ca}^{2+}$ -binding domain (Fukuda et al., 2002). Both the  $\beta$  and  $\gamma$  isoforms of Syt-7 were bound to  $\text{Ca}_v2.1$  channels in extracts of transfected tsA-201 cells to a similar extent as Syt-7 $\alpha$ , as indicated by co-immunoprecipitation with anti- $\text{Ca}_v2.1$  antibodies and immunoblotting with isoform-specific anti-Syt-7 antibodies (Fig. 8A–C, left). In complementary experiments,  $\text{Ca}_v2.1$  was co-immunoprecipitated from transfected tsA-201 cells with antibodies against Syt-7 $\beta$  and Syt-7 $\gamma$  (Fig. 8A–C, right). The consistent results in these two complementary immunoprecipitation protocols indicate that these protein interactions are specifically detected independent of the antibodies used for immunoblotting. Co-expression of Syt-7 $\alpha$ , Syt-7 $\beta$ , and Syt-7 $\gamma$  together with  $\text{Ca}_v2.1$  channels did not have significant effects on the peak amplitude of either  $\text{Ba}^{2+}$  or  $\text{Ca}^{2+}$  currents in comparison to expression of Syt-7 $\alpha$  alone (Extended Data Fig. 8-1). Together, these experiments indicate that all three Syt-7 isoforms bind to  $\text{Ca}_v2.1$  channels in transfected cells without significantly altering their level of functional expression.

We investigated the effects of co-expression with Syt-7 $\beta$  and Syt-7 $\gamma$  on facilitation of  $\text{Ca}_v2.1$  channels with  $10$  mM  $\text{CaCl}_2$  in the extracellular solution. Syt-7 $\beta$  increased the facilitation ratio of  $\text{Ca}_v2.1$  channels ( $p < 0.01$ ; Fig. 9A)




**Figure 9.** Syt-7β and Syt-7γ differentially modulate facilitation of Ca<sub>v</sub>2.1 channels. Inset, Pulse protocol. Currents recorded with 10 mM extracellular Ca<sup>2+</sup> and 0.5 mM EGTA in the intracellular recording solution were elicited by test pulses to +10 mV before (P1) and 5 ms after (P2) 10-mV conditioning prepulses of the indicated durations. **A**, Left, Syt-7β increases the facilitation ratio with increasing prepulse duration. Right, Syt-7β accelerates the onset of facilitation as a function of prepulse duration. **B**, Left, Syt-7γ increases the facilitation ratio with increasing prepulse duration. Right, Syt-7γ does not accelerate the onset of facilitation as a function of prepulse duration. Facilitation was obtained by normalizing the peak current from P2 to that from P1. Single-exponential fits of the data are shown. Data are represented as mean ± SEM additional experiments with different pulse protocols provide additional information on the effects of Syt-7β and Syt-7γ on facilitation of Ca<sub>v</sub>2.1 channels (Extended Data Fig. 9-1).

and accelerated the rate of facilitation, as demonstrated by the significant increase in the slope in cells co-transfected with Ca<sub>v</sub>2.1 plus Syt-7β (slope = 0.02 ± 0.004 ms<sup>-1</sup>; n = 15, p = 0.006) compared with control cells (slope = 0.009 ± 0.001 ms<sup>-1</sup>; n = 20; Fig. 9A, inset). However, these effects were substantially smaller than with co-expression of Syt-7α (Fig. 2). Co-expression of Syt-7γ also increased the peak level of facilitation (p < 0.01) to a lesser degree than Syt-7α (Fig. 9B), and it showed only a trend toward a significant increase in facilitation rate (slope = 0.03 ± 0.01 ms<sup>-1</sup>; n = 9, p = 0.16, ns) compared with control cells (slope = 0.01 ± 0.003 ms<sup>-1</sup>; n = 20; Fig. 9B, inset). To determine whether Syt-7β and Syt-7γ can compete effectively with Syt-7α, we co-expressed all three Syt-7 isoforms and measured Ca<sup>2+</sup>-dependent facilitation (Extended Data Fig. 9-1). We found

that co-expression Syt-7β and Syt-7γ together with Syt-7α reduced the strong increase in the rate and extent of facilitation observed with Syt-7α alone (Extended Data Fig. 9-1, red). These results are consistent with the conclusion that Syt-7β and Syt-7γ effectively compete for occupancy of the synprint site and alter modulation of Ca<sub>v</sub>2.1 by Syt-7α. Evidently, replacement of Syt-7α with either Syt-7β or Syt-7γ at the synprint site would reduce facilitation of Ca<sub>v</sub>2.1 channels.

In experiments testing the effect of a depolarizing prepulse on the voltage dependence of activation, neither Syt-7β (Fig. 10A-C) nor Syt-7γ (Fig. 10E-G) increased the maximum prepulse facilitation of Ca<sub>v</sub>2.1 at positive membrane potentials in contrast to Syt-7α. Similarly, neither Syt-7β (Fig. 10A-D) nor Syt-7γ (Fig. 10E-H) caused a significant shift in the voltage dependence of activation of Ca<sub>v</sub>2.1




**Figure 10.** Syt-7 $\beta$  and Syt-7 $\gamma$  differentially modulate prepulse facilitation of Ca<sub>v</sub>2.1 channels. Inset, Pulse protocol. Currents recorded with 10 mM extracellular Ca<sup>2+</sup> and 0.5 mM EGTA in the intracellular recording solution were elicited by test pulses to +10 mV before (P1) and 5 ms after (P2) 10-mV conditioning prepulses of the indicated durations. Facilitation was obtained by normalizing the peak current from P2 to that from P1. Single-exponential fits of the data are shown. **A**, Ca<sub>v</sub>2.1 alone. **B**, Ca<sub>v</sub>2.1 with Syt-7 $\beta$ . **C**, Overlay of panels **A**, **B**. **D**, V<sub>50</sub> values for results in panel **C**. **E**, Ca<sub>v</sub>2.1 alone. **F**, Ca<sub>v</sub>2.1 with Syt-7 $\gamma$ . **G**, overlay of panels **D**, **E**. **H**, V<sub>50</sub> values for panel **G**. Data are represented as mean  $\pm$  SEM. Additional experiments with different pulse protocols provide additional information on the effects of Syt-7 $\beta$  and Syt-7 $\gamma$  on the voltage dependence of facilitation of Ca<sub>v</sub>2.1 channels (Extended Data Fig. 10-1).

channels, unlike Syt-7 $\alpha$  (Fig. 10C,D). Co-expressing the three Syt-7 isoforms together caused a negative shift in the voltage dependence of activation following a depolarizing prepulse, in contrast to the positive shift in the voltage dependence of activation following a prepulse caused by co-expression of Syt-7 $\alpha$  alone (Extended Data Fig. 10-1, red). All of these voltage-dependent activation curves are monophasic (Figs. 7, 10; Extended Data Fig. 10-1), consistent with stoichiometric binding of each Syt-7 isoform to Ca<sub>v</sub>2.1 resulting in complete shifts of the activation curves. Together, these results suggest a dominant effect of Syt-7 $\beta$  and Syt-7 $\gamma$  on the voltage dependence of activation in paired-pulse protocols in the presence of all three Syt-7 isoforms.

In addition to their differential effects on Ca<sup>2+</sup>-dependent facilitation, co-expression of Syt-7 $\beta$  or Syt-7 $\gamma$  also had different effects on Ca<sup>2+</sup>-dependent inactivation of Ca<sub>v</sub>2.1 channels compared with Syt-7 $\alpha$  (Fig. 11). In the presence of 10 mM Ba<sup>2+</sup> as the permeant extracellular cation, Ca<sub>v</sub>2.1 channels activated rapidly and did not inactivate significantly in 200-ms depolarizing pulses when co-expressed with any of the Syt-7 isoforms (Fig. 11A). In contrast, in the presence of 10 mM Ca<sup>2+</sup> as permeant ion, Ca<sub>v</sub>2.1 channels inactivated with a time

constant of  $\sim$ 600 ms through their Ca<sup>2+</sup>/CaM-dependent inactivation mechanism (Fig. 11B, black). Strikingly, co-expression of Syt-7 $\alpha$  substantially slowed Ca<sup>2+</sup>-dependent inactivation (Fig. 11B, red), whereas co-expression of Syt-7 $\beta$  had a smaller effect (Fig. 11B, blue) and co-expression of Syt-7 $\gamma$  had no effect on Ca<sup>2+</sup>-dependent inactivation (Fig. 11B, green). These results indicate that replacement of Syt-7 $\alpha$  with either Syt-7 $\beta$  or Syt-7 $\gamma$  at the synprint site would decrease Ca<sup>2+</sup> entry in single depolarizations by preventing the inhibition of Ca<sup>2+</sup>-dependent inactivation of Ca<sub>v</sub>2.1 channels induced by Syt-7 $\alpha$  (Fig. 11), and at the same time would reduce prolonged Ca<sup>2+</sup> entry by decreasing the enhanced facilitation of Ca<sub>v</sub>2.1 channels caused by Syt-7 $\alpha$  during repetitive depolarizations (Fig. 9). This parallel modulation of Ca<sup>2+</sup> entry by single depolarizations plus trains of depolarizations would have a potent impact on synaptic transmission. Altogether, these results indicate that the two minor Syt-7 isoforms bind to Ca<sub>v</sub>2.1 channels in cellular context without altering functional expression of Ca<sub>v</sub>2.1. However, co-expression of Syt-7 $\beta$  and Syt-7 $\gamma$  can partially reverse the functional effects of Syt-7 $\alpha$  on facilitation (Extended Data Fig. 9-1), the voltage dependence of activation following a depolarizing prepulse (Extended





**Figure 11.** Syt-7 isoforms differentially modulate  $\text{Ca}^{2+}$ -dependent inactivation of  $\text{Ca}_v2.1$  channels.  $\text{Ca}_v2.1$  currents were elicited by depolarizing from a holding potential of  $-80$  mV to a test potential of  $+10$  mV. **A**, Time courses (200 ms) of  $I_{\text{Ba}^{2+}}$  with  $10$  mM  $\text{Ba}^{2+}$  as a permeant cation. **B**, Time courses (1000 ms) of  $I_{\text{Ca}^{2+}}$  with  $10$  mM  $\text{Ca}^{2+}$  as permeant ion. Syt-7 $\alpha$  and Syt-7 $\beta$  significantly slowed inactivation of the  $\text{Ca}_v2.1$  channel in the presence of  $10$  mM  $\text{Ca}^{2+}$ , whereas Syt-7 $\gamma$  had no effect. Data are represented as mean  $\pm$  SEM.

Data Fig. 10-1), and the rate of  $\text{Ca}^{2+}$ -dependent inactivation (Fig. 11). The differential actions of the three isoforms of Syt-7 provide a rich panoply of modulatory effects on  $\text{Ca}_v2.1$  channel activation, facilitation, and inactivation that would have a strong influence on synaptic transmission.

## Discussion

### Syt-7 modulates $\text{Ca}_v2.1$ channels through binding to the synprint site

In presynaptic nerve terminals,  $\text{Ca}_v2.1$  and  $\text{Ca}_v2.2$  channels associate with SNARE proteins and a large number of other presynaptic proteins (Khanna et al., 2007; Müller et al., 2010; Nanou and Catterall, 2018). SNARE proteins interact with the synprint site located in the intracellular loop between domains II and III (Sheng et al., 1994, 1996; Rettig et al., 1996), which is thought to play an important role in the incorporation of  $\text{Ca}_v2.1$  channels into the synaptic vesicle fusion machinery and regulation of their function (Mochida et al., 2003b; Szabo et

al., 2006).  $\text{Ca}^{2+}$  influx through  $\text{Ca}_v2.1$  channels is a crucial step in triggering  $\text{Ca}^{2+}$ -dependent exocytosis of neurotransmitter vesicles (Olivera et al., 1994; Dunlap et al., 1995). Previous studies showed that the fast  $\text{Ca}^{2+}$  sensor Syt-1 binds to the synprint site of both  $\text{Ca}_v2.1$  and  $\text{Ca}_v2.2$  channels (Sakurai et al., 1996; Sheng et al., 1996; Charvin et al., 1997; Kim and Catterall, 1997). These protein interactions are likely to modulate the rapid, synchronous component of neurotransmitter release mediated by Syt-1 (Bacaj et al., 2013).

In contrast to these extensive studies of SNARE proteins and the fast  $\text{Ca}^{2+}$  sensor Syt-1, the slow, high-affinity  $\text{Ca}^{2+}$  sensor Syt-7 is unique in mediating synaptic facilitation (Jackman et al., 2016; Nanou et al., 2016a; Turecek and Regehr, 2018) and asynchronous release (Bacaj et al., 2013; Turecek and Regehr, 2018), but its interactions with  $\text{Ca}_v2.1$  channels had not previously been investigated. Our results presented here show that Syt-7 binds to the synprint site of  $\text{Ca}_v2.1$  channels *in vivo* in mouse brain membranes, *in vitro* in transfected cells, and in solution in protein-interaction experiments. Unexpectedly, in contrast to Syt-1, our results provide evidence that Syt-7 modulates  $\text{Ca}^{2+}$ -dependent facilitation and inactivation (CDI) of  $\text{Ca}_v2.1$  channels, which are implicated in short-term forms of synaptic plasticity, including synaptic facilitation and the rapid phase of synaptic depression (Lee et al., 2000, 2002; Mochida et al., 2008; Nanou et al., 2016a, b, 2018). Direct interaction of Syt-7 and  $\text{Ca}_v2.1$  as shown here may contribute to short-term synaptic facilitation, in which both of these interacting protein partners are thought to play essential roles.

### Syt-7 isoforms differentially enhance facilitation of $\text{Ca}_v2.1$ channels

In cells expressing  $\text{Ca}_v2.1$  channels, we consistently observed  $\text{Ca}^{2+}$ -dependent facilitation of the  $\text{Ca}^{2+}$  current, as reported previously (Lee et al., 1999, 2000, 2003). In the presence of Syt-7 $\alpha$ , both the rate and extent of facilitation of  $\text{Ca}_v2.1$  channels were increased, and the rate of decay of facilitation was also accelerated. These results suggest that expression of Syt-7 $\alpha$  in presynaptic terminals *in vivo* would enhance  $\text{Ca}^{2+}$ -dependent facilitation and sharpen the time-dependent peak of facilitation of  $\text{Ca}_v2.1$  channels. Syt-7 $\beta$  and Syt-7 $\gamma$  also bound to the synprint site. However, compared with Syt-7 $\alpha$ , Syt-7 $\beta$ , and Syt-7 $\gamma$  had lesser effects on facilitation in response to voltage steps and did not shift the voltage dependence of prepulse facilitation. Differential expression of these Syt-7 isoforms could confer cell-specific regulation via interactions with  $\text{Ca}_v2.1$  channels and other regulatory targets.

### Syt-7 isoforms differentially modulate inactivation of $\text{Ca}_v2.1$ channels

In our depolarizing step protocols, none of the Syt-7 isoforms had any significant effect on the peak amplitude of  $\text{Ba}^{2+}$  current. However, our data show that Syt-7 $\alpha$  significantly increased  $\text{Ca}^{2+}$ -dependent inactivation of the  $\text{Ca}^{2+}$  channel, which would oppose facilitation. Syt-7 $\beta$  and Syt-7 $\gamma$  had lesser effects. The combination of increased



facilitation followed by increased inactivation induced by Syt-7 $\alpha$  would have the overall effect of sharpening the peak of the presynaptic calcium current to allow effective facilitation of repetitive rounds of neurotransmitter release. Syt-7 $\beta$  and Syt-7 $\gamma$  would bind to the synprint site of Ca $_v$ 2.1 channels but induce lesser functional effects.

### Comparison with regulation by CaM-like calcium sensor proteins

Our work characterizes an unexpected form of regulation of P/Q-type current conducted by Ca $_v$ 2.1 channels by the high affinity Ca $^{2+}$  sensor Syt-7 through a direct interaction with the synprint site. Interaction of Ca $_v$ 2.1 with Syt-7 may enhance facilitation of presynaptic Ca $^{2+}$  current and thereby play a role in triggering activation of the Ca $^{2+}$ -dependent exocytosis machinery, including the SNARE proteins. These effects would be dependent on the isoform of Syt-7 that is expressed in different cells and synapses. In previous experiments, CaM has been shown to regulate Ca $_v$ 2.1 channel activity, inducing increased facilitation and increased Ca $^{2+}$ -dependent inactivation, dependent on the local Ca $^{2+}$  concentration (Lee et al., 1999, 2000; DeMaria et al., 2001). Our results further show that Ca $^{2+}$ -dependent inactivation of Ca $_v$ 2.1 channels is modulated by Syt-7 in an isoform-dependent manner. In presynaptic nerve terminals, these changes in both the Ca $^{2+}$  entry in response to single action potentials plus trains of action potentials would substantially alter the encoding properties of synaptic transmission.

Other neuronal Ca $^{2+}$  sensor proteins related to CaM are expressed in the central nervous system, including Ca $^{2+}$  binding protein-1 (CaBP-1), visinin-like protein-2 (VILIP-2), and neuronal Ca $^{2+}$  sensor-1 (NCS-1). These Ca $^{2+}$  sensor proteins displace CaM from the C-terminal domain of Ca $_v$ 2.1 and modify short-term synaptic facilitation and rapid synaptic depression (Nanou and Catterall, 2018). It will be interesting to further investigate how these two distinct regulatory mechanisms mediated by Syt-7 and Ca $^{2+}$  sensor proteins converge on the Ca $_v$ 2.1 channel on the millisecond time frame of short-term synaptic plasticity.

In conclusion, our work characterizes a novel form of regulation of P/Q-type Ca $_v$ 2.1 channels by the high affinity Ca $^{2+}$  sensor Syt-7 through direct interaction with the synprint site. Ca $_v$ 2.1/Syt-7 interaction potentiates facilitation of Ca $^{2+}$  current and may play a role in triggering Ca $^{2+}$ -dependent exocytosis along with other SNARE proteins. Syt-7 also modulates Ca $^{2+}$ /CaM-dependent inactivation. Understanding the mechanism by which Syt-7 isoforms enhance facilitation and modulate inactivation of Ca $_v$ 2.1 channels in presynaptic terminals is a first step toward deciphering the complete picture of the role played by Syt-7 in the brain.

### References

- Bacaj T, Wu D, Yang X, Morishita W, Zhou P, Xu W, Malenka RC, Südhof TC (2013) Synaptotagmin-1 and synaptotagmin-7 trigger synchronous and asynchronous phases of neurotransmitter release. *Neuron* 80:947–959.
- Ben-Johny M, Yue DT (2014) Calmodulin regulation (calmodulation) of voltage-gated calcium channels. *J Gen Physiol* 143:679–692.
- Berridge MJ, Lipp P, Bootman MD (2000) Signal transduction. The calcium entry pas de deux. *Science* 287:1604–1605.
- Catterall WA, Few AP (2008) Calcium channel regulation and presynaptic plasticity. *Neuron* 59:882–901.
- Charvin N, Lévêque C, Walker D, Berton F, Raymond C, Kataoka M, Shoji-Kasai Y, Takahashi M, De Waard M, Seagar MJ (1997) Direct interaction of the calcium sensor protein synaptotagmin I with a cytoplasmic domain of the alpha1A subunit of the P/Q-type calcium channel. *EMBO J* 16:4591–4596.
- Christel C, Lee A (2012) Ca $^{2+}$ -dependent modulation of voltage-gated Ca $^{2+}$  channels. *Biochim Biophys Acta* 1820:1243–1252.
- DeMaria CD, Soong TW, Alseikhan BA, Alvania RS, Yue DT (2001) Calmodulin bifurcates the local Ca $^{2+}$  signal that modulates P/Q-type Ca $^{2+}$  channels. *Nature* 411:484–489.
- Dunlap K, Luebke JI, Turner TJ (1995) Exocytotic Ca $^{2+}$  channels in mammalian central neurons. *Trends Neurosci* 18:89–98.
- Erickson MG, Alseikhan BA, Peterson BZ, Yue DT (2001) Preassociation of calmodulin with voltage-gated Ca $^{2+}$  channels revealed by FRET in single living cells. *Neuron* 31:973–985.
- Fukuda M, Ogata Y, Saegusa C, Kanno E, Mikoshiba K (2002) Alternative splicing isoforms of synaptotagmin VII in the mouse, rat and human. *Biochem J* 365:173–180.
- Inchauspe CG, Martini FJ, Forsythe ID, Uchitel OD (2004) Functional compensation of P/Q by N-type channels blocks short-term plasticity at the calyx of held presynaptic terminal. *J Neurosci* 24:10379–10383.
- Jackman SL, Turecek J, Belinsky JE, Regehr WG (2016) The calcium sensor synaptotagmin 7 is required for synaptic facilitation. *Nature* 529:88–91.
- Khanna R, Zougman A, Stanley EF (2007) A proteomic screen for presynaptic terminal N-type calcium channel (Ca $_v$ 2.2) binding partners. *J Biochem Mol Biol* 40:302–314.
- Kim DK, Catterall WA (1997) Ca $^{2+}$ -dependent and -independent interactions of the isoforms of the alpha1A subunit of brain Ca $^{2+}$  channels with presynaptic SNARE proteins. *Proc Natl Acad Sci U S A* 94:14782–14786.
- Lee A, Scheuer T, Catterall WA (2000) Ca $^{2+}$ /calmodulin-dependent facilitation and inactivation of P/Q-type Ca $^{2+}$  channels. *J Neurosci* 20:6830–6838.
- Lee A, Wong ST, Gallagher D, Li B, Storm DR, Scheuer T, Catterall WA (1999) Ca $^{2+}$ /calmodulin binds to and modulates P/Q-type calcium channels. *Nature* 399:155–159.
- Lee A, Westenbroek RE, Haeseleer F, Palczewski K, Scheuer T, Catterall WA (2002) Differential modulation of Ca $_v$ 2.1 channels by calmodulin and Ca $^{2+}$ -binding protein 1. *Nat Neurosci* 5:210–217.
- Lee A, Zhou H, Scheuer T, Catterall WA (2003) Molecular determinants of Ca $^{2+}$ /calmodulin-dependent regulation of Ca $_v$ 2.1 channels. *Proc Natl Acad Sci U S A* 100:16059–16064.
- Mochida S, Westenbroek RE, Yokoyama CT, Itoh K, Catterall WA (2003a) Subtype-selective reconstitution of synaptic transmission in sympathetic ganglion neurons by expression of exogenous calcium channels. *Proc Natl Acad Sci U S A* 100:2813–2818.
- Mochida S, Westenbroek RE, Yokoyama CT, Zhong H, Myers SJ, Scheuer T, Itoh K, Catterall WA (2003b) Requirement for the synaptic protein interaction site for reconstitution of synaptic transmission by P/Q-type calcium channels. *Proc Natl Acad Sci U S A* 100:2819–2824.
- Mochida S, Few AP, Scheuer T, Catterall WA (2008) Regulation of presynaptic Ca $_v$ 2.1 channels by Ca $^{2+}$  sensor proteins mediates short-term synaptic plasticity. *Neuron* 57:210–216.
- Müller CS, Haupt A, Bildl W, Schindler J, Knaus H, Meissner M, Rammner B, Striessnig J, Flockner V, Fakler B, Schulte U (2010) Quantitative proteomics of the Ca $_v$ 2 channel nano-environments in the mammalian brain. *Proc Natl Acad Sci U S A* 107:14950–14957.
- Nanou E, Catterall WA (2018) Calcium channels, synaptic plasticity, and neuropsychiatric disease. *Neuron* 98:466–481.

- Nanou E, Sullivan JM, Scheuer T, Catterall WA (2016a) Calcium sensor regulation of the  $\text{Ca}_v2.1$   $\text{Ca}^{2+}$  channel contributes to short-term synaptic plasticity in hippocampal neurons. *Proc Natl Acad Sci U S A* 113:1062–1067.
- Nanou E, Yan J, Whitehead NP, Kim MJ, Froehner SC, Scheuer T, Catterall WA (2016b) Altered short-term synaptic plasticity and reduced muscle strength in mice with impaired regulation of presynaptic  $\text{Ca}_v2.1$   $\text{Ca}^{2+}$  channels. *Proc Natl Acad Sci U S A* 113:1068–1073.
- Nanou E, Lee A, Catterall WA (2018) Control of excitation/inhibition balance in a hippocampal circuit by calcium sensor protein regulation of presynaptic calcium channels. *J Neurosci* 38:4430–4440.
- Olivera BM, Miljanich GP, Ramachandran J, Adams ME (1994) Calcium channel diversity and neurotransmitter release: the omega-conotoxins and omega-agatoxins. *Annu Rev Biochem* 63:823–867.
- Rettig J, Sheng ZH, Kim DK, Hodson CD, Snutch TP, Catterall WA (1996) Isoform-specific interaction of the  $\alpha 1A$  subunits of brain  $\text{Ca}^{2+}$  channels with the presynaptic proteins syntaxin and SNAP-25. *Proc Natl Acad Sci U S A* 93:7363–7368.
- Sakurai T, Westenbroek RE, Rettig J, Hell J, Catterall WA (1996) Biochemical properties and subcellular distribution of the BI and  $\alpha 1A$  isoforms of alpha 1A subunits of brain calcium channels. *J Cell Biol* 134:511–528.
- Sheng ZH, Rettig J, Takahashi M, Catterall WA (1994) Identification of a syntaxin-binding site on N-type calcium channels. *Neuron* 13:1303–1313.
- Sheng ZH, Rettig J, Cook T, Catterall WA (1996) Calcium-dependent interaction of N-type calcium channels with the synaptic core-complex. *Nature* 379:451–454.
- Szabo Z, Obermair GJ, Cooper CB, Zamponi GW, Flucher BE (2006) Role of the synprint site in presynaptic targeting of the calcium channel  $\text{Ca}_v2.2$  in hippocampal neurons. *Eur J Neurosci* 24:709–718.
- Turecek J, Regehr WG (2018) Synaptotagmin 7 mediates both facilitation and asynchronous release at granule cell synapses. *J Neurosci* 38:3240–3251.
- Zamponi GW, Striessnig J, Koschak A, Dolphin AC (2015) The physiology, pathology, and pharmacology of voltage-gated calcium channels and their future therapeutic potential. *Pharmacol Rev* 67:821–870.
- Zühlke RD, Pitt GS, Deisseroth K, Tsien RW, Reuter H (1999) Calmodulin supports both inactivation and facilitation of L-type calcium channels. *Nature* 399:159–162.