



ORAL PRESENTATION

Open Access

Trends in hospitalization of preterm infants with intraventricular hemorrhage and hydrocephalus in the United States, 2000-2010

Eisha Anne Christian^{1*}, Diana Jin¹, Frank Attenello¹, Timothy Wen¹, Steven Cen¹, William J Mack¹, Mark D Krieger^{1,2}, J Gordon McComb^{1,2}

From Hydrocephalus 2015
Banff, Canada. 18-21 September 2015

Objective

Even with improved prenatal and neonatal care, intraventricular hemorrhage (IVH) occurs in approximately 25-30% of preterm infants, and a subset of these patients develop hydrocephalus. We aim to describe current trends in hospitalization of preterm infants with IVH and post-hemorrhagic hydrocephalus (PHH) using the Nationwide Inpatient Sample (NIS) and Kids Inpatient Database (KID).

Methods

The KID and NIS databases were combined to generate data for the years 2000 – 2010. All neonatal discharges with ICD9-CM codes for preterm birth with IVH alone or with IVH and hydrocephalus were included.

Results

There were 147,823 preterm neonates with IVH, and 9% of this group developed hydrocephalus during the same admission. Twenty-five percent and 28% of patients with Grades 3 and 4 IVH respectively developed hydrocephalus in comparison to 1% and 4% of patients with Grades 1 and 2 IVH. Thirty-eight percent of patients with PHH had permanent ventricular shunts inserted. Mortality rates were 4%, 10%, 18%, and 40% respectively for Grades 1-4 during initial hospitalization. Length of stay has been trending upward for both groups of IVH (49d in 2000, 56d in 2010) and PHH (59d in 2000, 70d in 2010). Average hospital cost per patient (inflation-adjusted) has also increased from \$201,578 to \$353,554 (IVH) and \$260,077 to \$495,697 (PHH) over 11 years.

* Correspondence: echristi@usc.edu

¹University of Southern California, Los Angeles, CA, USA

Full list of author information is available at the end of the article

Conclusion

The number of admissions of neonates with IVH has increased despite a decrease in the number of preterm births. Rates of hydrocephalus and mortality correlated closely with IVH grade. Incidence of hydrocephalus in preterm infants with IVH remained stable between 8-10%. Over an 11-year period, there was a progressive increase in hospital cost and length of stay for preterm neonates with IVH and PHH with a concurrent increase in the proportion of patients with congenital cardiac anomalies.

Authors' details

¹University of Southern California, Los Angeles, CA, USA. ²Children's Hospital Los Angeles, CA, USA.

Published: 18 September 2015

References

1. Olaya JE, Christian E, Ferman D, Luc Q, Sanger T, Krieger M, Liker M: Deep Brain Stimulation in Children and Young Adults with Secondary Dystonia: The Children's Hospital Los Angeles Experience. *Neurosurgical Focus* 2013, **35**(5):E7.
2. Karas PJ, Mikell CB, Christian E, Liker M, Sheth S: Deep Brain Stimulation: a mechanistic and clinical update. *Neurosurgical Focus* 2013, **35**(5):E1.
3. Christian E, Yu C, Apuzzo ML: Focused Ultrasound: Relevant History and Prospects for the Addition of Mechanical Energy to the Neurosurgical Armamentarium. *World Neurosurgery* 2014, **82**(3-4):354.
4. Christian E, Harris B, Wrobel B, Zada G: Endoscopic endonasal transsphenoidal surgery: implementation of an operative and perioperative checklist. *Neurosurgical Focus* 2014, **37**(4):E1.
5. Christian E, Huang C, Yen C, Acosta FA, Chen TC, Liu JC, Spoonamore M, Wang JC, Hsieh PC: A modified Gaines approach for lumbosacral traumatic spondylolisthesis: A historical review and case illustration. *J Spine* 2014, **3**(4).
6. Christian E, Imahiyerobo T, Nallappa S, Urata M, McComb JG, Krieger M: Intracranial hypertension after surgical correction for craniosynostosis: a systematic review. *Neurosurgical Focus* May 2015.

doi:10.1186/2045-8118-12-S1-O1

Cite this article as: Christian et al.: Trends in hospitalization of preterm infants with intraventricular hemorrhage and hydrocephalus in the United States, 2000-2010. *Fluids and Barriers of the CNS* 2015 **12**(Suppl 1):O1.