

CORRESPONDENCE

Open Access

# Classical dynamin DNM1 and DNM3 genes attain maximum expression in the normal human central nervous system

Antoni Romeu\* and Lluís Arola

## Abstract

Dynamin is a super-family of large GTPase proteins that polymerise during their biological activity. Dynamin polymers form around lipid tubes and contribute to the membrane fission and scission of nascent vesicles from parent membranes. Here we used the NCBI Gene Expression Omnibus (GEO) database and the BioGPS gene expression portal to study differential dynamin gene expression in normal human organs or tissues. From the GDS1096 and GDS596 dataset, we downloaded the relative expression levels of dynamin-related genes (presented as percentages), with respect to all of the other genes on the array (platform Affymetrix GPL96), which includes the best characterised human genes. The expression profiles of dynamin in the central nervous system (CNS) are clearly distinct from the expression profiles in the other organs or tissues studied. We found that the classical dynamin DNM1 and DNM3 genes reach their maximum expression levels (100% of maximal expression) in all normal human CNS tissues studied. This is in contrast to the expression profile in the other normal human organs or tissues studied, in which both dynamin DNM1 and DNM3 genes showed approximately 50% maximal expression. This data mining analysis supports the concept that there is a relationship between the synapse and the molecular function of dynamin, suggesting a new field of work in the study of neurodegenerative diseases.

**Keywords:** Dynamin, Central nervous system, Gene expression, Data mining

## Discussion


We have focused our attention on the expression profiling of dynamin genes in several normal human organs or tissues, using the NCBI Gene Expression Omnibus (GEO) database. We have chosen to study dynamin genes because dynamin is a super-family of GTP-binding mechanochemical proteins that are involved in fundamental processes, such as the scission of nascent vesicles from parent membranes and mitochondrial fusion and/or fission [1]. In addition, we believe that the dynamin super-family is a biological system suitable for studying differential gene expression in organs or tissues.

Specifically, we considered the group of genes coding the following protein families within the dynamin super-family [2]: (i) the classical dynamins and dynamin-like proteins that tubulate and sever membranes and are involved in clathrin-mediated endocytosis and other vesicular trafficking

processes (DNM1, DNM1L, DNM2, DNM3); (ii) the guanylate-binding proteins, which are induced by type II interferons and anti-inflammatory cytokines (GBP1 and GBP2); (iii) the proteins on the outside of the outer mitochondrial membrane that participate in mitochondrial fusion and contribute to the maintenance and function of the mitochondrial network (MFN1 and MFN2); (iv) proteins in the mitochondrial inter membrane space that are involved in mitochondrial fusion, and in which mutations in the gene are associated with optic atrophy type 1 (OPA1); and (v) Mx proteins, which are involved in resistance against viral infections (MX1 and MX2). Taken together, these proteins constitute a large set of GTPase proteins that polymerise within each group or family according to the dynamin molecular function [3]. These dynamin polymers (such as rings and helices) are formed around lipid tubes and contribute to membrane fission [4].

To address this issue, we relied on the independent studies by Su A.I. et al. [5] (GDS596) and Ge X. et al. [6] (GDS1096). We downloaded the relative expression levels


\* Correspondence: [antoni.romeu@urv.cat](mailto:antoni.romeu@urv.cat)  
Nutrigenomics Research Group, Department of Biochemistry and Biotechnology, Rovira i Virgili University, Tarragona, Spain



**Figure 1 Data retrieved from the dataset records GDS596 and GDS1096 of NCBI-GEO database.** The figure depicts the dynamin expression profiles of several normal human organs or tissues. The dynamin genes are listed on the x-axis, while the relative gene expression is shown on the y-axis. The points represent the gene expression level (presented as a percentage). Each point represents the mean of the gene expression level in normal organ or tissue samples that were common to both of the independent experimental datasets (GDS596 and GDS1096) of the NCBI-GEO database. According to the NCBI-GEO data analysis, all values within an array are rank ordered and then placed into percentile 'bins'. In other words, all the values of one hybridisation are sorted and then split into 100 groups. Thus, the points give an indication of where the expression of a given gene falls with respect to all genes on that array. In these records, the platform GPL96 (Affymetrix Human Genome U133A Array) were used, that includes over 1,000,000 unique oligonucleotide features covering more than 39,000 transcript variants, which in turn represent more than 33,000 of the best characterised human genes. See Table 1 for a detailed specification of the data.

of dynamin-related genes (presented as percentages) with respect to all other genes on the array from these NCBI-GEO database records. It is important to note that both NCBI-GEO data sets were created based on the same array platform (Affymetrix GPL96), which include all the best characterised human genes. Figure 1 shows the profiles of dynamin gene expression levels (expressed as

percentage and presented as beam shaped lines) in samples from normal human organs or tissues that are common to the NCBI-GEO records of both studies. From a strict visual analysis of these beam lines, the data can be separated into two consistent populations or sub-beam lines of expression profiles, one of which groups together the profiles corresponding to components of the central



**Figure 2 Dynamin expression profiling in representative components of normal human CNS tissues and organs.** Data retrieved from Figure 1. This figure depicts the expression profiling of the representative components of the CNS. The genes studied are denoted along the x-axis and the gene expression (presented as a percentage) on the y-axis. The Figure 1 legend explains how to interpret the NCBI-GEO profile charts.

**Table 1 Relative gene expression profiles from normal human organs and tissues (presented as percentatges)**

Organ-tissue	DNM1 (215116_s_at)	DNM1L (203105_s_at)	DNM2 (202253_s_at)	DNM3 (209839_at)	GBP1 (202269_x_at)	GBP2 (202748_at)	MFN1 (211801_x_at)	MFN2 (201155_s_at)	MX1 (202086_at)	MX2 (204994_at)	OPA1 (212213_x_at)
Amygdala	<b>99</b>	67	55	<b>96</b>	8	35	64	89	74	68	92
Brain foetal	<b>95</b>	55	50	<b>91</b>	15	31	65	87	68	67	85
Brain whole	<b>99</b>	52	64	<b>94</b>	1	35	54	92	78	70	88
Caudate nucleus	<b>98</b>	67	67	<b>97</b>	10	42	62	92	77	71	94
Cerebellum	<b>99</b>	47	64	<b>96</b>	2	35	60	93	71	66	79
Hippocampus	<b>98</b>	69	52	<b>97</b>	20	48	59	92	82	71	92
Hypothalamus	<b>98</b>	69	60	<b>97</b>	22	49	64	92	90	75	91
Spinal cord	<b>93</b>	56	65	<b>94</b>	46	51	59	91	84	63	91
Thalamus	<b>99</b>	71	55	<b>98</b>	17	38	64	93	82	70	91
Adrenal gland	<b>44</b>	42	72	<b>56</b>	43	77	53	88	78	73	74
Bone marrow	<b>47</b>	22	73	<b>47</b>	8	63	58	93	81	85	77
Heart	<b>52</b>	54	68	<b>46</b>	26	64	64	98	81	81	87
Kidney	<b>59</b>	51	78	<b>40</b>	39	67	55	88	75	71	74
Liver	<b>52</b>	27	68	<b>48</b>	56	55	50	91	83	81	76
Lung	<b>51</b>	35	76	<b>44</b>	61	75	62	88	86	84	72
Lung foetal	<b>67</b>	50	73	<b>40</b>	50	59	63	83	70	71	72
Ovary	<b>35</b>	35	49	<b>53</b>	43	83	61	88	71	73	84
Pancreas	<b>63</b>	46	59	<b>34</b>	25	56	72	87	74	81	74
Pituitary	<b>91</b>	65	55	<b>83</b>	29	49	58	88	81	70	87
Placenta	<b>38</b>	49	69	<b>35</b>	58	66	61	83	73	73	70
Prostate	<b>52</b>	50	69	<b>38</b>	52	68	54	86	75	71	79
Salivary gland	<b>57</b>	43	57	<b>46</b>	38	71	50	89	80	82	73
Skin	<b>75</b>	33	49	<b>57</b>	35	77	72	91	59	76	84
Testis	<b>52</b>	62	71	<b>47</b>	14	62	61	89	64	74	70
Thymus	<b>47</b>	41	72	<b>45</b>	61	74	62	86	88	88	70
Thyroid	<b>49</b>	42	75	<b>36</b>	80	70	63	89	85	79	83
Trachea	<b>64</b>	35	65	<b>45</b>	63	70	64	87	84	83	79
Uterus	<b>71</b>	55	60	<b>47</b>	81	82	65	82	72	77	82

Data are presented as the mean values of the dynamin gene expression percentatges calculated from the individual values of the samples common to both NCBI-GEO records (GDS596 and GDS1096). The GPL96 Affymetrix probe set ID is displayed in parentheses below the gene symbol. Where there is more than one entry for the same gene, we selected the entry for which the Affymetrix probe set ID refers to the mRNA or coding DNA sequence (cds). The CNS tissues and organs are shown in the top block of the table. DNM1 and DNM3 gene expression values are shown in bold. See the legend of Figure 1.

nervous system (CNS) (Figure 2). This data mining analysis allowed us characterise dynamin gene expression trends in the CNS. We found clear evidence that the classical dynamin DNMI and DNMI3 genes reach their maximum expression level (100% measurement score) in all of the normal human CNS tissues studied. Moreover, this results is in contrast to the expression profiles observed in the other normal human organs and tissues studied, in which both the dynamin DNMI and DNMI3 genes showed approximately 50%. A detailed breakdown of the data is presented in Table 1. To improve the rigor of our analysis, we used the BioGPS gene expression portal [7,8] to explore the expression profiles of human dynamin-related genes. We found that the classical dynamin DNMI and DNMI3 genes are also maximally expressed in normal human CNS components and at significantly lower level in others organs or tissues [9,10].

The CNS consists of the brain and spinal cord, where synapses are the dynamic structures through which all nervous system signals traverse. Dynamamin polymerisation in membrane fission is thought to play a significant role in the synapse (Gene Ontology GO:0003373) which is suggestive of the concept of a relationship between synapses and dynamamins [11]. Several animal model studies have shown that dynamin genes are highly expressed in neurons [12-15]. In conclusion, based on their high levels of expression in human CNS, DNMI and DNMI3 can be considered as CNS-specific dynamin genes, at least in the following human CNS components: amygdala, foetal brain, whole brain, caudate nucleus, cerebellum, cerebellum peduncles, cingulate cortex, globus pallidus, hippocampus, hypothalamus, medulla oblongata, occipital lobe, parietal lobe, pineal gland, pineal gland, pons, prefrontal cortex, spinal cord, subthalamic nucleus, temporal lobe, and thalamus. We suggest that the role of dynamamins in the CNS could be a potentially interesting area for further biochemical research in neurodegenerative diseases.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

AR carried out the downloading of gene expression data from the NCBI-GEO database, helped design the study, and contributed to writing the manuscript. LA wrote the manuscript and helped design the study. Both authors read and approved the final manuscript.

#### Acknowledgements

This study was supported by a grant from the "Ministerio de Economía y Competitividad" of the Spanish Government (AGL 2008-00387/ALI).

Received: 28 December 2013 Accepted: 24 March 2014

Published: 28 March 2014

#### References

1. Praefcke GJK, McMahon HT: The dynamin superfamily: universal membrane tubulation and fission molecules? *Nature Rev Mol Cell Biol* 2004, **5**:133-147.
2. *Dynamamin Hope Page*. <http://www.endocytosis.org/Dynamamin/>.

3. Hinshaw JE, Schmid SL: Dynamamin self-assembles into rings suggesting a mechanism for coated vesicle budding. *Nature* 1995, **374**:190-192.
4. Sever S, Chang J, Gu C: Dynamamin rings: not just for fission. *Traffic* 2013, **14**:1194-1199. doi:10.1111/tra.12116.
5. Su AI, Wiltshire T, Batalov S, Lapp H, Ching KA, Block D, Zhang J, Soden R, Hayakawa M, Kreiman G, Cooke MP, Walker JR, Hogenesch JB: A gene atlas of the mouse and human protein-encoding transcriptomes. *Proc Natl Acad Sci U S A* 2004, **101**:6062-6067.
6. Ge X, Yamamoto S, Tsutsumi S, Midorikawa Y, Ihara S, Wang SM, Aburatani H: Interpreting expression profiles of cancers by genome-wide survey of breadth of expression in normal tissues. *Genomics* 2005, **86**:127-141.
7. Wu C, Orozco C, Boyer J, Leglise M, Goodale J, Batalov S, Hodge CL, Haase J, Janes J, Huss JW 3rd, Su AI: BioGPS: an extensible and customizable portal for querying and organizing gene annotation resources. *Genome Biol* 2009, **10**:R130.
8. Wu C, MacLeod I, Su AI: BioGPS and MyGene.info: organizing online, gene-centric information. *Nucl Acids Res* 2013, **41**(D1):D561-D565.
9. DNMI1 default BioGPS gene report. <http://biogps.org/#goto=genereport&id=1759>.
10. DNMI3 default BioGPS gene report. <http://biogps.org/#goto=genereport&id=26052>.
11. Watanabe S, Rost BR, Camacho-Pérez M, Davis MW, Söhl-Kielczynski B, Rosenmund C, Jørgensen EM: Ultrafast endocytosis at mouse hippocampal synapses. *Nature* 2013, **504**:242-247. doi:10.1038/nature12809.
12. Cook T, Mesa K, Urrutia RJ: Three dynamamin-encoding genes are differentially expressed in developing rat brain. *J Neurochem* 1996, **67**:927-931.
13. Gray NW, Fourgeaud L, Huang B, Chen J, Cao H, Oswald BJ, Hemar A, McNiven MA: Dynamamin 3 is a component of the postsynapse, where it interacts with mGluR5 and Homer. *Curr Biol* 2003, **13**:510-515.
14. Ferguson SM, Brasnjo G, Hayashi M, Wolfel M, Collesi C, Giovedi S, Raimondi A, Gong LW, Ariel P, Paradise S, O'Toole E, Flavell R, Cremona O, Miesenböck G, Ryan TA, De Camilli P: A selective activity-dependent requirement for dynamamin 1 in synaptic vesicle endocytosis. *Science* 2007, **316**:570-574.
15. Raimondi A, Ferguson SM, Lou X, Armbruster M, Paradise S, Giovedi S, Messa M, Kono N, Takasaki J, Cappello V, O'Toole E, Ryan TA, De Camilli P: Overlapping role of dynamamin isoforms in synaptic vesicle endocytosis. *Neuron* 2011, **70**:1100-1114.

doi:10.1186/1756-0500-7-188

Cite this article as: Romeu and Arola: Classical dynamamin DNMI and DNMI3 genes attain maximum expression in the normal human central nervous system. *BMC Research Notes* 2014 **7**:188.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

