

Adherence of cardiologist physicians to the American Heart Association guideline in approach to risk factors of cardiovascular diseases: An experience from a teaching hospital

Zarrintaj Hosseinzadeh-Shanjani⁽¹⁾ , Soodabeh Hoveidamanesh⁽²⁾,
Mozhdeh Ramezani⁽³⁾, Farnoush Davoudi⁽³⁾, Marzieh Nojomi⁽⁴⁾ 

Short Communication

Abstract

BACKGROUND: Cardiovascular disease (CVD) is the leading cause of death globally and has enormous costs for healthcare systems. This disease has a strong association with lifestyle behaviors. Therefore, applying reliable and effective strategies for prevention and treatment of CVD is important. In this study, we aimed to evaluate the adherence of cardiologist physicians to the American Heart Association (AHA) guideline for prevention of CVD.

METHODS: Using a cross-sectional study, data were gathered for 208 patients using their medical records in the cardiology ward of a general teaching hospital. A physician systematically reviewed the medical records and completed the checklist in each domain. Adherence to the AHA guideline was evaluated in treating physician's choices and recommendations regarding these eight variables: hypertension (HTN), dietary intake, weight management, diabetes management, physical activity, blood lipid management, smoking, and aspirin prescription.

RESULTS: Medical records of 208 patients (109 men and 99 women) with the mean age of 62 ± 14 years were reviewed. The frequency of CVDs was 5.3% for coronary heart failure (HF) and 67.8% for the acute coronary syndrome (ACS). Cardiovascular risk factors of patients were HTN (53.8%), diabetes (34.6%), hyperlipidemia (17.3%), smoking (17.8%), and obesity (31.7%). We found a proportion of 59%, 15%, and 26% for high, moderate, and low adherence to AHA guideline, respectively.

CONCLUSION: Our study showed almost 60% high adherence to the AHA guideline by physicians in a teaching hospital. The most and the least adherence to the AHA guideline were for obesity and diabetes recommendations, respectively. More studies are needed to evaluate preventive guideline adherence in Iran. Establishing national preventive and therapeutic guidelines may increase the physicians' adherence to them.

Keywords: American Heart Association, Cardiovascular Diseases, Cardiologists, Guideline, Guideline Adherence

Date of submission: 20 Apr. 2018, *Date of acceptance:* 02 Oct. 2018

Introduction

Cardiovascular disease (CVD) is the leading cause of death globally and has huge costs for health care systems in many countries.¹ CVD, particularly coronary artery disease (CAD), is one of the most preventable non-communicable disorders that can be reduced by modifying behavioral risks.²

How to cite this article: Hosseinzadeh-Shanjani Z, Hoveidamanesh S, Ramezani M, Davoudi F, Nojomi M. **Adherence of cardiologist physicians to the American Heart Association guideline in approach to risk factors of cardiovascular diseases: An experience from a teaching hospital.** ARYA Atheroscler 2019; 15(1): 38-43.

1- Assistant Professor, Department of Community Medicine, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran
2- Community Medicine Specialist, Preventive Medicine and Public Health Research Center AND Burn Research Center, Iran University of Medical Sciences, Tehran, Iran

3- Assistant Professor, Preventive Medicine and Public Health Research Center AND Department of Community Medicine, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

4- Professor, Preventive Medicine and Public Health Research Center AND Department of Community Medicine, School of Medicine, Iran University of Medical Sciences AND Academy of Medical Sciences of Islamic Republic of Iran, Tehran, Iran

Correspondence to: Marzieh Nojomi, Email: mnojomi@iums.ac.ir

CVD causes 38% of all deaths in Iran annually, and about 3 million people aged 40 and above die every year from ischemic heart diseases (IHDs) in Iran.²

The evidence suggests disturbing increase in the prevalence of CVD risk factors such as diabetes, obesity, and the metabolic syndrome.³⁻⁵ High-risk behaviors such as cigarette smoking, unhealthy dietary habits, physical inactivity, and psychosocial factors have a strong association with CVD.⁶ Furthermore, hypertension (HTN), diabetes, and dyslipidemia are under the influence of the unhealthy lifestyle.⁷ World Health Organization (WHO) has announced that lifestyle modification can prevent 75% of CVDs.²

Recommendations of international guidelines are the most effective way of ensuring that patients receive optimal care.⁸ Physician adherence to cardiovascular preventive care guidelines is one of the critical steps to improve the outcomes of patients. Moreover, using guidelines can improve clinical practice.⁹ Another study showed that use of guideline-based therapies was associated with significant reduction in mortality and other major adverse outcomes.¹⁰

For prevention and management of CVD, the American College of Cardiology (ACC) and the American Heart Association (AHA) develop guidelines, standards, and policies to promote optimal patient care and cardiovascular health.¹¹ The proposed guideline is one of the most reliable guides that are widely used in CVD prevention. It consists of recommendations for the control of blood glucose, lipids, weight, and blood pressure, encouragement of physical activity, safe diet, and also aspirin usage.¹¹ According to the importance of CVD in population health and its socio-economic consequences, applying reliable and effective strategies for prevention and treatment of these diseases is important. In this study, we aimed to evaluate the adherence of Iranian physicians to the AHA guideline for CVD prevention.

Materials and Methods

In this cross-sectional study, data about physicians' adherence to the AHA guideline were collected from medical records of patients admitted to the cardiology ward of a general teaching hospital of Iran University of Medical Sciences in Tehran City, Iran. Data collection was conducted from April 4 to July 6, 2015. Eligible patients were those who were admitted to the cardiology ward with one of the following diagnoses: IHD, HTN, heart failure (HF),

or stroke. We excluded all records that were incomplete or free of CVD.

A data collection checklist was developed for the evaluation of the adherence to the AHA guideline and was completed for each patient using his/her medical records. The selection of checklist domains was performed by an expert panel consisting of experts in relevant medical specialties including community medicine, cardiology, and endocrinology, and eight risk factors were selected for inclusion in the checklist. Adherence to the AHA guideline was evaluated in treating physicians' choices and recommendations regarding these eight variables: HTN, dietary intake, weight management, diabetes management, physical activity, blood lipid management, smoking, and aspirin prescription.

A physician systematically reviewed the medical records and completed the checklists in each domain.

At first, she examined the number of existing risk-factors for each patient. Then, she compared the physicians' recommendations in each patient's medical record with the AHA guideline to determine the number of well-treated risk factors. The next step was to calculate the adherence to the AHA guideline by dividing the first number by the second. The total adherence to the AHA guideline was illustrated in percentage. Finally, we considered an adherence for less than 50%, 50%-80%, and more than 80% of risk factors as a low, moderate, and high adherence, respectively, according to expert opinions in the hospital.

Treatment goals for each variable were as follows:

- **HTN:** the target blood pressure was 130/80 for diabetes and 140/90 for the others. Moreover, prescribing proper antihypertensive medications from different classes in cases of high blood pressure was a desirable goal for HTN.

- **Dietary intake:** getting recommendations for an overall healthy eating pattern or following the recommendations by the patients.

- **Aspirin:** prescribing a low-dose aspirin for individuals at higher risk of coronary heart disease (CHD), especially those with the 10-year risk of CHD $\geq 10\%$.

- **Weight management:** achieving and maintaining a desirable weight [body mass index (BMI): 18.5–24.9 kg/m²]. Furthermore, it was a goal that patients received proper recommendations about their weight when their BMI was ≥ 25 kg/m² or their waist circumference at iliac crest level was equal or more than 40 inches in men or ≥ 35 inches in women.

• **Diabetes management:** reaching normal fasting plasma glucose (110 mg/dl) or near normal hemoglobin A1c (HbA1c) (7%).

• **Physical activity:** at least 30 minutes of moderate-intensity physical activity on most/all days of the week or recommending this.

• **Blood lipid management:** low-density lipoprotein cholesterol (LDL-C) < 160 mg/dl if at least one risk factor was present. LDL-C < 130 mg/dl if two risk factors were present and the 10-year risk of CHD < 20%. LDL-C < 100 mg/dl if two risk factors were present and the 10-year risk of CHD ≥ 20% or if the patient had diabetes.

• **Smoking:** complete cessation and no exposure to environmental tobacco smoke.

All statistical analyses were performed using the SPSS software (version 16, SPSS Inc., Chicago, IL, USA). Categorical variables were described as frequency and percentage. Also, continuous variables were reported by mean ± standard deviation (SD).

For calculating the total adherence to the AHA guideline, we divided the number of the well-treated risk factors (in accordance to the AHA guideline) by a total number of the existing risk factors in the patients. The final results were reported in percentage.

Study protocols and procedures have been approved by the review board of Iran University of Medical Sciences.

Results

208 patients (109 men and 99 women) with mean age of 62 ± 14 years were admitted during the study duration and their medical records were reviewed. Frequency of CVD and risk factors are shown in table 1.

Adherence to the AHA guideline was first evaluated for each risk factor separately, and the results are shown in table 2. These included: HTN, diabetes, obesity, hyperlipidemia, and smoking. Among the studied risk factors, obesity treatment

had the highest adherence to the AHA guideline (100%) followed by smoking (86.5%). The least adherence was for diabetes with only 31.9%.

Table 1. Baseline clinical characteristics of the study population

Type of CVD	Women (n)	Men (n)	Total [n (%)]
CHF	9	2	11 (5.3)
ACS	59	82	141 (67.8)
Arrhythmia	6	1	7 (3.4)
Type of cardiovascular risk factor			
HTN	69	43	112 (53.8)
Diabetes	40	32	72 (34.6)
Hyperlipidemia	23	13	36 (17.3)
Smoking	3	34	37 (17.8)
Obesity	39	27	66 (31.7)

CVD: Cardiovascular disease; ACS: Acute coronary syndrome; CHF: Congestive heart failure; HTN: Hypertension

Recommendation of healthy alternatives for lifestyle risk factors according to the AHA guideline was investigated as shown in table 3. These included: healthy diet recommendation, physical activity, and aspirin prescription. The results of adherence to the AHA guideline are shown in table 4.

According to the scoring, 59% of the records indicated a high degree of adherence by physicians in the cardiovascular unit of one teaching hospital; 15% of the files showed a medium adherence, and 26% of them illustrated low adherence by physicians.

Discussion

This study has been done by evaluation of 208 patients' medical records who were admitted to the cardiovascular unit of one general teaching hospital. Adherence to the AHA guideline by the treating cardiovascular specialists was evaluated. Our study showed 59% high, 15% moderate, and 26% low adherence to the guideline by physicians in this teaching hospital.

Table 2. Adherence to the American Heart Association (AHA) guideline based on the presence of cardiovascular risk factor in the study population

Risk factor	Patients with the risk-factor (n)	Non-adherence to the AHA guideline recommendation	Full adherence to the AHA guideline recommendation
		[n (%)]	[n (%)]
HTN	112	23 (20.5)	89 (79.5)
Smoking	37	5 (13.5)	32 (86.5)
Hyperlipidemia	36	15 (41.5)	21 (58.5)
Diabetes	72	49 (68.1)	23 (31.9)
Obesity	66	0 (0)	66 (100)

AHA: American heart association; HTN: Hypertension

Table 3. Adherence to preventive measures according to the American Heart association (AHA) guideline

Preventive measure	Non-adherence to the AHA guideline recommendation	Full adherence to the AHA guideline recommendation
	[n (%)]	[n (%)]
Healthy diet recommendation	115 (55.3)	93 (44.7)
Physical activity recommendation	118 (56.7)	90 (43.3)
Aspirin prescription	42 (20.2)	166 (79.8)

AHA: American heart association

Table 4. Adherence to the American Heart Association (AHA) guideline

Adherence	Percentage
High	59
Moderate	15
Low	26

In one study which was done in Shahid Modarres Hospital, Tehran City, 52.9% of physicians did not adhere to ACC/AHA guidelines in patients with unstable angina,¹² which is almost in accordance to our study findings (41%). The increase in our adherence result can be due to the fact that current health system pays more attention to the prevention of chronic CVDs. WHO has mentioned that 3/4 of all cardiovascular-related deaths can be prevented by changing one's lifestyle.²

Smoking is one of the most important behavioral risk factors for CVD.¹³ In this study, 17.8% of the patients were smoker, and smoking cessation was advised to 68.7% of them. There is a relationship between blood pressure levels and occurrence of CVD even in blood pressures that have been defined as normal. In this study, 53.8% of patients had HTN, among which 79.5% had been followed.

High cholesterol/triglyceride (TG) and low high-density lipoprotein (HDL) levels are associated with the occurrence of CVD.¹⁴ In developed countries, cholesterol level is increasing, which is mostly because of the low level of activity and increased fat consumption. In our 208 patients, 36 had hyperlipidemia, from which 21 received diet recommendations and 15 patients did not receive any recommendation. We know that each 1% decrease in blood cholesterol level can cause a 2%-5% decrease in CVD occurrence.¹⁴ In this study, 79.8% of patients were on aspirin that is an effective way to decrease CVD.

Other recommendations in the AHA guideline which have been used in this study were obesity, diabetes, physical activity, and diet. We found that 44.7% of patients were advised to eat more vegetables and fruits and decrease their dietary salt and fat intake. No dietary recommendation was

given to 55.3% of patients.

Changes in diet and lifestyle are vital in preventing CVD and cardiovascular deaths.¹⁵ That's why the change of lifestyle and having a healthy diet are among the most important recommendations of the 2006 AHA guideline. According to our study, less than half of the patients did not receive any lifestyle recommendation, noting an important gap between adherence to the therapeutic recommendations of the guideline and the preventive part of that. In Spain, the researchers studied the percentage of adherence to ACC/AHA guidelines and its effects on changing the survival of patients with atrial fibrillation (AF). They found that using the guideline increased the survival of patients with AF up to 3 years.¹⁶

A study in Croatia showed that although most specialists believed that guidelines were beneficial, only 56.9% used them.¹⁷ HTN was thought to be the most important risk factor by primary care physicians, but diabetes was thought as the most important risk factor by the cardiologist. As mentioned before, guideline-based treatment for patients can reduce mortality and major adverse outcomes.^{10,18} Furthermore, most of the physicians accepted guidelines but only half of them used guidelines, and their knowledge about the guidelines was not satisfactory.¹⁷ Also in a study, physicians' adherence to the guideline was significantly lower in nonteaching clinics.¹⁹ Thus, these reasons as well as the lack of time, resources, and knowledge are most likely the main causes of poor management of the risk factors.¹⁷

Our study had some limitations such as incomplete records and taking data from medical records of the patients. Conducting study in just one general teaching hospital with the probability of more adherence to the guideline is another limitation and reduces generalizability of findings.

The strength point of the current study was to assess the prevention and risk factors of CVDs besides the treatment approach. Majority of previous studies have just considered the treatment approach to CVDs. We suggest further studies to focus on adherence to especially preventive

guidelines using prospective studies with more supervision of gathering data.

Conclusion

Our study showed 59% high, 15% moderate, and 26% low adherence to the AHA guideline by physicians in the cardiology ward of one teaching hospital. The most adherence to the AHA guideline was about obesity (100%), smoking (86.5%), aspirin prescription (79.8%), and blood pressure (79.5%), respectively. The less adherence was about diabetes (31.9%).

More studies are needed to evaluate preventive guideline adherence in Iran. Establishing national preventive and therapeutic guidelines may increase the physicians' adherence to them and finally promote the health of people.

Acknowledgments

This study has been derived from a PhD dissertation (No. P/22128) approved by Iran University of Medical Sciences. The authors are grateful to the assistants in the current study.

Conflict of Interests

Authors have no conflict of interests.

References

1. Tarride JE, Lim M, DesMeules M, Luo W, Burke N, O'Reilly D, et al. A review of the cost of cardiovascular disease. *Can J Cardiol* 2009; 25(6): e195-e202.
2. Perk J, De Backer G, Gohlke H, Graham I, Reiner Z, Verschuren WM, et al. European guidelines on cardiovascular disease prevention in clinical practice (version 2012): The fifth joint task force of the European society of cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of nine societies and by invited experts). *Atherosclerosis* 2012; 223(1): 1-68.
3. Declining prevalence of no known major risk factors for heart disease and stroke among adults-United States, 1991-2001. *MMWR Morb Mortal Wkly Rep* 2004; 53(1): 4-7.
4. Mokdad AH, Ford ES, Bowman BA, Dietz WH, Vinicor F, Bales VS, et al. Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *JAMA* 2003; 289(1): 76-9.
5. American Heart Association. Heart disease and stroke statistics-2005. Dallas, TX: American Heart Association; 2005.
6. D'Agostino RB Sr, Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, et al. General cardiovascular risk profile for use in primary care: The Framingham Heart Study. *Circulation* 2008; 117(6): 743-53.
7. Huxley R, Barzi F, Woodward M. Excess risk of fatal coronary heart disease associated with diabetes in men and women: Meta-analysis of 37 prospective cohort studies. *BMJ* 2006; 332(7533): 73-8.
8. Komajda M, Cowie MR, Tavazzi L, Ponikowski P, Anker SD, Filippatos GS. Physicians' guideline adherence is associated with better prognosis in outpatients with heart failure with reduced ejection fraction: The QUALIFY international registry. *Eur J Heart Fail* 2017; 19(11): 1414-23.
9. Cabana MD, Kim C. Physician adherence to preventive cardiology guidelines for women. *Womens Health Issues* 2003; 13(4): 142-9.
10. Shah BR, O'Brien EC, Roe MT, Chen AY, Peterson ED. The association of in-hospital guideline adherence and longitudinal postdischarge mortality in older patients with non-ST-segment elevation myocardial infarction. *Am Heart J* 2015; 170(2): 273-80.
11. Smith SC Jr, Allen J, Blair SN, Bonow RO, Brass LM, Fonarow GC, et al. AHA/ACC guidelines for secondary prevention for patients with coronary and other atherosclerotic vascular disease: 2006 update: endorsed by the National Heart, Lung, and Blood Institute. *Circulation* 2006; 113(19): 2363-72.
12. Farahzadi M, Shafiee A, Bozorgi A, Mahmoudian M, Sadeghian S. Assessment of adherence to ACC/AHA guidelines in primary management of patients with NSTEMI in a referral cardiology hospital. *Crit Pathw Cardiol* 2015; 14(1): 36-8.
13. Wannamethee SG, Lowe GD, Shaper AG, Rumley A, Lennon L, Whincup PH. Associations between cigarette smoking, pipe/cigar smoking, and smoking cessation, and haemostatic and inflammatory markers for cardiovascular disease. *Eur Heart J* 2005; 26(17): 1765-73.
14. Astrup A, Dyerberg J, Elwood P, Hermansen K, Hu FB, Jakobsen MU, et al. The role of reducing intakes of saturated fat in the prevention of cardiovascular disease: Where does the evidence stand in 2010? *Am J Clin Nutr* 2011; 93(4): 684-8.
15. Ignarro LJ, Balestrieri ML, Napoli C. Nutrition, physical activity, and cardiovascular disease: An update. *Cardiovasc Res* 2007; 73(2): 326-40.
16. Diez-Manglano J, Gomes-Martin J, Al-Cheikh-Felices P, Perez SI, Diez-Angulo R, Clemente-Sarasa C. Adherence to guidelines and mortality in atrial fibrillation. *Int J Cardiol* 2014; 176(2): 430-6.
17. Reiner Z, Sonicki Z, Tedeschi-Reiner E. Physicians' perception, knowledge and awareness of cardiovascular risk factors and adherence to prevention guidelines: The PERCRO-DOC survey. *Atherosclerosis* 2010; 213(2): 598-603.

18. Nguyen T, Le KK, Cao HTK, Tran DTT, Ho LM, Thai TND, et al. Association between in-hospital guideline adherence and postdischarge major adverse outcomes of patients with acute coronary syndrome in Vietnam: A prospective cohort study. *BMJ Open* 2017; 7(10): e017008.

19. Cheng-Lai A, Snead J, Ng C, Verges C, Chung P. Comparison of adherence to the 2013 ACC/AHA cholesterol guideline in a teaching versus nonteaching outpatient clinic. *Ann Pharmacother* 2018; 52(4): 338-44.