



Original Article

The Diagnostic Value of Pulsed Wave Tissue Doppler Imaging in Asymptomatic Beta- Thalassemia Major Children and Young Adults; Relation to Chemical Biomarkers of Left Ventricular Function and Iron Overload

Seham M Ragab¹, Waleed M Fathy², Walaa FAbd El-Aziz³ and Rasha T Helal ²

Departments of Pediatrics¹, Clinical pathology² and Cardiology³, Faculty of Medicine, Menoufia University. Naser street, Shebeen El-koom, Menoufia, Egypt.

Abstract. Background: Cardiac iron toxicity is the leading cause of death among β -thalassaemia major (TM) patients. Once heart failure becomes overt, it is difficult to reverse.

Objectives: To investigate non-overt cardiac dysfunctions in TM patients using pulsed wave Tissue Doppler Imaging (TD I) and its relation to iron overload and brain natriuretic peptide (BNP).

Methods: Thorough clinical, conventional echo and pulsed wave TDI parameters were compared between asymptomatic 25 β -TM patients and 20 age and gender matched individuals. Serum ferritin and plasma BNP levels were assayed by ELISA.

Results: TM patients had significant higher mitral inflow early diastolic (E) wave and non significant other conventional echo parameters. In the patient group, pulsed wave TDI revealed systolic dysfunctions, in the form of significant higher isovolumetric contraction time (ICT), and lower ejection time (E T), with diastolic dysfunction in the form of higher isovolumetric relaxation time (IRT), and lower mitral annulus early diastolic velocity E` (12.07 ± 2.06 vs 15.04 ± 2.65 , $P = 0.003$) compared to the controls. Plasma BNP was higher in patients compared to the controls. Plasma BNP and serum ferritin had a significant correlation with each other and with pulsed wave conventional and TDI indices of systolic and diastolic functions. Patients with $E/E' \geq 8$ had significant higher serum ferritin and plasma BNP levels compared to those with ratio < 8 without a difference in Hb levels.

Conclusion: Pulsed wave TDI is an important diagnostic tool for latent cardiac dysfunction in iron-loaded TM patients and is related to iron overload and BNP.

Citation: Ragab S.M., Fathy W.M., El-Aziz W.F., Helal R.T. The Diagnostic value of pulsed wave tissue doppler imaging in asymptomatic beta- thalassemia major children and young adults; relation to chemical biomarkers of left ventricular function and iron overload. *Mediterr J Hematol Infect Dis* 2015, 7(1): e2015051, DOI: <http://dx.doi.org/10.4084/MJHID.2015.051>

Published: August 24, 2015

Received: May 17, 2015

Accepted: July 15, 2015

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Correspondence to: Dr. Seham Mohammed Ragab. Assistant Professor Of Pediatrics. Pediatric Department, Hematology and Oncology Unit, Faculty of Medicine, Minoufia University, Menouf^{7th}, EL-HadethaStreet, Minoufia, Egypt. Tel.:0020483663235 – Mobil: 00201220190474. Fax: 0020482322810. E-mail: sehambasel@gmail.com. sehambasel@yahoo.com. seham.ragab@med.menoufia.edu.eg

Introduction. Thalassemia is one of the most common genetic disorders. Thus, it is considered a global health problem. Worldwide, about 5% of the population carry globin variants. Beta (β)–Thalassemia is caused by the reduced synthesis of β -globin chains, which leaves an erythrocyte excess of unopposed α -chains resulting in

ineffective erythropoiesis and chronic hemolytic anemia.¹

According to the severity, β -thalassemias are classified into: transfusion dependent β thalassemias (TDT) or β -thalassemia major (TM), non transfusion dependent β -thalassemia (NTDT) or β - thalassemia

intermedia (TI) and β -thalassemia trait (asymptomatic carriers). β -TM is the severest form that develops during the first year of life and requires lifelong transfusion therapy for survival.²

Although improving survival, repeated blood transfusion regimen causes iron overload and iron toxicity in different organs including the heart.³

Despite the progress in iron chelation therapy, congestive heart failure due to iron accumulation.

Is still the leading cause of death in β -TM patients.^{4,5}

Iron overload in combination with other inflammatory and immunogenetic factors can cause left ventricular systolic dysfunction, dilatation and failure, whereas the sole iron overload may result in left ventricular diastolic dysfunction with myocardial restriction and subsequent pulmonary hypertension and right ventricular dilatation.⁶

Patients with TM may remain asymptomatic and global left ventricular (LV) function may be preserved until late in the disease process.^{7,8} So, early detection of myocardial dysfunction may be useful in the management plan.⁹

Echocardiography is an essential imaging modality for diagnosis of ventricular function,¹⁰ that allows exclusion of overt LV systolic dysfunction (left ventricular ejection fraction < 50%).¹¹

However, changes of segmental wall motion – the early sign of myocardial dysfunction in thalassemia patients - may be subtle and could be missed by conventional echocardiographic examination which may remain normal until late stages during this disease process.¹²

Tissue Doppler Imaging (TDI) is a relatively new Doppler ultrasound modality that records regional systolic and diastolic velocities within the myocardium. It allows quantitative measurement of both systolic and diastolic velocities directly from the ventricular myocardium with the determination of the extent of mitral annular displacement in systole and diastole.¹³

This new technique can show additional information compared with other echocardiography techniques, detecting even minor changes before the occurrence of abnormal indices of global ventricular dysfunction.¹⁴

Brain natriuretic peptide (BNP) is one of the natriuretic peptide system that is stored in the myocardial cells as pre- proBNP. It is secreted from the heart as a result of direct wall stress, caused by either stretch or pressure affecting cardiocytes. Once released, BNP has pronounced natriuretic, diuretic and vasodilating properties, working to dramatically reduce volume overload and hypertension.¹⁵ BNP level is useful for the diagnosis of left ventricular systolic and diastolic dysfunctions and is correlated with the severity and prognosis.^{16,17}

So, the aim of this work was to investigate the

utility of pulsed wave DTI to detect latent or non-overt cardiac dysfunctions in asymptomatic TM patients and its relation to the iron overload assayed by serum ferritin and to BNP as a biomarker of cardiac dysfunction.

Materials and Methods. This is a cross-sectional study, performed upon 45 subjects (patients and controls); 25 β -TM patients and 20 age and sex matched healthy individuals as controls.

The patient group included 25 multi-transfused β -TM patients (14 males and 11 females). Their ages ranged from 4 to 20 years with mean age of 12 ± 5.79 years and median of 12 years. These patients were kept on a regular blood transfusion regimen (every 3-4 weeks) since infancy to maintain pre-transfusion Hb above 7 gm/dl and post-transfusion Hb above 10gm/dl. They were on long-term chelation therapy for at least one year either by Deferoxamine (DFO) monotherapy, 30–50 mg/kg body weight by subcutaneous infusion with an infusion pump for 8–12 h, 5 days per week (16/25=64%) , oral Deferasirox monotherapy , 20-30 mg/kg/day,daily (4/25=16%) or combined therapy with DFO and Deferasirox (DFO at the dosage of 40 mg/kg/day for 3 days/week and daily Deferasirox at the dosage of 30 mg/Kg/day, 5/25=20%).

The study included cardiac asymptomatic TM patients with ejection fraction >55% and a normal resting 12-lead electrocardiogram (ECG).

Patients on cardiovascular treatment, with any cardiovascular complaints, documented arrhythmia, hypertension, renal disease, diabetes mellitus, congenital or rheumatic heart disease, use of medications altering myocardial functions or a history of smoking were excluded. Also, those who developed transfusion associated circulatory overload (TACO) or transfusion-related acute lung injury (TRALI) were excluded.

The control group consisted of 20 healthy age and gender-matched subjects (10 males and 10 females). Their ages ranged from 4 to 18 years with mean age of 10.9 ± 4.86 years and median of 10 years. They were free from acute (especially viral illness) or chronic illness (including cardiac diseases) with no family history of chronic hemolytic anemia. All controls had normal complete blood count (CBC), Hemoglobin (Hb) electrophoresis with normal ECG and conventional Echocardiographic findings. They had been randomly selected from children presented to our general outpatient clinic for routine check up especially for growth, or for non-specific complaints like non-specific abdominal pain.

This study had been carried out at the Hematology Unit, Pediatric Department in collaboration with Clinical Pathology and Cardiology Departments, Faculty of Medicine, Menoufia University, Egypt, in the period of time between January 2012 and

September 2013. The study was approved by the ethical committee of Menoufia Medical School, and informed consent was obtained from the patient or his or her legal guardian.

Methods: All participants in the study were subjected to a full history taking and comprehensive clinical examination including a cardiac examination.

For the patient group, a special emphasis was given to the age of the disease manifestations, time of the first blood transfusion, frequency of blood transfusion with calculation of red blood cells transfusion index (RBCsTI) during the last year, chelation therapy details, hepatic, renal, histories and history of splenectomy.

For all included children (patients and controls) weight and height were measured by the standard methods and plotted against age and sex specific centiles.

The participants were investigated by the following:

I- Conventional Echocardiography

Echocardiography, in the form of complete two-dimensional, continuous and pulsed wave echocardiographic examination, was done using ultrasound machine (vivid 9, General Electric Medical Systems, Horton, Norway), equipped with 5MHz variable frequency harmonic –phased array transducer with simultaneous ECG monitoring, performed without sedation, during normal respiration in the left lateral decubitus. Images were recorded in the standard parasternal long axis, apical four and two chamber views.

Conventional Echo-Doppler Measurements

Routine M-mode, two-dimensional continuous wave Doppler recordings were obtained for each subject. The left atrial (LA), aortic (AO) diameters and left ventricular (LV) internal cavity dimensions including left ventricular end systolic diameter (LVESD) and left ventricular end diastolic diameter (LVEDD) were determined. LV ejection fraction (EF) and LV fractional shortening (FS) were measured using Teichholz's M-mode formula.¹⁸

Transmittal flow patterns were obtained by pulsed-wave Doppler echocardiography from apical four-chamber view. The peak of early diastolic flow velocity (E), the peak of late diastolic flow velocity (A) and the ratio of E/A were measured.

II- Pulsed Wave Tissue Doppler Imaging (TDI) Measurements

The pulsed wave TDI was performed using the same machine. To display tissue velocities; from the apical 4 and 2-chamber views, the Doppler sample volume was placed at four different sites of the mitral annulus: anterior, lateral, septal and inferior walls in order to record major velocities. The following parameters were registered: mitral annulus systolic velocity (S'), mitral annulus early diastolic velocity (E'), mitral annulus late diastolic velocity (A') and

time intervals; isovolumetric contraction time (ICT), isovolumetric relaxation time (IRT) and ejection time (ET).

Then calculation of the mean E/E' (mitral inflow E wave/ E' mitral annulus velocity) ratio was done. According to the E/E' ratio, patients were classified into: patients with $E/E' \geq 15$ (diastolic dysfunction), patients with $E/E' \geq 8$ but less than 15 (suspected diastolic dysfunction) and those with $E/E' < 8$ (without diastolic dysfunction).^{19,20}

All pulsed-wave Doppler and PW-TDI parameters were measured at the end of expiration, at a sweep speed of 100 mm/s on three consecutive heart beats and the average for each was taken.

All data were obtained according to the recommendations of the American Society of Echocardiography.²¹

III-Laboratory investigations including:

1- Complete blood count (CBC): using Beckman 750, Int, U.S.A, Auto-counter after calibration.

2- Serum ferritin level was measured by Enzyme Linked Immune Sorbent Assay (ELISA) technique (ELISA, GenWay Biotech, Inc, NP 000137, Swiss) on Microplate reader (Bio-Rad 680 Hercules, California, USA). The mean yearly serum ferritin level in the previous year was considered (on the average of 4 determinations) for patients and at the time of sampling for the controls.

3- Plasma BNP: Three ml venous blood samples were drawn by sterile vein-puncture on EDTA tube. Blood samples were immediately centrifuged for 15 minutes at 3000 rpm; plasma samples were separated then were stored at -20°C until analysis. Plasma BNP level was measured by ELISA using kits supplied by Ray Biotech, Inc. GA 30092.

For the patient group, the echocardiographic examination and the blood sampling for CBC and BNP assay were performed on the fourth day following blood transfusion.²²

Statistical analysis: The data were processed on an IBM-PC compatible computer using SPSS version 16 (SPSS Inc., Chicago, IL, USA). Continuous parametric variables were presented as means \pm SD while for categorical variables numbers (%) were used. Chi-square test was used for qualitative variables. The difference between 2 groups was performed by student's t-test for parametric continuous variables and Man Whitney (U) test for non-parametric variables. Pearson correlation (r): was the test used to measure the association between two quantitative parametric variables and Spearman correlation coefficient was applied for non-parametric data. Receiver Operating Characteristic curve (ROC curve) analysis is a graph of sensitivity against 1- specificity at different cutoff points. The optimal cutoff point is that gives the highest sensitivity and specificity. Two-sided *p* value of < 0.05 was considered statistically significant.

Results. For the patient group, their ages at diagnosis ranged from 0.5–1.5 years with a mean of 0.77 ± 0.24 years. The mean age of first blood transfusion was 0.69 ± 0.21 years with a range of 0.5–1 year. The mean duration of transfusion treatment was 10.1 ± 5.11 years, that of the number of the transfusions /year was 11.2 ± 1.22 (median of 11 transfusions/year).

Comparison between the studied groups regarding clinical and laboratory data were represented in **Table 1**. The studied groups were matched regarding age, sex, the mean body weight, the mean height, and pulse rate. History of splenectomy was documented in 12 (48%) of TM patients. The patient group had a significantly lower post-transfusion Hb level with significantly higher mean yearly serum ferritin and

plasma BNP levels.

The conventional echocardiography parameters were presented in **Table 2**. Compared to the controls, TM patients had significant higher AO, LA, LVEDD, LVESD diameters and the mitral inflow early diastolic wave velocity (E). No significant difference was found between the studied groups regarding the left ventricular EF, left ventricular FS, the mitral inflow late diastolic wave velocity (A) or the E/A ratio.

As regard to the pulsed wave TDI parameters(**Table 3**), TM patients had significant higher ICT and lower ET compared to the controls in all tested sites of the mitral annulus as well as in the mean values of these parameters. The mean IRT and its values at the septal and the lateral walls of the mitral annulus were

Table 1. Comparison of the clinical and laboratory parameters between TM patients and controls.

Variables	Patients (25)	Controls (20)	P value
Age (years)	12.0 \pm 5. 79	10.9 \pm 4.86	0.56
Gender , males ,n(%)	14(56)	10(50)	1
Body weight (kg)	27.88 \pm 10.68	40.87 \pm 21.6	
< 5 th percentile , n(%)	6(24)	0(0)	0.19
Normal percentile range, n (%)	19(76)	20(100)	0.019
Height (cm)	129.28 \pm 21.63	137.53 \pm 31.53	0.23
< 5 th percentile ,n (%)	9(36)	0(0)	
Normal percentile range, n(%)	16(64)	20(100)	0.003
Pulse (beat /min)	90.32 \pm 6.24	83.2 \pm 11.22	0.064
Splenectomy,n (%)	12(48%)	-	-
RBCs TI(ml/kg/year)	109.75 \pm 30.39	-	-
Chelation ,n (%)			
DFO	16(64)	-	-
Deferasirox	4(16)		
DFO+Deferasirox	5(20)		
Pre transfusion Hb(g/dl)	7.28 \pm 1.03	12.2 \pm 0.58	<0.0001
Post transfusion Hb(g/dl)	10.38 \pm 0.41	12.2 \pm 0.58	<0.0001
Serum ferritin (ng/ml)	4126.12 \pm 3042.80	124.50 \pm 22.81	<0.0001
Median	4408	122	
Range	825-10200	96-170	
BNP (pg/ml)	29.28 \pm 14.30	8.80 \pm 1.81	<0.0001

Bold numerical values indicate significance.

Table 2. Comparison of conventional echocardiographic measures between TM patients and the controls.

Variables	Patients (25)	Controls (20)	P value
AO(mm)	22.92 \pm 4.22	19.40 \pm 3.61	0.02
LA(mm)	30.40 \pm 5.84	23.60 \pm 8.0	0.01
LVEDD(mm)	46.48 \pm 8.38	33.30 \pm 3.92	<0.0001
LVESD(mm)	29.16 \pm 5.83	20.80 \pm 2.48	<0.0001
EF(%)	66.24 \pm 5.57	69.50 \pm 3.34	0.11
FS (%)	37.48 \pm 6.65	38.20 \pm 2.44	0.31
E(cm/sec)	112.60 \pm 19.91	91.0 \pm 15.22	0.001
A(cm/sec)	75.20 \pm 22.65	69.60 \pm 14.07	0.56
E/A ratio	1.54 \pm 0.34	1.36 \pm 0.29	0.28

AO: aortic diameter; LA: left atrium diameter; LVEDD: left ventricular end diastolic diameter; LVESD: left ventricular end systolic diameter; EF: ejection fraction; FS: fraction shortening; E: mitral inflow early diastolic wave; A: mitral inflow late diastolic wave; E/A: mitral inflow early to late diastolic wave ratio. Bold numerical values indicate significance.

Table 3. Comparison of Pulsed wave TDI measures between TM patients and the controls.

	Septal wall	Lateral wall	Anterior wall	Inferior wall	The mean
S` (cm/s)					
Patients	7.90±0.87	8.92±2.21	7.90 ±0.88	8.84±1.34	8.28±0.9
Controls	8.56±1.98	9.10±1.37	8.04±2.17	8.20 ±1.62	8.6±1.26
P value	0.53	0.59	0.54	0.15	0.47
E` (cm/s)					
Patients	11.50±2.22	12.90±1.85	11.0±2.36	12.90 ±2.77	12.07 ±2.06
Controls	13.16±3.92	16.84±3.32	14.40±2.52	15.48±3.15	15.04±2.65
P value	0.14	0.001	0.002	0.04	0.003
A` (cm/s)					
Patients	7.90 ±1.45	7.70±0.67	7.10±1.20	7.80 ±1.40	7.45 ±0.92
Controls	7.92±2.63	7.76±2.52	7.4±1.19	9.16±2.09	8.2±2.02
P value	0.94	0.61	0.69	0.12	0.48
ICT(ms)					
Patients	66.88±8.98	69.96±11.02	67.16±14.40	64.6±17.79	67.15±10.89
Controls	45.0 ±8.26	58.20±29.76	41.30±5.12	45.90 ±9.53	47.85±8.97
P value	<0.0001	0.001	<0.0001	0.001	<0.0001
ET(ms)					
Patients	231.70±17.99	228.20±30.20	231.60±29.60	225.40 ±20.95	229.22±23.17
Controls	250.96±24.34	252.16±23.58	254.88±23.42	253.08±24.24	252.77±22.25
P value	0.02	0.03	0.04	0.008	0.02
IRT(ms)					
Patients	59.60±10.11	53.0±10.84	56.72±10.15	56.08±12.64	56.28±8.35
Controls	46.90 ±8.05	45.40±6.57	51.90±10.49	50.60 ±9.51	48.7±4.18
P value	0.001	0.04	0.14	0.49	0.008
E/E`					
Patients					8.0±2.09
Controls					7.42 ±0.3
P value					0.77

S`: mitral annulus systolic velocity ;E`: mitral annulus early diastolic velocity ;A`: mitral annulus late diastolic velocity; ICT: isometric contraction time; ET: ejection time; IRT: isovolumic relaxation time; E/E`: mitral inflow early diastolic wave E / mitral annulus early diastolic velocity E` ratio. Bold numerical values indicate significance

significantly higher in TM compared to the controls. The mean E` as well as its values at the lateral, anterior and inferior walls of the mitral annulus were significantly lower in TM patients compared to the controls. No significant difference was found in the S` or A` between the studied groups at any of the tested mitral annulus walls or in the mean values. TM patients had non-significant difference in E/E` ratio in comparison to the controls. Abnormal E/E` (mitral inflow E wave/E` mitral annulus velocity) ratio (≥ 15) was not found in any of TM patients. According to E/E`, TM patients were classified into those with E/E` < 8 (16 patients, 64%) and those with this ratio ≥ 8 but less than 15 (9 patients, 36%). Patients with E/E` ≥ 8 exhibited significant higher RBCs TI , mean yearly serum ferritin and plasma BNP levels compared to those with E/E` ratio < 8 without difference in post-transfusion Hb levels (**Figure 1 A and B**).

Univariate analysis among TM patients revealed that serum ferritin and plasma BNP were positively correlated with each other and each of them had significant positive correlation with the mean values of

E, ICT, and IRT; significant negative correlation with ET without significant correlation with E/A or E/E` ratios (**Table 4**).

The data, obtained from the Roc Curve, showed that the best sensitivity of 100% and specificity of 81.9% for the plasma BNP were at cutoff point of 28.5 pg/ml in ruling out diastolic dysfunction (E/E < 8). Negative predictive value was 100 % while positive predictive value was 75%.The area under the curve was 0.86, P = 0.003 (95% CI = 0.71 – 1.01) (**Figure 2A**).

For the mean yearly serum ferritin , the data obtained from the Roc Curve revealed that serum ferritin level at cutoff point of 4790.5 ng /ml had the best sensitivity of 88.9% and specificity of 81.2% in ruling out diastolic dysfunction (E/E < 8). Negative predictive value was 92.9% while positive predictive value was 72, P =0.007. The area under the curve was 0.83 (95% CI = 0.64 – 1.02) (**Figure 2B**).

Comparison of the pulsed wave TDI mean parameters regarding the age and serum ferritin categories (age of < 14 years and ≥ 14 years ; serum ferritin ≤ 2500 ng/ml and > 2500 ng/ml) revealed that




Figure 1a. Comparison of post transfusion Hb (g/dl), plasma BNP (pg/ml) and RBCs TI (ml/kg/year) in TM patients with E/E' ratio ≥ 8 compared to patients with E/E' < 8.

Figure 1b. Comparison of mean yearly serum ferritin (ng/ml) in TM patients with E/E' ratio ≥ 8 compared to patients with E/E' < 8.

Table 4 Correlations between BNP and serum ferritin with different conventional echo cardiography and TDI parameters .

	BNP (pg/ml)		Serum ferritin (ng/ml)	
	Correlation coefficient	P value	Correlation coefficient	P value
Serum Ferritin (ng/ml)	+ 0.63	<0.0001		
E (cm/sec)	+ 0.62	<0.0001	+ 0.59	<0.0001
E /A	+ 0.29	0.09	+ 0.12	0.48
ICT(ms)	+ 0.48	0.003	+ 0.57	<0.0001
ET (ms)	- 0.68	<0.0001	- 0.44	0.008
IRT(ms)	+ 0.45	0.007	+ 0.55	0.001
E/E'	+ 0.30	0.08	+ 0.25	0.14

E: mitral inflow early diastolic wave; A: mitral inflow late diastolic wave; E/A: mitral inflow early to late diastolic wave ratio; ICT: isometric contraction time; ET: ejection time; IRT: isovolumic relaxation time; E/E': mitral inflow early diastolic wave E / mitral annulus early diastolic velocity E' ratio. Bold numerical values indicate significance.




Figure 2 A. Receiver operator characteristic curve of BNP values for the identification of patients with E/E' ≥ 8 .

Figure 2 B. Receiver operator characteristic curve of serum ferritin values for the identification of patients with E/E' ≥ 8 .

Table 5. Comparison of Pulsed wave TDI mean measures among TM patients regarding the age and serum ferritin categories.

	Age < 14 years (14)	Age ≥14 years (11)	P value	Serum ferritin ≤2500 (ng/ml) (9)	Serum ferritin >2500 (ng/ml) (16)	P value
S` (cm/s)	8.79±0.99	8.45±1.45	0.5	8.7±1.4	8.42±1.00	0.6
E` (cm/s)	15.54±2.68	14.41±2.61	0.3	14.51±2.38	15.97±2.99	0.19
A` (cm/s)	8.54±2.34	7.77±1.52	0.36	9.00±2.42	7.75±1.67	0.14
ICT(ms)	67.66±9.81	67.32±11.37	0.94	65.69±10.73	68.53±10.25	0.52
ET(ms)	268.02±15.14	237.07±25.33	0.002	258.16±17.86	237.42±34.04	0.09
IRT(ms)	54.25±7.99	58.79±8.33	0.18	51.78±7.4	58.76±7.88	0.04
E/E`	7.41±1.41	8.76±2.61	0.11	7.12±0.74	8.5±2.45	0.14
E/E` <8	11(78.6)	5(45.5)		8(88.9)	8(50)	
E/E` ≥8	3(21.4)	6(54.5)	0.087	1(11.1)	8(50)	0.052

S`: mitral annulus systolic velocity; E`: mitral annulus early diastolic velocity; A`: mitral annulus late diastolic velocity; ICT: isometric contraction time; ET: ejection time; IRT: isovolumic relaxation time; E/E`: mitral inflow early diastolic wave E / mitral annulus early diastolic velocity E` ratio. Bold numerical values indicate significance.

the ET was significantly lower in TM patients ≥ 14 years compared to those < 14 years. TM patients with serum ferritin >2500 ng/ml had significantly higher IRT compared to those with serum ferritin level ≤ 2500 ng/ml. There was no significant difference in the age categories regarding E/E` ratio < 8 or ≥ 8, while there was a trend of prevalent E/E` ratio < 8 in those with serum ferritin ≤ 2500 ng/ml (P=0.052) (**Table 5**).

Discussion. Cardiac failure due to iron overload remains the most common cause of death in β-TM patients accounting for up to 71% of all deaths from this disease.³

Although intense iron chelation therapy can prevent and delay myocardial dysfunction, once dysfunction has become clinically evident, it is difficult to reverse.²³

In this study, M-mode conventional echocardiography revealed a significant elevation in LV dimensions (LVEDD and LVESD) in the studied TM patients compared to the controls, which is concordant with what was reported by other investigators.^{10,24,25} This cardiac dilatation is attributed to cardiac compensation and adaptation to the chronic high-output anemic state and hypoxia.²⁶

In accordance with previous studies,^{20,27} in our study asymptomatic TM patients had a mean values of LVEF and FS comparable to those of the control group suggesting preserved systolic function, as assessed by M-mode conventional echocardiography, till late in the disease. However, other studies reported a significantly lower LVEF in thalassemia patients in comparison with healthy age and sex-matched individuals.^{28,29}

Echocardiographic evaluation of diastolic functions has been traditionally performed by measurement of trans-mitral flow parameters including the early (E) and late (A) diastolic filling velocities and the E/A ratio with conventional pulsed wave Doppler. The trans-mitral E wave is related to the time course of

active LV relaxation that generates a pressure gradient from the left atrium through the LV inflow tract.³⁰

In this study, mitral E velocity was higher in patients than in controls without a significant difference in the A velocity or E/A ratio. Similar results were reported in other studies.^{14,24,29,31} Absence of a significant difference in E/A ratio in TM patients compared to the controls could be due to the exclusion of patients with heart failure symptoms. It could be that the E/A alone is not sufficient to diagnose diastolic dysfunction.²⁹

Nevertheless, different patterns of abnormality were documented by other researchers,^{27,32} who found a significant reduction in both E and A velocities in TM patients than controls without significant alteration in E/A ratio. Moreover, E/A ratio was found to be increased in thalassemia patients in the study done by Garadah et al.,³³ denoting restrictive diastolic dysfunction. In this regard, out of our studied 25 TM patients, only 2 were found to have E/A ratio > 2 (denoting restrictive filling pattern - grade 3 diastolic dysfunction) while the others had this ratio between 1 and 2 (normal value). It has been postulated that myocardial iron deposition in some TM patients may not directly affect left ventricular contractility, but it may rather cause left ventricular myocardial restriction causing this restrictive diastolic dysfunction.³⁴

In fact, LV diastolic function, as measured by conventional pulsed Doppler transmitral flow recordings, is limited by its dependence on age and loading conditions.²²

So, elevated E velocity among our studied TM patients who were anemic compared to the controls without higher mean E/A ratio reflects an increase in the preload state due to chronic anemia.³⁵ However, considering the 2 patients with E/A ratio > 2, iron deposition was the contributing factor that dominated the preload state resulting in this restrictive pattern.

Cardiac iron deposition in TM patients was found to

be patchy and more in myocytes rather than the interstitium.³⁶ This contributes to regional wall motion abnormalities found in these patients.³⁷

Pulsed wave TDI can detect regional systolic and diastolic myocardial dysfunctions earlier than global dysfunction in thalassemia patients.²³ It has the advantage that the measured velocities have been reported to be independent of loading conditions.³⁸

In the current study, pulsed wave TDI of TM patients revealed presence of systolic dysfunction in the form of significantly higher mean and regional ICT and lower ET values compared to the controls with non-significant lower regional and mean values of the systolic velocity S' . TM patients ≥ 14 years old had significant lower mean ET compared to those below 14 years without any difference in other parameters, meaning that systolic dysfunction was more evident by increasing age.

Significant reduction in the lateral annulus ejection time of TM patients (age ≤ 16 years) compared with healthy subjects was reported by Iarussi et al.¹⁴ Furthermore, Abdelmokter and Azer²⁸ and Garadah et al.³³ had reported significant lower tissue Doppler systolic velocity in the β -TM group compared to controls.

In terms of diastolic dysfunction indices, the studied TM patients had significantly lower early diastolic velocity E' and higher IRT.

Decreased E' is one of the earliest markers of diastolic dysfunction and is present in all stages of diastolic dysfunction.¹³ Reduced E' velocity in TM patients was found by other researchers who attributed this to myocardial stiffness.^{28,33}

Prolonged IRT had been previously reported in TM patients with normal systolic function.^{27,39} It was suggested that prolonged IRT to be the earliest sign of diastolic dysfunction in these patients and to reflect iron-induced impairment in left ventricular relaxation.³⁹

Combining trans-mitral flow velocity with annular velocity (E/E') has been proposed as a tool for assessing LV filling pressures since it is influenced by both trans-mitral driving pressure and myocardial relaxation.⁴⁰ Because E' velocity remains reduced and mitral E velocity increases with higher filling pressure, the ratio between trans-mitral E and E' (E/E' ratio) correlates well with LV filling pressure or pulmonary capillary wedge pressure (PCWP). The subjects have diastolic dysfunction when PCWP is ≥ 20 mm Hg if E/E' is ≥ 15 , a normal function if E/E' is < 8 ^{19,38} and a suspected diastolic dysfunction having a E/E' ratio between 8 and 15 (grey-intermediate zone).^{19,20}

The use of E/E' ratio has provided independent and incremental diagnostic and prognostic information in some major cardiac diseases, including heart failure.¹³ It has attracted attention for assessing LV diastolic function due to its load independent nature, its un-

affection by elevated LA pressure and linear correlation with LV end diastolic pressure.²⁷

E/E' ratio has a peculiar diagnostic importance for diastolic dysfunction among TM patients.⁴¹

In this regard, our study results revealed that the mean E/E' of the TM patients was higher compared to the controls yet the difference did not reach a significant level ($P > 0.05$). None of the studied TM patients had $E/E' > 15$ but in 9 out of the studied 25 patients (36%), E/E' ratio was in the gray zone, between 8 and 15, while the other 16 (64%) patients had normal E/E' ratio of < 8 . Regarding E/E' ratio, both patient categories had comparable post-transfusion Hb level with higher serum ferritin and BNP in those with a high ratio. The non-significant elevation of E/E' ratio is in accordance with Agha et al.³¹ but not with Kremastinos et al.²⁰ and Parale et al.²⁷ who found that there was a significant elevation in E/E' in the TM patients compared to the control group. This difference regarding the E/E' ratio with the 2 mentioned studies^{20,27} could be related to the difference in the age range, the iron load and the use of chelation therapy. Actually, our included children were younger than those of the first study (12.0 ± 5.79 years versus, 27.2 ± 12.5 years for our patients and their patients respectively). Also all involved patients in our study were under regular chelation therapy with lower mean yearly serum ferritin level than those of the second study (4126.12 ± 3042.80 ng/ml versus 8370.85 ± 2660.35 ng/ml respectively) who did not receive any chelation therapy before.

It is worth mentioning that, our results indicate the absence of global LV diastolic dysfunction that is in accordance with what was reported before.^{10,22}

So, among our non-symptomatic TM patients having a normal global systolic function by conventional echocardiography, the pulsed wave TDI had detected combined latent systolic and diastolic dysfunctions denoting the importance of this technique. By this, pulsed wave TDI detected combined systolic and diastolic dysfunctions in thalassemia patients as in previous reports.^{23,28}

Considered as a surrogate marker for heart failure,^{16,17} BNP was assayed in this work. Our studied TM patients exhibited significantly higher BNP compared to the controls being higher in those with suspected diastolic dysfunction ($E/E' \geq 8$). Its level demonstrated significant correlations with systolic (positive correlation with ICT and negative correlation with ET) and diastolic (positive correlation with both E wave and IRT) dysfunction indices, with a positive correlation trend with E/E' ratio ($P = 0.08$). BNP also had good correlation with serum ferritin as an indicator of iron overload. Elevated BNP in TM patients and its association with diastolic dysfunction was documented in previous studies.^{20,41}

Nikolidakis et al.⁴² found that BNP levels were

statistically higher in the severely iron-loaded thalassemia group (with MRI T2* < 24 ms). BNP levels inversely correlated with myocardial T2 relaxation time values. The authors adopted a cut-off value of 29 pg/ml (sensitivity of 88%, specificity of 58%, negative predictive value of 94% and positive predictive value of 37%) for the identification of patients with severe myocardial hemosiderosis using magnetic resonance imaging as the comparative “gold” standard.

In accordance, our results revealed a BNP level of 28.5 pg/ml in ruling out diastolic dysfunction (E/E' < 8) and to discriminate patients with E/E' ≥ 8 (suspected diastolic dysfunction) indicating the importance of BNP in prediction of cardiac iron load and its induced diastolic dysfunction before conventional indices of systolic function are affected.

Iron overload is the main determinant of cardiac morbidity in TM patients.^{4,5}

Although not completely reliable, serum ferritin is the widely accepted parameter of iron overload in clinical practice.⁴³ Single ferritin measurement may be misleading and it does not reflect long-term ferritin levels or correlate with cardiac iron levels.⁴⁴ However, the yearly trends in ferritin levels may reflect the direction of body iron loading and long term elevations in ferritin predict cardiac mortality.⁴⁵ By this, the mean yearly serum ferritin during the last year of our enrolled TM patients, had a significant relation to pulsed wave conventional and TDI systolic and diastolic dysfunction indices. It also had significant positive correlation with BNP, the chemical marker of cardiac dysfunction. In addition it was significantly higher among patients with E/E' ratio ≥ 8 than those with this ratio of < 8, finding that was coupled with significant higher RBC s consumption in these patients. However, in some previous studies, no relation was found between diastolic dysfunction and serum ferritin level.^{28,33}

Serum ferritin level of > 2500 ng/ml had been suggested to indicate increased risk of cardiac affection.³ The studied TM patients with serum ferritin > 2500 ng/ml had significantly higher IRT compared to those with serum ferritin level ≤ 2500 ng/ml with a

trend of prevalent E/E' ratio < 8 (normal diastolic function) in those with serum ferritin < 2500 ng/ml (P = 0.052). However, using the ROC curve, serum ferritin level of 4790.5 ng/ml was the cutoff value to predict E/E' ratio ≥ 8. In this regards, El Beshlawy et al.⁴⁶ had reported that there was a low prevalence of myocardial siderosis as measured by cardiovascular magnetic resonance (MRI T2*) in the Egyptian TM patients in spite of very high serum ferritin and high liver iron concentration (LIC) . The authors postulated that, the possibility of a genetic component for the resistance to cardiac iron loading in this population should be considered. This could reflect genetic susceptibility to cardiac iron toxicity in different populations and needs further large-scale studies.

In summary, cardiac dysfunction is a common morbidity among iron-loaded TM patients despite regular chelation therapy. In this study cardiac non-symptomatic TM patients with normal global systolic functions by conventional Echo-Doppler measurements demonstrated abnormal left ventricular systolic and diastolic indices using the pulsed-wave TDI study which were correlated with plasma BNP and serum ferritin. This confirms the importance of this diagnostic modality in TM patients especially in developing countries where wide use of cardiovascular magnetic resonance (MRI T2*) is limited by its expensiveness.²³ BNP cut-off value for prediction of diastolic dysfunction (E/E' ≥ 8) was more or less equal to the previously documented one for prediction of MRI T2* < 24 ms (28.5 pg/ml vs 29 pg/ml) and this raise the concern about the importance of this simple un-expensive test in iron-loaded TM patients.

Conclusions. Asymptomatic TM children under regular chelation therapy may have latent diastolic and or systolic dysfunctions that could not be detected by conventional echocardiography but could be highlighted by TDI. Hence, application of pulsed-wave TDI in these patients is appropriate. Integrated use of echocardiography, pulsed-wave TDI and BNP level for an accurate assessment of cardiac functions is highly recommended to help identifying subjects at risk and facilitates early intervention.

References:

1. Rund D, Rachmilewitz E. Beta-thalassemia. *N Engl J Med.* 2005;353: 1135–1146. <http://dx.doi.org/10.1056/NEJMra050436> PMID:16162884
2. Thein SL. Genetic modifiers of β-thalassemia. *Haematologica .* 2005 May; 90(5):651-660.
3. Olivieri NF, Nathan DG, MacMillan JH, Wayne AS, Liu PP, McGee A, Martin M, Koren G, Cohen AR. Survival in medically treated patients with homozygous beta-Thalassemia. *N Engl J Med.* 1994 Sep 1;331(9):574-578. <http://dx.doi.org/10.1056/NEJM199409013310903> PMID:8047081
4. Walker JM. The heart in thalassaemia. *Eur Heart J.* 2002 23:102–105. <http://dx.doi.org/10.1053/euhj.2001.2850> PMID:11785990
5. Borgna-Pignatti C, Cappellini MD, De Stefano P, Del Vecchio GC, Forni GL, Gamberini MR, Ghilardi R, Piga A, Romeo MA, Zhao H, Cnaan A. Cardiac morbidity and mortality in deferoxamine or deferiprone-treated patients with thalassemia major. *Blood.* 2006 May 1;107(9):3733-3737. <http://dx.doi.org/10.1182/blood-2005-07-2933> PMID:16373663
6. Kremastinos DT, Farmakis D, Aessopos A, Hahalis G, Hamodraka E, Tsiapras D, Keren A. Beta-thalassemia cardiomyopathy: history, present considerations, and future perspectives. *Circ Heart Fail.* 2010 May;3(3):451-458. <http://dx.doi.org/10.1161/CIRCHEARTFAILURE.109.913863> PMID:20484195
7. Pennell DJ . T2* magnetic resonance and myocardial iron in thalassemia. *Ann N Y Acad Sci.* 2005;1054:373-378. <http://dx.doi.org/10.1196/annals.1345.045> PMID:16339685
8. Cohen AR, Galanello R, Pennell DJ, Cunningham MJ, Vichinsky E. Thalassemia. *Hematology Am Soc Hematol Educ Program*

- 2004;14-34.
9. Hahalis G, Manolis AS, Gerasimidou I, Alexopoulos D, Sitafidis G, Kourakli A. Right ventricular diastolic function in beta-thalassaemia major: Echocardiographic and clinical correlates. *Am Heart J*. 2001;141:428-434. <http://dx.doi.org/10.1067/mhj.2001.113077> PMID:11231441
 10. Hamdy AM. Use of strain and tissue velocity imaging for early detection of regional myocardial dysfunction in patients with beta thalassaemia. *Eur J Echocardiogr*. 2007 Mar;8(2):102-109. <http://dx.doi.org/10.1016/j.euje.2006.02.004> PMID:16564231
 11. Mottram PM, Marwick TH. Assessment of diastolic function: what the general cardiologist needs to know. *Heart*. 2005 May;91(5):681-695. <http://dx.doi.org/10.1136/hrt.2003.029413> PMID:15831663 PMCID:PMC1768877
 12. Leonardi B, Margossian R, Colan SD, Powell AJ. Relationship of magnetic resonance imaging estimation of myocardial iron to left ventricular systolic and diastolic function in thalassaemia. *JACC Cardiovasc Imaging*. 2008 Sep;1(5):572-578. <http://dx.doi.org/10.1016/j.jcmg.2008.04.005> PMID:19356483
 13. Yu CM, Sanderson JE, Marwick TH, Oh JK. Tissue Doppler imaging a new prognosticator for cardiovascular diseases. *J Am Coll Cardiol*. 2007 May 15;49(19):1903-1914. <http://dx.doi.org/10.1016/j.jacc.2007.01.078> PMID:17498573
 14. Iarussi D, Di Salvo G, Pergola V, Coppolino P, Tedesco MA, Ratti G, Esposito L, Calabrò R, Ferrara M. Pulsed doppler tissue imaging and myocardial function in thalassaemia major. *Heart Vessels*. 2003 Mar;18(1):1-6. <http://dx.doi.org/10.1007/s003800300000> PMID:12644874
 15. Daniels LB, Maisel AS. Natriuretic Peptides. *J Am Coll Cardiol*. 2007 Dec 18;50(25):2357-2368. <http://dx.doi.org/10.1016/j.jacc.2007.09.021> PMID:18154959
 16. Sagnella GA. Measurement and importance of plasma brainnatriuretic peptide and related peptides. *Ann Clin Biochem*. 2001 Mar;38(Pt 2):83-93. <http://dx.doi.org/10.1258/0004563011900317> PMID:11269760
 17. Kremastinos DT, Tsiapras DP, Kostopoulou AG, Hamodraka ES, Chaidaroglou AS, Kapsali ED. NT-proBNP levels and diastolic dysfunction in β -thalassaemia major patients. *Eur J Heart Fail*. 2007 May;9(5):531-536. <http://dx.doi.org/10.1016/j.ejheart.2006.11.004> PMID:17317307
 18. Teichholz LE, Kreulen T, Herman MV, Gorlin R. Problems in echocardiographic volume determinations: Echocardiographic – angiographic correlations in the presence or absence of asynergy. *Am J Cardiol*. 1976 Jan;37(1):7-11. [http://dx.doi.org/10.1016/0002-9149\(76\)90491-4](http://dx.doi.org/10.1016/0002-9149(76)90491-4)
 19. Ommen SR, Nishimura RA, Appleton CP, Miller FA, Oh JK, Redfield MM, Tajik AJ. Clinical utility of Doppler echocardiography and Tissue Doppler imaging in the estimation of left ventricular filling pressures: a comparative simultaneous Doppler-catheterization study. *Circulation*. 2000 Oct 10;102(15):1788-1794. <http://dx.doi.org/10.1161/01.CIR.102.15.1788> PMID:11023933
 20. Kremastinos DT, Hamodraka E, Parissis J, Tsiapras D, Dima K, Maisel A. Predictive value of B- type natriuretic peptides in detecting latent left ventricular diastolic dysfunction in beta-thalassaemia major. *Am Heart J*. 2010 Jan;159(1):68-74. <http://dx.doi.org/10.1016/j.ahj.2009.10.025> PMID:20102869
 21. Pellikka PA, Nagueh SF, Elhendy AA, Kuehl CA, Sawada SG; American Society of Echocardiography. American Society of Echocardiography recommendations for performance, interpretation, and application of stress echocardiography. *J Am Soc Echocardiogr*. 2007;20:1021-1041. <http://dx.doi.org/10.1016/j.echo.2007.07.003> PMID:17765820
 22. Kremastinos DT, Tsiapras DP, Tsetsos GA, Rentoukas EI, Vretou HP, Toutouzas PK. Left ventricular diastolic Doppler characteristics in beta-thalassaemia major. *Circulation*. 1993 Sep;88(3):1127-1135. <http://dx.doi.org/10.1161/01.CIR.88.3.1127> PMID:8353874
 23. Vogel M, Anderson LJ, Holden S, Deanfield JE, Pennell DJ, Walker JM. Tissue doppler echocardiography in patients with beta thalassaemia detects early myocardial dysfunction related to myocardial iron overload. *Eur Heart J*. 2003 Jan;24(1):113-119. [http://dx.doi.org/10.1016/S0195-668X\(02\)00381-0](http://dx.doi.org/10.1016/S0195-668X(02)00381-0)
 24. Aypar EI, Alehan D, Hazirolan T, Gümriük F. The efficacy of tissue Doppler imaging in predicting myocardial iron load in patients with beta-thalassaemia major: correlation with T2* cardiovascular magnetic resonance. *Int J Cardiovasc Imaging*. 2010 Apr;26(4):413-429. <http://dx.doi.org/10.1007/s10554-010-9591-6> PMID:20127175
 25. Uçar T, Ileri T, Atalay S, Uysal Z, Tutar E, Ertem M. Early detection of myocardial dysfunction in children with beta-thalassaemia major. *Int J Cardiovasc Imaging*. 2009 Apr;25(4):379-386. <http://dx.doi.org/10.1007/s10554-008-9404-3> PMID:19107572
 26. Ehlers KH, Levin AR, Markenson AL, Marcus JR, Klein AA, Hilgartner MW, Engle MA. Longitudinal study of cardiac function in thalassaemia major. *Ann N Y Acad Sci*. 1980;344:397-404. <http://dx.doi.org/10.1111/j.1749-6632.1980.tb33678.x> PMID:6930879
 27. Parale GP, Pawar SS, Tapare VS. Assessment of LV diastolic function in patients with beta-thalassaemia major with special reference to E/E ann ratio. *J Pediatr Hematol Oncol*. 2009 Jan;31(1):69-73. <http://dx.doi.org/10.1097/MPH.0b013e31818ab138> PMID:19125094
 28. Abdelmuktader AM, Azer HY. Usefulness of pulsed wave tissue doppler imaging in assessment of left ventricular functions in children with beta-thalassaemia major. *Indian J Pediatr*. 2013 Sep;80(9):721-725. <http://dx.doi.org/10.1007/s12098-013-1020-0> PMID:23604609
 29. Noori NM, Mehralzadeh S. Echocardiographic evaluation of systolic and diastolic heart function in patients suffering from beta-thalassaemia major aged 5-10 years at the Zahedan Research Center for Children and Adolescent Health. *Anatolian journal of cardiology*. 2010 Apr;10(2):150-153.
 30. Nishimura RA, Tajik AJ. Evaluation of diastolic filling of left ventricle in health and disease: Doppler echocardiography is the clinician's Rosetta Stone. *J Am Coll Cardiol*. 1997 Jul;30(1):8-18. [http://dx.doi.org/10.1016/S0735-1097\(97\)00144-7](http://dx.doi.org/10.1016/S0735-1097(97)00144-7)
 31. Agha HM, Beshlawy A, Hamdy M, Sobehi A, El Zahrae F, Abd El Satar IA, AbdelMassih A, Said F, Abd El Aziz O, El Tagui M, Pennell DJ. Early detection of right ventricular diastolic dysfunction by pulsed tissue Doppler echocardiography in iron loaded beta thalassaemia patients. *Pediatr Cardiol*. 2015 Mar;36(3):468-474. <http://dx.doi.org/10.1007/s00246-014-1035-y> PMID:25293426
 32. Seliem MA, Al-Saad HI, Bou-Holaigah IH, Khan MN, Palileo MR. Left ventricular diastolic dysfunction in congenital chronic anaemias during childhood as determined by comprehensive echocardiographic imaging including acoustic quantification. *Eur J Echocardiogr*. 2002 Jun;3(2):103-110. <http://dx.doi.org/10.1053/euje.2001.0122> PMID:12114094
 33. Garadah T, Kassab S, Mahdi N, Abu-Taleb A and Jamsheer A. Pulsed and Tissue Doppler Echocardiographic Changes in Patients with Thalassaemia Major. *Clinical Medicine Insights: Cardiology* 2010;3 :1-8.
 34. Kremastinos DT. Heart failure in beta-thalassaemia. *Congest Heart Fail*. 2001;7:312-314. <http://dx.doi.org/10.1111/j.1527-5299.2001.00259.x> PMID:11828176
 35. Vaccari M, Crepaz R, Fortini M, Gamberini MR, Scarcia S, Pitscheider W, Bosi G. Left ventricular remodeling, systolic function, and diastolic function in young adults with beta thalassaemia intermedia. *Chest*. 2002 Feb;121(2):506-512. <http://dx.doi.org/10.1378/chest.121.2.506> PMID:11834665
 36. Fitchett DH, Coltart DJ, Littler WA, Leyland MJ, Trueman T, Gozzard DI, Peters TJ. Cardiac involvement in secondary haemochromatosis: a catheter biopsy study and analysis of myocardium. *Cardiovasc Res*. 1980 Dec;14(12):719-724. <http://dx.doi.org/10.1093/cvr/14.12.719> PMID:7260965
 37. Lattanzi F, Bellotti P, Picano E, Chiarella F, Mazzarisi A, Melevendi C, Forni G, Landini L, Distanto A, Vecchio C. Quantitative ultrasonic analysis of myocardium in patients with thalassaemia major and iron overload. *Circulation*. 1993 Mar;87(3):748-754. <http://dx.doi.org/10.1161/01.CIR.87.3.748> PMID:8443895
 38. Sohn DW, Chai IH, Lee DJ, Kim HC, Kim HS, Oh BH, Lee MM, Park YB, Choi YS, Seo JD, Lee YW. Assessment of mitral annulus velocity by Doppler tissue imaging in evaluation of left ventricular diastolic function. *J Am Coll Cardiol*. 1997 Aug;30(2):474-480. [http://dx.doi.org/10.1016/S0735-1097\(97\)88335-0](http://dx.doi.org/10.1016/S0735-1097(97)88335-0)
 39. Gharzuddine WS1, Kazma HK, Nuwayhid IA, Bitar FF, Koussa SF, Moukarbel GV, Taher AT. Doppler characterization of left ventricular diastolic function in beta-thalassaemia major. Evidence for an early stage of impaired relaxation. *Eur J Echocardiogr*. 2002 Mar;3(1):47-51. <http://dx.doi.org/10.1053/euje.2001.0114> PMID:12067534
 40. Nagueh SF, Lakkis NM, Middleton KJ, Spencer WH 3rd, Zoghbi WA, Qui-ones MA. Doppler estimation of left ventricular filling pressures in patients with hypertrophic cardiomyopathy.

- Circulation. 1999 Jan 19;99(2):254-261.
<http://dx.doi.org/10.1161/01.CIR.99.2.254> PMID:9892592
41. Chrysohoou C, Greenberg M, Pitsavos C, Panagiotakos DB, Ladis V, Barbetseas J, Brili S, Singh S, Stefanadis C. Diastolic function in young patients with beta-thalassemia major: an echocardiographic study. *Echocardiography*. 2006 Jan;23(1):38-44.
<http://dx.doi.org/10.1111/j.1540-8175.2006.00148.x>
PMid:16412181
42. Nikolidakis S, Flessa C, Nikolaou N, Gotsis ED, Koutouzis M, Polyhronaki E, Vrettou E, Kyriakides ZS. Brain natriuretic peptide as marker of myocardial iron load in beta-thalassemia. *Int J Cardiol*. 2007 Jun 12;118(3):408-409.
<http://dx.doi.org/10.1016/j.ijcard.2006.07.052> PMID:17045671
43. Lo L, Singer ST. Thalassemia; current approach to an old disease. *Pediatr Clin North Am*. 2002 Dec;49(6):1165-91, v.
[http://dx.doi.org/10.1016/S0031-3955\(02\)00088-3](http://dx.doi.org/10.1016/S0031-3955(02)00088-3)
44. Anderson LJ1, Holden S, Davis B, Prescott E, Charrier CC, Bunce NH, Firmin DN, Wonke B, Porter J, Walker JM, Pennell DJ. Cardiovascular T2 star (T2*) magnetic resonance for the early diagnosis of myocardial iron overload. *Eur Heart J*. 2001 Dec;22(23):2171-2179. <http://dx.doi.org/10.1053/euhj.2001.2822>
PMid:11913479
45. Pennell DJ, Udelson JE, Arai AE, Bozkurt B, Cohen AR, Galanello R, Hoffman TM, Kiernan MS, Lerakis S, Piga A, Porter JB, Walker JM, Wood J; American Heart Association Committee on Heart Failure and Transplantation of the Council on Clinical Cardiology and Council on Cardiovascular Radiology and Imaging. Cardiovascular function and treatment in b-Thalassemia major: a consensus statement from the american heart association. *Circulation*. 2013 Jul 16;128(3):281-308.
<http://dx.doi.org/10.1161/CIR.0b013e31829b2be6>
PMid:23775258
46. El Beshlawy A, El Tagui M, Hamdy M, El Ghamrawy M, Azim KA, Salem D, Said F, Samir A, St Pierre T, Pennell DJ. Low prevalence of cardiac siderosis in heavily iron loaded Egyptian thalassemia major patients. *Ann Hematol*. 2014 Mar;93(3):375-379.
<http://dx.doi.org/10.1007/s00277-013-1876-0>
PMid:23949317