



Draft Genome Sequence of the Feruloyl Esterase-Producing Strain *Lactobacillus fermentum* CRL1446, a Probiotic for Malnutrition

María C. Abeijón Mukdsi,^a Lucila Saavedra,^a María P. Gauffin Cano,^a  Elvira M. Hebert,^a  Roxana B. Medina^a

^aCentro de Referencia para Lactobacilos (CERELA)–CONICET, San Miguel de Tucumán, Tucumán, Argentina

ABSTRACT We report here the draft genome sequence of *Lactobacillus fermentum* CRL1446 (2,148,781 bp, 51.4% G+C content). This strain exhibits feruloyl esterase activity and important technological and probiotic properties. Because of its proven beneficial effects *in vivo*, it represents an interesting candidate for the development of functional foods or pharmabiotics for malnutrition.

Lactobacillus fermentum CRL1446 is a strain isolated from an artisanal goat milk cheese produced in northwest Argentina. It presents optimal technological properties for the development of fermented dairy products, mainly cheeses, due to its capacity to generate flavor compounds (1). Furthermore, this strain displays probiotic properties and feruloyl esterase (FE) activity, which is capable of releasing antioxidant ferulic acid from its nonbioavailable esterified forms naturally present in vegetable foods (2). Free ferulic acid can be absorbed in the gut and can exert several health benefits on the host (3).

The administration of *L. fermentum* CRL1446 to mice subjected to normal and malnutrition diets (caloric restriction and high-fat diets) increased intestinal FE activity and enhanced oxidative status. Moreover, it reduced glycemia, cholesterolemia, and triglyceridemia and positively modulated intestinal microbiota, as evidenced by a marked bifidogenic effect (4–7). This noninflammatory strain, which is able to modulate the profile of adipokines *in vitro* (8), also reduced body weight gain and leptin levels in diet-induced obese mice (7). Due to these functional properties, *L. fermentum* CRL1446 could be used as a probiotic for reverting or improving alterations related to malnutrition.

Here, we present a draft genome sequence of *L. fermentum* CRL1446 that was sequenced using a whole-genome shotgun strategy on an Illumina HiSeq 2500 sequencer. Paired reads with lengths of 250 bp were obtained, corresponding to a 670-fold coverage. Quality-filtered reads were assembled using the NGEN assembler (DNASTar version 12.2.0), resulting in 56 contigs. The Rapid Annotations using Subsystems Technology (RAST) server and the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) were used for functional annotation of predicted genes (9).

The 2,148,781-bp genome is composed of a single circular chromosome with a G+C content of 51.4%. The chromosome contains 2,187 coding sequences and 74 RNA genes, as predicted by PGAP (15 rRNAs, 56 tRNAs, and 3 noncoding RNA genes). There are 316 subsystems represented in the chromosome, which comprise only 49% of the sequences assigned. According to the RAST analysis, 1,599 protein-encoding genes were assigned to known functions.

In silico studies revealed that *L. fermentum* CRL1446 carries eight genes encoding proteins predicted as esterases/lipases (E.C.3.1.1.x), which belong to the α/β fold hydrolase family and display the classical serine nucleophilic motif (GxSxG) described for some carboxyl esterases (10). One of these predicted proteins (GenBank accession

Received 20 March 2018 Accepted 24 March 2018 Published 24 May 2018

Citation Abeijón Mukdsi MC, Saavedra L, Gauffin Cano MP, Hebert EM, Medina RB. 2018. Draft genome sequence of the feruloyl esterase-producing strain *Lactobacillus fermentum* CRL1446, a probiotic for malnutrition. *Genome Announc* 6:e00225-18. <https://doi.org/10.1128/genomeA.00225-18>.

Copyright © 2018 Abeijón Mukdsi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to María C. Abeijón Mukdsi, cabeijon@cerela.org.ar, or Roxana B. Medina, rmedina@cerela.org.ar.

number PNV58462) shares 100% identity with a FE from *L. fermentum* SD1210 (GenBank accession number AOR52356).

Further in-depth analysis of the *L. fermentum* CRL1446 genome will help to exploit the biotechnological and probiotic potential of this strain and elucidate the mechanisms underlying the beneficial effects observed *in vivo*. This information could support the use of *L. fermentum* CRL1446 for developing novel functional foods, dietary supplements, or pharmabiotics directed to malnutrition and other oxidative stress-related ailments.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [POTQ00000000](https://www.ncbi.nlm.nih.gov/nuclink/POTQ00000000). The version described in this paper is the first version, POTQ01000000.

ACKNOWLEDGMENTS

This work was supported by grants from CONICET (PIP 215) and ANPCyT (PICT 2015-2916).

REFERENCES

1. Abeijón Mukdsi MC, Maillard MB, Medina RB, Thierry A. 2018. Ethyl butanoate is synthesised both by alcoholysis and esterification by dairy lactobacilli and propionibacteria. *LWT-Food Sci Technol* 89:38–43. <https://doi.org/10.1016/j.lwt.2017.10.012>.
2. Abeijón Mukdsi MC, Argañaraz Martínez E, Perez Chaia A, Medina R. 2016. Feruloyl esterase activity is influenced by bile, probiotic intestinal adhesion and milk fat. *Benef Microbes* 7:597–607. <https://doi.org/10.3920/BM2015.0197>.
3. Palani Swamy SK, Govindaswamy V. 2015. Therapeutical properties of ferulic acid and bioavailability enhancement through feruloyl esterase. *J Funct Foods* 17:657–666. <https://doi.org/10.1016/j.jff.2015.06.013>.
4. Abeijón Mukdsi MC, Gauffin Cano MP, González SN, Medina RB. 2012. Administration of *Lactobacillus fermentum* CRL1446 increases intestinal feruloyl esterase activity in mice. *Lett Appl Microbiol* 54:18–25. <https://doi.org/10.1111/j.1472-765X.2011.03166.x>.
5. Abeijón Mukdsi MC, Haro C, González SN, Medina RB. 2013. Functional goat milk cheese with feruloyl esterase activity. *J Funct Foods* 5:801–809. <https://doi.org/10.1016/j.jff.2013.01.026>.
6. Russo M, Fabersani E, Abeijón Mukdsi MC, Ross R, Fontana C, Benítez Páez A, Gauffin Cano P, Medina RB. 2016. *Lactobacillus fermentum* CRL1446 ameliorates oxidative and metabolic parameters by increasing intestinal feruloyl esterase activity and modulating microbiota in caloric-restricted mice. *Nutrients* 8:415. <https://doi.org/10.3390/nu8070415>.
7. Fabersani Marrades ME. 2016. PhD thesis. National University of Tucumán, San Miguel de Tucumán, Argentina.
8. Fabersani E, Abeijón-Mukdsi MC, Ross R, Medina R, González S, Gauffin-Cano P. 2017. Specific strains of lactic acid bacteria differentially modulate the profile of adipokines *in vitro*. *Front Immunol* 8:266. <https://doi.org/10.3389/fimmu.2017.00266>.
9. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
10. Bornscheuer UT. 2002. Microbial carboxyl esterases: classification, properties and application in biocatalysis. *FEMS Microbiol Rev* 26:73–81. <https://doi.org/10.1111/j.1574-6976.2002.tb00599.x>.