

## Anesthesiologists and the High Risk of Exposure to COVID-19

### To the Editor

We read the study by Tompkins and Kerchberger,<sup>1</sup> personal protective equipment (PPE) for Care of Pandemic Influenza Patients: A Training Workshop for the Powered Air-Purifying Respirators (PAPR), with great interest. Since the 2009 Pandemic Influenza A (H1N1 subtype) pandemic of the influenza virus, the Centers for Disease Control and Prevention (CDC) guidelines state that health care providers (HCP), when performing or participating in procedures that are associated with pathogen generating aerosols (aerosol-generating procedures [AGPs]), should wear an N95 filter mask and consider PAPR. AGPs include performance of the following procedures in patients with acute respiratory diseases<sup>2</sup>: intubation and related procedures, cardiopulmonary resuscitation, bronchoscopies, surgeries, and autopsies.

Regarding preventive measures for the Coronavirus Disease 2019 (COVID-19) virus in Mexico, the official norms NOM-017-STPS-2008 for PPE selection for health care personnel, highlight 2 important considerations: (1) PPE are all the set of elements and devices, that are specifically designed to protect the HCP against accidents and illnesses that could be caused by agents during the performance of their daily basis activities as well as in the emergency care; and (2) the occupational risk related to the exposure of the HCP must be identified and analyzed.<sup>3</sup>

The Joint Commission International (JCI) emphasizes that current status of PPE supplies remains inadequate to achieve minimum standards in most hospitals. The JCI has been calling for action at all government levels to address the shortage and protect those who work heroically to care for infected patients with COVID-19. We can confirm that in Mexico, there

have been “hospital outbreaks” with 329 HCPs who have been infected with COVID-19 throughout the country. Planning an adequate distribution of PPE to health workers and developing appropriate strategies in clinics can diminish the impact of this pandemic on HCPs.

**Nayely García-Méndez, PhD**

Department of Hemodynamic  
General Hospital of the National Medical Center “La Raza”  
Mexican Institute of Social Security (IMSS)  
Mexico City, Mexico  
PhD Program in Medical Sciences  
Universidad de La Frontera  
Temuco, Chile

ayeyigmendez@comunidad.unam.mx

**Juan Lagarda Cuevas, MD**

Department of Anesthesia  
New York University  
American British Cowdray  
Medical Center Mexico City  
Mexico City, Mexico

**Tamara Otzen, PhD**

**Carlos Manterola, PhD**  
Center of Excellence in Morphological  
and Surgical Studies (CEMyQ)  
Universidad de La Frontera  
Temuco, Chile

### REFERENCES

1. Tompkins BM, Kerchberger JP. Personal protective equipment for care of pandemic powered air purifying respirator. *Anes Analg*. 2010;111:933–945.
2. Organización Mundial de la Salud. Prevención y control de infección en enfermedades respiratorias agudas con tendencia epidémica y pandémica durante la atención sanitaria Pautas. Available at: [www.paho.org/es/documentos/prevencion-control-infeccion-enfermedades-respiratorias-agudas-con-tendencia-epidemica](http://www.paho.org/es/documentos/prevencion-control-infeccion-enfermedades-respiratorias-agudas-con-tendencia-epidemica). Accessed April 9, 2020.
3. Norma Oficial Mexicana NOM-017-SSA2-2012, Para la vigilancia epidemiológica. Available at: <https://www.gob.mx/stps/documentos/norma-oficial-mexicana-nom-017-stps-2008-equipo-de-proteccion-personal-seleccion-uso-y-manejo-en-los-centros-de-trabajo>. Accessed April 9, 2020.

DOI: 10.1213/ANE.0000000000004919