



Pericyte Contractile Responses to Endothelin-1 and A β Peptides: Assessment by Electrical Impedance Assay

Elliott Hibbs*, Seth Love and J. Scott Miners

Dementia Research Group, Clinical Neurosciences, University of Bristol, Bristol, United Kingdom

Pericytes are vascular mural cells that contract and relax in response to vasoactive stimuli to regulate neurovascular coupling and cerebral blood flow. Pericytes are damaged and degenerate in Alzheimer's disease (AD). We previously showed that the level of the regulatory vasoconstrictor, endothelin-1 (EDN1), is elevated in AD cerebral cortex and upregulated by amyloid-beta (A β). We have used electrical impedance analysis to monitor the contractile and proliferative response of cultured human fetal and adult brain-derived pericytes to EDN1 in real-time. EDN1 caused transient, dose-dependent contraction of fetal and adult brain pericytes that was mediated by EDN1 type A receptors and increased the subsequent proliferation of fetal but not adult cells. The contractile responses to EDN1 were weaker in the adult pericytes. The EDN1-mediated contractile response of fetal pericytes was unchanged after exposure to A β_{1-40} or A β_{1-42} (0.1–10 μ M) for 1 h but both contraction and subsequent relaxation were significantly impaired upon exposure to A β for 24 h. These data suggest that chronic exposure to A β interferes with EDN1-mediated pericyte contractility, potentially contributing to neurovascular uncoupling and reduced cerebral blood flow in AD.

OPEN ACCESS

Edited by:

Robert Lindsay Medcalf,
Monash University, Australia

Reviewed by:

Brad A. Sutherland,
University of Tasmania, Australia
Daniel Beard,
The University of Newcastle, Australia

*Correspondence:

Elliott Hibbs
eh15249@bristol.ac.uk

Keywords: A β peptide, Alzheimer's disease, electrical impedance, endothelin-1, pericyte

INTRODUCTION

Pericytes are vascular mural cells that are particularly abundant within the brain. They are expressed within the capillary network but also reside on pre- and post-capillary vessels. Pericytes are critical components of the neurovascular unit and dynamically regulate cerebral blood flow (CBF) in response to local metabolic demand (Joyce et al., 1985; Neuhaus et al., 2017; Nortley et al., 2019). In stroke, pericytes die in rigor, clamping shut capillary lumina, contributing to ischemic injury (Hall et al., 2014; Heyba et al., 2019). Recent studies indicate that pericytes are damaged and degenerate in Alzheimer's disease (AD) and vascular dementia (VaD) (Sagare et al., 2013; Kisler et al., 2017; Ding et al., 2020). This is thought to play a major role in blood-brain barrier (BBB) breakdown and cerebral hypoperfusion, and to affect clearance of A β in AD (Daneman et al., 2010).

Brain ischemia has long been recognized as the principal pathological process in VaD but recent studies have shown that cerebral hypoperfusion and BBB leakiness are early and substantial contributors to disease progression and cognitive decline in AD (Ding et al., 2020). We previously showed that the enzymes responsible for synthesis of the vasoconstrictor peptide endothelin-1 (EDN1) were upregulated in AD (Palmer et al., 2009, 2013) and associated with elevated EDN1

Specialty section:

This article was submitted to
Non-Neuronal Cells,
a section of the journal
Frontiers in Cellular Neuroscience

Received: 15 June 2021

Accepted: 28 July 2021

Published: 20 August 2021

Citation:

Hibbs E, Love S and Miners JS
(2021) Pericyte Contractile
Responses to Endothelin-1 and A β
Peptides: Assessment by Electrical
Impedance Assay.
Front. Cell. Neurosci. 15:723953.
doi: 10.3389/fncel.2021.723953

level in the cerebral cortex. The level of EDN1 correlated strongly with biochemical evidence of cerebral hypoperfusion in AD and with the level of A β , which we found to increase the production of EDN1 by neurons and endothelial cells *in vitro* (Palmer et al., 2012). EDN1 induces contraction of smooth muscle cells, causing constriction of penetrating arteries and arterioles and reducing blood flow in the cerebral cortex and deep within the brain (Wynne et al., 2009). A recent study confirmed that capillaries were constricted in proximity to A β plaques in human biopsy brain tissue and that mechanistically, reactive oxygen species in response to oligomeric A β mediated EDN1-induced pericyte contraction and consequent narrowing of capillaries in brain cortical slices, and was likely to contribute to cerebral hypoperfusion in AD (Nortley et al., 2019).

In vitro studies of pericyte function have been performed with bovine retinal pericytes (Yamagishi et al., 1993) using semi-quantitative methods, such as the “wrinkling assay” [pericyte contraction causes folds when the cells are grown on a silicone membrane (Durham et al., 2014)], and with brain-derived human pericytes isolated from fetal brain tissue (ScienCell, United States) (Persidsky et al., 2016; Neuhaus et al., 2017). The fetal cells express pericyte markers such as platelet-derived growth factor receptor- β (PDGFR β) and chondroitin sulfate proteoglycan-4 (NG2) and are negative for endothelial markers and non-vascular markers including GFAP and IBA1 (Hall et al., 2014). The cells respond to vasoactive metabolites and contract and migrate in culture (Neuhaus et al., 2017; Nortley et al., 2019; Kemp Scott et al., 2020). They have been shown to express α -smooth muscle actin (α -SMA) (Nehls and Drenckhahn, 1991).

Fetal pericytes are responsive to multiple vasoactive peptides in the brain, and contract in response to EDN1 as previously shown using an electrical impedance assay (Neuhaus et al., 2017). It remains unclear whether the pathophysiological responses of fetal brain pericytes reflect those in the adult brain. EDN1 also evokes pericyte-mediated constriction of capillaries in human and rat cortical slices in response to oligomeric A β (Nortley et al., 2019). In the present study, we have assessed EDN1-mediated contraction of fetal and adult brain-derived pericytes by use of an electrical impedance assay and investigated whether the contractile response is modified by exposure of pericytes to A β peptides.

MATERIALS AND METHODS

Pericyte Culture

Human brain vascular pericytes of fetal origin (fHBVP; ScienCell, United States) and human brain vascular pericytes of adult origin (aHBVP; Cell Systems, United States) were grown in pericyte medium (ScienCell, United States) at 37°C in 5% CO₂. All experiments were performed using cells at passages of <8. Cells were cultured in poly-L-lysine-coated flasks (10 mg/mL) until 80% confluent and then sub-cultured following the manufacturer's protocol. All work was performed in a Class II microbiological safety cabinet (Holten, United Kingdom) under sterile conditions.

Immunofluorescent Labeling of Pericytes

Immunofluorescence was used to assess the expression and distribution of cell-specific markers. Cells were cultured on poly-L-lysine coated coverslips and left to grow for 24 h at 37°C in a 5% CO₂ incubator. The medium was aspirated, and cells were washed gently three times with PBS before being fixed in 4% paraformaldehyde (PFA; Sigma-Aldrich, United States) for 10 min at 37°C in a 5% CO₂ incubator. PFA was removed and cells were washed three more times in PBS and left in PBS at 4°C until immunolabelled. At this point, PBS was removed, cold methanol added, and the cells incubated at -20°C for 10 min, washed with PBS 3 times for 1 min each and then blocked with 5% donkey serum (Sigma-Aldrich, United States) for 30 min at room temperature (RT). The cells were incubated with primary antibody overnight at 4°C: α -SMA (1:100, Abcam United Kingdom, rabbit polyclonal), PDGFR β (1:500, R&D Systems United States, biotinylated goat polyclonal), EDNRA (1:500, Abcam United Kingdom, rabbit polyclonal), and EDNRB (1:1000, Abcam United Kingdom, rabbit polyclonal). The cells were washed again in PBS three times for 1 min each followed by addition of a donkey anti-rabbit IgG Alexa Fluor 488 (Invitrogen, United Kingdom) secondary antibody (1:500) for α -SMA and streptavidin Alexa Fluor 488 conjugate (Invitrogen, United Kingdom) for PDGFR β for 1 h at RT in the dark. Samples were washed in PBS three times for 1 min each and coverslips were placed onto glass microscope slides and mounted using Vectashield mounting medium with DAPI (Vector Labs, United States). Slides were stored at 4°C in the dark before being imaged by confocal microscopy (Nikon Eclipse 80i).

EDN1-Mediated Contraction of Pericytes Assessed Using an Electrical Impedance Assay (xCELLigence)

Pericytes (fHBVP and aHBVP) were seeded at 5000 cells/well onto pre-coated poly-L-lysine impedance plates (E-plate VIEW[®] 96 or E Plate insert 16[®], ACEA Biosciences, United States) and allowed to adhere for 1 h at RT within the hood and then overnight in the incubator at 37°C, 5% CO₂. Cells were then starved in serum-free medium for 2 h. EDN1 (Abcam, United Kingdom) was added *in situ* and the cell index (see below) was measured using either the xCELLigence[®] RTCA S16 or SP96-well plates (ACEA Biosciences, United States) maintained at 37°C in a 5% CO₂ incubator. The cell index was recorded in real-time throughout the duration of the experiment. Addition of serum-free medium was used as a vehicle control. The relative impedance is presented as a “cell index” (an arbitrary value derived from the impedance software) which reflects the contact surface area between cells and the culture dish and the resistance to electrical current (see **Supplementary Figure 1**); the average normalized cell index for a particular well is generated by normalizing the raw cell index data with reference to the mean index across duplicate vehicle control wells. Upon addition of EDN1, parameters for the xCELLigence software were set to record in 5 s intervals for a total duration of 2 h (contraction-relaxation phase) and then 1 h intervals for a further 48 h (proliferation phase). Data analyses were performed using RTCA

Data Analysis Software 1.0 (ACEA Biosciences, United States). The change in slope (1/h) of the cell index curve, to reflect the rate of responses, was calculated using the built-in function of the data analysis software. Measurements were recorded in duplicate wells for each condition; “*n*” refers to the number of independent experimental repeats.

To assess the effect of receptor blockers on EDN1-mediated contraction, BQ123 (Tocris, United Kingdom) and BQ788 (Sigma-Aldrich, United States) were added in serum-free medium, 1 h before the addition of EDN1.

To assess the effect of A β _{1–40} (rPeptide, United States) and A β _{1–42} (rPeptide, United States) on EDN1-mediated contraction, freshly solubilized A β peptides prepared in 35% acetonitrile and diluted in serum-free medium were added either 1 or 24 h prior to the addition of EDN1.

Statistical Software

All statistical analyses were performed using GraphPad Prism 8.1.2 software package. Unpaired two-tailed *t*-tests or ANOVA with Dunnett's multiple comparisons *post hoc* tests were selected and used for comparisons between groups as appropriate. *P*-values of <0.05 were considered statistically significant.

RESULTS

Pericytes Isolated From Fetal and Adult Human Brain Express Pericyte Markers

fHBVP (ScienCell, United States) and aHBVP (Cell-Systems, United States) expressed the canonical pericyte marker PDGFR β (Figures 1A,B). Pericytes expressed α -SMA (Figures 1C,D) and both endothelin-1 type A (EDNRA) (Figures 1E,F) and type B (EDNRB) receptors (Figures 1G,H).

EDN1-Induced Contraction and Proliferation of Fetal Brain-Derived Pericytes

Electrical impedance assays (xCELLigence) were used to assess pericyte contraction, relaxation, and proliferation in response to EDN1, as previously reported (Neuhaus et al., 2017). Immediately after addition of EDN1, pericyte contraction caused a rapid decrease in cell index (relative impedance), followed by a relaxation phase (~10 min) – a classic G-protein coupled receptor response – and subsequent proliferation, leading to a progressive rise in cell index over 48–72 h (Figure 2).

The rate of contraction of fHBVPs increased with the concentration of EDN1 (1 pM–1 μ M) as illustrated in Figure 2A. The initial rapid decrease in cell index over control values (cells in pericyte medium only) was significant after exposure to 1 nM (*p* = 0.0039), 0.01 μ M (*p* = 0.0003), 0.1 μ M (*p* < 0.0001), or 1 μ M (*p* < 0.0001) EDN1 (*n* = 12) (Figure 2B). Contraction was followed by an increase in cell index (i.e., relaxation) as illustrated in Figure 2C. The rate of increase was related to the previous rate of decrease and was significantly higher in cells that had been exposed to 0.1 nM (*p* = 0.0085), 1 nM (*p* < 0.0001), 0.01 μ M

(*p* < 0.0001), 0.1 μ M (*p* < 0.0001), or 1 μ M (*p* < 0.0001) EDN1 (*n* = 12) (Figure 2D).

To monitor the effects of EDN1 on proliferation, the impedance was monitored for a further 48–72 h (Figure 2E), during which impedance rose reflecting an increase in the number of cells covering the plate. The rate of increase was calculated and indicated that proliferation of fHBVP was significantly greater than in untreated control cells when exposed to 0.1 nM (*p* = 0.0290), 1 nM (*p* = 0.0008), 0.01 μ M (*p* = 0.0038), 0.1 μ M (*p* = 0.0041), or 1 μ M (*p* = 0.0004) EDN1 (*n* = 12) (Figure 2F). We similarly showed proliferation of fHBVPs in response to EDN1 using a standardized BRDU assay (Supplementary Figure 2).

Pretreatment with BQ123, an EDNRA antagonist, inhibited EDN1-mediated pericyte contraction in a dose-dependent manner (Figure 3A). A significant reduction in the rate of contraction induced by EDN1 (100 nM) was observed in cells pre-treated with BQ123 at 1 μ M (*p* = 0.0158) or 10 μ M (*p* = 0.0001) (*n* = 7) (Figure 3B). Pre-incubation with BQ788, an EDNRB receptor antagonist, did not significantly alter EDN1-induced pericyte contraction (*n* = 7) (Figures 3C,D).

Adult Brain-Derived Pericytes Contract in Response to EDN1 via EDNRA but Do Not Proliferate


EDN1 induced contraction of aHBVP. The rate of initial decline of cell index (i.e., contraction) was dose-related (1 pM–1 μ M), as illustrated in Figure 4A. The rate of contraction was significantly higher in adult cells treated with 0.1 μ M (*p* = 0.0008) or 1 μ M (*p* < 0.0001) EDN1 than in untreated cells (*n* = 4) (Figure 4B). The initial contraction was followed by a slower increase in cell index (i.e., relaxation), as shown in Figure 4C; the increase in cell index was most rapid after 0.1 μ M EDN1 exposure (*p* = 0.0159) (*n* = 4) (Figure 4D).

Impedance was monitored for a further 48 h to assess the effect of EDN1 on adult pericyte proliferation. The cell index increased slowly (Figure 4E) and was not affected by exposure to EDN1 (*n* = 4) (Figure 4F).

Pre-treatment with BQ123 inhibited EDN1-mediated contraction of aHBVP (Figure 5A). A significant reduction in the rate of contraction induced by EDN1 (100 nM) was seen in pericytes pre-treated with 0.1 μ M (*p* = 0.0154), 1 μ M (*p* = 0.0241), or 10 μ M (*p* = 0.0016) BQ123 compared to cells treated with EDN1 alone (*n* = 7) (Figure 5B). Pre-treatment with BQ788 did not significantly influence EDN1-induced adult pericyte contraction (Figures 5C,D).

EDN1-Induced Contractile Response of Fetal Pericytes Was Impaired by Exposure to A β Peptides

To investigate the acute effects of A β on EDN1-induced contraction of fHBVP we pre-incubated cells with A β _{1–40} or A β _{1–42} (10–0.1 μ M) 1 h before the addition of EDN1 (100 nM). Neither contraction nor relaxation was significantly altered with exposure to A β _{1–40} or A β _{1–42} (Figures 6A–H).



We next determined if longer exposure to A β peptides for 24 h impacted fHBVP contractility. Pre-treatment with 1 μ M A β_{1-40} for 24 h significantly impaired EDN1-induced pericyte contraction ($p = 0.0249$) (**Figures 7A,B**). The subsequent increase in cell index (reflecting relaxation) was also significantly slower in cells treated with 1 μ M A β_{1-40} ($p = 0.0409$) (**Figures 7C,D**). The contractile response to EDN1 tended also to be lower after pre-treatment with A β_{1-42} but the difference was not statistically significant ($n = 6$) (**Figures 7E-H**).

Exposure to A β_{1-40} and A β_{1-42} for 24 h at concentrations above 1 μ M caused pericyte death, indicated by the incorporation of ethidium-1 dye. Pericyte death was minimal at A β_{1-40} and A β_{1-42} levels below 1 μ M (**Supplementary Figure 3**).


It was not possible to assess the effects of prolonged exposure of aHBVP to EDN1 because of deteriorating viability and responsiveness of the adult cells after 24 h exposure to A β_{1-40} or A β_{1-42} .

DISCUSSION

In this study, we have used electrical impedance assays (xCELLigence) to characterize the contractile and proliferative responses of fHBVP and aHBVP to EDN1. EDN1-induced contraction of fHBVP and aHBVP was mediated specifically by activation of EDNRA. The contractile responses were stronger in the fetal than adult cells, and the former proliferated after EDN1-induced contraction whereas the latter did not. Exposure to


A β_{1-40} for 24 h significantly abrogated the contractile response to EDN1 in fHBVP, and subsequent relaxation. In the absence of EDN1, neither A β_{1-40} nor A β_{1-42} induced cell contraction.

Pericytes respond to vasoactive stimuli; ATP (Hørlyck et al., 2021) and EDN1 (Chakravarthy et al., 1992) induce pericyte contraction whereas adenosine and nitric oxide cause pericyte relaxation (Neuhaus et al., 2017). In human and rat cortical slices, noradrenaline constricts, and glutamate dilates capillaries in proximity to pericyte cell bodies (Peppiatt et al., 2006; Hall et al., 2014; Nortley et al., 2019). EDN1 was previously reported to cause pericyte contraction of bovine retinal pericytes (Chakravarthy et al., 1992), associated with the release of cytosolic Ca²⁺ and reorganization of actin filaments (Dehouck et al., 1997). More recently, EDN1 was shown to induce strong pericyte-mediated contraction reducing the lumina of capillaries in rat cortical slices – an effect mediated by the EDNRA receptor (Neuhaus et al., 2017; Nortley et al., 2019). Neuhaus et al. (2017) demonstrated the utility of xCELLigence to first measure contraction, relaxation and proliferation of human fetal brain-derived pericytes in response to EDN1 and various vascular stimuli. We have confirmed that EDN1 induces contraction of fHBVP, across a similar dose range to that previously reported (1 nM–1 μ M), reflecting the normal concentration of EDN1 in human plasma (3.55 ± 1.78 μ g/ml) (Kostov et al., 2016). The contractile response was dose-dependent and was typical of a classical G-protein coupled response (Billington and Penn, 2003). We have extended the previous findings and demonstrated that both fHBVP and aHBVP are responsive to EDN1, and that



the contraction is mediated through activation of the EDNRA receptor. Exposure to EDN1 alone does not cause pericyte toxicity, with longer exposure causing cell proliferation between 48 and 72 h (Neuhaus et al., 2017).

Fetal pericytes were more responsive to EDN1. The minimum concentration of EDN1 needed to elicit contraction in aHBVP was 0.1 μM –100 times greater than needed for contraction of fHBVP. aHBVP also took longer to relax following contraction,




as demonstrated by the slower increase in cell index following contraction compared to that for fHBVP. Impaired relaxation of aHBVP could potentially affect neurovascular coupling and contribute to cerebral hypoperfusion in normal aging. Unlike fHBVP, which proliferated in response to EDN1 as reported previously (Neuhaus et al., 2017), aHBVP did not. Whether this reflects a genuine physiological difference between pericytes obtained from fetal and adult brains or is specific to this commercial source of pericytes (Cell-Systems), remains to be determined. Further studies to characterize the morphology and physiology of fetal and adult brain-derived pericytes and their anatomical subtypes should provide important insights into age- and disease-related pericyte dysfunction.

Soluble A β peptides, consisting of monomeric and oligomeric species cause a slow constriction of brain capillaries, mediated by pericyte contraction, reducing the diameter of the vessel lumen by up to 25% in human and rat cortical sections (Nortley


et al., 2019). Transgenic rats treated with A β_{1-40} in the lateral ventricles showed an increase in EDNRA expression in the cerebral cortex, hippocampus, and brain stem (Briyal et al., 2011). We therefore wondered whether pre-exposure to A β had an impact on EDN1-induced contraction of fHBVP and aHBVP *in vitro*.

Short-term exposure had no effect. However, 24 h exposure of fHBVP to A β_{1-40} , at concentrations reported in the human brain (~6 nM) (Nortley et al., 2019), significantly altered the contractile response of pericytes to EDN1. Higher concentrations of A β (in the micromolar range) impaired the contractile responses of pericytes but this may have been partly because of cell toxicity. Previous studies found that exposure to 5–25 μM A β_{1-40} over 3–7 days caused degenerative changes (Verbeek et al., 1997; Sagare et al., 2013) whereas 1.4 μM for 3 h did not cause pericyte death in rat cortical slices (Nortley et al., 2019). We have not found evidence of pericyte



degeneration at concentrations of $A\beta_{1-40}$ or $A\beta_{1-42}$ below 1 μM over a 24 h period. Relaxation of fHBVP was also impaired after exposure to $A\beta_{1-40}$. The deleterious impact of Aβ on pericyte responsiveness – slowing and reducing


the contractile responses as well as subsequent relaxation – is likely to interfere with the rapid adaptation of capillary blood flow to changes in neuronal metabolic demand, and suggests a mechanism whereby elevated Aβ may contribute to




neurovascular uncoupling and cerebral hypoperfusion in AD. It remains to be determined the extent to which the aggregation states of A β (monomeric, oligomeric, and fibrillar), influences pericyte contractile function and the impact of A β on pericyte responses to vasodilators such as NO, adenosine and PGE₂, also merits further study.

Limitations of this study include the reliance on single sources and batches of human fetal and adult pericytes. Pericytes are likely to show heterogeneity of responsiveness *in vivo*, because of multiple genetic and epigenetic factors. Although the fetal pericytes have been used in numerous studies and express canonical pericyte receptors including PDGFR β , NG2, and desmin (Neuhaus et al., 2017), we are still unsure of the precise CNS source of these cells and our observations need to be confirmed by study of pericyte function *in vivo*. Recent transcriptomic studies, such as Vanlandewijck et al. (2018) suggest that these cells may represent vascular fibroblast-like

cells rather than pericytes (*personal communication*) especially after long term culture. This needs further analysis, although we would note that our studies were all on cells that had been passaged fewer than 8 times. We have used a reductionist approach to examine pericytes in isolation whereas *in vivo*, the actions of EDN1 may potentially be modulated locally by neurons, endothelial cells, astrocytes, and microglia. Lastly, we have relied on electrical impedance measurements as proxy indicators of contraction and proliferation. EDN1 was previously shown to mediate proliferation in a study that used both xCELLigence and traditional trypan-blue cell counting (Neuhaus et al., 2017), and we have used a BRDU assay to confirm pericyte proliferation in response to EDN1 (**Supplementary Figure 2**). Other methods, such as measuring the relative change in cell area by confocal microscopy (Hørlyck et al., 2021), are likely to be useful in future *in vivo* studies to validate our findings.





In conclusion, we have shown that fetal and adult brain-derived pericytes contract in response to activation of EDNRA by EDN1. Adult pericytes were less sensitive than fetal pericytes to EDN1 and remained contracted for longer. The contractile response of fetal pericytes was impaired after chronic exposure to A β . These differences may be relevant to disease pathogenesis and should be considered in the design of future studies.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding author.

AUTHOR CONTRIBUTIONS

EH performed all the experiments, analyzed and interpreted the data, and prepared the first draft of the manuscript. All authors contributed to design of the study, revised and edited subsequent drafts of the manuscript, contributed to the article, and approved the submitted version.

FUNDING

This work was supported in part by grant MR/N0137941/1 for the GW4 BIOMED DTP, awarded to the Universities of Bath, Bristol, Cardiff, and Exeter from the Medical Research Council (MRC)/UKRI and the University of Bristol who covered the article processing fees.

REFERENCES

- Billington, C. K., and Penn, R. B. (2003). Signaling and regulation of G protein-coupled receptors in airway smooth muscle. *Respir. Res.* 4:2.
- Briyal, S., Philip, T., and Gulati, A. (2011). Endothelin-A receptor antagonists prevent amyloid- β -induced increase in ETA receptor expression, oxidative stress, and cognitive impairment. *J. Alzheimers Dis.* 23, 491–503. doi: 10.3233/jad-2010-101245
- Chakravarthy, U., Gardiner, T. A., Anderson, P., Archer, D. B., and Trimble, E. R. (1992). The effect of endothelin 1 on the retinal microvascular pericyte. *Microvasc. Res.* 43, 241–254. doi: 10.1016/0026-2862(92)90022-h
- Daneman, R., Zhou, L., Kebede, A. A., and Barres, B. A. (2010). Pericytes are required for blood-brain barrier integrity during embryogenesis. *Nature* 468, 562–566. doi: 10.1038/nature09513
- Dehouck, M. P., Vigne, P., Torpier, G., Breittmayer, J. P., Cecchelli, R., and Frelin, C. (1997). Endothelin-1 as a mediator of endothelial cell-pericyte interactions in bovine brain capillaries. *J. Cereb. Blood Flow Metab.* 17, 464–469. doi: 10.1097/00004647-199704000-00012
- Ding, R., Hase, Y., Ameen-Ali, K. E., Ndung'u, M., Stevenson, W., Barsby, J., et al. (2020). Loss of capillary pericytes and the blood-brain barrier in white matter in poststroke and vascular dementias and Alzheimer's disease. *Brain Pathol.* 30, 1087–1101. doi: 10.1111/bpa.12888
- Durham, J. T., Surks, H. K., Dulmovits, B. M., and Herman, I. M. (2014). Pericyte contractility controls endothelial cell cycle progression and sprouting: insights into angiogenic switch mechanics. *Am. J. Physiol. Cell Physiol.* 307, C878–C892.

ACKNOWLEDGMENTS

We are grateful to the members of the Dementia Research Group for the support and assistance in setting up the impedance software.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fncel.2021.723953/full#supplementary-material>

Supplementary Figure 1 | (A) Graphical representation of cell contraction on impedance plates. Contraction reduces the cell surface area in contact with culture dish electrodes represented by a reduction in cell index. **(B)** Example impedance trace readout from the xCELLigence software. Image **(B)** taken and adapted from ACEA Biosciences Inc.

Supplementary Figure 2 | Fetal human brain-derived vascular pericytes (fHBVPs) were seeded in a 96-well plate at 5,000 cells per well and left to settle overnight. Pericytes were incubated with purified recombinant EDN1 (diluted in serum free medium) for 72 h ($n = 1$). A BRDU incorporation assay (Abcam, United Kingdom) was used according to the manufacturer's instructions to assess pericyte proliferation.

Supplementary Figure 3 | (A) Fetal human brain-derived vascular pericytes (fHBVPs) and **(B)** adult human brain derived-vascular pericytes (aHBVP) were seeded in black clear bottomed 96-well plates at 5,000 cells per well and left to settle overnight. Pericytes were then incubated with recombinant A β peptides (rPeptide, diluted in 35% acetonitrile as a stock and diluted in serum-free medium) for 24 h. Cell toxicity was assessed using an ethidium dye, in the live/dead mammalian cell assay (Invitrogen, United Kingdom), according to the manufacturer's guidelines ($n = 2$). The percentage of cells that had died was calculated with reference to the untreated control wells. The bars represent the mean values and SD.

- Hall, C. N., Reynell, C., Gesslein, B., Hamilton, N. B., Mishra, A., Sutherland, B. A., et al. (2014). Capillary pericytes regulate cerebral blood flow in health and disease. *Nature* 508, 55–60. doi: 10.1038/nature13165
- Heyba, M., Al-Abdullah, L., Henkel, A. W., Sayed, Z., Malatiali, S. A., and Redzic, Z. B. (2019). Viability and contractility of rat brain pericytes in conditions that mimic stroke; an in vitro study. *Front. Neurosci.* 13:1306. doi: 10.3389/fnins.2019.01306
- Hörlyck, S., Cai, C., Helms, H. C. C., Lauritzen, M., and Brodin, B. (2021). ATP induces contraction of cultured brain capillary pericytes via activation of P2Y-type purinergic receptors. *Am. J. Physiol. Heart Circ. Physiol.* 320, H699–H712.
- Joyce, N. C., Haire, M. F., and Palade, G. E. (1985). Contractile proteins in pericytes. I. Immunoperoxidase localization of tropomyosin. *J. Cell Biol.* 100, 1379–1386. doi: 10.1083/jcb.100.5.1379
- Kemp Scott, S., Aguera Kalia, N., Cha, B., and Davis George, E. (2020). Defining endothelial cell-derived factors that promote pericyte recruitment and capillary network assembly. *Arterioscler. Thromb. Vasc. Biol.* 40, 2632–2648. doi: 10.1161/ATVBAHA.120.314948
- Kisler, K., Nelson, A. R., Rege, S. V., Ramanathan, A., Wang, Y., Ahuja, A., et al. (2017). Pericyte degeneration leads to neurovascular uncoupling and limits oxygen supply to brain. *Nat. Neurosci.* 20, 406–416. doi: 10.1038/nn.4489
- Kostov, K., Blazhev, A., Atanasova, M., and Dimitrova, A. (2016). Serum concentrations of endothelin-1 and matrix metalloproteinases-2,-9 in prehypertensive and hypertensive patients with type 2 diabetes. *Int. J. Mol. Sci.* 17:1182. doi: 10.3390/ijms17081182
- Nehls, V., and Drenckhahn, D. (1991). Heterogeneity of microvascular pericytes for smooth muscle type alpha-actin. *J. Cell Biol.* 113, 147–154. doi: 10.1083/jcb.113.1.147

- Neuhaus, A. A., Couch, Y., Sutherland, B. A., and Buchan, A. M. (2017). Novel method to study pericyte contractility and responses to ischaemia in vitro using electrical impedance. *J. Cereb. Blood Flow Metab.* 37, 2013–2024. doi: 10.1177/0271678x16659495
- Nortley, R., Korte, N., Izquierdo, P., Hirunpattarasilp, C., Mishra, A., Jaunmuktane, Z., et al. (2019). Amyloid beta oligomers constrict human capillaries in Alzheimer's disease via signaling to pericytes. *Science* 365:eaav9518. doi: 10.1126/science.aav9518
- Palmer, J. C., Baig, S., Kehoe, P. G., and Love, S. (2009). Endothelin-converting enzyme-2 is increased in Alzheimer's disease and up-regulated by Abeta. *Am. J. Pathol.* 175, 262–270. doi: 10.2353/ajpath.2009.081054
- Palmer, J. C., Barker, R., Kehoe, P. G., and Love, S. (2012). Endothelin-1 is elevated in Alzheimer's disease and upregulated by amyloid-beta. *J. Alzheimers Dis.* 29, 853–861. doi: 10.3233/jad-2012-111760
- Palmer, J. C., Tayler, H. M., and Love, S. (2013). Endothelin-converting enzyme-1 activity, endothelin-1 production, and free radical-dependent vasoconstriction in Alzheimer's disease. *J. Alzheimers Dis.* 36, 577–587. doi: 10.3233/jad-130383
- Peppiatt, C. M., Howarth, C., Mobbs, P., and Attwell, D. (2006). Bidirectional control of CNS capillary diameter by pericytes. *Nature* 443, 700–704. doi: 10.1038/nature05193
- Persidsky, Y., Hill, J., Zhang, M., Dykstra, H., Winfield, M., Reichenbach, N. L., et al. (2016). Dysfunction of brain pericytes in chronic neuroinflammation. *J. Cereb. Blood Flow Metab.* 36, 794–807. doi: 10.1177/0271678x15606149
- Sagare, A. P., Bell, R. D., Zhao, Z., Ma, Q., Winkler, E. A., Ramanathan, A., et al. (2013). Pericyte loss influences Alzheimer-like neurodegeneration in mice. *Nat. Commun.* 4:2932.
- Vanlandewijck, M., He, L., Mäe, M. A., Andrae, J., Ando, K., Del Gaudio, F., et al. (2018). A molecular atlas of cell types and zonation in the brain vasculature. *Nature* 554, 475–480. doi: 10.1038/nature25739
- Verbeek, M. M., de Waal, R. M., Schipper, J. J., and Van Nostrand, W. E. (1997). Rapid degeneration of cultured human brain pericytes by amyloid beta protein. *J. Neurochem.* 68, 1135–1141. doi: 10.1046/j.1471-4159.1997.68031135.x
- Wynne, B. M., Chiao, C.-W., and Webb, R. C. (2009). Vascular smooth muscle cell signaling mechanisms for contraction to angiotensin ii and endothelin-1. *J. Am. Soc. Hypertens. JASH* 3, 84–95. doi: 10.1016/j.jash.2008.09.002
- Yamagishi, S., Hsu, C. C., Kobayashi, K., and Yamamoto, H. (1993). Endothelin 1 mediates endothelial cell-dependent proliferation of vascular pericytes. *Biochem. Biophys. Res. Commun.* 191, 840–846. doi: 10.1006/bbrc.1993.1293

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Hibbs, Love and Miners. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.