

World Journal of Biological Chemistry

Submit a Manuscript: http://www.f6publishing.com

World J Biol Chem 2017 August 26; 8(3): 163-174

DOI: 10.4331/wjbc.v8.i3.163

ORIGINAL ARTICLE

ISSN 1949-8454 (online)

Basic Study Control of nuclear-cytoplasmic shuttling of Ankrd54 by PKC δ

Amy L Samuels, Alison Louw, Reza Zareie, Evan Ingley

Amy L Samuels, Alison Louw, Evan Ingley, Cell Signalling Group, Harry Perkins Institute of Medical Research and Centre for Medical Research, the University of Western Australia, Nedlands, WA 6009, Australia

Reza Zareie, Proteomics International Laboratories Ltd, Nedlands, WA 6009, Australia

Reza Zareie, Proteowa Pty Ltd, SABC, Murdoch University, Murdoch, WA 6150, Australia

Evan Ingley, Cell Signalling Group, School of Veterinary and Health Sciences, Murdoch University, Murdoch, WA 6150, Australia

Author contributions: Samuels AL and Louw A contributed equally to the manuscript and designed, supported and performed experiments, and analyzed data; Zareie R designed and performed experiments, and analysed data; Ingley E designed and supported the research, designed and undertook experiments, analyzed data and contributed to writing the manuscript.

Supported by the National Health and Medical Research Council, Nos. 403987, 513714 and 634352; the Medical Research Foundation of Royal Perth Hospital; and the Cancer Council of Western Australia; Evan Ingley received support from the Cancer Council of Western Australia, the Harry Perkins Institute of Medical Research; Sock-it-to-Sarcoma and the Hollywood Private Hospital Research Foundation; and the MACA Ride to Conquer Cancer.

Conflict-of-interest statement: Each author declares no conflict of interest.

Data sharing statement: All data are available upon request.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http://creativecommons.org/ licenses/by-nc/4.0/

Manuscript source: Unsolicited manuscript

Correspondence to: Dr. Evan Ingley, Professor, Cell Signalling Group, School of Veterinary and Health Sciences, Murdoch University, 90 South Street, Murdoch, WA 6150, Australia. evan.ingley@mudcoh.edu.au Telephone: +61-8-61510738 Fax: +61-8-61510701

Received: January 25, 2017 Peer-review started: February 1, 2017 First decision: March 8, 2017 Revised: April 28, 2017 Accepted: May 12, 2017 Article in press: May 15, 2017 Published online: August 26, 2017

Abstract

AIM

To identify and characterize the effect of phosphorylation on the subcellular localization of Ankrd54.

METHODS

HEK293T cells were treated with calyculin A, staurosporin or phorbol 12-myristate 13-acetate (PMA). Cells were transfected with eGFP-tagged Ankrd54 with or without Lyn tyrosine kinase (wild-type, *Y397F* mutant, or *Y508F* mutant). The subcellular localization was assessed by immunofluorescence imaging of cells, immunoblotting of subcellular fractionations. The phosphorylation of Ankrd54 was monitored using Phos-tagTM gel retardation. Phosphorylated peptides were analysed by multiplereaction-monitoring (MRM) proteomic analysis.

RESULTS

Activation of PKC kinases using PMA promoted nuclear export of Ankrd54 and correlated with increased Ankrd54 phosphorylation, assayed using Phos-tagTM gel retardation. Co-expression of an active form of the PKC δ



isoform specifically promoted both phosphorylation and cytoplasmic localization of Ankrd54, while PKC δ , Akt and PKA did not. Alanine mutation of several serine residues in the amino-terminal region of Ankrd54 (Ser14, Ser17, Ser18, Ser19) reduced both PMA induced cytoplasmic localization and phosphorylation of Ankrd54. Using MRM proteomic analysis, phosphorylation of the Ser18 residue of Ankrd54 was readily detectable in response to PMA stimulation. PMA stimulation of cells co-expressing Ankrd54 and Lyn tyrosine kinase displayed increased co-immunoprecipitation and enhanced co-localization in the cytoplasm.

CONCLUSION

We identify phosphorylation by PKC δ as a major regulator of nuclear-cytoplasmic shuttling of Ankrd54, and its interaction with the tyrosine kinase Lyn.

Key words: Ankrd54; PKCδ; Lyn; Btk; Phosphorylation

© **The Author(s) 2017.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Ankrd54 is a nuclear-cytoplasmic shuttling adaptor that interacts with Lyn, Btk, Txk, and HCLS1. Activation of PKC kinases promoted nuclear export and phosphorylation of Ankrd54, and increased interaction and cytoplasmic co-localization with Lyn. Co-expression of an active form of the PKC δ isoform specifically promoted both phosphorylation and cytoplasmic localization of Ankrd54. Alanine mutation of several serine residues in the amino-terminal region of Ankrd54 reduced both phorbol 12-myristate 13-acetate induced cytoplasmic localization and phosphorylation of Ankrd54. These results identify PKC δ as a major regulator of nuclear-cytoplasmic shuttling and interaction of Ankrd54 with Lyn, through its phosphorylation of at least the Ser18 residue.

Samuels AL, Louw A, Zareie R, Ingley E. Control of nuclearcytoplasmic shuttling of Ankrd54 by PKC8. *World J Biol Chem* 2017; 8(3): 163-174 Available from: URL: http://www.wjgnet. com/1949-8454/full/v8/i3/163.htm DOI: http://dx.doi.org/10.4331/ wjbc.v8.i3.163

INTRODUCTION

Protein nuclear-cytoplasmic compartmentalization and the shuttling of proteins between the nucleus and cytoplasm is regulated at multiple levels and is vitally important for many cellular processes (*e.g.*, transcription). During interphase, when the nuclear envelope exists, the nuclear pore complex (NPC) controls the nuclear entry and exit of many proteins and protein complexes^[1]. While proteins/ complexes less than approximately 40 kDa in size can typically diffuse through the NPC opening, larger proteins rely on an active process to traverse the NPC and enter/ exit the nucleus. The primary control mechanism for most proteins passing through the NPC is the exposure of a nuclear localization sequence (NLS) and/or a nuclear

export sequence (NES)^[2]. The classical NLS is enriched in basic residues and mediates interaction with an importin α/β complex, which then guides the NLS containing cargo through the NPC, and it is released into the nucleus upon RanGTP binding^[3]. Proteins exiting the nucleus typically display a leucine-rich NES which interacts with Crm1 in the presence of RanGTP and mediates their export through the NPC. Many proteins that actively shuttle between the nucleus and cytoplasm contain both NLS and NES motifs^[4].

Posttranslational modifications of nuclear-cytoplasmic shuttling proteins have been shown to regulate the exposure and interaction of NLS and NES motifs with the importin α/β and Crm1 complexes, respectively^[5]. Phosphorylation appears to be a major regulator and mediator of both NLS and NES activity, with examples of phosphorylation promoting nuclear import (*e.g.*, STATs and Erk1/2) as well as nuclear export (*e.g.*, NFAT)^[6-8].

Previously we identified Ankrd54 (also called Liar) as a novel nuclear-cytoplasmic (containing functional NLS and NES motifs) shuttling adaptor of the Src family tyrosine kinase Lyn and other signalling molecules (e.g., HCLS1), that regulates down-stream signalling in erythropoietin-responsive erythroid cells^[9]. Ankrd54 was also identified as an adaptor for the Tec family tyrosine kinases Btk and Txk^[10]. Both these studies illustrated that Ankrd54 interacts with the SH3 domains of these kinases/signalling molecules via its ankyrin repeats and this interaction can regulate the nuclear-cytoplasmic compartmentalization of Lyn/Btk^[9,10]. While we identified that erythropoietin stimulation of erythroid cells promoted nuclear speckle accumulation of both endogenous Ankrd54 and Lyn^[9], the molecular mechanism controlling Ankrd54 nuclear-cytoplasmic shuttling remained unclear. Consequently, we investigated the role of phosphorylation in regulating Ankrd54 localization to nuclear/cytoplasmic compartments and identified an amino-terminal serine cluster that is required for phorbol ester (PMA)-stimulated cytoplasmic accumulation of Ankrd54, mediated by the serine/threonine kinase PKC δ isoform. Intriguingly, tyrosine phosphorylation of PKC δ by Src family kinases activates the serine/threonine kinase, which could then promote cytosolic accumulation of Ankrd54 and cytoplasmic interaction with Lyn.

MATERIALS AND METHODS

Cell culture, antibodies and reagents

HEK293T cells were maintained in Delbocco's modified Eagle's medium (DMEM) with 10% fetal calf serum (FCS). Polyclonal rabbit antibodies were raised against murine Ankrd54 using the purified full-length protein fused to GST, as the antigen, and high titer serum was purified on protein-A beads as described previously^[9]. Additional antibodies used for immunoblotting were anti-Lyn (Santa-Cruz Biotechnology, sc-15 and sc-7274); anti- β -actin (AC-15, ab6276, Abcam, Cambridge, United Kingdom); anti-Histone H3 (H0164, Sigma-Aldrich); and anti-14-3-3 ζ (ab51129, Abcam). TRITC-labeled phalloidin (P1591, Sigma-Aldrich) was used to visualize filamentous actin (F-actin). Phorbol 12-myristate 13-acetate (PMA, P8139), staurosporine (S4400) and leptomycin B (L2913) were purchased from Sigma-Aldrich, calyculin A (#9902) was purchased from Cell Signaling Technologies, and Phos-tag[™]-Acrylamide TM (ALL-107) was purchased from Wako Laboratory Chemicals (Japan).

Plasmid construction

All plasmid constructs were generated by site-directed mutagenesis using oligonucleotides (sequences available upon request), subcloning in frame into the appropriate vector and confirmation by sequencing. Murine Lyn, LynY397F and LynY508F expressing plasmids have previously been described^[11]. Full-length murine Ankrd54 cDNA was generated by incorporation of a 5'RACE product and a cDNA isolated from a cDNA library using site-directed mutagenesis as previously described^[9]. eGFP-tagged Ankrd54 was generated by subcloning murine Ankrd54 into the mammalian expression vector pEGFPC1 (Invitrogen, Carlsbad, CA, United States). This eGFP-Ankrd54 expression construct was then used as a substrate for site-directed mutagenesis to generate the NLS (Ankrd54KKRL111AALA, Ankrd54RR98AA, Ankrd54D98-114) and NES (Ankrd54LSL289ASA) mutants as previously described^[9]. The S14A/S17A/ S18A/S19A mutant was generated by the same procedure using oligonucleotides that simultaneously introduced these serine-alanine mutations in one reaction. The mCherry-SH3 domain fusion was generated by subcloning the SH3 domain from murine Lyn cDNA into the pmCherry-C1 (Invitrogen) vector as an inframe C-terminal fusion of mCherry. The Akt mammalian expression vectors were a kind gift from Professor Brian A. Hemmings (FMI, Switzerland) and have been previously described^[12]. The PKC mammalian expression vectors for PKC α and PKC δ isoforms were a generous gift from Professor Jae-Won Soh (Inha University, South Korea) and have been described previously^[13,14]. Expression vectors for the catalytic domain of cAMP-dependent protein kinase (PKA) were a kind gift from Professor G. Stanley McKnight (University of Washington, Seattle, WA, United States) and have previously been described^[15].

Cell lysis, fractionation, immunoblotting, and immunoprecipitation

Cells (HEK293T) were transiently transfected using Lipofectamine 2000 (Invitrogen) and harvested 24-48 h post transfection. Protein lysates were prepared in 1% nonidet P40, 0.1% SDS, 150 mmol/L NaCl, 50 mmol/L Tris-HCl pH 8.0, 1 mmol/L EDTA, 1 mmol/L EGTA, 25 mmol/L sodium fluoride, 25 mmol/L β -glycerophosphate, 1 mmol/L Na₃VO₄, 1 × protease inhibitor cocktail (Complete, ROCHE) or 1 mmol/L benzamidine/1 mmol/L phenylmethylsulphoyl fluoride (PMSF)/1 µg/mL aprotinin, for 30 min on ice, followed by centrifugation at 10000 × g for 20 min. Protein concentration was estimated using the BioRad D_c protein assay according to manufacturer's instructions (BioRad), using bovine serum albumin as a

standard.

For immunoprecipitation, 5 mg of total protein from clarified cell lysates were incubated with specific antibodies (10 μ g) for 2 h at 4 °C, collected with protein G-Sepharose beads (Sigma-Aldrich) for 16 h (4 °C) before washing in lysis buffer, and subsequent analysis by SDS-PAGE.

For separation of nuclear and cytoplasmic compartments subcellular fractionation was performed. Cells were washed (twice) with cold PBS, and lysed in cell lysis buffer (20 mmol/L Tris-HCl pH 8, 0.5% NP40, 10 mmol/L NaCl, 3 mmol/L MqCl₂ supplemented with protease inhibitors) for 30 min on ice with vigorous pipetting every 5 min. Nuclei were sedimented by centrifugation at 5000 $\times g$ for 5 min at 4 °C and the supernatant (cytoplasmic fraction) was collected. The nuclear pellet was washed 3 times in cell lysis buffer to remove cytoplasmic traces and the nuclei were lysed in nuclear lysis buffer (50 mmol/L Tris-HCl, 1% SDS) on ice for 20 min followed by brief sonication to ensure nuclei rupture. Nuclear debri was collected by centrifugation at 12500 \times g for 10 min at $4 ^{\circ}$ C and the supernatant (nuclear fraction) was collected. Nuclear and cytoplasmic fractions were stored at -80 $^{\circ}C$ prior to analysis by SDS-PAGE and Western blotting.

After probing with specific primary antibodies, proteins were revealed using either secondary antibody coupled to horseradish peroxidase (GE Healthcare or Cell Signaling Technology) and detection by enhanced chemiluminescence (GE Healthcare), or with fluorescently labeled secondary antibodies and an Odyssey scanner (LI-COR Biosciences, Lincoln, NE, United States). Quantitation of immunoblot analysis was performed using on Odyssey scanned blots using Image Studio Light (V3.1.4, LI-COR Biosciences).

Immunofluorescent microscopy

Glass coverslips (No. 1.5H, Marienfeld) were coated with either poly-L-lysine (Sigma-Aldrich) or serum before seeding cells (HEK293T). At the end of specific cell treatments of specific time points, coverslips were washed with PBS (37 °C) then fixed in 4% paraformaldehyde/ PBS for 15 min at 37 $^\circ\!\!\mathbb{C}$, followed by permeabilization with 0.5% triton X-100 in PBS (5 min). Coverslips were then either processed immediately or stored at -80 $^{\circ}\mathrm{C}$ with minimal PBS media. Coverslips were blocked for 2 h at 25 $^{\circ}$ C in 3% BSA/PBS/Tween-20 (0.1%), then incubated with primary antibody diluted in blocking buffer at 4 ℃ for 16 h, then washed 3 times in PBS/Tween-20 (15 min) before incubation with fluorophore-conjugated secondary antibodies (Alexa-Fluor-labeled-488/594, Life-Technologies), with Hoechst 33342 at 0.01 µg/mL, for 1 h at 25 $^\circ\!\!\!\mathrm{C}$, then washed in PBS/Tween-20 before mounting in Vectashield (H-1000, Vector Labs). For some experiments TRITC-labeled phalloidin (P1951, Sigma-Aldrich) at 50 µg/mL was added to the secondary antibody/Hoechst solution to visualize filamentous actin (F-actin). For experiments where only transfected fluorescent protein was being analysed then permeabilized cells were incubated directly with Hoechst with or without



TRITC-phalloidin. For cells co-transfected with eGFP-Ankrd54 and Lyn expressing plasmids, Lyn was detected using anti-Lyn antibodies (Sant-Cruz Biotechnology, sc-15) and Alexa-Fluor-546 donkey anti-rabbit IgG (ThermoFisher Scientific). Fluorescent microscopy was performed using three different instruments: An MRC 1024 UV laser scanning confocal microscope (Bio-Rad) using a Nikon Diaphot 300 microscope fitted with a 40 \times /1.15 oil objective and analysed using Confocal Assistant (v4.02, Bio-Rad); an × 60 plan-Apo 1.40 oil objective on an inverted Elcips-Ti microscope (Nikon, Japan), fitted with a CoolSNAP HQ² CCD camera (Photometrics), images analysed using NIS Elements 4.x (Nikon); or on a DeltaVision Elite system and analysed with softWoRx suite 2.0 (GE Healthcare). Generally, images were processed as z-stacks (deconvolution software used for the non-confocal systems) and maximum image projections generated. Quantitation of subcellular localization was performed on 50-100 individual cells for each experiment by analyzing 10 nuclear and 10 cytoplasmic sub-compartments per individual cell. Ratios of nuclear: cytoplasmic amounts of fluorescence (eGFP channel) were then collated. For statistical analysis experiments were repeated as three biological replicates and analysed by students *t*-test or ANOVA with two-tailed analysis, with orthogonal comparisons (using Prism v7.0b.154, Graphpad). Data are presented as mean \pm SD, *P* < 0.05, *P* < 0.01, ns *P* > 0.05. The statistical methods of this study were reviewed by Michael Phillips from the Harry Perkins Institute of Medical Research.

Multiple reaction monitoring mass spectrometry

Protein bands obtained following SDS-PAGE were reduced and alkylated using DTT and iodoacetamide before digestion with chymotrypsin at pH approximately 8.0. Peptides were extracted as previously described^[16] and analysed by electrospray ionisation mass spectrometry using an Ultimate 3000 nano HPLC system (Dionex) coupled to a 4000 Q TRAP mass spectrometer (Applied Biosystems). Peptides were resolved using a C18 PepMap100 column (LC Packings) running a linear gradient of water/acetonitrile/with 0.1% formic acid. Multiple reaction monitoring (MRM) transitions for the putative phosphopeptides (flanking Ser18 of ANKKRD54) were generated using the Skyline software^[17]. Both singly-phosphorylated and nonphosphorylated forms of the peptides, with one optional missed cleavage, were targeted. A non-phosphorylated peptide (sequence: QQDVEPRDEL, corresponding to amino acid 72-81) was also included in the MRM experiment as an internal normalization control. Spectra were imported back into the Skyline software for interpretation. Peak heights from each sample were divided by the relative peak height of the internal normalization control in the same sample to obtain normalized quantities. Normalized values associated with each hypermass were summed to obtain an overall quantity indicator for that hypermass. The overall hypermass quantities were then divided over the

maximum observed for that hypermass to obtain the overall percentage for the hypermass.

RESULTS

Ankrd54 contains a functional NLS and NES when expressed as an eGFP fusion, co-localizes with active Lyn and interacts with the SH3 domain of Lyn

We have previously shown that myc/HA-tagged and untagged Ankrd54 contain functional NLS and NES sequences^[9]. For this study, we generated an N-terminal eGFP fusion of Ankrd54 and confirmed that the NLS and NES were still functional in this fusion construct (Figure 1A). Further, the Crm1 inhibitor Leptomycin B promoted strong nuclear accumulation of eGFP-Ankrd54 (Figure 1B) as has been observed with myc/HA-tagged Ankrd54 previously^[9]. Interaction of eGFP-Ankrd54 and Lyn was also analogous to that previously delineated with the strongest co-localization seen with constitutively active (Y508F) Lyn and minimal with inactive (Y397F) Lyn (Figure 1C-E).

In erythroid cells Lyn and Ankrd54 both co-localize to nuclear speckles after erythropoietin stimulation^[9], while when overexpressed in HEK293 cells active Lyn and Ankrd54 co-localize in the cytoplasm/plasma-membrane (Figure 1E). To ascertain if Ankrd54 could co-localize with Lyn in the nucleus *via* their Ankyrin repeat-SH3 interaction, we co-expressed an mCherry fusion of the SH3 domain of Lyn with eGFP-tagged Ankrd54 (Figure 1F). Here eGFP-Ankrd54 was predominantly nuclear and localized to nuclear speckles. The SH3 domain of Lyn localized to both the cytoplasm and nucleus with stronger accumulation in the nucleus and specifically within Ankrd54 co-localized nuclear speckles (Figure 1F).

Modulation of phosphorylation via CalyculinA and staurosporine regulates nuclear-cytoplasmic shuttling of Ankrd54

Having established that the eGFP fusion of Ankrd54 retained functional NLS and NES, localization dynamics and Lyn/SH3 domain interactions, we then sought to delineate what signals could influence its nuclearcytoplasmic shuttling. Modulation of phosphorylation using the phosphatase inhibitor CalyculinA and the kinase inhibitor Staurosporine produced profound effects on the subcellular localization of eGFP-Ankrd54 (Figure 1G and H). Immunofluorescence microscopy revealed that activation of kinases through inhibition of phosphatases with CalyculinA promoted cytoplasmic accumulation of eGFP-Ankrd54, while inhibition of kinases with Staurosporine significantly increased nuclear accumulation (Figure 1G). Biochemical subcellular fractionation and immunoblot analysis also showed that CalyculinA significantly promoted cytoplasmic accumulation of eGFP-Ankrd54 (Figure 1H). Further, addition of CalyculinA was unable to influence the compartmentalization of either NLS or NES deletion mutants of eGFP-Ankrd54, which accumulate in the





Figure 1 Ankrd54 shuttles between nuclear and cytoplasmic compartments and phosphorylation regulates this subcellular compartmentalization. A: Localization analysis of eGFP-tagged Ankrd54 (top panel), eGFP-Ankrd54 with the NLS deleted (Ankrd54 Δ NLS, middle panel), and eGFP-Ankrd54 with the NES deleted (Ankrd54 Δ NES, bottom panel) in HEK293 cells. Delineation of the nucleus by Hoechst staining of the DNA is on the right, eGFP fluorescence is on the left. Nuclear and cytoplasmic localization of Ankrd54 was enumerated (graph at right), ^a*P* < 0.05; B: Wild-type eGFP-Ankrd54 expressing HEK293 cells were analysed after treatment with Leptomycin B (+LB) for 2 h (0.4 ng/mL), eGFP fluorescence (left) DNA counter staining (right), and quantitation of nuclear/cytoplasmic localization (graph at right), ^a*P* < 0.05; C: Localization analysis of co-expressed eGFP-Ankrd54 (green) and Lyn wild-type (LWT, red), with DNA counterstained (blue). Merged and individual channels are shown; D: Localization analysis of co-expressed eGFP-Ankrd54 (green) and kinase inactive Lyn (L397, red), with DNA counterstained (blue). Merged and individual channels are shown; F: Localization analysis of eGFP-Ankrd54 (green) and kinase inactive Lyn (L598, red), with DNA counterstained (blue). Merged and individual channels are shown; F: Localization analysis of eGFP-Ankrd54 and an mCherry-fusion of the SH3 domain of Lyn, with DNA counterstained. Merged image (left panel) illustrates co-localizing nuclear puncta (arrow heads); G: Effect of CalyculinA (50 nmol/L, 60 min) or Staurosporine (100 nmol/L, 60 min) on eGFP-Ankrd54 NUS, aubcellular fractionation analysis of eGFP-Ankrd54 (WT), eGFP-Ankrd54ΔNLS (ΔNLS) and eGFP-Ankrd54ΔNES (ΔNES), without (left) and with (right) the addition of CalyculinA (50 nmol/L, 60 min). Nuclear (N) and cytoplasmic (C) fractions of transfected HEK293 cells were immunobloted using anti-Ankrd54, anti-Histone (H3), anti-14-3-3ζ, and anti-β-actin antibodies. Quantitation of nuclear/cytoplasmic localization of Ankrd54, a

155

cytoplasm or nucleus, respectively (Figure 1H).

Identification of PKC δ as mediating phosphorylation and cytoplasmic accumulation of Ankrd54

Having established that global phosphorylation is important for NES mediated cytoplasmic accumulation of Ankrd54 (Figure 1G and H), we then investigated the direct effect of phosphorylation on Ankrd54 subcellular localization. Analysis of phosphorylation databases (PhosphoSitePlus[®], Cell Signaling Technologies; PhosphoNET, Kinexus Bioinformatics Corporation, Canada; and PHOSIDA, Max Planck Institute of Biochemistry, Germany), indicated that Ankrd54 is a phosphoprotein with several residues identified in mass spectrometric data. Utilizing Phos-Tag[™] SDS-PAGE to detect phosphorylation-mediated mobility shifted Ankrd54, we discovered that CalyculinA promoted phosphorylation of Ankrd54 in the cytoplasmic fraction (Figure 2A). Using this Phos-Tag mobility shift assay we then analysed the ability of co-expressing candidate kinases (predicted from database searches, PhosphoNET, PHOSIDA) and kinase activators for their ability to mimic the CalyculinA effect. While co-expression of constitutively active forms of Akt or the catalytic subunit of PKA failed to promote a bandshift of Ankrd54, the addition of the PKC activator PMA did produce a clear phospho-Ankrd54 shifted band (Figure 2B and C). Consequently, we then tested the ability of two widely expressed PMA activated PKC isoforms (a conventional PKC, PKC α ; and a novel PKC, PKC δ) to promote cytoplasmic phosphorylation of Ankrd54 (Figure 2D and E). This clearly demonstrated that PKC δ could promote phosphorylation of cytoplasmic Ankrd54, especially when expressed as a constitutively active form (Δ NPS, N-terminal pseudo-substrate domain deletion), while PKCδ failed to promote Ankrd54 phosphorylation (Figure 2D and E). Further, quantitation of the PKCoNPS promoted phospho-Ankrd54 revealed that PKC δ also promoted cytoplasmic accumulation of Ankrd54 (Cytoplasmic: nuclear ratio; Ankrd54 1.0:1.5; Ankrd54 + PKC&ANPS 1.0:0.3). In addition, the CalyculinA mediated bandshift of cytoplasmic Ankrd54 was enhanced with NLS deleted, but strongly reduced in NES deleted mutants of Ankrd54 (Figure 2F). Consequently, we then ascertained the effect of co-expressing active PKC δ (PKC $\delta\Delta$ NPS) with Ankrd54 on the subcellular localization of Ankrd54 by immunofluorescent microscopy (Figure 2G). This revealed, in agreement with the biochemical analysis (Figure 2E), that PKC δ significantly promotes cytoplasmic accumulation of Ankrd54.

Utilizing predictive algorithms (NetPho3.1)^[18,19] and interpretation of potential site of PKC phosphorylation (Figure 2H), identified four regions of potential phosphorylation; an amino-terminal cluster of serines (Ser14, Ser17, Ser18, Ser19), Thr103, Ser116 and Ser271. The C-terminal site Ser271, close to the NES motif, showed low probability of PKC phosphorylation due to a lack of N-terminal basic residues within the site. The Thr103 and Ser116 sites are located within the NLS motif (Thr103) and immediately C-terminal (Serh116) to the NLS motif, and the N-terminal cluster (Ser14, Ser17, Ser18, Ser19) all have potential for PKCmedicated phosphorylation.

Mass spectrometric analysis of Ser18 of Ankrd54 correlates its phosphorylation status with PMA stimulation

While we identified several potential PKC phosphorylation sites, only the N-terminal Ser14/Ser17/Ser18/Ser19 cluster has been robustly detected in mass-spectrometry datasets (PhosphositePlus®, PhosphoNET, PHOSIDA), in addition to several phosphorylated residues located close to the P-loop (Thr31, Ser38, Ser44, Ser54, Ser58, Ser63), with the Thr103, Ser116 and Ser271 motifs not robustly detected in these global phosphorylation analysis datasets. Consequently, we focused our initial attention on the amino-terminal Ser14/Ser17/Ser18/ Ser19 cluster and its phosphorylation in response to PMA treatment. Using MRM of singly phosphorylated peptides encompassing the N-terminal serine cluster (Table 1) we identified this region as showing strong correlation of its phosphorylation with exposure of cells to PMA. Significantly, this included an inverse correlation of PMA stimulation with the non-phosphorylated peptide.

Mutation of amino-terminal serine cluster (Ser14, Ser17, Ser18, Ser19) mitigates PMA stimulated cytoplasmic accumulation of Ankrd54

Stimulation of eGFP-Ankrd54 transfected HEK293 cells with PMA promotes significant cytoplasmic accumulation of Ankrd54 by 1 h post phorbol ester addition, assayed by immunofluorescent microscopy (Figure 3A). Consequently, we used this time point of PMA stimulation to analyse the effect of mutagenesis of the N-terminal serine cluster (S14A, S17A, S18A, S19A) on cytoplasmic accumulation of Ankrd54 (Figure 3B-E). While wild-type Ankrd54 (Figure 3B) showed significant cytoplasmic accumulation after PMA stimulation in addition to phosphorylation (assayed by Phos-Tag[®] SDS-PAGE), the S14/17/18/19A mutant failed to show significant alteration to its predominantly nuclear subcellular localization, and failed to show a robust phosphorylation mobility shift (Figure 3C). Importantly, direct in-gel comparison of wild-type and the SA mutant of Ankrd54 showed a strong reduction in the band shift upon PMA stimulation of mutant Ankrd54 (Figure 3C, far right panel). Interestingly, mutagenesis of the NES motif (promoting nuclear localization) also disrupted the PMAmediated cytoplasmic accumulation and phosphorylation of Ankrd54 (Figure 3D), while the NLS motif mutant (predominantly cytoplasmic) showed no significant further cytoplasmic accumulation upon PMA stimulation, but did show strong PMA-mediated phosphorylation (Figure 3E).

Modulation of phosphorylation by PMA stimulation regulates nuclear-cytoplasmic shuttling of Ankrd54 and its interaction with Lyn

Stimulation of eGFP-Ankrd54 and Lyn co-transfected HEK293 cells with PMA for 1 h promotes interaction



Samuels AL et al. PKC controls Ankrd54 localization



Figure 2 Phosphorylation by PKCô regulates subcellular compartmentalization of Ankrd54. A: Immunoblot analysis of nuclear (N) and cytoplasmic (C) fractionation of eGFP-Ankrd54 using Phos-Tag™ SDS-PAGE (pTag), with (+CalA) and without (-CalA) CalyculinA treatment (50 nmol/L, 60 min). Transfected HEK293 cells were immunoblotted using anti-Ankrd54, anti-Histone (H3), anti-14-3-3ζ, and anti-β-actin antibodies. Mobility shifted Ankrd54, due to phosphorylation, is indicated (pAnkrd54); B: Immunoblot analysis of nuclear (N) and cytoplasmic (C) fractionation of eGFP-Ankrd54 using Phos-TagTM SDS-PAGE (pTag), co-transfected with PKA expression plasmids, or with addition of PMA (200 nmol/L, 60 min). The wild-type (PKA-Cat-WT) catalytic subunit of PKA, or a dominant active mutant (PKA-Cat-RQ) were induced to express by the addition of Zn. Transfected HEK293 cells were immunoblotted using anti-Ankrd54, anti-Histone (H3), anti-14-3-3ζ, and anti-β-actin antibodies. Mobility shifted Ankrd54, due to phosphorylation, is indicated (pAnkrd54); C: Immunoblot analysis of nuclear (N) and cytoplasmic (C) fractionation of eGFP-Ankrd54 using Phos-Tag™ SDS-PAGE (pTag), co-transfected with Akt expression plasmids. Plasmids expressed either wild-type Akt (Akt-WT), a kinase inactive mutant (Akt-K/A), an oncogenic myristoylated/palmitylated Akt (m/p-Akt), or an activation phospho-mimic (Akt-S/D). Transfected HEK293 cells were immunoblotted using anti-Ankrd54, anti-Histone (H3), anti-14-3-3ζ, and anti-β-actin antibodies. Mobility shifted Ankrd54, due to phosphorylation, is indicated (pAnkrd54); D: Immunoblot analysis of nuclear (N) and cytoplasmic (C) fractionation of eGFP-Ankrd54 using Phos-Tag™ SDS-PAGE (pTag), co-transfected with HA-tagged PKCα expression plasmids. Plasmids expressed either wild-type PKCa (HA-PKAaWT), a kinase inactive mutant (HA-PKCaKR), or an activation mutant (HA-PKCaANPS), Transfected HEK293 cells were immunoblotted using anti-Ankrd54, anti-HA (PKCα), anti-Histone (H3), anti-14-3-3ζ, and anti-β-actin antibodies. Mobility shifted Ankrd54, due to phosphorylation, is indicated (pAnkrd54). Quantitation of cytoplasmic:nuclear ratios are displayed; E: Immunoblot analysis of nuclear (N) and cytoplasmic (C) fractionation of eGFP-Ankrd54 using Phos-Tag™ SDS-PAGE (pTag), co-transfected with HA-tagged PKCδ expression plasmids. Plasmids expressed either wildtype PKCδ (HA-PKAδWT), a kinase inactive mutant (HA-PKCδKR), or an activation mutant (HA-PKCδΔNPS). Transfected HEK293 cells were immunoblotted using anti-Ankrd54, anti-Ha (PKCδ), anti-Histone (H3), anti-14-3-3ζ, and anti-β-actin antibodies. Mobility shifted Ankrd54, due to phosphorylation, is indicated (pAnkrd54). Quantitation of cytoplasmic:nuclear ratios are displayed; F: Immunoblot analysis of cytoplasmic fractionation of eGFP-Ankrd54 using Phos-TagTM SDS-PAGE (pTag), with and without the addition of CalyculinA (50 nmol/L, 60 min). Transfected HEK293 cells were immunoblotted using anti-Ankrd54 and anti-β-actin antibodies. Mobility shifted Ankrd54, due to phosphorylation, is indicated (pAnkrd54); G: Localization analysis of eGFP-Ankrd54 with and without co-expression of dominant active PKCô (PKCôANPS) in HEK293 cells. Delineation of the nucleus by Hoechst staining of the DNA is on the right, eGFP fluorescence (green), PKCô (HA-tag, red), with merged image (left). Nuclear and cytoplasmic localization of eGFP-Ankrd54 was enumerated (graph at right), ^aP < 0.05; H: Prediction of phosphorylated residues in Ankrd54. Top, schematic of Ankrd54 domain structure showing location of P-loop (yellow), NLS (red), ankyrin repeats (green) and NES (purple). Middle, graph of potential phosphorylated residues, predicted by NetPhos3.1. Below, amino acid sequences surrounding top predicted motifs as indicated.

WJBC www.wjgnet.com



Figure 3 Amino terminal serine cluster (Ser14, Ser17, Ser18, Ser19) of Ankrd54 is required for phorbol 12-myristate 13-acetate stimulated nuclear export. A: Localization analysis of eGFP-tagged Ankrd54 (green) in HEK293 cells stimulated with PMA (200 nmol/L) for the indicated times. Delineation of the nucleus by Hoechst staining (blue), and F-actin with TRITC-phalloidin (red). Nuclear and cytoplasmic localization of Ankrd54 was enumerated (graph at right), ${}^{b}P < 0.01$, ns: Not significant (P > 0.05); B-E: Localization analysis of eGFP-tagged Ankrd54 by immunofluorescence of cells (left panels) and by Phos-TagTM SDS-PAGE (pTag) analysis (right panel), in HEK293 cells with and without PMA treatment (200 nmol/L, 60 min). Delineation of the nucleus by Hoechst staining (blue), and F-actin with TRITCphalloidin (red). Nuclear and cytoplasmic localization of Ankrd54 was enumerated (graph in centre), ${}^{b}P < 0.01$. Right panel, immunoblot analysis of nuclear (N) and cytoplasmic (C) fractionation of transfected HEK293 cells were immunoblotted using anti-Ankrd54, anti-Histone (H3), anti-14-3-3 ζ , and anti- β -actin antibodies. Mobility shifted Ankrd54, due to phosphorylation, is indicated (pAnkrd54). Cell immunofluorescence and immunoblot analysis was undertaken for cells expressing wild-type (B), serine-alanine (S14A, S17A, S18A, S19A) mutated (C), NLS deleted (D) and NES deleted (E) constructs of eGFP-tagged Ankrd54. For panel (C) a direct in-gel comparison of wild-type and SA mutant Ankrd54, before and after PMA stimulation is shown in the far-right panel. PMA: Phorbol 12-myristate 13-acetate.

WJBC www.wjgnet.com

| Target peptide | m/z | Charge | Transition | Normalized quantity | | Overall percentage | |
|------------------------|---------|--------|--|---------------------|---------|--------------------|------|
| | | state | | + PMA | -PMA | +PMA | -PMA |
| Non-phosphorylated S18 | 936.078 | 3+ | AATGGGADDESRSGRSSSDGECAVAPEPL.y4+ | 4.2E+02 | 1.9E+03 | 26% | 91% |
| peptide | | | AATGGGADDESRSGRSSSDGECAVAPEPL.y2+ | 8.5E+02 | 2.5E+03 | | |
| | | | AATGGGADDESRSGRSSSDGECAVAPEPL.y15+++ | 2.6E+02 | 1.1E+03 | | |
| | 702.310 | 4+ | AATGGGADDESRSGRSSSDGECAVAPEPL.y4+ | ND | ND | ND | ND |
| | | | AATGGGADDESRSGRSSSDGECAVAPEPL.y2+ | ND | ND | | |
| | 826.871 | 4+ | AATGGGADDESRSGRSSSDGECAVAPEPLAEAGGL.y6+ | 2.0E+03 | 3.2E+03 | 61% | 100% |
| | | | AATGGGADDESRSGRSSSDGECAVAPEPLAEAGGL.y2+ | 2.0E+03 | 3.2E+03 | | |
| Normalization Control | 614.794 | 2 | QQDVEPRDEL.y8+ | 1 | 1 | 94% | 100% |
| (Q72-L81) | | | QQDVEPRDEL.y6+ | 1 | 1 | | |
| | | | QQDVEPRDEL.y5+ | 1 | 1 | | |
| Singly-phosphorylated | 962.733 | 3+ | AATGGGADDESRSGRSSSDGECAVAPEPL(p).y4+ | 1.3E+04 | 7.0E+03 | 70% | 44% |
| S18 peptide | | | AATGGGADDESRSGRSSSDGECAVAPEPL(p).y2+ | 1.1E+03 | 1.6E+03 | | |
| | | | AATGGGADDESRSGRSSSDGECAVAPEPL(p).y15+++ | | | | |
| | 722.302 | 4+ | AATGGGADDESRSGRSSSDGECAVAPEPL(p).y4+ | 5.5E+03 | 2.4E+03 | 71% | 38% |
| | | | AATGGGADDESRSGRSSSDGECAVAPEPL(p).y2+ | 5.8E+02 | 9.0E+02 | | |
| | 846.862 | 4+ | AATGGGADDESRSGRSSSDGECAVAPEPLAEAGGL(p).y6+ | ND | ND | 100% | 83% |
| | | | AATGGGADDESRSGRSSSDGECAVAPEPLAEAGGL(p).y2+ | 1.1E+03 | 8.8E+02 | | |

Table 1 Multiple reaction monitoring of Ser18 of Ankrd54 with phorbol 12-myristate 13-acetate stimulation

Singly-phosphorylated and nonphosphorylated forms of peptides encompassing the Ser18 residue (S18) were analysed and compared to the non-phosphorylated normalization control peptide of Ankrd54, before and after phorbol 12-myristate 13-acetate stimulation (1 μ mol/L, 1 h), of immunoprecipitated Ankrd54. ND: Not detected.

of Lyn and Ankrd54, assayed by immunoblotting of Ankrd54 immunoprecipitates (Figure 4A). Further, immunofluorescent microscopy (Figure 4B) of co-transfected cells illustrated strong co-localization in the cytoplasmic compartment of cells co-expressing Ankrd54 and Lyn, when the cells were stimulated with PMA for 1 h.

DISCUSSION

Here we describe the identification of an N-terminal serine cluster that can be phosphorylated by PKC δ to promote cytoplasmic accumulation of the nuclearcytoplasmic shuttling adaptor Ankrd54 (binding partner of Lyn and Btk tyrosine kinases, and other signalling proteins)^[9,10,20,21] (summarized in Figure 5). These are important findings that help place the pathway context in which Ankrd54 complexes shuttle between the nuclear and cytoplasmic compartments, as being controlled (in part) by PKC δ (Figure 5).

Many of the binding partners of Ankrd54 have important functions both within the cytoplasm as well as the nucleus and enact these subcellular-specific functions in response to many different stimuli. Further, recent findings have identified the nuclear speckles that Ankrd54 can localize to^[9] as being PML (promyelocytic leukaemia) nuclear bodies and paraspeckles^[22], and the strong interaction of the SH3 domain of Lyn and Ankrd54 has been independently confirmed^[23,24]. PML nuclear bodies have a plethora of functions including cell cycle arrest, apoptosis, senescence and transcription and are regulated by signals that feed into these pathways. Paraspeckles function as regulators of gene expression, in response to cellular stress and differentiation, by controlling nuclear retention of RNA (containing doublestranded RNA regions) which have undergone adenosineto-inosine editing^[25-27]. The identification of Ankrd54 as a component of these two sub-nuclear structures, which contain many nuclear-cytoplasmic shuttling molecules, suggests the pathways that Ankrd54 (and its binding partners) is a part of are related to those that these two nuclear bodies/speckles are intimately involved with, *i.e.*, cell cycle, senescence, apoptosis, differentiation and transcription. Indeed, strong nuclear accumulation of Ankrd54 promotes apoptosis, and altering Ankrd54 levels and subcellular localization influences viability and differentiation pathways^[9,20].

PKC δ , a member of the novel (activated by diacylglycerol but not calcium) PKC subfamily has an intimate link with Src family kinases (SFK), including Lyn, as well as nuclear-cytoplasmic shuttling/signalling and the control of apoptosis in response to DNA damage^[28-31]. Activation of PKC δ involves diacylglycerol binding as well as tyrosine phosphorylation by SFK (as well as Abl and Syk), which can lead to activated PKC δ localizing to the plasma-membrane, mitochondria or the nucleus^[30]. Nuclear and mitochondrial localized PKC_δ appear associated with induction of apoptosis^[29,32]. Differential tyrosine phosphorylation of PKCo by SFKs/Abl mediates activation and either translocation to the nucleus or plasma-membrane^[29,31]. Interestingly, when active Lyn is co-expressed with Ankrd54 (Figure 1C-E)^[9], or Btk and Ankrd54 are co-expressed^[10], both accumulate in the cytoplasmic/plasma-membrane fraction. Consequently, Lyn/Btk tyrosine phosphorylated and activated $\text{PKC}\delta$ could be the driver of the cytoplasmic accumulation of Ankrd54 and interaction with Lyn. In addition, when an active form of PKC δ is co-expressed with Ankrd54 both localize outside of the nucleus (Figure 2G). This could also indicate that the activation of PKC δ within this context promotes a viability/anti-apoptotic response, due

WJBC | www.wjgnet.com

Samuels AL et al. PKC controls Ankrd54 localization



Figure 4 Phorbol 12-myristate 13-acetate stimulation of cells co-expressing Ankrd54 and Lyn promotes their co-immunoprecipitation and co-locations. A: Immunoblot analysis of Ankrd54 immunoprecipitates from cells transfected with eGFP-Ankrd54 and/or Lyn with or without PMA stimulation (200 mmol/L, 1 h). Immunoprecipitates and cell lysates were immunoblotted using anti-Ankrd54, anti-Lyn (sc-15, sc-7274), anti-pY-4G10, and anti- β -actin antibodies, ^bP < 0.01; B: Localization analysis of eGFP-tagged Ankrd54 (green) and Lyn (sc-15, alexaFlour546) by immunofluorescence of HEK293 cells with and without PMA treatment (200 nmol/L, 1 h). Delineation of the nucleus by Hoechst staining (blue). Scale bar = 10 μ m; PMA: Phorbol 12-myristate 13-acetate.



Figure 5 Schematic of proposed model of PKCδ regulation of Ankrd54 nuclear-cytoplasmic shuttling. Model of potential mechanism by which PKCδ regulated the cytoplasmic accumulation of Ankrd54. Ankrd54 shuttles between the nuclear and cytoplasmic compartments, promoted by its NLS and NES motifs. Ankrd54 can interact with Lyn, Btk, Txk, HCLS1, LASP1, Vav1, Hip55 and ESE2L. Nuclear Ankrd54 can localize to PML nuclear bodies and paraspeckles (Para-sp). Lyn, and potentially Btk/Txk, can phosphorylate and activate PKCδ, which can then phosphorylate Ankrd54, promoting cytoplasmic accumulation.

to the subcellular localization of PKC δ .

The mechanism by which phosphorylation of Ankrd54 by PKC δ enhances cytoplasmic accumulation of Ankrd54 and its interaction with Lyn remains to be determined. The sites of phosphorylation do not fit the consensus for 14-3-3 binding and so it is unlikely to be *via* 14-3-3 sequestration^[33]. Alternatively, it may interfere with the NLS function and/or enhance the NES function. However, it would be worth exploring the composition of cytoplasmic localized Ankrd54 to determine potential interactions (that may be phosphorylation dependent)

that mediate its cytoplasmic accumulation, as well as determining the influence of nuclear/cytoplasmic localized Ankrd54 on the subcellular distribution of its known binding partners. Further, studies on rates of nuclear/cytoplasmic shuttling for mutants of the N-terminal serine cluster could also identify whether there is a direct effect upon the ability of Ankrd54 to traverse the nuclear pore, *i.e.*, to influence the function of the NLS and/or NES.

In this manuscript, we have identified a mechanism that controls the subcellular localization of the nuclear-

cytoplasmic adaptor Ankrd54 and promotes its interaction with Lyn, through PKC δ mediated phosphorylation of an N-terminal serine cluster (Figure 5). Consequently, it will be important to define the consequences of this PKC δ mediated phosphorylation of Ankrd54 on its other binding partners, their subcellular compartmentalization as well as its influence on their biological function.

ACKNOWLEDGMENTS

We thank Irma Larma and Kevin Li for technical assistance.

COMMENTS

Background

Protein nuclear-cytoplasmic compartmentalization and the shuttling of proteins between the nucleus and cytoplasm is regulated at multiple levels and is vitally important for many cellular processes. Phosphorylation is a major regulator of subcellular compartmentalization of proteins. Ankrd54 shuttles between the nucleus and cytoplasm and here the authors investigated the involvement of phosphorylation on subcellular compartmentalization of this protein.

Research frontiers

The regulation of nuclear-cytoplasmic transport and subcellular localization of signalling molecules such as Ankrd54 is critical for their functions in both normal and diseased cells. The identification of the kinases important for the regulation of subcellular localization of Ankrd54 was the objective of the authors.

Innovations and breakthroughs

This is the first discovery of the involvement of PKC kinases as being important in the nuclear-cytoplasmic shuttling of Ankrd54.

Applications

Knowing that PKCs can regulate the subcellular localization of Ankrd54 may find applications in identifying way to regulate its important interacting partners (*i.e.*, Lyn, Btk, Txk, HCLS1) in health and disease.

Terminology

Regulated subcellular localization: The localization of a protein inside a cell can have dramatic impact on its function, and in many disease incorrect location of proteins leads to specific pathologies. Further, many proteins have altered localization after specific stimulation the promotes of inhibits their function. A proteins localization within a cell is consequently critical for its function and is often regulated by post-translation modification, with phosphorylation being the most common.

Peer-review

In this manuscript, the authors characterized phosphorylation of Ankrd54, a nuclear-cytoplasmic shuttling adaptor protein. They provided good evidence that phosphorylation plays a role in regulating subcellular localization of Ankrd54. Specifically, phosphorylation of the N-terminal region of the protein promotes its accumulation in cytoplasm. They also demonstrate enhanced phosphorylation of Ankrd54 by activated version of PKC-delta, implying it as a potential kinase responsible for Ankrd54 phosphorylation. Using cutting-edge mass spec analysis, they nicely demonstrated that phosphorylation at Ser18 is responding to phorbol 12-myristate 13-acetate stimulation. Overall, the work is very carefully done and the data are with high quality.

REFERENCES

- D'Angelo MA, Hetzer MW. Structure, dynamics and function of nuclear pore complexes. *Trends Cell Biol* 2008; 18: 456-466 [PMID: 18786826 DOI: 10.1016/j.tcb.2008.07.009]
- 2 Pemberton LF, Paschal BM. Mechanisms of receptor-mediated

nuclear import and nuclear export. *Traffic* 2005; **6**: 187-198 [PMID: 15702987 DOI: 10.1111/j.1600-0854.2005.00270.x]

- 3 Lange A, Mills RE, Lange CJ, Stewart M, Devine SE, Corbett AH. Classical nuclear localization signals: definition, function, and interaction with importin alpha. *J Biol Chem* 2007; 282: 5101-5105 [PMID: 17170104 DOI: 10.1074/jbc.R600026200]
- 4 Fabbro M, Henderson BR. Regulation of tumor suppressors by nuclear-cytoplasmic shuttling. *Exp Cell Res* 2003; 282: 59-69 [PMID: 12531692 DOI: 10.1016/S0014-4827(02)00019-8]
- 5 Nardozzi JD, Lott K, Cingolani G. Phosphorylation meets nuclear import: a review. *Cell Commun Signal* 2010; 8: 32 [PMID: 21182795 DOI: 10.1186/1478-811X-8-32]
- 6 Meyer T, Vinkemeier U. Nucleocytoplasmic shuttling of STAT transcription factors. *Eur J Biochem* 2004; 271: 4606-4612 [PMID: 15606747 DOI: 10.1111/j.1432-1033.2004.04423.x]
- 7 Khokhlatchev AV, Canagarajah B, Wilsbacher J, Robinson M, Atkinson M, Goldsmith E, Cobb MH. Phosphorylation of the MAP kinase ERK2 promotes its homodimerization and nuclear translocation. *Cell* 1998; 93: 605-615 [PMID: 9604935 DOI: 10.1016/ S0092-8674(00)81189-7]
- 8 Kehlenbach RH, Dickmanns A, Gerace L. Nucleocytoplasmic shuttling factors including Ran and CRM1 mediate nuclear export of NFAT In vitro. *J Cell Biol* 1998; 141: 863-874 [PMID: 9585406 DOI: 10.1083/jcb.141.4.863]
- 9 Samuels AL, Klinken SP, Ingley E. Liar, a novel Lyn-binding nuclear/ cytoplasmic shuttling protein that influences erythropoietin-induced differentiation. *Blood* 2009; 113: 3845-3856 [PMID: 19064729 DOI: 10.1182/blood-2008-04-153452]
- 10 Gustafsson MO, Hussain A, Mohammad DK, Mohamed AJ, Nguyen V, Metalnikov P, Colwill K, Pawson T, Smith CI, Nore BF. Regulation of nucleocytoplasmic shuttling of Bruton's tyrosine kinase (Btk) through a novel SH3-dependent interaction with ankyrin repeat domain 54 (ANKRD54). *Mol Cell Biol* 2012; **32**: 2440-2453 [PMID: 22527282 DOI: 10.1128/MCB.06620-11]
- 11 Tilbrook PA, Ingley E, Williams JH, Hibbs ML, Klinken SP. Lyn tyrosine kinase is essential for erythropoietin-induced differentiation of J2E erythroid cells. *EMBO J* 1997; 16: 1610-1619 [PMID: 9130706 DOI: 10.1093/emboj/16.7.1610]
- 12 Andjelković M, Alessi DR, Meier R, Fernandez A, Lamb NJ, Frech M, Cron P, Cohen P, Lucocq JM, Hemmings BA. Role of translocation in the activation and function of protein kinase B. *J Biol Chem* 1997; 272: 31515-31524 [PMID: 9395488 DOI: 10.1074/jbc.272.50.31515]
- 13 Soh JW, Lee EH, Prywes R, Weinstein IB. Novel roles of specific isoforms of protein kinase C in activation of the c-fos serum response element. *Mol Cell Biol* 1999; 19: 1313-1324 [PMID: 9891065]
- 14 Soh JW, Weinstein IB. Roles of specific isoforms of protein kinase C in the transcriptional control of cyclin D1 and related genes. *J Biol Chem* 2003; 278: 34709-34716 [PMID: 12794082 DOI: 10.1074/jbc. M302016200]
- 15 Riggle KM, Riehle KJ, Kenerson HL, Turnham R, Homma MK, Kazami M, Samelson B, Bauer R, McKnight GS, Scott JD, Yeung RS. Enhanced cAMP-stimulated protein kinase A activity in human fibrolamellar hepatocellular carcinoma. *Pediatr Res* 2016; 80: 110-118 [PMID: 27027723 DOI: 10.1038/pr.2016.36]
- 16 Bringans S, Eriksen S, Kendrick T, Gopalakrishnakone P, Livk A, Lock R, Lipscombe R. Proteomic analysis of the venom of Heterometrus longimanus (Asian black scorpion). *Proteomics* 2008; 8: 1081-1096 [PMID: 18246572 DOI: 10.1002/pmic.200700948]
- 17 MacLean B, Tomazela DM, Shulman N, Chambers M, Finney GL, Frewen B, Kern R, Tabb DL, Liebler DC, MacCoss MJ. Skyline: an open source document editor for creating and analyzing targeted proteomics experiments. *Bioinformatics* 2010; 26: 966-968 [PMID: 20147306 DOI: 10.1093/bioinformatics/btq054]
- 18 Blom N, Gammeltoft S, Brunak S. Sequence and structure-based prediction of eukaryotic protein phosphorylation sites. *J Mol Biol* 1999; 294: 1351-1362 [PMID: 10600390 DOI: 10.1006/jmbi. 1999.3310]
- 19 Blom N, Sicheritz-Pontén T, Gupta R, Gammeltoft S, Brunak S. Prediction of post-translational glycosylation and phosphorylation of proteins from the amino acid sequence. *Proteomics* 2004; 4: 1633-1649 [PMID: 15174133 DOI: 10.1002/pmic.200300771]

- 20 Barber DL. Truth or dare: role of Liar in EPO-dependent signaling. Blood 2009; 113: 3650-3651 [PMID: 19372263 DOI: 10.1182/ blood-2009-01-198861]
- 21 Ingley E. Integrating novel signaling pathways involved in erythropoiesis. *IUBMB Life* 2012; 64: 402-410 [PMID: 22431075 DOI: 10.1002/iub.1024]
- 22 Fong KW, Li Y, Wang W, Ma W, Li K, Qi RZ, Liu D, Songyang Z, Chen J. Whole-genome screening identifies proteins localized to distinct nuclear bodies. *J Cell Biol* 2013; 203: 149-164 [PMID: 24127217 DOI: 10.1083/jcb.201303145]
- 23 Zhu J, Larman HB, Gao G, Somwar R, Zhang Z, Laserson U, Ciccia A, Pavlova N, Church G, Zhang W, Kesari S, Elledge SJ. Protein interaction discovery using parallel analysis of translated ORFs (PLATO). *Nat Biotechnol* 2013; **31**: 331-334 [PMID: 23503679 DOI: 10.1038/nbt.2539]
- 24 Larman HB, Liang AC, Elledge SJ, Zhu J. Discovery of protein interactions using parallel analysis of translated ORFs (PLATO). *Nat Protoc* 2014; 9: 90-103 [PMID: 24336473 DOI: 10.1038/nprot.2013.167]
- 25 Bond CS, Fox AH. Paraspeckles: nuclear bodies built on long noncoding RNA. J Cell Biol 2009; 186: 637-644 [PMID: 19720872 DOI: 10.1083/jcb.200906113]
- 26 Fox AH, Lamond AI. Paraspeckles. Cold Spring Harb Perspect Biol 2010; 2: a000687 [PMID: 20573717 DOI: 10.1101/cshperspect. a000687]
- 27 Hennig S, Kong G, Mannen T, Sadowska A, Kobelke S, Blythe A, Knott GJ, Iyer KS, Ho D, Newcombe EA, Hosoki K, Goshima N, Kawaguchi T, Hatters D, Trinkle-Mulcahy L, Hirose T, Bond CS, Fox AH. Prion-like domains in RNA binding proteins are essential

for building subnuclear paraspeckles. *J Cell Biol* 2015; **210**: 529-539 [PMID: 26283796 DOI: 10.1083/jcb.201504117]

- 28 Kajimoto T, Sawamura S, Tohyama Y, Mori Y, Newton AC. Protein kinase C {delta}-specific activity reporter reveals agonist-evoked nuclear activity controlled by Src family of kinases. *J Biol Chem* 2010; 285: 41896-41910 [PMID: 20959447 DOI: 10.1074/jbc. M110.184028]
- 29 Humphries MJ, Ohm AM, Schaack J, Adwan TS, Reyland ME. Tyrosine phosphorylation regulates nuclear translocation of PKCdelta. *Oncogene* 2008; 27: 3045-3053 [PMID: 18059334 DOI: 10.1038/ sj.onc.1210967]
- 30 Reyland ME, Jones DN. Multifunctional roles of PKCδ: Opportunities for targeted therapy in human disease. *Pharmacol Ther* 2016; 165: 1-13 [PMID: 27179744 DOI: 10.1016/j.pharmthera.2016.0 5.001]
- 31 Xue ZH, Zhao CQ, Chua GL, Tan SW, Tang XY, Wong SC, Tan SM. Integrin alphaMbeta2 clustering triggers phosphorylation and activation of protein kinase C delta that regulates transcription factor Foxp1 expression in monocytes. *J Immunol* 2010; **184**: 3697-3709 [PMID: 20190138 DOI: 10.4049/jimmunol.0903316]
- 32 Qi X, Mochly-Rosen D. The PKCdelta -Abl complex communicates ER stress to the mitochondria - an essential step in subsequent apoptosis. *J Cell Sci* 2008; 121: 804-813 [PMID: 18285444 DOI: 10.1242/jcs.024653]
- 33 Yaffe MB, Rittinger K, Volinia S, Caron PR, Aitken A, Leffers H, Gamblin SJ, Smerdon SJ, Cantley LC. The structural basis for 14-3-3: phosphopeptide binding specificity. *Cell* 1997; 91: 961-971 [PMID: 9428519 DOI: 10.1016/S0092-8674(00)80487-0]

P- Reviewer: Jia J, Wang Y S- Editor: Song XX L- Editor: A E- Editor: Lu YJ







Published by Baishideng Publishing Group Inc

7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA Telephone: +1-925-223-8242 Fax: +1-925-223-8243 E-mail: bpgoffice@wjgnet.com Help Desk: http://www.f6publishing.com/helpdesk http://www.wjgnet.com

