

Vitamin K1 As A Potential Molecule For Reducing Single-Walled Carbon Nanotubes-Stimulated α -Synuclein Structural Changes And Cytotoxicity

This article was published in the following Dove Press journal:
International Journal of Nanomedicine

Amitis Naskhi,^{1,*} Sanaz Jabbari,^{1,*}
Goran Qader Othman,² Falah
Mohammad Aziz,³ Abbas Salihi,^{3,4}
Majid Sharifi,⁵ Soyar Sari,¹ Keivan
Akhtari,⁶ Shang Ziyad Abdulqadir,³
Asaad AB Alasady,⁷ Osama K
Abou-Zied,⁸ Anwarul
Hasan,^{9,10} Mojtaba Falahati⁵

¹Department of Cellular and Molecular Biology, Faculty of Advanced Science and Technology, Tehran Medical Sciences, Islamic Azad University, Tehran, Iran; ²Department of Medical Laboratory, Health Technical College, Erbil Polytechnic University, Erbil, Iraq; ³Department of Biology, College of Science, Salahaddin University-Erbil, Kurdistan Region, Iraq; ⁴Department of Medical Analysis, Faculty of Science, Tishk International University, Erbil, Iraq; ⁵Department of Nanotechnology, Faculty of Advanced Science and Technology, Tehran Medical Sciences, Islamic Azad University, Tehran, Iran; ⁶Department of Physics, University of Kurdistan, Sanandaj, Iran; ⁷Anatomy, Histology, and Embryology Unit, College of Medicine, University of Duhok, Kurdistan Region, Iraq; ⁸Department of Chemistry, Faculty of Science, Sultan Qaboos University, Muscat, Sultanate of Oman; ⁹Department of Mechanical and Industrial Engineering, College of Engineering, Qatar University, Doha 2713, Qatar; ¹⁰Biomedical Research Center, Qatar University, Doha 2713, Qatar

*These authors contributed equally to this work

Correspondence: Mojtaba Falahati
Department of Nanotechnology, Faculty of
Advanced Science and Technology, Tehran
Medical Sciences, Islamic Azad University,
Tehran, Iran
Email mojtaba.falahati@alumni.ut.ac.ir

Anwarul Hasan
Department of Mechanical and Industrial
Engineering, College of Engineering, Qatar
University, Doha 2713, Qatar
Email hasan.anwarul.mit@gmail.com

Aims: Different kinds of vitamins can be used as promising candidates to mitigate the structural changes of proteins and associated cytotoxicity stimulated by NPs. Therefore, the structural changes of α -syn molecules and their associated cytotoxicity in the presence of SWCNTs either alone or co-incubated with vitamin K1 were studied by spectroscopic, bioinformatical, and cellular assays.

Methods: Intrinsic and ThT fluorescence, CD, and Congo red absorption spectroscopic approaches as well as TEM investigation, molecular docking, and molecular dynamics were used to explore the protective effect of vitamin K1 on the structural changes of α -syn induced by SWCNTs. The cytotoxicity of α -syn/SWCNTs co-incubated with vitamin K1 against SH-SY5Y cells was also carried out by MTT, LDH, and caspase-3 assays.

Results: Fluorescence spectroscopy showed that vitamin K1 has a significant effect in reducing SWCNT-induced fluorescence quenching and aggregation of α -syn. CD, Congo red adsorption, and TEM investigations determined that co-incubation of α -syn with vitamin K1 inhibited the propensity of α -syn into the structural changes and amorphous aggregation in the presence of SWCNT. Docking studies determined the occupation of preferred docked site of SWCNT by vitamin K1 on α -syn conformation. A molecular dynamics study also showed that vitamin K1 reduced the structural changes of α -syn induced by SWCNT. Cellular data exhibited that the cytotoxicity of α -syn co-incubated with vitamin K1 in the presence of SWCNTs is less than the outcomes obtained in the absence of the vitamin K1.

Conclusion: It may be concluded that vitamin K1 decreases the propensity of α -syn aggregation in the presence of SWCNTs and induction of cytotoxicity.

Keywords: α -Syn, single-walled carbon nanotube, aggregation, vitamin K1, cytotoxicity

Introduction

Due to the unique and extraordinary mechanical, electronic, optical, geometric, and biological properties of CNTs,¹ they have been widely used in biological activities such as drug delivery, tissue engineering, cancer treatment, and gene therapy,² recently. In general, CNTs are one-dimensional molecules with one or more walls of graphene that in the form of SWCNTs with a diameter of 0.4 to 3.5 nm can cross the lipid membrane.³ Despite considerable attention in both fields of pharmacological⁴ and toxicological^{5,6} profiles, it is still vital to study the critical effects of CNTs on bimolecular structures and cell integrity. Several reports have shown that CNT causes chronic brain inflammation, multiple sclerosis, an increase of autism risk, a decrease of the IQ in children, neurodegenerative diseases, and

microglia activation.^{7–10} Despite the mentioned reports, dynamic and atomic interactions between SWCNTs and biomolecules are still unknown.¹⁰ Because it has been reported that SWCNTs alter the secondary structure of specific proteins in various forms^{11,12} and accelerate their aggregation based on the conditions of each assay.¹³ The fibrillation of some proteins including A β , α -syn, and tau proteins is linked to a group of neurological diseases such as Alzheimer's and Parkinson's,^{1,14} which have been affected by CNTs.¹³ Generally, CNTs specifically SWCNTs affect protein aggregation by stimulating nucleation and polymerization growth.¹⁵ However, surface changes such as functionalization of SWCNTs can reduce their cytotoxicity.¹⁶ In this regard, Liu et al.¹⁶ revealed that the use of hydroxylating agents on SWCNTs could significantly reduce the amyloid and amorphous protein aggregation, which are very suitable for Alzheimer's treatment.

α -Syn is a soluble protein with 140 amino acids, which are found mainly in the nervous system and red blood cells.¹⁷ Despite the initial recognition of α -syn in the pre-synaptic nerve terminal, its natural role remains unknown.¹⁸ The most common functions for α -syn are the regulation of soluble N-ethylmaleimide-sensitive factor (NSF) attachment protein receptor, production of dopamine and control of synaptic vesicle recycling.¹⁹ The fact is that increase in α -syn aggregation, which can be due to gene amplification, nucleotide polymorphism, and ageing,²⁰ result in neuropathology such as Parkinson's and Alzheimer's diseases.²¹ In this field, Alarcón-Arís et al²² exhibited that reducing the level of α -syn and eliminating their aggregation increases the potential treatment of Parkinson's disease. However, due to the effect of soluble α -syn on the function and development of neurons, it is not possible to ultimately decrease the level of these molecules.²³ Nevertheless, many compounds have been reported to reduce α -syn protein aggregation, including rifampicin,²⁴ dopamine analog compounds,²⁵ quercetin,²⁶ vitamin K,²⁷ gallic acid,²⁸ catepsin D,²⁹ and curcumin compounds.³⁰

Different types of vitamins, specifically fat-soluble vitamin (A, E, D, and K), have shown to reduce the aggregation of the protein.³¹ In this line, experiments have determined that vitamin K (1,4-naphthoquinones) has a wide range of activities on neurological disorders such as Hsp90 inhibition,³² inhibition of monoamine oxidase activity,³³ and inhibition of protein aggregation.³¹ In this regard, da Silva et al²⁷ reported an inhibitory effect of vitamin K against α -syn fibrillation. Hence, in this paper, we examined

the effect of SWCNTs on structural changes and subsequent aggregation of α -syn either alone or co-incubated with vitamin K1. Also, the cytotoxicity of α -syn/SWCNT complex co-incubated with or without vitamin K1 against neuron-like cells (SH-SY5Y) was explored.

Materials And Methods

Materials

α -Syn, ThT, Congo red, vitamin K1, DMEM: F12, FBS, L-glutamine, penicillin, streptomycin, NGF, and MTT were purchased from Sigma (St. Louis, MO, USA). LDH Assay Kit (ab102526) and Annexin V-FITC Apoptosis Staining Kit (ab14085) were purchased from Abcam (Cambridge, CB4 0FL, UK). Caspase-3 activity kit (E13183) was obtained from Thermo Fisher Scientific, Massachusetts, USA.

Sample Preparation

A stock solution of α -syn was prepared in 20 mM phosphate buffer pH 7.4, and protein concentration with an extinction coefficient of $5120 \text{ M}^{-1}\text{cm}^{-1}$ was calculated by a UV-visible (vis) spectrophotometer (Varian, Carry 100 Bio, Australia) at 280 nm. SWCNT was dissolved in DMSO (0.5%, v/v) and sonicated for 20 min using a sonicator probe (Misonix- S3000, USA). Vitamin K1 was also dissolved in DMSO (0.5%, v/v). In the present study, α -syn (50 μM) either alone or with SWCNT (10 $\mu\text{g}/\text{mL}$) was co-incubated with a similar molar ratio of vitamin K1 for 24 hrs. In the future studies, the concentration-dependent inhibition spectra of vitamin K1 against α -syn structural changes induced by nanomaterial can be investigated by spectroscopic assays.

Intrinsic Fluorescence Assay

Intrinsic fluorescence assay was performed with a fluorescence spectrophotometer (Varian, Cary eclipse, Australia). Aliquots of α -syn samples either alone or with vitamin K1 incubated with the single dose of SWCNT for 24 hrs were then removed and diluted ten-fold with 20 mM phosphate buffer pH 7.4 at final concentrations of 5 μM . The excitation was fixed at 270 nm (slit width: 10 nm), and intensity was recorded from 290 to 370 nm (slit width: 10 nm). The resulting spectra were corrected against buffer blank, SWCNT solution, vitamin solution, and inner filter effects.

ThT Fluorescence Assay

Protein samples (5 μM) mixed with ThT to achieve the final ThT concentration of 10 μM . Samples were then

incubated in the dark for 15 min. The excitation was fixed at 440 nm (slit width: 10 nm), and intensity was recorded from 450 to 550 nm (slit width: 10 nm). The resulting spectra were corrected against the blanks.

Congo Red Binding Assay

The protein and Congo red concentrations were mixed at 5 and 20 μM , respectively, and incubated in the dark for 30 mins. The absorbance spectra (400–650 nm) were then recorded with a UV-vis spectrophotometer (Varian, Carry 100 Bio, Australia).

Far-UV CD Assay

The CD signals of α -syn with or without SWCNT either alone or co-incubated with vitamin K1 were recorded using an AVIV 215 spectropolarimeter (Aviv, 215, Lakewood, NJ, USA). Signals as ellipticity changes $[\theta]$ were scanned in the range of 190–250 nm and their blanks (SWCNT and vitamin K1) were subtracted from the protein spectrum.

TEM

TEM images were taken on a Zeiss microscope (EM10C, 100 kV, Germany). The aggregation formation was determined by applying 10 μL of α -syn (50 μM) in the absence and presence of vitamin K1 with SWCNT on a 200-mesh copper grid. Grids were then dried at room temperature for 30 mins.

Simulation Methods

A (6, 6) armchair CNT model of lengths of about 7 nm was constructed and HEX 6.3 software was used to perform docking.³⁴ The structure of the human α -syn protein (PDB ID: 1XQ8) was downloaded from Brookhaven Protein Data Bank. The structure of vitamin K1 was obtained from Avogadro software (Libavogadro Library, Pittsburgh, PA, USA).³⁵ Water molecules and ions were replaced with hydrogen atoms. The protein was fixed to be rigid, and the effect of solvent molecules on docking was ignored. The molecular dynamics simulations were done using the Forcite code and the Dreiding force field.³⁶ The smaller CNT model and α -syn in the absence and presence of vitamin K1 were surrounded by 1000 water molecules, and the geometries were optimized. A time step of 1 fs and a total simulation time of 500 ps were used in the simulation.

Cell Culture

SH-SY5Y cells obtained from Royan institute (Tehran, Iran) were cultured in DMEM: F12 medium in a humidified

atmosphere with 5% (v/v) CO_2 /air at 37°C containing 10% (v/v) FBS and 100 U/mL penicillin and streptomycin.

Cell Viability Assay

MTT assay was explored to assay the cell viability in the presence of different species of α -syn molecules. For cellular assays, the aliquots of α -syn either alone or co-incubated with vitamin K1 in the presence or absence of SWCNT for 24 hrs were then removed and added to the cell culture medium with a final concentration of 5 μM in the 96-well plates. Briefly, cells were seeded at 10^4 cells/well with the differentiation medium, treated with an aliquot of α -syn/SWCNT complex with or without vitamin K1 for 24 hrs, followed by addition of MTT (concentration of 0.5 mg/mL), incubated for 4 hrs at 37°C, and followed by removal of supernatant and addition of DMSO (200 μL). Afterward, the absorbance was read at 570 nm using an ELISA reader (Expert 96, Asys Hitch, Ec Austria).

LDH Release Assay

LDH release assay was performed using the LDH Cytotoxicity Detection kit (ab102526, UK). After treatment, 50 μL of supernatant was added to the cell culture medium and mixed with 50 μL reaction mixture (30 mins) followed by the addition of 50 μL of stop solution. The optical density was read at 470 nm using an ELISA reader (Expert 96, Asys Hitch, Ec Austria).

Caspase-3 Activity Assay

Caspase-3 activity was carried out by employing the caspase-3 activity kit (E13183, Thermo Fisher Scientific). Briefly, cells after treatment were homogenized in reaction buffer followed by determination of protein concentration by Bradford assay. Then, 2mM caspase-3 substrate was added and incubated for 2 hrs at 37°C. The absorbance was finally read at 405 nm using an ELISA reader (Expert 96, Asys Hitch, Ec Austria).

Statistical Analysis

The statistical analysis was done by performing one-way ANOVA for three independent experiments. The significance of outcomes was explored as $P \leq 0.05$.

Results

CNT Characterization

SWCNT characterization was fully carried out in our previous study.¹⁴ Briefly, it was shown that SWCNT shows

microns long with uniform outer diameters of around 1–2 nm.¹⁴ Also, the hydrodynamic radius of SWCNT was around 40–60 nm.¹⁴

Intrinsic Fluorescence Study

To explore the structural changes of α -syn upon interaction with SWCNT, tyrosine fluorescence emission spectrum was acquired for the α -syn in the presence of a single dose of SWCNT after 24 hrs incubation. It was detected that SWCNT could interact with α -syn monomers, eventually causing a quenching impact on the intrinsic fluorescence of α -syn. However, when α -syn co-incubated with vitamin K1 for 24 hrs, the quenching effect of SWCNT on the intrinsic fluorescence of protein was reduced. Although SWCNTs have induced some pronounced structural change in α -syn, these effects were decreased in the presence of vitamin K1 (Figure 1). The higher quenching effect of SWCNT in the absence of vitamin K1 may indicate the better adsorption of α -syn on the surface of SWCNT. It could be suggested that vitamin K1 binds α -syn and mask the binding site of SWCNT on the protein surface, which may explain the inhibiting effect of vitamin K1 on α -syn structural changes induced by SWCNT.

ThT Study

Aggregation of α -syn induced by SWCNT and the inhibitory effect of vitamin K1 can be explored by using ThT fluorescence study. The interaction of SWCNTs with α -syn for 24 hrs substantially significantly enhanced α -syn aggregation as determined by ThT fluorescence measurements (Figure 2). However, ThT fluorescence intensity of the α -syn/SWCNT complex was reduced when protein




Figure 1 Tyrosine fluorescence signals of α -syn with or without vitamin K1 in the presence of SWCNTs.




Figure 2 ThT fluorescence signal of α -syn with or vitamin K1 in the presence of SWCNTs.

samples were co-incubated with vitamin K1 for 24 hrs. These data indicate that the SWCNTs can enhance the aggregated formation of α -syn. This effect is dependent on the presence of the inhibitor as checked by the presence of vitamin K1 (Figure 2). Indeed, ThT fluorescence intensity of α -syn was varied in a more significant manner by SWCNT relative to α -syn/vitamin K1 complex.

Congo Red Absorption Study

Congo red is known as an important dye, which is vastly employed to determine the presence of amyloid fibrils.³⁷ Indeed, binding to amyloid fibrils results in an increased absorbance and red-shift in the maximum optical density of Congo red probe. As shown in the Congo red absorption spectra of α -syn/SWCNT (Figure 3), a significant enhancement in optical density is detected at 493 nm. However, this increase is not accompanied by a detectable red-shift from 493 nm to higher wavelength, determining the formation of non-ordered fibrillar species in the α -syn sample upon interaction with SWCNT. However, the Congo red absorbance (493 nm) decreased after co-incubation of α -syn with vitamin




Figure 3 Congo red absorption band of α -syn with or vitamin K1 in the presence of SWCNTs.

K1 in the presence of SWCNT, revealing the reduction in aggregation propensity of α -syn molecules. Therefore, ThT fluorescence and Congo red absorbance outcomes revealed that the formation of protein aggregates in the presence of SWCNTs exhibited nonfibrillar features and vitamin K1 reduces the formation of these aggregated species.

CD Study

CD is considered a widely used spectroscopic technique to explore the secondary structural changes of biomolecules.^{14,37} The CD spectra of α -syn monomer demonstrated the typical band of predominantly random coil conformation. As shown in Figure 4, the aggregation of α -syn while incubating with SWCNT at 25°C for 24 hrs caused a reduction in random coil structure. However, no considerable concomitant enhancement in the β -sheet fraction was observed, indicating the aggregation of protein into an amorphous morphology. Also, we observed a less reduction in single minima around 195 nm in α -syn samples co-incubated with K1 in the presence of SWCNT compared to the samples without vitamin K1. These data indicated that structural transition in α -syn by SWCNT was decreased upon co-incubation with vitamin K1.

TEM Analysis

To further study the morphology of α -syn aggregation in the presence of SWCNT and the efficacy of vitamin K1 on α -syn aggregation inhibition, TEM study was done to detect the morphological characteristics of α -syn/SWCNT samples in the absence and presence of vitamin K1 after 24 hrs of incubation. As displayed in Figure 5A, SWCNT shows an outer diameter of around 1–2 nm with a worm-like shape.




Figure 4 CD signals of α -syn with or without vitamin K1 in the presence of SWCNTs.

The micrograph of α -syn in the presence of SWCNT shows large quantities of branched nonfibrillar aggregates of \sim 5 nm in diameter and several μ m in length, indicating amorphous aggregated species (Figure 5B). However, as compared to the α -syn/SWCNT sample, α -syn sample co-incubated with 50 μ M vitamin K1 exhibits less aggregated species and are sparsely populated in the presence of SWCNT (Figure 5C). All in all, intrinsic and ThT fluorescence spectroscopy, Congo red binding spectroscopy, CD spectroscopy, and TEM data indicated that SWCNTs induced amorphous aggregation of α -syn, whereas co-incubation of α -syn with vitamin K1 leads to suppression of these nonfibrillar aggregates induced by SWCNT.

Molecular Docking Study

Molecular docking was run to determine the kind of interactions involved between vitamin K1 and α -syn that control the protein aggregation. The molecular docking was performed with a model of CNT and α -syn in the absence or presence of a stable conformer of vitamin K1. The resulting binding energy for CNT and α -syn was -286.32 E-value. To investigate the α -syn affinity to CNTs in the presence of vitamin K1, a successive molecular docking was performed with human α -syn as receptor and 30 molecules of vitamin K1 as ligands. Then, a molecular docking study was performed by the obtained complex and CNT model. The obtained binding energy was -220.63 E-value which reveals a lower affinity of CNT to interact with α -syn/SWCNT complex relative to free α -syn. The docked site was visualized by using CHIMERA (www.cgl.ucsf.edu/chimera) and PyMOL (<http://pymol.sourceforge.net/>) tools. The docked complexes for α -syn with SWCNT either alone or with K1 are shown in Figure 6A or B, respectively. As can be seen in Figure 7A, in the absence of vitamin K1 the nearest residues are Lys-21, Ala-17, Ala-18, Gly-14, Ala-11, Lys-10, while in the presence of vitamin K1, the preferred sites of α -syn for CNT are occupied by vitamin K1 molecules (Figure 6B) which can be the essential factor in decreasing the affinity of CNT to protein/vitamin complex. In this case, the CNT interacts with Val-63, Thr-64, Gly-67 (Figure 7B). Therefore, it may be concluded that vitamin K1 prevented aggregation of α -syn after interaction with SWCNT through non-covalent interaction.

Molecular Dynamics Study

The conformation of α -syn either alone (Figure 8A) or with vitamin K1 (Figure 8B) in the beginning and after 500 ps evolution without (Figure 8C) and with vitamin K1




Figure 5 TEM images of SWCNT (A) α -syn/SWCNT (B) and α -syn/vitamin K1/SWCNT (C).




Figure 6 Binding mode of SWCNT to α -syn in the absence (A) and the presence of vitamin K (B).

(Figure 8D) has been shown in Figure 8. As can be observed, the α -syn upon interaction with CNT in the absence of vitamin K1 tends to attach the CNT wall and be crowded at the nanotube surface which leads to folding in the protein chain. However, the presence of vitamin K1 inhibited the SWCNT-induced structural changes of α -syn by interacting with residues involved in aggregation formation. The molecular dynamics data is in good agreement with experimental spectroscopy outcomes.

MTT Assay

As indicated by MTT assay, SWCNT (1 μ g/mL) or α -syn (5 μ M) showed no cytotoxicity against SH-SY5Y cells




Figure 7 α -syn residues surrounding SWCNT in the absence (A) and presence of vitamin K1 (B) are displayed with numbers within 4 Å.

after 24 hrs (Figure 9). However, cell viability was markedly decreased to 62.1% \pm 10.69% (*P<0.05) after a 24-hrs exposure to 5 μ M α -syn/SWCNT complex. Afterward, when cells were treated with an aliquot of α -syn/SWCNT




Figure 8 The structure of α -syn/SWCNT in the initial (A), α -syn/SWCNT/vitamin K1 in the initial (B), α -syn/SWCNT after 500 ps evolution (C), and α -syn/SWCNT/vitamin K1 after 500 ps evolution (D).

complex co-incubated with vitamin K1 for 24 hrs, cell toxicity was dramatically attenuated ($\#P < 0.05$) (Figure 9). Indeed, treatment of cells with α -syn/vitamin K1/SWCNT complex for 24 hrs significantly elevated the cell viability to 78.88 ± 6.80 . Vitamin K1 did not possess remarkable SH-SY5Y cytotoxic capacities with a concentration of $5 \mu\text{M}$ (data not shown).

LDH Assay

To further investigate the cytotoxic effect of α -syn/SWCNT complex with or without vitamin K1, the release of LDH, another indicator of cytotoxicity, was explored. As displayed in Figure 10, a remarkable increase in LDH release ($*P < 0.05$) was detected after 24 hrs exposures to $5 \mu\text{M}$ α -syn/SWCNT complex. However, treatment of the cells with an aliquot of α -syn/SWCNT complex co-incubated with vitamin K1 for 24 hrs attenuated this enhancement markedly ($\#P < 0.05$). Our data clearly showed that α -syn/SWCNT complex-induced membrane leakage in SH-SY5Y cells was attenuated in the presence of vitamin K1.

Caspase-3 Assay

Caspase-3 as the key apoptotic executive protein can be activated by both extrinsic and intrinsic pathways. Figure 11 exhibits that treatment of SH-SY5Y cells with α -syn/SWCNT complex for 24 hrs significantly increased the activity of caspase-3 ($*P < 0.05$), and this enhancement was remarkably reduced by an aliquot of α -syn/SWCNT complex co-incubated with vitamin K1 for 24 hrs ($\#P < 0.05$).

Discussion

Although some reports have shown that oxidative stress or post-translational changes can play a vital role in oligomerization and α -syn aggregation,^{38,39} further investigations have demonstrated that α -syn aggregation causing neurological problems such as Alzheimer's and Parkinson's diseases are exacerbated by nano-material such as CNTs.⁸ Therefore, understanding the role of CNTs in α -syn aggregation and introduction of some protective agents is crucial for the development of new therapies, which this study examines the protective effect of vitamin K1 against α -syn aggregation and their associated cytotoxicity in SH-SY5Y cell model induced by SWCNT. CNTs are basically insoluble in aqueous solutions and are not suitable for use in




Figure 9 SH-SY5Y cell viability after being exposed to aliquots of α -syn (5 μ M), SWCNT (1 μ gml⁻¹), α -syn/SWCNT complex formed in the absence and presence of vitamin K1. *P \leq 0.05 statistically different from the control group, #P \leq 0.05 statistically different from the α -syn/SWCNT complex group.

medical programs due to biological activity at their surfaces, such as increasing the aggregation of tau and α -syn proteins.¹⁵ Similar to our results, Cavallo et al,⁴⁰ Avti et al,⁴¹ and Yu⁴² reported that the use of SWCNT at concentrations of below 10 μ g/mL does not create cytotoxicity. Also, in agreement with our results, Zeinabad et al¹⁴ described that the use of SWCNTs did not affect LDH activity and cell membrane potential. Whereas, Pichardo et al,⁴³ Toyokuni et al,⁴⁴ and Syama et al⁴⁵ exhibited that the use of SWCNTs could induce cytotoxicity even at a concentration of 0.1 μ g/mL. The difference in reports is due to the variation in the SWCNTs production method, the difference in surface factors, the type of target cells, length and diameter of the SWCNTs, the type of cytotoxicity assays, and even the media used in cellular studies. However, when CNTs interact with proteins, their effects on structural changes of proteins and associated cytotoxicity are not comprehensively understood. In this regard, the results of Ge et al,⁴⁶ Ebrahim-Habibi et al,⁴⁷ and Raghavendra et al⁴⁸ displayed that the interaction of proteins with SWCNTs would be dictated by functional groups on nanotube surface, which will be very effective in controlling the cytotoxicity of SWCNTs. On the other hand, the results of Du et al⁴⁹ demonstrated that SWCNTs caused the aggregation

of lysozyme protein with a change in the protein structure. It was also found that lysozyme aggregation in multiwall CNTs was more severe than SWCNTs. However, changing surface factors from hydroxylates to carboxylates reduces the cytotoxicity of SWCNTs by reducing the accumulation of proteins. Also, SWCNTs result in more lysozyme aggregation (300-fold) compared to albumin protein corona based on multilayer accumulation at the CNT surface.⁴⁹ Analogously, Xie et al⁵⁰ demonstrated that in the presence of the hydroxylating agents on the SWCNT surface, they could significantly reduce the level of protein aggregation in the solution. It was determined that the reduction in protein aggregation is due to increased solubility of SWCNTs in the presence of the hydroxyl groups. Previously using molecular simulations, Li et al⁵¹ and Jana et al⁵² estimated that the hydrophobic interactions between SWCNTs and soluble proteins are considered as main sources in the initiation of protein aggregation. In line with the impact of carbon nanomaterial on the protein aggregation, it has been reported that graphene accelerates the α -syn aggregation and increases the cytotoxicity caused by the tendency of protein to aggregation.⁵³ The presence of some small molecules like vitamins also can play a potential role in the interaction of proteins and NPs. In this study,




Figure 10 SH-SY5Y cell LDH release after being exposed to aliquots of α -syn (5 μ M), SWCNT (1 μ g mL⁻¹), α -syn/SWCNT complex formed in the absence and presence of vitamin K1. *P \leq 0.05 statistically different from the control group, #P \leq 0.05 statistically different from the α -syn/SWCNT complex group.

the protective effects of vitamin K1 against α -syn aggregation stimulated by SWCNTs were explored by fluorescence, CD, UV-vis, TEM, and bioinformatical investigations. It was shown that minor changes occur in the content of the α -syn structure co-incubated with vitamin K1 after interaction with SWCNTs, while the α -syn structure without vitamin K1 showed a significant structural change and subsequent aggregation after addition of SWCNT. Increasing absorption of α -syn on the SWCNT surface results in high levels of disorder in protein structure and function compared to the control group. More data are needed to evaluate the inhibitory effects of vitamin K1 against aggregation triggered by SWCNTs. Besides, the cytotoxicity of α -syn co-incubated with vitamin K1 in the presence of SWCNT on SH-SY5Y cells was also determined by MTT, LDH, and caspase-3 assays. Our results are consistent with reports by da Silva et al,²⁷ Xie et al,⁵⁰ Du et al,⁴⁹ and Pang⁵⁴ which showed that by controlling carbon nanomaterial with auxiliary compounds, it is possible to inhibit the protein aggregation and associated cytotoxicity induced by CNTs.

Despite minimal information on the use of vitamins to control the toxicity of CNTs against protein aggregation, similar to our finding, da Silva et al²⁷ illustrated that vitamin K not only prevented the α -syn aggregation, but also reduced the population of formed oligomers. In addition, they reported that vitamin K, by attaching to the lysine residues of α -syn, causes the formation of amorphous aggregation from the oligomer species.²⁷ Likewise, it was determined that the use of vitamin E along with SWCNTs can prevent the reduction of intracellular protein content and significantly reduce the cytotoxicity of SWCNTs.^{55,56} It has been revealed that the connection between SWCNTs and proteins can increase the active sites for fibril formation and decrease the soluble protein content. The deposition and changing the intracellular protein activity by SWCNTs resulted in increased cell death or apoptosis and exacerbation of pulmonary inflammation in vivo. In another form, Alam et al³¹ reported that the presence of vitamin K could prevent the aggregation of lysozyme and A β peptide which improves




Figure 11 SH-SY5Y cell caspase-3 assay after being exposed to aliquots of α -syn (5 μ M), SWCNT (1 μ g mL⁻¹), α -syn/SWCNT complex formed in the absence and presence of vitamin K1. *P \leq 0.05 statistically different from the control group, #P \leq 0.05 statistically different from the α -syn/SWCNT complex group.

cellular viability in the SH-SY5Y cell line. In addition to vitamin E and K, vitamin A has also been reported as a potential inhibitory molecule against α -syn aggregation.⁵⁷ However, the biological properties of α -syn in the presence of vitamin A are not fully investigated and may be related to the high hydrophobicity of these molecules.

In general, it should be emphasized that intensive care should be considered to apply CNTs in medical applications,^{58–63} especially for the treatment of neurological diseases.

Conclusion

Herein, vitamin K1 was revealed to considerably prevent the formation of amorphous aggregation of α -syn in the presence of SWCNT and its relevant cytotoxicity. Our data suggested that vitamin K1 binds to α -syn monomer and mask the binding site of SWCNT. Furthermore, cytotoxicity experiments revealed that the amorphous aggregates formed by SWCNT in the absence of vitamin K1 are more toxic compared to non-aggregated species produced in the

presence of vitamin K1. The data obtained may provide in-depth insight into probable mechanisms of aggregation inhibition by vitamins and an applicable suggestion about designing unique inhibitors.

Abbreviations

α -syn, α -synuclein; CNTs, carbon nanotubes; CD, circular dichroism; DMEM: F12, Dulbecco's minimum essential medium and Ham's F12 (1:1); FBS, fetal bovine serum; LDH, lactate dehydrogenase; NP, nanoparticle; NGF, nerve growth factor; SWCNTs, single-walled carbon nanotubes; ThT, thioflavin T; TEM, transmission electron microscopy; MTT, 3-(4, 5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide.

Acknowledgement

This article was made possible by the grant NPRP10-120-170-211 from Qatar National Research Fund (a part of Qatar Foundation). All statements here are the sole responsibility of the authors.

Disclosure

The authors declare no conflicts of interest in this work.

References

- Butnaru D, Chapman J. The impact of self-replicating proteins on inflammation, autoimmunity and neurodegeneration—an untraveled path. *Autoimmun Rev*. 2019. doi:10.1016/j.autrev.2018.09.009
- Mohseni-Dargah M, Akbari-Birgani S, Madadi Z, Saghatchi F, Kaboudin B. Carbon nanotube-delivered iC9 suicide gene therapy for killing breast cancer cells in vitro. *Nanomedicine*. 2019;14(8):1033–1047.
- Tagmatarchis N, Prato M. Functionalization of carbon nanotubes via 1, 3-dipolar cycloadditions. *J Mater Chem*. 2004;14(4):437–439. doi:10.1039/b314039c
- Ali-Boucetta H, Kostarelos K. Pharmacology of carbon nanotubes: toxicokinetics, excretion and tissue accumulation. *Adv Drug Del Rev*. 2013;65(15):2111–2119. doi:10.1016/j.addr.2013.10.004
- Zhao X, Lu D, Hao F, Liu R. Exploring the diameter and surface dependent conformational changes in carbon nanotube-protein corona and the related cytotoxicity. *J Hazard Mater*. 2015;292:98–107. doi:10.1016/j.jhazmat.2015.03.023
- Zhao X, Hao F, Lu D, Liu W, Zhou Q, Jiang G. Influence of the surface functional group density on the carbon-nanotube-induced α -chymotrypsin structure and activity alterations. *ACS Appl Mater Interfaces*. 2015;7(33):18880–18890. doi:10.1021/acsami.5b05895
- Bussy C, Al-Jamal KT, Boczkowski J, et al. Microglia determine brain region-specific neurotoxic responses to chemically functionalized carbon nanotubes. *ACS Nano*. 2015;9(8):7815–7830. doi:10.1021/acs.nano.5b02358
- Migliore L, Uboldi C, Di Bucchianico S, Coppedè F. Nanomaterials and neurodegeneration. *Environ Mol Mutagen*. 2015;56(2):149–170. doi:10.1002/em.21931
- Costa PM, Bourgoignon M, Wang JT, Al-Jamal KT. Functionalised carbon nanotubes: from intracellular uptake and cell-related toxicity to systemic brain delivery. *J Control Release*. 2016;241:200–219. doi:10.1016/j.jconrel.2016.09.033
- Kafa H, Wang JT-W, Rubio N, et al. Translocation of LRP1 targeted carbon nanotubes of different diameters across the blood-brain barrier in vitro and in vivo. *J Control Release*. 2016;225:217–225. doi:10.1016/j.jconrel.2016.01.031
- Luo J, Wärmländer SKTS, Yu C-H, Muhammad K, Gräslund A, Pieter Abrahams J. The A β peptide forms non-amyloid fibrils in the presence of carbon nanotubes. *Nanoscale*. 2014;6(12):6720–6726. doi:10.1039/c4nr00291a
- Park M, Park J, Lee J, Ju S-Y. Scaling of binding affinities and cooperativities of surfactants on carbon nanotubes. *Carbon*. 2018;139:427–436. doi:10.1016/j.carbon.2018.07.003
- Allegri M, Perivoliotis DK, Bianchi MG, et al. Toxicity determinants of multi-walled carbon nanotubes: the relationship between functionalization and agglomeration. *Toxicol Rep*. 2016;3:230–243. doi:10.1016/j.toxrep.2016.01.011
- Zeinabad HA, Zarrabian A, Saboury AA, Alizadeh AM, Falahati M. Interaction of single and multi wall carbon nanotubes with the biological systems: tau protein and PC12 cells as targets. *Sci Rep*. 2016;6:26508–26518. doi:10.1038/srep26508
- Li C, Mezzenga R. The interplay between carbon nanomaterials and amyloid fibrils in bio-nanotechnology. *Nanoscale*. 2013;5(14):6207–6218. doi:10.1039/c3nr01644g
- Liu F, Wang W, Sang J, Jia L, Lu F. Hydroxylated single-walled carbon nanotubes inhibit A β 42 fibrillogenesis, disaggregate mature fibrils, and protect against A β 42-induced cytotoxicity. *ACS Chem Neurosci*. 2019;10(1):588–598. doi:10.1021/acschemneuro.8b00441
- Marques O, Outeiro TF. Alpha-synuclein: from secretion to dysfunction and death. *Cell Death Dis*. 2012;3:350–360. doi:10.1038/cddis.2012.94
- Wales P, Pinho R, Lázaro DF, Outeiro TF. Limelight on alpha-synuclein: pathological and mechanistic implications in neurodegeneration. *J Parkinsons Dis*. 2013;3(4):415–459. doi:10.3233/JPD-130216
- Valdinocci D, Radford RAW, Siow SM, Chung RS, Pountney DL. Potential modes of intercellular α -synuclein transmission. *Int J Mol Sci*. 2017;18(2):469–480. doi:10.3390/ijms18020469
- Book A, Guella I, Candido T, et al. A meta-analysis of α -Synuclein multiplication in familial parkinsonism. *Front Neurol*. 2018;9:1–10. doi:10.3389/fneur.2018.00001
- Brundin P, Dave KD, Kordower JH. Therapeutic approaches to target alpha-synuclein pathology. *Exp Neurol*. 2017;298:225–235. doi:10.1016/j.expneurol.2017.10.003
- Alarcón-Aris D, Recasens A, Galofré M, et al. Selective α -synuclein knockdown in monoamine neurons by intranasal oligonucleotide delivery: potential therapy for parkinson's disease. *Mol Ther*. 2018;26(2):550–567. doi:10.1016/j.yjmt.2017.11.015
- Pujols J, Peña-Díaz S, Lázaro DF, et al. Small molecule inhibits α -synuclein aggregation, disrupts amyloid fibrils, and prevents degeneration of dopaminergic neurons. *Proc Natl Acad Sci*. 2018;115(41):10481–10486. doi:10.1073/pnas.1804198115
- Li J, Zhu M, Rajamani S, Uversky VN, Fink AL. Inhibits R. α -synuclein fibrillation and disaggregates fibrils. *Chem Biol*. 2004;11(11):1513–1521. doi:10.1016/j.chembiol.2004.08.025
- Li H-T, Lin D-H, Luo X-Y, et al. Inhibition of α -synuclein fibrillization by dopamine analogs via reaction with the amino groups of α -synuclein. *Febs J*. 2005;272(14):3661–3672. doi:10.1111/j.1742-4658.2005.04792.x
- Zhu M, Han S, Fink AL. Oxidized quercetin inhibits α -synuclein fibrillization. *Biochim Biophys Acta*. 2013;1830(4):2872–2881. doi:10.1016/j.bbagen.2012.12.027
- da Silva FL, Coelho Cerqueira E, de Freitas MS, Gonçalves DL, Costa LT, Follmer C. Vitamins K interact with N-terminus α -synuclein and modulate the protein fibrillization in vitro. Exploring the interaction between quinones and α -synuclein. *Neurochem Int*. 2013;62(1):103–112. doi:10.1016/j.neuint.2012.10.001
- Ardah MT, Paleologou KE, Lv G, et al. Structure activity relationship of phenolic acid inhibitors of α -synuclein fibril formation and toxicity. *Front Aging Neurosci*. 2014;6(197):1–10. doi:10.3389/fnagi.2014.00001
- McGlinchey RP, Lee JC. Cysteine cathepsins are essential in lysosomal degradation of α -synuclein. *Proc Natl Acad Sci*. 2015;112(30):9322–9327. doi:10.1073/pnas.1500937112
- Jha NN, Ghosh D, Das S, et al. Effect of curcumin analogs on α -synuclein aggregation and cytotoxicity. *Sci Rep*. 2016;6:28511–28520. doi:10.1038/srep28511
- Alam P, Chaturvedi SK, Siddiqi MK, et al. Vitamin k3 inhibits protein aggregation: implication in the treatment of amyloid diseases. *Sci Rep*. 2016;6:26759–26765. doi:10.1038/srep26759
- Hadden MK, Hill SA, Davenport J, Matts RL, Blagg BS. Synthesis and evaluation of Hsp90 inhibitors that contain the 1, 4-naphthoquinone scaffold. *Biorg Med Chem*. 2009;17(2):634–640. doi:10.1016/j.bmc.2008.11.064
- Cerqueira EC, Netz PA, Diniz C, do Canto VP, Follmer C. Molecular insights into human monoamine oxidase (MAO) inhibition by 1, 4-naphthoquinone: evidences for menadione (vitamin K3) acting as a competitive and reversible inhibitor of MAO. *Biorg Med Chem*. 2011;19(24):7416–7424. doi:10.1016/j.bmc.2011.10.049
- Ritchie DW, Venkatraman V. Ultra-fast FFT protein docking on graphics processors. *Bioinformatics*. 2010;26(19):2398–2405. doi:10.1093/bioinformatics/btq444
- Hanwell MD, Curtis DE, Lonie DC, Vandermeersch T, Zurek E, Hutchison GR. Avogadro: an advanced semantic chemical editor, visualization, and analysis platform. *J Cheminform*. 2012;4(1):17–25. doi:10.1186/1758-2946-4-17

36. Mayo SL, Olafson BD, Goddard WA. Dreiding—A generic force-field for molecular simulations. *J Phys Chem.* 1990;94:8897–8909. doi:10.1021/j100389a010
37. Mahdavi Mehr M, Meratan AA, Ghobeh M, Ghasemi A, Saboury AA, Nemat-Gorgani M. Inhibition of HEWL fibril formation by taxifolin: mechanism of action. *PLoS One.* 2017;12(11):0187841–0187845. doi:10.1371/journal.pone.0187841
38. Breydo L, Wu JW, Uversky VN. α -Synuclein misfolding and Parkinson's disease. *Biochim Biophys Acta Mol Basis Dis.* 2012;1822(2):261–285. doi:10.1016/j.bbadis.2011.10.002
39. Wan OW, Chung KK. The role of alpha-synuclein oligomerization and aggregation in cellular and animal models of Parkinson's disease. *PLoS One.* 2012;7(6):38545–38554. doi:10.1371/journal.pone.0038545
40. Cavallo D, Fanizza C, Ursini CL, et al. Multi-walled carbon nanotubes induce cytotoxicity and genotoxicity in human lung epithelial cells. *J Appl Toxicol.* 2012;32(6):454–464. doi:10.1002/jat.2711
41. Avti PK, Caparelli ED, Sitharaman B. Cytotoxicity, cytocompatibility, cell-labeling efficiency, and in vitro cellular magnetic resonance imaging of gadolinium-catalyzed single-walled carbon nanotubes. *J Biomed Mater Res A.* 2013;101(12):3580–3591. doi:10.1002/jbm.a.34643
42. Yu IJ. Single-Wall Carbon Nanotubes (SWCNT) Induce Cytotoxicity and Genotoxicity Produced by Reactive Oxygen Species (ROS) Generation in Phytohemagglutinin (PHA)-stimulated male human peripheral blood lymphocytes AU - Kim, Jin Sik. *J Toxicol Environ Health A.* 2014;77(19):1141–1153. doi:10.1080/15287394.2014.917062
43. Pichardo S, Gutiérrez-Praena D, Puerto M, et al. Oxidative stress responses to carboxylic acid functionalized single wall carbon nanotubes on the human intestinal cell line Caco-2. *Toxicol In Vitro.* 2012;26(5):672–677. doi:10.1016/j.tiv.2012.03.007
44. Toyokuni S, Jiang LI, Kitaura R, Shinohara H. Minimal inflammogenicity of pristine single-wall carbon nanotubes. *Nagoya J Med Sci.* 2015;77(1–2):195–202.
45. Syama S, Mohanan PV. Safety and biocompatibility of graphene: A new generation nanomaterial for biomedical application. *Int J Biol Macromol.* 2016;86:546–555. doi:10.1016/j.ijbiomac.2016.01.116
46. Ge C, Du J, Zhao L, et al. Binding of blood proteins to carbon nanotubes reduces cytotoxicity. *Proc Natl Acad Sci USA.* 2011;108(41):16968–16973. doi:10.1073/pnas.1105270108
47. Ebrahim-Habibi MB, Ghobeh M, Mahyari FA, Rafii-Tabar H, Sasanpour P. An investigation into non-covalent functionalization of a single-walled carbon nanotube and a graphene sheet with protein G: A combined experimental and molecular dynamics study. *Scientific Reports.* 2019;9(1):1273.
48. Raghavendra AJ, Fritz K, Fu S, Brown JM, Podila R, Shannahan JH. Variations in biocorona formation related to defects in the structure of single walled carbon nanotubes and the hyperlipidemic disease state. *Sci Rep.* 2017;7(1):8382–8390. doi:10.1038/s41598-017-08896-w
49. Du P, Zhao J, Mashayekhi H, Xing B. Adsorption of bovine serum albumin and lysozyme on functionalized carbon nanotubes. *J Phys Chem C.* 2014;118(38):22249–22257. doi:10.1021/jp5044943
50. Xie L, Lin D, Luo Y, Li H, Yang X, Wei G. Effects of hydroxylated carbon nanotubes on the aggregation of A β 16–22 peptides: a combined simulation and experimental study. *Biophys J.* 2014;107(8):1930–1938. doi:10.1016/j.bpj.2014.08.034
51. Li H, Luo Y, Derreumaux P, Wei G. Carbon nanotube inhibits the formation of β -sheet-rich oligomers of the Alzheimer's amyloid- β (16–22) peptide. *Biophys J.* 2011;101(9):2267–2276. doi:10.1016/j.bpj.2011.09.046
52. Jana AK, Sengupta N. Adsorption mechanism and collapse propensities of the full-length, monomeric A β 1–42 on the surface of a single-walled carbon nanotube: a molecular dynamics simulation study. *Biophys J.* 2012;102(8):1889–1896. doi:10.1016/j.bpj.2012.03.036
53. Mohammadi S, Nikkhal M, Hosseinkhani S. Investigation of the effects of carbon-based nanomaterials on A53T alpha-synuclein aggregation using a whole-cell recombinant biosensor. *Int J Nanomedicine.* 2017;12:8831–8840. doi:10.2147/IJN.S144764
54. Pang R, Li M, Zhang C. Degradation of phenolic compounds by laccase immobilized on carbon nanomaterials: diffusional limitation investigation. *Talanta.* 2015;131:38–45. doi:10.1016/j.talanta.2014.07.045
55. Shvedova AA, Kisin ER, Murray AR, et al. Vitamin E deficiency enhances pulmonary inflammatory response and oxidative stress induced by single-walled carbon nanotubes in C57BL/6 mice. *Toxicol Appl Pharmacol.* 2007;221(3):339–348. doi:10.1016/j.taap.2007.03.018
56. Singh RP, Sharma G, Sonali, et al. Vitamin E TPGS conjugated carbon nanotubes improved efficacy of docetaxel with safety for lung cancer treatment. *Colloids Surf B Biointerfaces.* 141; 2016:429–442. doi:10.1016/j.colsurfb.2016.02.011
57. Ono K, Yamada M. Vitamin A potentially destabilizes preformed α -synuclein fibrils in vitro: implications for Lewy body diseases. *Neurobiol Dis.* 2007;25(2):446–454. doi:10.1016/j.nbd.2006.10.010
58. Yoosefian M, Jahani M. A molecular study on drug delivery system based on carbon nanotube for the novel norepinephrine prodrug, Droxidopa. *J Mol Liq.* 2019;15(284):258–264. doi:10.1016/j.molliq.2019.04.016
59. Yoosefian M, Rahmanifar E, Etminan N. Nanocarrier for levodopa Parkinson therapeutic drug: comprehensive benserazide analysis. *Artif Cells Nanomed Biotechnol.* 2018;46(sup1):434–446. doi:10.1080/21691401.2018.1430583
60. Yoosefian M, Pakpour A, Etminan N. Nanofilter platform based on functionalized carbon nanotubes for adsorption and elimination of Acrolein, a toxicant in cigarette smoke. *Appl Surf Sci.* 2018;444:598–603. doi:10.1016/j.apsusc.2018.03.108
61. Yoosefian M, Etminan N. Leucine/Pd-loaded (5, 5) single-walled carbon nanotube matrix as a novel nanobiosensors for in silico detection of protein. *Amino Acids.* 2018;50(6):653–661. doi:10.1007/s00726-018-2552-4
62. Yoosefian M. A high efficient nanostructured filter based on functionalized carbon nanotube to reduce the tobacco-specific nitrosamines, NNK. *Appl Surf Sci.* 2018;15(434):134–141. doi:10.1016/j.apsusc.2017.10.166
63. Yoosefian M, Etminan N. The role of solvent polarity in the electronic properties, stability and reactivity trend of a tryptophane/Pd doped SWCNT novel nanobiosensor from polar protic to non-polar solvents. *RSC Adv.* 2016;6(69):64818–64825. doi:10.1039/C6RA14006H

International Journal of Nanomedicine

Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Journal Citation Reports/Science Edition, EMBASE, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-nanomedicine-journal>