



ESC

European Society
of Cardiology

Heart rate and blood pressure monitoring in heart failure

Piotr Ponikowski^{1,2}, Ilaria Spoletini³, Andrew J.S. Coats³, Massimo F. Piepoli⁴, and Giuseppe M.C. Rosano^{3*}

¹Department of Heart Diseases, Wroclaw Medical University, Wroclaw, Poland;

²Centre for Heart Diseases, Military Hospital, Wroclaw, Poland;

³Department of Medical Sciences, Centre for Clinical and Basic Research, IRCCS San Raffaele Pisana, via della Pisana, 235, 00163 Rome, Italy; and

⁴Heart Failure Unit, Guglielmo da Saliceto Hospital, Piacenza, Italy

KEYWORDS

Heart failure;
Blood pressure;
Heart rate;
Monitoring

It has been long known that incessant tachycardia and severe hypertension can cause heart failure (HF). In recent years, it has also been recognized that more modest elevations in either heart rate (HR) or blood pressure (BP), if sustained, can be a risk factor both for the development of HF and for mortality in patients with established HF. Heart rate and BP are thus both modifiable risk factors in the setting of HF. What is less clear is the question whether routine systematic monitoring of these simple physiological parameters to a target value can offer clinical benefits. Measuring these parameters clinically during patient review is recommended in HF management in most HF guidelines, both in the acute and chronic phases of the disease. More sophisticated systems now allow long-term automatic or remote monitoring of HR and BP and whether this more detailed patient information can improve clinical outcomes will require prospective RCTs to evaluate. In addition, analysis of patterns of both HR and BP variability can give insights into autonomic function, which is also frequently abnormal in HF. This window into autonomic dysfunction in our HF patients can also provide further independent prognostic information and may in itself be target for future interventional therapies. This article, developed during a consensus meeting of the Heart Failure Association of the ESC concerning the role of physiological monitoring in the complex multi-morbid HF patient, highlights the importance of repeated assessment of HR and BP in HF, and reviews gaps in our knowledge and potential future directions.

Introduction

Monitoring heart rate (HR) and blood pressure (BP) is an established part of routine clinical practice in heart failure (HF) patients. The role of an elevated HR and BP in determining cardiovascular risk is well-known since the Framingham study.¹ Hypertension is the most prevalent

risk factor for HF in the developed world, and long known to be a major and remediable risk factor for HF. The optimal target for BP control in primary prevention has been evaluated in many large-scale antihypertensive trials, but the optimal target BP or HR within an established HF population is less clear. The impact of elevated HRs on adverse outcomes and mortality in HF has been extensively studied,^{2,3} especially in sinus rhythm. For HR, we have compelling evidence that it is beneficial to target a HR close to 60 b.p.m., but for BP the optimal target remains unclear.

*Corresponding author. Tel: +39 06 5225 2409, Fax: +39 06 5225 2465, Email: giuseppe.rosano@gmail.com

Increased HR has a complex interaction with BP, in the short-term BP increases cause reflex bradycardia via the arterial baroreflex system and yet the same factors that drive an increased BP also frequently drive an increased HR, such as increased sympathetic tone. The development of systems to continuously measure both BP and HR can offer novel ways for us to measure HR and BP behaviour and to do so in intense detail.⁴ It may well be that the complex interplay of these control systems of BP and HR and of respiration may offer even more powerful ways to measure prognostic risk within the HF population⁵ and devise novel interventions.⁶ Much of this HR/BP interaction behaviour is linked to the complex autonomic nervous system and its reflex control systems, which themselves are now being evaluated as a therapeutic target in HF.⁷ It may well be that measurement of BP and HR by continuous or semi-continuous monitoring may offer a window for us to use this new research to improve our care of HF patients^{8,9} and also to offer novel targets for intervention in HF. Thus, both BP and HR, and also their interactions, carry potentially important clinical value in the management of HF, and may in future offer us new therapeutic targets and approaches to managing our HF patients.

Benefits of heart rate and blood pressure monitoring in heart failure

Monitoring of HR and BP can contribute to improved outcomes in HF management if it leads to more effective use of guideline-directed medical therapy. In the case of resting HR in sinus rhythm, there is an inverse relation between this parameter and life expectancy.¹⁰ Increased resting sinus HR is a risk factor for the development of HF and sudden cardiac death^{11,12} and HR reduction a beneficial predictive factor in HF.¹³ It has been demonstrated that HR reduction by 10 beats saves almost 5 kg ATP per day.¹⁴ In fact, chronic HF is characterized by higher energy expenditure and blood supply to the myocardium, with impaired vascular resistance and ventricular loading and mechanical dyssynchrony.

The BEAUTIFUL trial showed the prognostic relevance of HR in HF with reduced ejection fraction patients, showing that those with higher HR had greater occurrence of cardiovascular death and HF hospitalization.¹⁵ Similarly, the SHIFT trial found that the risk of cardiovascular death and HF hospitalization increased by 3% with every beat increase from baseline HR and by 16% for every 5 b.p.m. increase.¹⁶ The CHARM programme¹⁷ revealed the prognostic importance of temporal changes in resting HR in HF. In fact, change in HR over time predicted the outcome in HF patients. This further highlights the need for frequent outpatient monitoring of HR. Indeed, HR is a prognostic factor in HF especially in the early post-discharge period where it is related to symptomatic improvement and prognosis.¹⁸ The EVEREST trial clarified that elevated 1 and 4-week post-discharge HR predicted mortality in patients with HF and left ventricular systolic dysfunction in sinus rhythm.¹⁹ Monitoring HR in these post-discharge intervals is, therefore, recommended.

Finally, a meta-analysis²⁰ demonstrated that the magnitude of HR reduction is associated with the survival benefit of beta-blockers in HF, whereas the dose of beta-blocker is not. This implies that HR reduction in beta-blocker treatment for chronic HF mainly contributes to the clinical benefit in this treatment.^{21,22} In contrast to the case for sinus rhythm in atrial fibrillation there is no clear association between HR and prognosis at least over the range of HR from 60 to 100 b.p.m.²³

As for HR, BP is a key prognostic variable to be monitored over time, being directly related to HF development and progression.¹ Chronic hypertension has the potential to determine structural cardiac changes and ventricular dysfunction. For this reason, the lifetime risk for HF doubles in those with BP >160/100 vs. <140/90 mmHg in both sexes.²⁴ Further, there is increased recognition of the harmful consequences of pre-hypertensive chronic states (<140 mmHg).²⁵ There is no clear cut-off point delimitating the need for antihypertensive therapy, as the association between systolic BP and HF risk is continuous.^{25,26} More complex BP behaviour may also be important to monitor in HF syndromes such as the interaction of BP and HR variability (HRV) discussed elsewhere in this article and BP response to simple tests.²⁷

Guidelines-based indications for in-hospital heart rate and blood pressure monitoring

According to the ESC guidelines,²⁸ standard non-invasive monitoring of HR and BP is recommended (Class I) and should be performed from the initial presentation of HF.

For in-hospital monitoring, the 2018 guidance from the HFA/ESC²⁹ indicated that assessments of HR and BP should be repeated at intervals. The first monitoring should occur during hospitalization for acute HF, in order to provide optimal management and vaso-active drug usage.^{30,31} Baseline/initial monitoring usually is performed in the emergency department at the point of initial admission. From then on clinical status monitoring should be performed based on the changing clinical variables of the patient. This usually happens in intensive or critical care unit settings. Repeated monitoring of HR and BP is made in the general ward or prior to discharge.²⁹ Finally, serial assessments of HR and BP should be part of a multidisciplinary after-discharge patient monitoring programme (remote monitoring).

Remote patient monitoring: implantable devices

Using implantable devices, HR and BP can be easily monitored over time, in order to prevent decompensation and consequent rehospitalizations.^{4,32-34} Often, before an episode of decompensation, a rapid increase in mean HR occurs, and it can be simply detected by an implantable device. Measurement of short-term HRV can similarly prevent sudden cardiac death in chronic HF.²⁹ Among currently available sensors, HR derivatives are able to assess mean HR, nocturnal HR, HRV, HRV footprint, and HR during exercise.

For BP monitoring, haemodynamic sensors allow the detection of right ventricular pressure (Chronicle IHM),³⁵ RV dP/dt_{max} , left atrial pressure (HeartPOD),³⁶ and pulmonary artery pressure (Champion).³⁷ There is encouraging evidence showing the effectiveness and safety of implantable device measurements and their correlation with serial invasive measurements. Positive results come from meta-analyses,³⁸⁻⁴¹ whereas few clinical trials to date have demonstrated that implementation of an implantable device improves patient outcome, a notable exception being the CardioMEMS device. This represents an important knowledge gap to be fulfilled.

Conclusions

Repeated assessment of HR and BP over time is recommended for the care of patients with HF and provides prognostic information. Serial comprehensive inpatient and outpatient monitoring is crucial to the optimal management of acute and chronic HF patients to improve outcomes and to prevent adverse events.⁴²⁻⁴⁶

However, most of the guideline recommendations related to monitoring are based on consensus of opinion of the experts and/or small studies, retrospective studies and/or registries (level of evidence C).⁴⁷ For this reason, additional research is warranted.

Conflict of interest: none declared.

References

- Levy D, Larson MG, Vasan RS, Kannel WB, Ho KK. The progression from hypertension to congestive heart failure. *JAMA* 1996;**275**: 1557-1562.
- Hori M, Okamoto H. Heart rate as a target of treatment of chronic heart failure. *J Cardiol* 2012;**60**:86-90.
- Rastogi A, Novak E, Platts AE, Mann DL. Epidemiology, pathophysiology and clinical outcomes for heart failure patients with a mid-range ejection fraction. *Eur J Heart Fail* 2017;**19**:1597-1605.
- Merchant FM, Dec GW, Singh JP. Implantable sensors for heart failure. *Circ Arrhythm Electrophysiol* 2010;**3**:657-667.
- Patel VN, Pierce BR, Bodapati RK, Brown DL, Ives DG, Stein PK. Association of Holter-derived heart rate variability parameters with the development of congestive heart failure in the cardiovascular health study. *JACC Heart Fail* 2017;**5**:423-431.
- Lachowska K, Bellwon J, Narkiewicz K, Gruchala M, Hering D. Long-term effects of device-guided slow breathing in stable heart failure patients with reduced ejection fraction. *Clin Res Cardiol* 2019;**108**: 48-60.
- Pearson MJ, Smart NA. Exercise therapy and autonomic function in heart failure patients: a systematic review and meta-analysis. *Heart Fail Rev* 2018;**23**:91-108.
- Yamada S, Yoshihisa A, Sato Y, Sato T, Kamioka M, Kaneshiro T, Oikawa M, Kobayashi A, Suzuki H, Ishida T, Takeishi Y. Utility of heart rate turbulence and T-wave alternans to assess risk for readmission and cardiac death in hospitalized heart failure patients. *J Cardiovasc Electrophysiol* 2018;**29**:1257-1264.
- Au-Yeung WM, Reinhall PG, Bardy GH, Brunton SL. Development and validation of warning system of ventricular tachyarrhythmia in patients with heart failure with heart rate variability data. *PLoS One* 2018;**13**:e0207215.
- Levine HJ. Rest heart rate and life expectancy. *J Am Coll Cardiol* 1997;**30**:1104-1106.
- Hasenfuss G. Benefit of heart rate reduction in heart failure. *Curr Heart Fail Rep* 2010;**7**:156-158.
- Pokorney SD, Al-Khatib SM, Sun JL, Schulte P, O'Connor CM, Teerlink JR, Armstrong PW, Ezekowitz JA, Starling RC, Voors AA, Velazquez EJ, Hernandez AF, Mentz RJ. Sudden cardiac death after acute heart failure hospital admission: insights from ASCEND-HF. *Eur J Heart Fail* 2018;**20**:525-532.
- Reil JC, Böhm M. BEAUTIFUL results-the slower, the better? *Lancet* 2008;**372**:779-780.
- Ferrari R, Ceconi C, Guardigli G. Pathophysiological role of heart rate: from ischaemia to left ventricular dysfunction. *Eur Heart J Suppl* 2008;**10**(Suppl F):F7-F10.
- Fox K, Ford I, Steg PG, Tendera M, Robertson M, Ferrari R; BEAUTIFUL Investigators. Heart rate as a prognostic risk factor in patients with coronary artery disease and left-ventricular systolic dysfunction (BEAUTIFUL): a subgroup analysis of a randomised controlled trial. *Lancet* 2008;**372**:817-821.
- Böhm M, Komajda M, Borer JS, Ford I, Maack C, Tavazzi L, Moyne A, Swedberg K; SHIFT Investigators. Duration of chronic heart failure affects outcomes with preserved effects of heart rate reduction with ivabradine: findings from SHIFT. *Eur J Heart Fail* 2018;**20**: 373-381.
- Vazir A, Claggett B, Jhund P, Castagno D, Skali H, Yusuf S, Swedberg K, Granger CB, McMurray JJ, Pfeffer MA, Solomon SD. Prognostic importance of temporal changes in resting heart rate in heart failure patients: an analysis of the CHARM program. *Eur Heart J* 2015;**36**: 669-675.
- Metra M, Gheorghiadu M, Bonow RO, Dei Cas L. Postdischarge assessment after a heart failure hospitalization: the next step forward. *Circulation* 2010;**122**:1782-1785.
- Greene SJ, Vaduganathan M, Wilcox JE, Harinstein ME, Maggioni AP, Subacius H, Zannad F, Konstam MA, Chioncel O, Yancy CW, Swedberg K, Butler J, Bonow RO, Gheorghiadu M; EVEREST Trial Investigators. The prognostic significance of heart rate in patients hospitalized for heart failure with reduced ejection fraction in sinus rhythm: insights from the EVEREST (Efficacy of Vasopressin Antagonism in Heart Failure: outcome Study With Tolvaptan) trial. *JACC Heart Fail* 2013; **1**:488-496.
- McAlister FA, Wiebe N, Ezekowitz JA, Leung AA, Armstrong PW. Meta-analysis: beta-blocker dose, heart rate reduction, and death in patients with heart failure. *Ann Intern Med* 2009;**150**:784-794.
- Agewall S. Cardiovascular pharmacotherapy and real-world data. *Eur Heart J Cardiovasc Pharmacother* 2018;**4**:65-66.
- Agewall S. Cardiovascular pharmacotherapy. *Eur Heart J Cardiovasc Pharmacother* 2018;**4**:1.
- Kotecha D, Flather MD, Altman DG, Holmes J, Rosano G, Wikstrand J, Packer M, Coats AJS, Manzano L, Böhm M, van Veldhuisen DJ, Andersson B, Wedel H, von Lueder TG, Rigby AS, Hjalmarsen A, Kjekshus J, Cleland JGF; Beta-Blockers in Heart Failure Collaborative Group. Heart rate and rhythm and the benefit of beta-blockers in patients with heart failure. *J Am Coll Cardiol* 2017;**69**: 2885-2896.
- Lloyd-Jones DM, Larson MG, Leip EP, Beiser A, D'Agostino RB, Kannel WB, Murabito JM, Vasan RS, Benjamin EJ, Levy D. Lifetime risk for developing congestive heart failure: the Framingham Heart Study. *Circulation* 2002;**106**:3068-3072.
- Georgiopoulou VV1, Kalogeropoulos AP, Butler J. Dilemmas of blood pressure management for heart failure prevention. *Circ Heart Fail* 2011;**4**:528-533.
- Lip GYH, Coca A, Kahan T, Boriani G, Manolis AS, Olsen MH, Oto A, Potpara TS, Steffel J, Marín F, de Oliveira Figueiredo MJ, de Simone G, Tzou WS, En Chiang C, Williams B. Hypertension and cardiac arrhythmias: executive summary of a consensus document from the European Heart Rhythm Association (EHRA) and ESC Council on Hypertension, endorsed by the Heart Rhythm Society (HRS), Asia-Pacific Heart Rhythm Society (APHRS), and Sociedad Latinoamericana de Estimulación Cardíaca y Electrofisiología (SOLEACE). *Eur Heart J Cardiovasc Pharmacother* 2017;**3**:235-250.
- Bronzwaer AGT, Bogert LWJ, Westerhof BE, Piek JJ, Daemen M, van Lieshout JJ. Abnormal haemodynamic postural response in patients with chronic heart failure. *ESC Heart Fail* 2017;**4**:146-153.
- Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, González-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano

- GM, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P; Authors/Task Force Members; Document Reviewers. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC). Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2016;18:891-975.
29. Harjola VP, Parissis J, Brunner-La Rocca HP, Celutkienė J, Chioncel O, Collins SP, De Backer D, Filippatos GS, Gayat E, Hill L, Lainscak M, Lassus J, Masip J, Mebazaa A, Miró Ò, Mortara A, Mueller C, Mullens W, Nieminen MS, Rudiger A, Ruschitzka F, Seferovic PM, Sionis A, Vieillard-Baron A, Weinstein JM, de Boer RA, Crespo-Leiro MG, Piepoli M, Riley JP. Comprehensive in-hospital monitoring in acute heart failure: applications for clinical practice and future directions for research. A statement from the Acute Heart Failure Committee of the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). *Eur J Heart Fail* 2018;20:1081-1099.
 30. Hoedemaker NPG, Damman P, Ottervanger JP, Dambrink JHE, Gosselink ATM, Kedhi E, Kolkman E, de Winter RJ, van't Hof AWJ. Trends in optimal medical therapy prescription and mortality after admission for acute coronary syndrome: a 9-year experience in a real-world setting. *Eur Heart J Cardiovasc Pharmacother* 2018;4:102-110.
 31. Cheema B, Ambrosy AP, Kaplan RM, Senni M, Fonarow GC, Chioncel O, Butler J, Gheorghiu M. Lessons learned in acute heart failure. *Eur J Heart Fail* 2018;20:630-641.
 32. Fudim M, O'Connor CM, Dunning A, Ambrosy AP, Armstrong PW, Coles A, Ezekowitz JA, Greene SJ, Metra M, Starling RC, Voors AA, Hernandez AF, Michael Felker G, Mentz RJ. Aetiology, timing and clinical predictors of early vs. late readmission following index hospitalization for acute heart failure: insights from ASCEND-HF. *Eur J Heart Fail* 2018;20:304-314.
 33. Nagueh SF. Non-invasive assessment of left ventricular filling pressure. *Eur J Heart Fail* 2018;20:38-48.
 34. Donal E, Galli E, Fraser AG. Non-invasive estimation of left heart filling pressures: another nail in the coffin for E/e'? *Eur J Heart Fail* 2017;19:1661-1663.
 35. Ho C. Implantable hemodynamic monitoring (the Chronicle IHM system): remote telemonitoring for patients with heart failure. *Issues Emerg Health Technol* 2008;111:1-4.
 36. Walton AS, Krum H. The HeartPOD implantable heart failure therapy system. *Heart Lung Circ* 2005;14 Suppl 2:S31-S33.
 37. Givertz MM, Stevenson LW, Costanzo MR, Bourge RC, Bauman JG, Ginn G, Abraham WT; CHAMPION Trial Investigators. Pulmonary artery pressure-guided management of patients with heart failure and reduced ejection fraction. *J Am Coll Cardiol* 2017;70:1875-1886.
 38. Klersy C, De Silvestri A, Gabutti G, Regoli F, Auricchio A. A meta-analysis of remote monitoring of heart failure patients. *J Am Coll Cardiol* 2009;54:1683-1694.
 39. Inglis SC, Clark RA, McAlister FA, Ball J, Lewinter C, Cullington D, Stewart S, Cleland JG. Structured telephone support or telemonitoring programmes for patients with chronic heart failure. *Cochrane Database Syst Rev* 2010;8:CD007228.
 40. Clark RA, Inglis SC, McAlister FA, Cleland JG, Stewart S. Telemonitoring or structured telephone support programmes for patients with chronic heart failure: systematic review and meta-analysis. *BMJ* 2007;334:942.
 41. Clarke M, Shah A, Sharma U. Systematic review of studies on telemonitoring of patients with congestive heart failure: a meta-analysis. *J Telemed Telecare* 2011;17:7-14.
 42. Pareek A, Chandurkar N, Dharmadhikari S. Congestive heart failure: more common as well as an important cardiovascular outcome. *Eur Heart J Cardiovasc Pharmacother* 2017;3:98.
 43. Bhambhani V, Kizer JR, Lima JAC, van der Harst P, Bahrami H, Nayor M, de Filippi CR, Enserro D, Blaha MJ, Cushman M, Wang TJ, Gansevoort RT, Fox CS, Gaggin HK, Kop WJ, Liu K, Vasan RS, Psaty BM, Lee DS, Brouwers FP, Hillege HL, Bartz TM, Benjamin EJ, Chan C, Allison M, Gardin JM, Januzzi JL Jr, Levy D, Herrington DM, van Gilst WH, Bertoni AG, Larson MG, de Boer RA, Gottdiener JS, Shah SJ, Ho JE. Predictors and outcomes of heart failure with mid-range ejection fraction. *Eur J Heart Fail* 2018;20:651-659.
 44. Agostoni P, Paolillo S, Mapelli M, Gentile P, Salvioni E, Veglia F, Bonomi A, Corrà U, Lagioia R, Limongelli G, Sinagra G, Cattadori G, Scardovi AB, Metra M, Carubelli V, Scrutinio D, Raimondo R, Emdin M, Piepoli M, Magrì D, Parati G, Caravita S, Re F, Cicoira M, Minà C, Corrae M, Frigerio M, Bussotti M, Oliva F, Battaia E, Belardinelli R, Mezzani A, Pastormerlo L, Guazzi M, Badagliacca R, Di Lenarda A, Passino C, Sciomer S, Zambon E, Pacileo G, Ricci R, Apostolo A, Palermo P, Contini M, Clemenza F, Marchese G, Gargiulo P, Binno S, Lombardi C, Passantino A, Filardi PP. Multiparametric prognostic scores in chronic heart failure with reduced ejection fraction: a long-term comparison. *Eur J Heart Fail* 2018;20:700-710.
 45. Perk J. The 2016 version of the European Guidelines on Cardiovascular Prevention. *Eur Heart J Cardiovasc Pharmacother* 2017;3:9-10.
 46. Kim JS, Park S, Yan P, Jeffers BW, Cerezo C. Effect of inter-individual blood pressure variability on the progression of atherosclerosis in carotid and coronary arteries: a post hoc analysis of the NORMALISE and PREVENT studies. *Eur Heart J Cardiovasc Pharmacother* 2017;3:82-89.
 47. Agewall S. Adherence to guidelines and registry data. *Eur Heart J Cardiovasc Pharmacother* 2017;3:183-184.