


RESEARCH ARTICLE

Open Access



# Association of in-hospital resource utilization with post-acute spending in Medicare beneficiaries hospitalized for acute myocardial infarction: a cross-sectional study

Sudhakar V. Nuti<sup>1,2</sup>, Shu-Xia Li<sup>2</sup>, Xiao Xu<sup>2,3</sup>, Lesli S. Ott<sup>2,10</sup>, Tara Lagu<sup>4,5</sup>, Nihar R. Desai<sup>2,6</sup>, Karthik Murugiah<sup>2,6</sup>, Michael Duan<sup>7</sup>, John Martin<sup>7</sup>, Nancy Kim<sup>8</sup> and Harlan M. Krumholz<sup>2,6,9\*</sup> 

## Abstract

**Background:** Efforts to decrease hospitalization costs could increase post-acute care costs. This effect could undermine initiatives to reduce overall episode costs and have implications for the design of health care under alternative payment models.

**Methods:** Among Medicare fee-for-service beneficiaries aged  $\geq 65$  years hospitalized with acute myocardial infarction (AMI) between July 2010 and June 2013 in the Premier Healthcare Database, we studied the association of in-hospital and post-acute care resource utilization and outcomes by in-hospital cost tertiles.

**Results:** Among patients with AMI at 326 hospitals, the median (range) of each hospital's mean per-patient in-hospital risk-standardized cost (RSC) for the low, medium, and high cost tertiles were \$16,257 (\$13,097–\$17,648), \$18,544 (\$17,663–\$19,875), and \$21,831 (\$19,923–\$31,296), respectively. There was no difference in the median (IQR) of risk-standardized post-acute payments across cost-tertiles: \$5014 (4295–6051), \$4980 (4349–5931) and \$4922 (4056–5457) for the low ( $n = 90$ ), medium ( $n = 98$ ), and high ( $n = 86$ ) in-hospital RSC tertiles ( $p = 0.21$ ), respectively. In-hospital and 30-day mortality rates did not differ significantly across the in-hospital RSC tertiles; however, 30-day readmission rates were higher at hospitals with higher in-hospital RSCs: median = 17.5, 17.8, and 18.0% at low, medium, and high in-hospital RSC tertiles, respectively ( $p = 0.005$  for test of trend across tertiles).

**Conclusions:** In our study of patients hospitalized with AMI, greater resource utilization during the hospitalization was not associated with meaningful differences in costs or mortality during the post-acute period. These findings suggest that it may be possible for higher cost hospitals to improve efficiency in care without increasing post-acute care utilization or worsening outcomes.

**Keywords:** Medicare, Costs, Bundled payments, Post-acute, Health policy

\* Correspondence: [harlan.krumholz@yale.edu](mailto:harlan.krumholz@yale.edu)

<sup>2</sup>Center for Outcomes Research and Evaluation, Yale-New Haven Hospital, 1 Church Street, Suite 200, New Haven, CT 06510, USA

<sup>6</sup>Section of Cardiovascular Medicine, Department of Internal Medicine, Yale School of Medicine, 789 Howard Avenue, FMP 330, New Haven, CT 06520, USA

Full list of author information is available at the end of the article



## Background

Although United States (U.S.) inpatient costs are declining over time [1], post-hospitalization costs have more than doubled since 2001 [2] and account for a large portion of U.S. health care spending growth [3]. As a result, the post-acute care period has become a focus of programs designed to reduce waste and control costs [4, 5]. The move towards payment models that extend beyond hospitalizations to the entire episode of a patient's illness [6] has only increased the interest in understanding the relationship between resource utilization during a hospitalization and utilization during the post-acute care period.

Although prior studies on episodes of care have shown that post-acute care is a substantial contributor to spending variation and therefore could be a target for cost savings [3, 7–9], there have been few studies on the relationship between in-hospital and post-acute resource utilization [10]. Such information would be valuable, as it is plausible that cost reductions in the hospitalization period could affect spending in the post-acute period. For example, hospitals that provide higher intensity care and spend more during the hospitalization might have a beneficial effect on patients after they are discharged, reducing the amount of post-acute care resources required. In contrast, hospitals that spend less during the hospitalization may discharge patients who require greater post-acute care utilization. Alternatively, it is also possible that there is no appreciable relationship between in-hospital and post-acute resource utilization. Knowledge about the relationship between early and late use of resources may help inform strategies to reduce resource utilization in the post-acute period while maintaining or improving health outcomes.

To examine these issues, we studied Medicare fee-for-post-discharge payment and 30-day outcomes measures are constructed [11]. We examined the association between in-hospital and post-discharge resource utilization, compared patient outcomes across hospitals stratified according to their in-hospital costs, and assessed patterns of in-hospital and post-acute resource utilization across hospitals to understand how hospitals use their resources.

## Methods

### Data sources

We used the Premier Healthcare Database (PHD) by Premier Inc., Charlotte, North Carolina, which contains highly specific information on in-hospital administrative and operational data from a large network of hospitals and ancillary providers nationwide, representing more than 330 million discharges. PHD contains a date-stamped log of all billed items and resource utilization during hospitalizations (e.g., medications and laboratory, diagnostic, and therapeutic services), diagnosis and procedure codes, and

patient and hospital characteristics [12, 13]. Patient-level data were de-identified in accordance with the Health Insurance Portability and Accountability Act.

We also utilized the Centers for Medicare & Medicaid Services (CMS) publicly reported measures on hospital-specific, 30-day risk-standardized payment (RSP), 30-day risk-standardized mortality rate (RSMR), and 30-day risk-standardized readmission rate (RSRR) for 2010–2013 for AMI [11]. Of note, the 30-day RSP and RSMR measures start at patient admission. In addition, we obtained hospital-specific data on average per-patient payment for various post-acute care settings after discharge from the index hospitalization for AMI such as emergency department, readmission, outpatient physician visits, and skilled nursing facility. The publicly reported data were linked to PHD by hospital ID. Notably, some hospitals in our cohort were not linkable to the CMS public reporting data as they did not have a CMS hospital ID, shared a hospital ID with another hospital, or did not report a given measure. In brief, the RSP measure is determined by calculating payments during the 30-day AMI episode of care after removing payment adjustments for policy initiatives and geographic factors to better reflect variation in resource utilization related to clinical care. Payments were then risk-standardized based on patient clinical characteristics in the AMI hospitalization and the 12 months before hospitalization [14, 15].

The study was conducted under a collaborative contract with Premier, Inc. regarding protection of privacy of hospitals and other providers, which permits the linking of hospital data. We do not report any results that would reveal hospital identities. The Yale University Human Investigation Committee exempted this study protocol as defined by the Office of Human Research Protections.

### Study population

We included hospitals that were revascularization-capable, had intensive care units (ICUs)/cardiac care units (CCUs), and had 25 or more eligible AMI hospitalizations between July 2010 and June 2013. The revascularization-capable hospitals are defined as those that performed  $\geq 5$  percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG) procedures for AMI from 2010 to 2013.

An eligible AMI hospitalization is defined as one in which the patient was 65 years or older at admission with a principal diagnosis of new AMI, i.e., International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes 410.xx (except for 410.x2 (AMI, subsequent episode-of-care)) and was not transferred from or to another acute care facility, did not involve a heart transplant or left ventricular assist device implantation, and had a length of stay greater than 1 day (unless death occurred). The age restriction was chosen to be consistent with CMS publicly reported mortality, readmission, and payment measures.

### Hospital characteristics

Hospital characteristics were available from PHD, including: number of beds, teaching hospital status, service of a rural or urban population, and the U.S census region where a hospital is located.

### Hospital costs and outcomes

We measured in-hospital cost for each hospitalization, the methods of which are detailed elsewhere [13]. Briefly, we used hospital cost accounting data available in the Premier database and removed geographic variation in input prices by applying average unit cost across all hospitals for each service item by year. All cost estimates were adjusted to 2013 U.S. dollars. To reduce the influence of outliers, we Winsorized cost data within each year at the 0.5th and 99.5th percentiles. Hospital-level risk-standardized in-hospital RSC (hereafter in-hospital RSC) was calculated by adjusting for patient case-mix using hierarchical generalized linear models [13]. Candidate risk factors include patient age, comorbidities [16], prior revascularization procedures (CABG and PCI), and hospital transfer-in and transfer-out rate for all patients with AMI during the study period. Stepwise selection was used to select final risk-adjustors.

To assess resource utilization after hospitalization for AMI, we also estimated a risk-standardized, post-acute payment measure (post-acute RSP) for each hospital. Because PHD is de-identified, we could not link a patient's hospitalization with his/her post-discharge care information and hence could not perform direct risk adjustment for the post-acute payment measure. Therefore, we used a "pseudo" risk-standardization approach via prorating. Specifically, for each hospital, using CMS data, we first calculated its average per patient post-acute care payment (not risk-standardized) and the proportion of its total 30-day episode of care payment (not risk-standardized) that occurred after discharge from the index hospitalization. Then we multiplied the hospital's CMS publicly reported 30-day RSP (risk-standardized) by this proportion, and used this estimate as our measure for post-acute RSP. For example, for a hospital, its 30-day RSP is \$15,000, and its unadjusted average per patient in-hospital and post-acute payments are \$18,000 and \$6000, respectively. We then calculate the post-acute RSP for this hospital as  $15,000 * [6000 / (18,000 + 6000)] = \$3750$ .

In addition, we assessed in-hospital mortality for each hospitalization. Hospital-level risk-standardized, in-hospital mortality (in-hospital RSMR) was estimated using hierarchical generalized linear models with adjustment for patient case-mix. We also linked hospitals in our sample to 2010–2013 CMS hospital-specific public reporting data on 30-day RSMRs and 30-day RSRRs, as PHD is de-identified and we could not measure post-discharge outcomes for individual hospitalizations.

### Hospital practice

For each hospitalization, we examined whether the patient was admitted to an ICU, the length of ICU/CCU stay in days, and the overall length of hospital stay; all hospitals in our sample had ICUs and CCUs. We also examined whether patients received any interventional cardiac therapies, such as cardiac catheterization, CABG surgery, PCI procedures, or intra-aortic balloon pump (IABP). Receipt of revascularization was determined based on ICD-9-CM procedure codes and hospital charge codes for PCI or CABG services (Additional file 1: Appendix A). Receipt of IABP was based on ICD-9-CM code 37.61. Use of ICU care, number of ICU days, and cardiac catheterizations were identified based on hospital charge codes (see Additional file 1: Appendix B for list of codes).

### Statistical analysis

Hospitals were divided into tertiles based on in-hospital RSC, namely low, medium, and high risk-standardized cost tertiles. We utilized frequency (proportion) to characterize categorical variables, and median (interquartile range: IQR) or mean (standard deviation: SD) to characterize continuous variables. Differences in patient and hospital characteristics across in-hospital RSC tertiles were assessed using chi-square tests for categorical variables and Kruskal-Wallis tests for continuous variables. In addition to assessing the post-acute RSP by in-hospital RSC tertiles, as a sensitivity analysis we assessed the Pearson correlation between the in-hospital RSC and post-acute RSP as another approach.

We also calculated and compared mean per-patient cost in various in-hospital cost departments (e.g., cardiac procedures, ICU/CCU, laboratory) and per-patient payment in different post-acute care settings (e.g., emergency department, skilled nursing facility (SNF), readmission) by in-hospital risk-standardized cost tertiles. We used Wilcoxon analyses to test whether there is overall association between mean cost in each cost department or post-acute care setting and in-hospital RSC tertiles. We used the Dwass, Steel, Critchlow-Fligner (DSCF) method for multiple comparison analysis [17–19]. We also assessed the association between in-hospital ICU/CCU use and post-discharge SNF payment by regressing logarithmic hospital average SNF payment over hospital ICU/CCU use rate (proportion of patients using ICU/CCU) and average days in ICU/CCU, because they are settings with higher, variable resource utilization [20]. We conducted a sensitivity analysis to determine whether vital status contributed to differences in post-acute payments. *P*-values < 0.05 were deemed statistically significant. Analyses were performed using SAS version 9.4.

Our study has limitations. First, it is a cross-sectional study, so we cannot draw causal inferences regarding whether change in in-hospital RSCs would affect post-

hospital payments. Future studies with longitudinal data would help provide additional insights. Second, our findings may not be broadly generalizable because we studied Medicare fee-for-service beneficiaries  $\geq 65$  years of age at 326 hospitals within the PHD. Third, there are limitations in using claims data for measurement of risk factors and risk standardization, but we utilized established risk models for adjustment that have been validated for claims [14, 21, 22]. Fourth, due to the de-identified nature of our data, we could not track resource utilization or outcomes for individual patients after discharge from the initial hospitalization. Therefore, we relied on CMS publicly reported measures to approximate longer-term patient outcomes and post-discharge resource utilization. We also had to use hospital-level post-acute payment data instead of patient-level linked in-hospital and post-acute cost data. Although this approach may not accurately reflect patient outcomes in our analytic sample or the exact costs of post-acute care, it should provide a reasonable approximation to help understand patterns of care and resource utilization. Furthermore, we were not able to link all of the Premier hospitals to CMS measures, which may affect the external validity of the study. Finally, we measured resource utilization via hospital costs for inpatient care and payments for post-acute care. While we recognize that they represent 2 different perspectives (provider vs. payer) of the burden on our healthcare system, they are a fair representation of the resources consumed in the respective settings.

## Results

### Variation in Total in-hospital RSC

Initially, there were 123,002 AMI patients in 326 hospitals in our cohort; after excluding non-Medicare patients and hospitals with less than 25 AMI patients during the study period, 82,008 patients hospitalized with AMI at 326 eligible hospitals remained (Additional file 1: Figure S1). There were 108 hospitals (24,799 patients with AMI) in the low in-hospital RSC tertile, 109 hospitals (28,198 patients) in the medium in-hospital RSC tertile, and 109 hospitals (29,011 patients) in the high in-hospital RSC tertile (Table 1, Additional file 1: Table S1). The median (range) of each hospital's mean per-patient in-hospital RSC for the low-, medium-, and high-cost tertiles was \$16,257 (\$13,097–\$17,648), \$18,544 (\$17,663–\$19,875), and \$21,831 (\$19,923–\$31,296), respectively.

### Variation in components of in-hospital resource utilization

In terms of utilization, patients in higher in-hospital RSC hospitals appeared to receive more intensive care during the inpatient stay (Table 2). The proportion of patients receiving ICU care was 44.0, 46.7, and 58.6%, respectively, at

low, medium, and high in-hospital RSC hospitals ( $p < 0.001$ ); and the IQRs for number of days in ICUs were 1–3, 1–4, and 2–5 ( $p < 0.001$ ), respectively. Similarly, patients in the higher in-hospital RSC hospitals were more likely to receive interventional therapies, such as proportion of patients receiving cardiac catheterization, CABG, PCI, and IABP ( $p < 0.001$  for all). Median length of stay in days (IQR) was 4 (2–6), 4 (3–7), and 4 (3–8) across in-hospital RSC tertiles ( $p < 0.001$ ).

The departments with the greatest resource utilization measured by per person mean spending were ICU/CCU, other room and board, supply, and cardiac procedures for all 3 in-hospital RSC tertiles (Fig. 1a, Additional file 1: Tables S2 and S3, Additional file 1: Figures S2–S4). However, departments that varied the most across low, medium, and high in-hospital RSC tertiles were ICU/CCU, supplies, and pharmacy. The high in-hospital RSC hospitals spent notably more on ICU/CCU and pharmacy than other hospitals: \$2339, \$2876 and \$4285 per person for ICU/CCU spending and \$1408, \$1572, and \$2578 per person for pharmacy spending, for the low, medium, and high in-hospital RSC hospital tertiles, respectively. Mean per-patient spending on cardiac procedures, however, was similar among the 3 tertiles (\$2008, \$2153, and \$2074, respectively).

### Variation in hospital characteristics and outcomes

There were no significant differences in hospital characteristics across the 3 in-hospital RSC tertiles (Tables 1 and 2). There was also no significant difference in in-hospital RSMRs (Table 3). Of the 326 eligible hospitals in our cohort, 276 and 277 were successfully linked to the CMS 30-day RSMR and RSRR measures, respectively. There was no statistically significant difference in 30-day RSMRs among the in-hospital RSC tertiles, but 30-day RSRRs differed significantly across the in-hospital RSC tertiles ( $p = 0.03$ ). Hospitals in tertiles with higher in-hospital RSC had higher RSRRs: the medians (IQRs) of RSRRs were 17.5% (16.8–18.4%), 17.8% (17.1–18.6%), and 18.0% (17.5–18.6%), respectively, for low, medium, and high in-hospital RSC tertiles, and the trend test treating in-hospital RSC tertile ranks as numeric variables had a  $p$ -value of 0.005.

### Relationship between in-hospital and post-acute care resource utilization

Of the 274 hospitals that were linked to the CMS AMI payment measure, there was no statistically significant difference in risk-standardized post-acute payments across in-hospital RSC tertiles: median (IQR) = \$5014 (4295–6051), \$4980 (4349–5931) and \$4922 (4056–5457) for the low ( $n = 90$ ), medium ( $n = 98$ ), and high ( $n = 86$ ) in-hospital RSC tertiles, respectively ( $p = 0.21$ ) (Table 3). However, there was a modest yet statistically significant negative correlation between in-hospital RSC and

**Table 1** Characteristics of hospitals in different tertiles of risk-standardized in-hospital cost

Hospital Characteristics	Overall	Risk-Standardized In-Hospital Cost Tertile			p-value
		Low (\$13,097-17,648)	Medium (\$17,663-19,875)	High (\$19,923-31,296)	
Total no. of hospitals	326	108	109	109	
AMI 3-year volume					0.06
< 146 (low)	33.4	42.6	25.7	32.1	
146–284 (medium)	33.1	30.6	39.5	29.4	
> 284 (high)	33.4	26.9	34.9	38.5	
Beds					0.09
< 257 (low)	33.1	43.5	26.6	29.4	
257–409 (medium)	33.4	27.8	37.6	34.9	
> 409 (high)	33.4	28.7	35.8	35.8	
Teaching hospitals					0.60
Yes	34.7	35.2	31.2	37.6	
No	65.3	64.8	68.8	62.4	
Census regions					0.38
Midwest	24.9	31.5	22.9	20.2	
Northeast	13.5	12.0	11.9	16.5	
South	42.3	36.1	44.0	46.8	
West	19.3	20.4	21.1	16.5	
Population served					0.37
Urban	85.3	82.4	89.0	84.4	
Rural	14.7	17.6	11.0	15.6	

**AMI** Acute myocardial infarction

risk-standardized post-acute care payments (correlation coefficient =  $-0.13$ ,  $p = 0.04$ , Additional file 1: Figure S5). On average, every \$1000 increase in in-hospital RSC was associated with \$50 less post-acute care payments ( $p = 0.04$ ).

Three settings displayed the highest mean per-patient post-acute care payments: readmission, SNFs, and

non-acute inpatient care (Additional file 1: Table S5, Fig. 1b); these findings were consistent in a sensitivity analysis stratified by vital status (Additional file 1: Table S4, Additional file 1: Figures S6a and b).


For the low, medium and high in-hospital RSC tertiles, respectively, the mean per-patient post-acute payments were \$1534, \$1633, and \$1596 for readmissions ( $p =$

**Table 2** Utilization characteristics of patients hospitalized with acute myocardial infarction by hospital risk-standardized in-hospital cost levels

Utilization Characteristics	Overall	Risk-Standardized In-Hospital Cost Tertile			p-value
		Low (\$13,097-17,648)	Medium (\$17,663-19,875)	High (\$19,923-31,296)	
Total no. of hospitals	326	108	109	109	
Selected inpatient care during index episode					
Index episode length of stay, median (IQR), days	4 (3–7)	4 (2–6)	4 (3–7)	4 (3–8)	< 0.001
ICU admission during index episode, %	50.1	44.0	46.7	58.6	< 0.001
ICU days if admitted, median (IQR)	2 (1–4)	2 (1–3)	2 (1–4)	3 (2–5)	< 0.001
Interventional cardiac therapies (%)					
Cardiac catheterization	63.9	61.4	64.9	64.9	< 0.001
CABG surgery	7.8	6.9	8.0	8.5	< 0.001
Percutaneous coronary intervention	41.4	40.1	41.2	42.8	< 0.001
Intra-aortic balloon pump	4.7	4.0	4.7	5.3	< 0.001

CABG Coronary artery bypass graft, ICU Intensive care unit, IQR Interquartile range





**Fig. 1 a and b.** Resource Utilization by In-Hospital Cost Tertiles. **a** Relative Contribution of Service Categories to In-Hospital Costs by Hospitals in Different In-Hospital Cost Tertiles. Green bars indicate the low in-hospital cost tertile, blue bars the medium in-hospital cost tertile, and red bars the high in-hospital cost tertile. ICU: intensive care unit; CCU: coronary care unit; OR: operating room. **b** Relative Contribution of Service Categories to Post-Acute Payments by Hospitals in Different In-Hospital Cost Tertiles. Green bars indicate the low in-hospital cost tertile, blue bars the medium in-hospital cost tertile, and red bars the high in-hospital cost tertile

**Table 3** Outcomes and post-acute care costs of patients hospitalized with acute myocardial infarction by hospital risk-standardized in-hospital cost levels

Variable	Total	Inpatient Cost Level			p-value
		Low (\$13,097-17,648)	Medium (\$17,663-19,875)	High (\$19,923-31,296)	
<b>Outcomes</b>					
In-hospital mortality rate	8.2 (7.9–8.8)	8.2 (7.8–8.7)	8.3 (7.8–8.8)	8.2 (7.8–8.7)	0.90
30-day risk-standardized mortality rate (N = 276) <sup>a</sup> , median (IQR)	14.7 (13.7–15.7)	14.9 (13.9–15.7)	14.6 (13.7–15.6)	14.7 (13.6–15.5)	0.42
30-day risk-standardized readmission rate (N = 277) <sup>a</sup> , median (IQR)	17.8 (17.1–18.5)	17.5 (16.8–18.4)	17.8 (17.1–18.6)	18.0 (17.5–18.6)	0.03
Hospital risk-standardized post-acute care payment (N = 274) (\$), median (IQR)	4964 (4250–5931)	5014 (4295–6051)	4980 (4349–5931)	4922 (4056–5457)	0.21

<sup>a</sup>Data provided for the number of hospitals (N) that could be linked to Centers for Medicare & Medicaid Services measures. By cost level, for risk-standardized mortality rates, there were 89, 99, and 88 hospitals; for risk-standardized readmission rates, 90, 99, and 88 hospitals; for risk-standardized post-acute care payment, 90, 98, and 86 hospitals  
 Statistics are percent unless stated otherwise

0.001), \$1535, \$1542, and \$1303 for SNFs ( $p < 0.001$ ), and \$533, \$649, and \$714 for the non-acute inpatient setting ( $p < 0.001$ ). The highest in-hospital RSC tertile had lower SNF and home health agency use and payments, with spending in other post-acute settings similar across cost tertiles (Additional file 1: Tables S5 and S6, Additional file 1: Figures S7-S9).

#### **Relationship between in-hospital ICU use and post-acute care resource utilization on SNF**

Given the variation in both ICU utilization and SNF post-acute payments across in-hospital RSC tertiles, we assessed the association between ICU use rate and the logarithm of mean payments for SNF utilization among the 274 hospitals. For every 10% increase in use of ICUs, there was a 0.7% decrease in SNF payment ( $p < 0.003$ ). Moreover, for each additional day in ICU, there was a 13% decrease in SNF payment ( $p < 0.002$ ).

#### **Discussion**

In our study of patients hospitalized with AMI, greater resource utilization during the hospitalization was not associated with substantially lower resource utilization during the post-acute period. Hospitals with higher in-hospital RSC spent more on ICU/CCU stays, pharmacy, and supplies, and had lower payments for SNFs and home health agencies. Across in-hospital RSC tertiles, hospitals had similar in-hospital and 30-day mortality rates, but those in the highest in-hospital RSC tertile had higher 30-day readmission rates. Although patients at hospitals with greater inpatient resource utilization were more likely to receive cardiac procedures, there were no marked differences in other treatment characteristics across in-hospital RSC tertiles.

Prior research on bundled and episode-of-care payments shows that post-acute care is a substantial contributor to spending variation [3, 7–9]. Our work extends this literature as the first large, contemporary study to assess the association of resource utilization patterns between in-hospital and post-acute care. A previous study analyzing patient-level data showed that among Medicare beneficiaries with AMI, patient disease severity at hospital discharge, discharging hospital for-profit ownership, and discharging hospital provision of home health services were important predictors of post-acute service utilization; however, the study did not have access to detailed in-hospital cost data and the ability to study in-hospital RSCs or post-acute spending, or conduct a hospital-level analysis [23]. Another study of patients with heart failure and pneumonia showed that those initially seen in lower cost hospitals had lower spending associated with re-admission within 6 months of the index hospitalization, but the study did not assess spending across the various post-acute care settings [24].

Several studies have assessed the association between hospital spending and outcomes, but the results have been mixed [25–31]. Our study adds to the literature that shows no association between increased hospital resource utilization and better outcomes. In fact, readmission was highest in hospitals with higher in-hospital resource utilization. Further research is needed to determine whether spending reduction is a means to improve outcomes.

There are several possible explanations for our findings. There may be a core set of services that are necessary to ensure that people have better outcomes and lower post-acute utilization that are provided by all hospitals, and the increased resource utilization by high-cost hospitals may be in lower yield areas. Second, there may be certain resource utilization “habits” in higher spending hospitals that inflate costs and could be curtailed. For example, one study in the Premier Healthcare Database has shown that some hospitals may perform more angiographies that ultimately do not result in higher rates of revascularization, which may partly explain the greater utilization in those hospitals [32]. Moreover, studies have shown that PCI procedures are associated with increased readmission rates [33], with many of these readmissions being preventable [34], which may contribute to the higher readmission rates in our study. Nonetheless, the prior cited study using the PHD found no differences in readmissions [32], with hospitals that do more being unlikely to have a lower threshold of readmission, and there is a large range of use that is not associated with outcomes [20, 35]. We could not, with this design, explain why higher costs may be associated with higher rates of re-admission. If the relationship is true, and not confounded, it may be related to the fact that excess tests, procedures, and medications increase the likelihood of adverse events. In addition, longer length of stay and intensity of the stay, including use of the intensive care unit, could provide more exposure to the allostatic stress of the hospitalization, leading to greater downstream vulnerability [36]. It may be that there is a sweet spot of intensity for each patient, and places that tend to do more may compromise downstream recovery. It is also possible that higher cost hospitals have stricter definitions of AMI and therefore have higher in-hospital utilization rates and, given their severity, patients may be more likely to be readmitted, though the magnitude of differences at the hospital level suggests that differences in severity alone are unlikely to explain our findings. Finally, there may be other determinants of post-acute care utilization that were not captured in our study, an important area for future research. For example, through linking patients in the in-hospital and post-acute settings, one might explore which particular categories of in-hospital expenditures are associated with the risk for greater utilization of

post-discharge services – and whether they are markers for higher risk of utilization or something that mediates the needs for greater utilization. In particular, there is a need for studies surrounding how the length and intensity of the hospitalization relates to the recovery after discharge.

Reducing unnecessary post-acute resource utilization is a goal as the government and health care industry try to contain costs and improve the value of care, and new models of payment that focus on the entire episode of patient care, such as the Bundled Payments for Care Improvement Initiative [6], are an important first step. Our study suggests the absence of a strong association between higher in-hospital resource utilization and reductions in post-discharge resource utilization. Furthermore, post-hospitalization outcomes were not better. These findings suggest that greater resource utilization at the higher cost hospitals may fail to provide a meaningful benefit for patients and the health system. Moving forward, higher spending hospitals may target areas of higher cost during a hospitalization without increasing post-acute care utilization or worsening outcomes. The insight, however, requires the ability to benchmark.

## Conclusions

In our study of a large number of hospitals across the country, greater resource utilization for patients hospitalized with AMI during the inpatient stay was not associated with substantially lower 30-day post-acute care resource utilization or better outcomes. In the context of novel care and payment models that extend across the entire episode of care, where entities will be rewarded for reducing spending, our findings suggest that it may be possible for higher cost hospitals to improve efficiency in care without increasing post-acute care utilization or jeopardizing outcomes of care.

## Additional file

**Additional file 1: Appendix A.** List of ICD-9 and Standard Charge Codes for Identifying Percutaneous Coronary Intervention and Coronary Artery Bypass Graft Procedures. Appendix B. Standard Charge Codes for Identifying ICU/CCU and Catheterization. **Table S1.** Characteristics of Patients Hospitalized with Acute Myocardial Infarction by Hospital Risk-Standardized In-Hospital Cost Levels. **Table S2.** In-Hospital Resource Utilization: Mean Per-Patient Cost in Each Department. **Table S3.** In-Hospital Resource Utilization: Mean per Patient Number of Items. **Table S4.** Patient Costs by In-Hospital Cost Ranks and Vital Status. **Table S5.** Mean Per Patient Post-Acute-Care Payment at Different Care Settings. **Table S6.** Percentage of Patients Using Various Post-Acute-Care Services by Cost Groups. **Figure S1.** Flowchart of Exclusion Criteria. **Figure S2.** Number of Items Ordered during a Hospitalization by In-Hospital Cost Tertile. **Figure S3.** Mean In-hospital Resource Utilization per User by In-Hospital Cost Tertiles. **Figure S4.** Relative Contribution of Service Categories to In-Hospital Costs by In-Hospital Cost Tertiles. **Figure S5.** Correlation Between Hospital Risk-Standardized In-Hospital Cost (\$) and Hospital Risk-Standardized Post-Acute Payments (\$). **Figure S6.a** In-Hospital Spending Among Survivors. **b** In-Hospital Spending Among the Expired. **Figure S7.** Percentage of Patients Using Different Post-Acute-Care Services by In-Hospital Cost Tertiles. **Figure S8.** Mean User Post-

Acute Payments by In-hospital Cost Tertiles. **Figure S9.** Relative Contribution of Service Categories to Post-Acute Payments by In-Hospital Cost Tertiles. (DOCX 178 kb)

## Acknowledgements

Not applicable.

## Funding

Dr. Lagu is supported by grant K01HL114745 from the National Heart, Lung, and Blood Institute, Bethesda, MD. Dr. Desai is supported by grant K12 HS023000–05 from the Agency for Healthcare Research and Quality. No funder had a role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

## Availability of data and materials

The data that support the findings of this study are available from Premier Inc. and the Centers for Medicare & Medicaid Services but restrictions apply to the availability of these data, which were used under specific data use agreements for the current study, and so are not publicly available. For questions regarding the data, please contact Shu-Xia Li ([shu-xia.li@yale.edu](mailto:shu-xia.li@yale.edu)).

## Authors' contributions

Each author has read and approved the final version and all made important contributions: SVN: Conception and design, analysis and interpretation, writing the article, critical revision of the article, final approval. SXL: Conception and design, analysis and interpretation, data collection, writing the article, critical revision of the article, final approval. XX: Analysis and interpretation, writing the article, critical revision of the article, final approval. LSO: Analysis and interpretation, data collection, writing the article, critical revision of the article, final approval. TL: Analysis and interpretation, writing the article, critical revision of the article, final approval. NRD: Analysis and interpretation, writing the article, critical revision of the article, final approval. KM: Analysis and interpretation, writing the article, critical revision of the article, final approval. MD: Analysis and interpretation, writing the article, critical revision of the article, final approval. JM: Analysis and interpretation, writing the article, critical revision of the article, final approval. NK: Analysis and interpretation, writing the article, critical revision of the article, final approval. HMK: Conception and design, analysis and interpretation, writing the article, critical revision of the article, final approval, overall study responsibility.

## Ethics approval and consent to participate

The study was conducted under a collaborative contract with Premier, Inc. regarding protection of privacy of hospitals and other providers, which permits the linking of hospital data. For use of Premier data, the Yale University Human Investigation Committee exempted this study protocol as defined by the Office of Human Research Protections because data was de-identified. For CMS data, institutional review board approval, including waiver of the requirement of participant informed consent, was provided by the Yale University Human Investigation Committee.

## Consent for publication

Not applicable.

## Competing interests

Dr. Krumholz is a recipient of research agreements from Medtronic and Johnson & Johnson (Janssen), through Yale, to develop methods of clinical trial data sharing; was the recipient of a grant from Medtronic and the Food and Drug Administration, through Yale, to develop methods for post-market surveillance of medical devices; received payment from the Arnold & Porter Law Firm for work related to the Sanofi clopidogrel litigation and from the Ben C. Martin Law Firm for work related to the Cook IVC filter litigation; chairs a cardiac scientific advisory board for UnitedHealth; is a participant/participant representative of the IBM Watson Health Life Sciences Board; is a member of the Advisory Boards for Element Science and for Facebook, and the Physician Advisory Board for Aetna; and is the founder of Hugo, a personal health information platform. Drs. Krumholz, Li, Xu, and Desai work under contract to the Centers for Medicare & Medicaid Services to develop



and maintain performance measures that are publicly reported. Dr. Lagu has received consulting fees from the Institute for Healthcare Improvement for her input on a project to help health systems achieve disability competence. The other authors do not have disclosures to report.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Author details

<sup>1</sup>Yale School of Medicine, 333 Cedar Street, New Haven, CT 06516, USA. <sup>2</sup>Center for Outcomes Research and Evaluation, Yale-New Haven Hospital, 1 Church Street, Suite 200, New Haven, CT 06510, USA. <sup>3</sup>Department of Obstetrics, Gynecology, and Reproductive Sciences, Yale School of Medicine, 310 Cedar Street, LSOG 205B, New Haven, CT 06510, USA. <sup>4</sup>Tufts University School of Medicine, 145 Harrison Ave, Boston, MA, USA. <sup>5</sup>Department of Medicine, Baystate Medical Center, 759 Chestnut Street, Springfield, MA 01199, USA. <sup>6</sup>Section of Cardiovascular Medicine, Department of Internal Medicine, Yale School of Medicine, 789 Howard Avenue, FMP 330, New Haven, CT 06520, USA. <sup>7</sup>Premier, Inc, 13034 Ballantyne Corporate Place, Charlotte, NC 28277, USA. <sup>8</sup>Section of General Internal Medicine, Department of Internal Medicine, Yale School of Medicine, 333 Cedar Street, New Haven, CT 06520, USA. <sup>9</sup>Department of Health Policy and Management, Yale School of Public Health, 60 College Street, New Haven, CT, USA. <sup>10</sup>Present address: Remedy Partners, 1120 Boston Post Road, Suite 3, Darien, CT 06820, USA.

Received: 19 September 2017 Accepted: 18 March 2019

Published online: 25 March 2019

## References

- Krumholz HM, Nuti SV, Downing NS, Normand SL, Wang Y. Mortality, hospitalizations, and expenditures for the Medicare population aged 65 years or older, 1999–2013. *JAMA*. 2015;314(4):355–65.
- Tian W. An all-payer view of hospital discharge to postacute care, 2013: statistical brief #205. Healthcare cost and utilization project (HCUP) statistical briefs. Rockville: Agency for Healthcare Research and Quality (US). 2016; <https://www.ncbi.nlm.nih.gov/books/NBK373736/>. Accessed 26 Aug 2018.
- Chandra A, Dalton MA, Holmes J. Large increases in spending on postacute care in medicare point to the potential for cost savings in these settings. *Health Aff (Millwood)*. 2013;32(5):864–72.
- Interim Report of the Committee on Geographic Variation in Health Care Spending and Promotion of High-Value Health Care: Preliminary committee observations. Washington, DC: The National Academies Press; 2013.
- Newhouse JP, Garber AM. Geographic variation in Medicare services. *N Engl J Med*. 2013;368(16):1465–8.
- Press MJ, Rajkumar R, Conway PH. Medicare's new bundled payments: design, strategy, and evolution. *JAMA*. 2016;315(2):131–2.
- Tsai TC, Joynt KE, Wild RC, Orav EJ, Jha AK. Medicare's bundled payment initiative: most hospitals are focused on a few high-volume conditions. *Health Aff (Millwood)*. 2015;34(3):371–80.
- Birkmeyer JD, Gust C, Baser O, Dimick JB, Sutherland JM, Skinner JS. Medicare payments for common inpatient procedures: implications for episode-based payment bundling. *Health Serv Res*. 2010;45(6):1783–95.
- Miller DC, Gust C, Dimick JB, Birkmeyer N, Skinner J, Birkmeyer JD. Large variations in Medicare payments for surgery highlight savings potential from bundled payment programs. *Health Aff (Millwood)*. 2011;30(11):2107–15.
- Mechanic R. Post-acute care—the next frontier for controlling Medicare spending. *N Engl J Med*. 2014;370(8):692–4.
- Centers for Medicare & Medicaid Services. Hospital Compare. <https://data.medicare.gov/data/hospital-compare>. Accessed 26 Aug 2018.
- Chen SI, Dharmarajan K, Kim N, et al. Procedure intensity and the cost of care. *Circ Cardiovasc Qual Outcomes*. 2012;5(3):308–13.
- Xu X, Li SX, Lin H, et al. "Phenotyping" hospital value of care for patients with heart failure. *Health Serv Res*. 2014;49(6):2000–16.
- Kim N, Bernheim SM, Ott LS, et al. An administrative claims measure of payments made for medicare patients for a 30-day episode of care for acute myocardial infarction. *Med Care*. 2015;53(6):542–9.
- Hospital-level, risk-standardized payment associated with a 30-day episode of care for AMI (Version 1.0): 2012 Measure Methodology Report. <http://www.qualitynet.org/dcs/ContentServer?c=Page&pagename=QnetPublic/Page/QnetTier4&cid=1228773321331>. Accessed 26 Aug 2018.
- Elixhauser A, Steiner C, Harris DR, Coffey RM. Comorbidity measures for use with administrative data. *Med Care*. 1998;36(1):8–27.
- Hollander M, Wolfe D. Nonparametric statistical methods, 2nd Edition. New York: Wiley; 1999.
- Dwass M. Some k-sample rank-order tests. In: Olkin I, editor. Contributions to probability and statistics. California: Stanford University Press; 1960. p. 198–202.
- Steel RG. A rank sum test for comparing all pairs of treatments. *Technometrics*. 1960;2(2):197–207.
- Safavi KC, Dharmarajan K, Kim N, et al. Variation exists in rates of admission to intensive care units for heart failure patients across hospitals in the United States. *Circulation*. 2013;127(8):923–9.
- Krumholz HM, Lin Z, Drye EE, et al. An administrative claims measure suitable for profiling hospital performance based on 30-day all-cause readmission rates among patients with acute myocardial infarction. *Circ Cardiovasc Qual Outcomes*. 2011;4(2):243–52.
- Krumholz HM, Wang Y, Mattera JA, et al. An administrative claims model suitable for profiling hospital performance based on 30-day mortality rates among patients with an acute myocardial infarction. *Circulation*. 2006; 113(13):1683–92.
- Bronskill SE, Normand SL, McNeil BJ. Post-acute service use following acute myocardial infarction in the elderly. *Health Care Financ Rev*. 2002;24(2):77–93.
- Chen LM, Jha AK, Guterman S, Ridgway AB, Orav EJ, Epstein AM. Hospital cost of care, quality of care, and readmission rates: penny wise and pound foolish? *Arch Intern Med*. 2010;170(4):340–6.
- Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, Pinder EL. The implications of regional variations in Medicare spending. Part 2: health outcomes and satisfaction with care. *Ann Intern Med*. 2003;138(4):288–98.
- Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, Pinder EL. The implications of regional variations in Medicare spending. Part 1: the content, quality, and accessibility of care. *Ann Intern Med*. 2003;138(4):273–87.
- Hussey PS, Wertheimer S, Mehrotra A. The association between health care quality and cost: a systematic review. *Ann Intern Med*. 2013;158(1):27–34.
- Lagu T, Rothberg MB, Nathanson BH, Pekow PS, Steingrub JS, Lindenaer PK. The relationship between hospital spending and mortality in patients with sepsis. *Arch Intern Med*. 2011;171(4):292–9.
- Romley JA, Jena AB, Goldman DP. Hospital spending and inpatient mortality: evidence from California: an observational study. *Ann Intern Med*. 2011;154(3):160–7.
- Romley JA, Jena AB, O'Leary JF, Goldman DP. Spending and mortality in US acute care hospitals. *Am J Manag Care*. 2013;19(2):e46–54.
- Stukel TA, Fisher ES, Alter DA, et al. Association of hospital spending intensity with mortality and readmission rates in Ontario hospitals. *JAMA*. 2012;307(10):1037–45.
- Safavi KC, Li SX, Dharmarajan K, et al. Hospital variation in the use of noninvasive cardiac imaging and its association with downstream testing, interventions, and outcomes. *JAMA Intern Med*. 2014;174(4):546–53.
- Khawaja FJ, Shah ND, Lennon RJ, et al. Factors associated with 30-day readmission rates after percutaneous coronary intervention. *Arch Intern Med*. 2012;172(2):112–7.
- Wasfy JH, Strom JB, Waldo SW, et al. Clinical preventability of 30-day readmission after percutaneous coronary intervention. *J Am Heart Assoc: Cardiovasc Cerebrovasc Dis*. 2014;3(5):e001290.
- Chen R, Strait KM, Dharmarajan K, et al. Hospital variation in admission to intensive care units for patients with acute myocardial infarction. *Am Heart J*. 2015;170(6):1161–9.
- Krumholz HM. Post-hospital syndrome—an acquired, transient condition of generalized risk. *N Engl J Med*. 2013;368(2):100–2.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

