

RESEARCH ARTICLE

Minimum inhibitory concentrations of commonly used antibiotics against *Helicobacter Pylori*: A multicenter study in South China

Xueping Huang^{1,2}✉, Yuan Liu³✉, Zhihui Lin^{1,2}, Baihe Wu³, Gaohui Nong⁴, Yushan Chen⁵, Yuping Lu^{1,2}, Xinhua Ji⁶, Xiang Zhou⁶, Biao Suo⁷, Qiuzhao Chen⁷, Jinqi Wei^{3*}

1 Department of Gastroenterology, Fujian Provincial Hospital, Fuzhou, Fujian, China, **2** Department of Gastroenterology, Shengli Clinical Medical College of Fujian Medical University, Fuzhou, Fujian, China, **3** Department of Gastroenterology, Fifth Affiliated Hospital of Sun Yat-sen University, Zhuhai, Guangdong, China, **4** Department of Microbiology, Zhuhai Health School, Zhuhai, Guangdong, China, **5** Department of Gastroenterology, Fuzhou First Hospital, Fuzhou, Fujian, China, **6** Department of Gastroenterology, Pucheng Hospital, Nanping, Fujian, China, **7** Department of Gastroenterology, Xiamen Hospital of Traditional Chinese Medicine, Xiamen, Fujian, China

✉ These authors contributed equally to this work.

* 491589958@qq.com



OPEN ACCESS

Citation: Huang X, Liu Y, Lin Z, Wu B, Nong G, Chen Y, et al. (2021) Minimum inhibitory concentrations of commonly used antibiotics against *Helicobacter Pylori*. A multicenter study in South China. PLoS ONE 16(9): e0256225. <https://doi.org/10.1371/journal.pone.0256225>

Editor: Muhammad Hussain Siddique, Government College University Faisalabad, PAKISTAN

Received: February 12, 2021

Accepted: August 2, 2021

Published: September 2, 2021

Copyright: © 2021 Huang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Considering patients' privacy and related regulations in China, we chose not to make the database public to everyone. If a researcher wants to use our raw data for scientific research purposes, he or she could apply for use with our ethics committee (fjslec@163.com).

Funding: This work was supported by the Youth Foundation of Fujian Provincial Health Commission (No. 2016-1-5), the Qihang Foundation of Fujian Medical University (No. 2016QH108) for Xueping

Abstract

Aim

To determine the minimum inhibitory concentrations (MICs) of commonly used antibiotics against *Helicobacter Pylori* (*H. pylori*) in South China and compare their resistance rates by using EUCAST breakpoints and other breakpoints.

Methods

Patients who had not previously received *H. pylori* treatment in clinical centers in South China were enrolled in this study from 2017 to 2020. Gastric biopsies were obtained for *H. pylori* culture. The MICs of amoxicillin (AMX), clarithromycin (CLA), metronidazole (MTZ), levofloxacin (LEV), tetracycline (TET) and furazolidone (FZD) were tested by broth microdilution method and assessed by two different breakpoints. ATCC43504 standard strain served as a control.

Results

A total of 208 *H. pylori* strains were isolated from patients' biopsy samples. The MICs of AMX, CLA, MTZ, LEV, TET and FZD for *H. pylori* were 0.0156-256mg/L (MIC₅₀ 0.125mg/L, MIC₉₀ 4mg/L), 0.0156- >256 mg/L (MIC₅₀ 0.0312mg/L, MIC₉₀ 64mg/L), 0.0156- >256mg/L (MIC₅₀ 8mg/L, MIC₉₀ 256mg/L), 0.0156-256mg/L (MIC₅₀ 0.25mg/L, MIC₉₀ 16mg/L), 0.0156-256mg/L (MIC₅₀ 0.0625mg/L, MIC₉₀ 4mg/L), and 0.0156- >256mg/L (MIC₅₀ 0.0312mg/L, MIC₉₀ 2mg/L), respectively. The MICs of AMX, CLA, MTZ, LEV, TET and FZD for ATCC43504 strain were 0.25mg/L, 0.0625mg/L, 64mg/L, 0.5mg/L, 1mg/L and 0.25mg/L, respectively. The resistance rate of FZD was 11.05%. The overall resistance rates

Huang, and Foundation of Fujian Provincial Hospital (No. 2014YNZD04) for Zhihui Lin.

Competing interests: The authors have declared that no competing interest exists.

according to EUCAST breakpoints and other breakpoints were 57.21% and 14.90% for AMX ($p < 0.001$), 38.94% and 38.94% for CLA ($p = 1$), 39.42% and 50.96% for MTZ ($p < 0.001$), 12.98% and 10.58% for TET ($p = 0.025$), 35.10% and 35.10% for LEV ($p = 1$), respectively.

Conclusions

Our results demonstrate that AMX, FZD, and TET, but not MTZ, CLR or LEV, showed good anti-*H. pylori* activity *in vitro* in South China. When different breakpoints were used, similar results were found with CLA, and LEV, but not with AMX, MTZ, or TET.

Introduction

Helicobacter pylori (*H. pylori*) is a spiral-shaped, gram-negative, and microaerophilic bacterium, which specifically colonizes the gastric epithelia and infects approximately half of the population worldwide [1]. In general, the infection rate of *H. pylori* is higher in developing countries compared to developed countries [2]. *H. pylori* is an important etiology of global health problems because it can cause chronic gastritis, dyspepsia, peptic ulcers, gastric malignancies, and extragastric diseases [3, 4]. Eradication of *H. pylori* can effectively alleviate symptoms in functional dyspepsia, heal peptic ulcers, and improve atrophic gastritis and intestinal metaplasia [5]. Specifically, since 1994, *H. pylori* has been identified as a class-A carcinogen by the International Agency for Research on Cancer (IARC) [6]. Eradication of *H. pylori* has a preventive effect on gastric cancers [7] and has been proven to be efficient in treating nearly 75% of early mucosa-associated lymphoid tissue (MALT) lymphoma [8]. Therefore, early treatment of *H. pylori* is recommended for all symptomatic patients [9].

The first-line treatment for the eradication of *H. pylori* consists of 1 proton pump inhibitor (PPI) and 2 antibiotics (amoxicillin, clarithromycin or metronidazole) [1]. Unfortunately, the eradication rate using the above-mentioned treatment plan has declined over the last decade to less than 90% [10, 11], mainly due to a rapid increase in antibiotic resistance, especially against clarithromycin and metronidazole [12, 13]. The prevalence of *H. pylori* antibiotic resistance significantly varies from country to country and between regions within the same country [14, 15]. Therefore, it is essential to conduct local surveillance of antibiotic resistance.

Before implementation of this study, there were limited data regarding the minimum inhibitory concentrations (MICs) of *H. pylori* resistance to antibiotics in South China. As to breakpoints proposed for commonly used antibiotics against *H. pylori*, there are different guidelines including European Committee on Antimicrobial Susceptibility Testing (EUCAST), British Society for Antimicrobial Chemotherapy (BSAC), Clinical and Laboratory Standards Institute (CLSI), et al. In this study, we aimed to investigate the MICs of commonly used antibiotics to *H. pylori* isolated from patients in South China and compare the resistance rates when using different breakpoints to provide guidance on prescription of antibiotics.

Methods

Patients

Patients who had dyspeptic symptoms and received gastroscopy in the following 4 hospitals in South China from January 4th, 2017 to October 1st, 2020 were enrolled in this study: Fujian Provincial Hospital (Fuzhou, Fujian, China), the Fifth Affiliated Hospital of Sun Yat-Sen

University (Zhuhai, Guangdong, China), Xiamen Hospital of Traditional Chinese Medicine (Xiamen, Fujian, China), and Pucheng Hospital (Nanping, Fujian, China). Exclusion criteria were as follows: (1) patients who had used bismuth salts, proton pump inhibitors, antibiotics, or H2 receptor antagonists a month before entrance of the study; (2) patients who had undergone surgery; (3) patients who had malignant tumors; (4) patients who had contraindications for gastroscopy.

Isolation, identification, and antibiotic susceptibility tests of *H. pylori*

One gastric mucosal specimen was taken from the antrum of the stomach in each patient during gastroscopy to detect the presence of *H. pylori* with a rapid urease test. If the rapid urease test was positive, another specimen from the antrum was collected, stored, and then cultured in the liquid medium for *H. pylori* containing brain heart immersion broth and 10% calf serum (Zhuhai Yeoman Bioengineering Products Factory). Organisms were identified as *H. pylori* if isolates demonstrated curved gram-negative rods along with positive urease, catalase, and oxidase reactions. The MICs of *H. pylori* to commonly used antibiotics including amoxicillin (AMX), clarithromycin (CLA), metronidazole (MTZ), levofloxacin (LEV), tetracycline (TET) and furazolidone (FZD) were determined by broth microdilution method. The standard strain ATCC43504 (provided by Shanghai Bioplus Biotechnology Company) served as a control. The culture suspension was inoculated onto plates with turbidity adjusted to McFarland standard of 2.0. The antibiotic powders of AMX, CLA, and LEV were purchased from BioDuly Biotechnology Co.Ltd. (Nanjing, China), and MTZ, TET and FZD were from Meilun Biotechnology Co. Ltd. (Dalian, China).

The resistance were defined by EUCAST [16] and other breakpoints [17–19] as follows: > 0.12 mg/L and ≥ 2 mg/L for AMX, > 0.5 mg/L and ≥ 1 mg/L for CLA, > 8 mg/L and ≥ 8 mg/L for MTZ, > 1 mg/L and ≥ 2 mg/L for LEV, > 1 mg/L and ≥ 4 mg/L for TET, respectively. The resistance breakpoint to FZD was set at > 2 mg/L [20].

Statement of ethics

This study was approved by the independent Ethics Committees of the Fujian Provincial Hospital (Approval No. K2016-11-028) and performed following the World Medical Association Declaration of Helsinki. Written informed consent was obtained from each patient. Authors had access to information that could identify individual participant during and after data collection.

Statistical analysis

Data were analyzed using SPSS 22.0 software (IBM, Armonk, NY, USA). Percentages were used to describe the antibiotic resistance rates of *H. pylori* isolates. The Chi-Square test was used to compare the drug resistance rates calculated by two breakpoint systems. A P-value of < 0.05 was considered statistically significant.

Results

Patient characteristics

A total of 208 patients were included in this study. The percentages of male and female were 58.2% ($n = 121$) and 41.8% ($n = 87$), respectively. The mean age was 45 years old (range: 18–77 years old). Chronic superficial gastritis, the predominant endoscopic finding, was present in 32.2% of patients ($n = 67$), chronic atrophic gastritis in 26.4% ($n = 55$), peptic ulcers in 23.1%

of patients (n = 48), and erosive gastritis in 18.3% (n = 38). Patient characteristics and endoscopic data are summarized in [Table 1](#).

The MICs of commonly used antibiotics to *H. pylori*

A total of 208 *H. pylori* strains were isolated from the biopsy samples of gastric mucosa. The MICs of AMX, CLA, MTZ, LEV, TET and FZD for the *H. pylori* isolates were 0.0156-256mg/L (MIC₅₀ 0.125mg/L, MIC₉₀ 4mg/L), 0.0156- >256 mg/L (MIC₅₀ 0.0312mg/L, MIC₉₀ 64mg/L), 0.0156- >256mg/L (MIC₅₀ 8mg/L, MIC₉₀ 256mg/L), 0.0156-256mg/L (MIC₅₀ 0.25mg/L, MIC₉₀ 16mg/L), 0.0156-256mg/L (MIC₅₀ 0.0625mg/L, MIC₉₀ 4mg/L), and 0.0156- >256mg/L (MIC₅₀ 0.0312mg/L, MIC₉₀ 2mg/L), respectively ([Fig 1](#)). The MICs of AMX, CLA, MTZ, LEV, TET and FZD for ATCC43504 strain were 0.25mg/L, 0.0625mg/L, 64mg/L, 0.5mg/L, 1mg/L and 0.25mg/L, respectively ([Table 2](#)).

The overall resistance rates of *H. pylori* between two different breakpoints

The resistance rate of *H. pylori* to FZD was 11.05%. The overall resistance rates according to EUCAST breakpoints and other breakpoints were 57.21% and 14.90% for AMX, respectively ($p < 0.001$), 38.94% and 38.94% for CLA, ($p = 1$), 39.42% and 50.96% for MTZ ($p < 0.001$), 12.98% and 10.58% for TET ($p = 0.025$), 35.10% and 35.10% for LEV ($p = 1$) ([Table 3](#)).

The multiple resistance rates of *H. pylori* between two different breakpoints

When comparing the use of EUCAST breakpoints to other breakpoints, monoresistance rates were 21.63% and 25.96% ($p = 0.163$), double resistance rates were 17.31% and 12.50% ($p = 0.121$), triple resistance rates were 10.10% and 12.50% ($p = 0.405$), and multiple resistance rates were 19.24% and 14.42% ($p = 0.006$), respectively ([Table 4](#)).

Impact of diseases on antibiotic resistance of *H. pylori* in South China

A significant difference was detected in the resistance of *H. pylori* to LEV among different disease groups ($p < 0.05$). The MIC of LEV was highest in erosive gastritis ($\log_2 \mu = 0.47$), followed by chronic superficial gastritis ($\log_2 \mu = -1.16$), chronic atrophic gastritis ($\log_2 \mu = -1.42$), and peptic ulcers ($\log_2 \mu = -1.81$). Additionally, the MICs of erosive gastritis were also highest among the 4 disease groups for the other 5 antibiotics, although they were not statistically different ($p > 0.05$) ([Fig 2](#)).

Table 1. Patient characteristics and endoscopic data.

	Descriptive statistics
Age, mean (range), y	45 (18–77)
Sex, n (%)	
Male	121 (58.2%)
female	87 (41.8%)
Endoscopic data, n (%)	
erosive gastritis	38 (18.3%)
peptic ulcers	48 (23.1%)
chronic superficial gastritis	67 (32.2%)
chronic atrophic gastritis	55 (26.4%)

<https://doi.org/10.1371/journal.pone.0256225.t001>




Fig 1. MIC distributions of common used antibiotics in *H. pylori* in South China. Arrows indicates the EUCAST (↓) or other (⇓) resistance breakpoint. A is for amoxicillin (AMX), B is for clarithromycin (CLA), C is for metronidazole (MTZ), D is for levofloxacin (LEV), E is for tetracycline (TET), F is for furazolidone (FZD).

<https://doi.org/10.1371/journal.pone.0256225.g001>

Impact of age on antibiotic resistance of *H. pylori* in South China

Patients were divided into two groups according to age (<45 years old vs. ≥45 years old). No significant difference in the resistance of *H. pylori* to all tested antibiotics including AMX, CLA, MTZ, LEV, TET, and FZD between the two age groups ($p > 0.05$) (Fig 3).

Table 2. Minimum inhibitory concentrations (MICs) of *H. pylori* to commonly used antibiotics (mg/L).

Antibiotics*	MICs	MIC ₅₀	MIC ₉₀	MICs of ATCC43504
AMX	0.0156 ~ 256	0.125	4	0.25
CLA	0.0156 ~ >256	0.0312	64	0.0625
MTZ	0.0156 ~ >256	8	256	64
LEV	0.0156 ~ 256	0.25	16	0.5
TET	0.0156 ~ 256	0.0625	4	1
FZD	0.0156 ~ >256	0.0312	2	0.25

*AMX: amoxicillin, CLA: clarithromycin, MTZ: metronidazole, LEV: levofloxacin, FZD: furazolidone, TET: tetracycline.

<https://doi.org/10.1371/journal.pone.0256225.t002>

Table 3. Differences in resistance rates of *H. pylori* using different breakpoints.

Antibiotics ¹	EUCAST ² breakpoints		Other ³ breakpoints		X ²	p
	Isolate (n)	Resistance rate (%)	Isolate (n)	Resistance rate (%)		
AMX	119	57.21	31	14.90	88	<0.001
CLA	81	38.94	81	38.94	0	1
MTZ	82	39.42	106	50.96	24	<0.001
FZD	-	-	23	11.05	-	-
TET	27	12.98	22	10.58	5	0.025
LEV	73	35.10	73	35.10	0	1

1. AMX: amoxicillin, CLA: clarithromycin, MTZ: metronidazole, LEV: levofloxacin, FZD: furazolidone, TET: tetracycline.

2. EUCAST: European Committee on Antimicrobial Susceptibility Testing.

3. Other breakpoints were as follows: ≥ 2 mg/L for AMX, ≥ 1 mg/L for CLA, ≥ 8 mg/L for MTZ, ≥ 2 mg/L for LEV, ≥ 4 mg/L for TET, and > 2 mg/L for FZD.

<https://doi.org/10.1371/journal.pone.0256225.t003>

Impact of gender on antibiotic resistance of *H. pylori* in South China

There was no significant difference in resistance to AMX, CLA, MTZ, LEV, TET and FZD among *H. pylori* strains isolated between the two gender groups ($p > 0.05$) (Fig 4).

Discussion

Resistance of *H. pylori* to the commonly used antibiotics has posed an increasing threat to the standard therapeutic regimens worldwide [11]. Therefore, it is extremely important to continuously monitor the resistance of *H. pylori* in various regions to find effective antibiotics to eradicate *H. pylori*.

Various methods have been reported to assess antibiotic susceptibility of *H. pylori in vitro*, including Kirby-Bauer, agar dilution, and broth dilution methods. Kirby-Bauer antibiotic test is semi-quantitative and unreliable. Agar and broth dilution methods, the most commonly used methods, are quantitative and can provide more accurate results [21]. Agar dilution is recognized as the gold standard of susceptibility testing. However, this method is labor-intensive and expensive. Epsilon test (E-test) is an agar diffusion method and demonstrates an excellent correlation with the agar dilution method [22, 23]. However, antibiotics has to be supplied by specified manufacturers, making this method costly for screening. The broth dilution method is also highly accurate and comparable to agar dilution [24], and can test more than one antibiotic at one time. Broth microdilution method uses microdilution plates with a capacity of 500 μ L/well to perform the broth dilution. The results of the ampicillin and clarithromycin disk diffusion assays for *H. pylori* showed 100% correlation with those of broth microdilution [25]. The E-test tends to overestimate the resistance of metronidazole for *H.*

Table 4. Differences in multiple antibiotic resistance rates of *H. pylori* using different breakpoints.

	EUCAST [#] breakpoints		Other breakpoints [*]		X ²	p
	Isolates (n)	Resistance rate (%)	Isolate (n)	Resistance rate (%)		
Monoresistance	45	21.63	54	25.96	0.190	0.163
Double resistance	36	17.31	26	12.50	2.382	0.121
Triple resistance	21	10.10	26	12.50	0.696	0.405
Multiple resistance	40	19.24	30	14.42	6.750	0.006

[#]EUCAST: European Committee on Antimicrobial Susceptibility Testing.

^{*} Other breakpoints were as follows: ≥ 2 mg/L for AMX, ≥ 1 mg/L for CLA, ≥ 8 mg/L for MTZ, ≥ 2 mg/L for LEV, ≥ 4 mg/L for TET, and > 2 mg/L for FZD.

<https://doi.org/10.1371/journal.pone.0256225.t004>




Fig 2. Impact of disease on different antibiotic resistance of *H. pylori* in South China. A is for amoxicillin (AMX), B is for clarithromycin (CLA), C is for furazolidone (FZD), D is for levofloxacin (LEV), E is for metronidazole (MTZ), F is for tetracycline (TET). CSG: chronic superficial gastritis; CAG: chronic atrophic gastritis; EG: erosive gastritis; PU: peptic ulcer.

<https://doi.org/10.1371/journal.pone.0256225.g002>




Fig 3. Impact of age on different antibiotic resistance of *H. pylori* in South China. A is for amoxicillin (AMX), B is for clarithromycin (CLA), C is for furazolidone (FZD), D is for levofloxacin (LEV), E is for metronidazole (MTZ), F is for tetracycline (TET).

<https://doi.org/10.1371/journal.pone.0256225.g003>




Fig 4. Impact of gender on different antibiotic resistance of *H. pylori* in South China. A is for amoxicillin (AMX), B is for clarithromycin (CLA), C is for furazolidone (FZD), D is for levofloxacin (LEV), E is for metronidazole (MTZ), F is for tetracycline (TET).

<https://doi.org/10.1371/journal.pone.0256225.g004>

pylori. Therefore, the results of metronidazole using E-test should be confirmed by agar dilution or broth microdilution [25, 26]. All discrepancies occurred when the E-test MIC values were at the range of 8 and 32 mg/L [25]. Therefore, the broth microdilution method was used in our study to monitor the drug resistance rates.

Breakpoints were used to classify the results of antimicrobial susceptibility by agar dilution or broth dilution methods into sensitive, intermediate, and resistant to a specific antibiotic. The CLSI only recommends breakpoints for clarithromycin to treat *H. pylori* and proposes agar dilution method as the standard for antimicrobial susceptibility testing. Different studies use different breakpoints, for example, the breakpoints of amoxicillin ranges from 0.125 to 2 µg/ml [27–30], making it hard to compare the results of different studies. Alarcón et al. [31] conducted antimicrobial susceptibility in *H. pylori* clinical isolates by using EUCAST breakpoints and compared to other breakpoints. They found similar results lied in the most frequently tested antibiotics, e.g. metronidazole, clarithromycin, tetracycline, and levofloxacin, with exception of amoxicillin and rifampicin. Our results agreed with these findings. The overall resistance rate for amoxicillin was 60.98% by using EUCAST breakpoints and 12.80% by using other breakpoints in our study. There were distinct differences between the two breakpoints for simple AMX, MTZ, and TET resistance. However, there was no significant difference for CLA and LEV. In addition to CLSI, EUCAST, and BSAC breakpoint systems, a breakpoint system appropriate for China, especially for AMX, MTZ and TET, is desired.

Resistance rates of CLA, MTZ, and LEV for *H. pylori* were all high in South China. While the MIC₉₀ of AMX, TET, and FZD were low, the MIC₉₀ of MTZ was high, 128 times higher than that of FZD and 64 times higher than those of AMX and TET. Our results indicate that FZD, AMX, and TET remain good choices for the eradication of *H. pylori* in South China, consistent with the results observed in other regions in China and abroad [20, 28, 32].

However, CLA, MTZ, and LEV should be avoided in empiric treatment of *H. pylori* in South China. CLA resistance has been indicated to account for the sharply decreased eradication rate by 70% (from 87.8% to 18.3%) in the regimen of PPI-clarithromycin-amoxicillin and 47% (from 97% to 50%) in the regimen of PPI-clarithromycin-metronidazole [33]. Resistance to CLA and MTZ has compromised the therapeutic efficacy. Malfertheiner et al. [1] recommended to use the bismuth quadruple as one of the first-line medications in areas with high dual resistance to CLA and MTZ.

MTZ is the classic nitroimidazole antibiotic. It reaches a high concentration in the stomach and its bactericidal activity is not affected by low pH in the stomach. *H. pylori* resistance to MTZ and CLA has been globally reported with regional differences and has been on the rise. While the resistance rate of *H. pylori* to MTZ is relatively low in Southern Europe, Taiwan, and Japan (less than 20%) [34–36], it is high in Africa (75.8%) [13] and China (78.2%) [37]. The main reasons for the high MTZ resistance are its previous prescription for parasitic or gynecological infections [38], extensive availability, and affordability. In majority of countries, the resistance rate of CLA is greater than 20% [32, 33, 37]. From 1997 to 2008, the resistance rate of CLA in Japan gradually increased from 8.7% to 34.5% [11].

We found that the MICs of *H. pylori* were highest in patients with erosive gastritis. This may be due to the resistant *H. pylori* isolates are virulent strains and often cause gastric erosion. Our study failed to detect significant difference in the resistance rate of *H. pylori* to tested antibiotics in different age and gender groups.

There were at least two limitations in our study. Firstly, drug resistance and sensitivity determined by antibiotic susceptibility test *in vitro* cannot fully represent the condition *in vivo*. And secondly, the selected samples from our clinical centers could not represent the whole population in South China.

Conclusions

Our results demonstrate that AMX, FZD, and TET, but not MTZ, CLR or LEV, showed good anti-*H. pylori* activity *in vitro* in South China. When different breakpoints were used, similar results were found with CLA, and LEV, but not with AMX, MTZ, or TET.

Supporting information

S1 Data.
(XLSX)

Author Contributions

Conceptualization: Xueping Huang, Zhihui Lin, Gaohui Nong, Jinqi Wei.

Data curation: Xueping Huang, Yuan Liu, Zhihui Lin, Yushan Chen.

Formal analysis: Xueping Huang.

Funding acquisition: Xueping Huang.

Investigation: Yuan Liu, Yushan Chen, Yuping Lu, Xinhua Ji, Xiang Zhou, Biao Suo, Qiuzhao Chen.

Methodology: Yuan Liu, Baihe Wu, Gaohui Nong.

Project administration: Jinqi Wei.

Resources: Yuan Liu, Yushan Chen.

Software: Baihe Wu.

Supervision: Zhihui Lin, Jinqi Wei.

Writing – original draft: Xueping Huang.

References

1. Malfertheiner P, Megraud F, O'Morain CA. Management of *Helicobacter pylori* infection—the Maastricht V/Florence Consensus Report. *Gut*. 2017; 66 (1):6–30. <https://doi.org/10.1136/gutjnl-2016-312288> PMID: 27707777.
2. Fock KM, Ang TL. Epidemiology of *Helicobacter pylori* infection and gastric cancer in Asia. *Journal of gastroenterology and hepatology*. 2010; 25 (3):479–486. <https://doi.org/10.1111/j.1440-1746.2009.06188.x> PMID: 20370726.
3. Sanchez Delgado J, Garcia-Iglesias P, Tito L, Puig I. Update on the management of *Helicobacter pylori* infection. A position paper from the Catalan Society of Digestology. *Gastroenterologia y hepatologia*. 2018; 41 (4):272–280 <https://doi.org/10.1016/j.gastrohep.2017.12.009> PMID: 29426621.
4. Smith SM, O'Morain C, McNamara D. Antimicrobial susceptibility testing for *Helicobacter pylori* in times of increasing antibiotic resistance. *World Journal of Gastroenterology*. 2014; 20 (29):9912–9921. <https://doi.org/10.3748/wjg.v20.i29.9912> PMID: 25110421.
5. Jung Mook, Kang Nayoung, Kim Cheol, et al. Predictive factors for improvement of atrophic gastritis and intestinal metaplasia after *Helicobacter pylori* eradication: a three-year follow-up study in Korea. *Helicobacter* 2012; 17 (2):86–95. <https://doi.org/10.1111/j.1523-5378.2011.00918.x> PMID: 22404438.
6. Vainio H. Schistosomes, liver flukes, and *Helicobacter pylori*. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Lyon, 7–14 June 1994. IARC monographs on the evaluation of carcinogenic risks to humans 1994; 61:1–241. PMID: 7715068
7. Sugimoto M, Murata M, Yamaoka Y. Chemoprevention of gastric cancer development after *Helicobacter pylori* eradication therapy in an East Asian population: Meta-analysis. *World Journal of Gastroenterology*. 2020; 26(15):1820–1840. <https://doi.org/10.3748/wjg.v26.i15.1820> PMID: 32351296.
8. Zullo A, Hassan C, Cristofari F, Andriani A, Francesco V.D, Ierardi E, et al. Effects of *Helicobacter pylori* eradication on early-stage gastric mucosa-associated lymphoid tissue lymphoma. *Clinical gastroenterology and hepatology*. 2010; 8 (2):105–110. <https://doi.org/10.1016/j.cgh.2009.07.017> PMID: 19631287.
9. Sugano K, Tack J, Kuipers EJ, Graham DY, El-Omar EM, Miura S, et al. Kyoto global consensus report on *Helicobacter pylori* gastritis. *Gut*. 2015; 64 (9):1353–1367. <https://doi.org/10.1136/gutjnl-2015-309252> PMID: 26187502.
10. Chung JW, Lee GH, Han JH, Jeong JY and Kim JH. The trends of one-week first-line and second-line eradication therapy for *Helicobacter pylori* infection in Korea. *Gastroenterology*. 2009; 136(105):A-24–A-24. [https://doi.org/10.1016/S0016-5085\(09\)60113-X](https://doi.org/10.1016/S0016-5085(09)60113-X). PMID: 21510323.
11. Sasaki M, Ogasawara N, Utsumi K, Kawamura N, Kamiya T, Kataoka H, et al. Changes in 12-Year First-Line Eradication Rate of *Helicobacter pylori* Based on Triple Therapy with Proton Pump Inhibitor, Amoxicillin and Clarithromycin. *Journal of clinical biochemistry and nutrition*. 2010; 47 (1):53–58. <https://doi.org/10.3164/jcbs.10-10> Epub 2010 Jun 17. PMID: 20664731.
12. De Francesco V, Giorgio F, Hassan C, Manes G Zullo A. Worldwide h. pylori antibiotic resistance: a systematic review. *Journal of gastrointestinal and liver diseases*.2010; 19 (4):409–414. [10.1111/j.1440-1746.2010.06571.x](https://doi.org/10.1111/j.1440-1746.2010.06571.x) PMID: 21188333.
13. Jaka H, Rhee JA, Östlu Linda, Smart L, Peck R, Mueller A, et al. The magnitude of antibiotic resistance to *Helicobacter pylori* in Africa and identified mutations that confer resistance to antibiotics: systematic review and meta-analysis. *BMC infectious diseases* 2018; 18 (1):193. <https://doi.org/10.1186/s12879-018-3099-4> PMID: 29699490.
14. Graham DY. *Helicobacter pylori* update: gastric cancer, reliable therapy, and possible benefits. *Gastroenterology*.2015; 148 (4):719–731 e713. <https://doi.org/10.1053/j.gastro.2015.01.040> PMID: 25655557.
15. Thung I, Aramin H, Vavinskaya V, Gupta S, Park JY, Crowe SE, et al. Review article: the global emergence of *Helicobacter pylori* antibiotic resistance. *Alimentary pharmacology & therapeutics*. 2016; 43 (4):514–533. <https://doi.org/10.1111/apt.13497> PMID: 26694080.
16. European Committee on Antimicrobial Susceptibility Testing. Breakpoint tables for interpretation of MICs and zone diameters. EUCAST Clinical Breakpoint Table v. 5.0, valid from 2015-01-01; <http://www.eucast.org/clinical/breakpoints>.

17. Clinical and Laboratory Standards Institute M100-S19: Performance Standards for Antimicrobial Susceptibility Testing; 19th Informational Supplement. Wayne, PA, USA: Clinical and Laboratory Standards Institute, 2009.
18. Su P, Li Y, Li H, Zhang J, Lin L, Wang Q, et al. Antibiotic resistance of *Helicobacter pylori* isolated in the Southeast Coastal Region of China. *Helicobacter* 2013; 18(4):274–279. <https://doi.org/10.1111/hel.12046> PMID: 23418857.
19. King A. Recommendations for susceptibility tests on fastidious organisms and those requiring special handling. *Journal of Antimicrobial Chemotherapy*. 2001; 48 Suppl. 1: 77–80. https://doi.org/10.1093/jac/48.suppl_1.77 PMID: 11420340.
20. Bai P, Zhou LY, Xiao XM, Luo Y, Ding Y. Susceptibility of *Helicobacter pylori* to antibiotics in Chinese patients. *Journal of Digestive Disease*. 2015; 16(8):464–470. <https://doi.org/10.1111/1751-2980.12271> PMID: 26147515.
21. Wiegand I, Hilpert K, Hancock RE. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nature Protocols*. 2008; 3(2):163–175. <https://doi.org/10.1038/nprot.2007.521> PMID: 18274517.
22. Piccolomini R., Bonaventura GD., Catamo G., Carbone F, Neri M. Comparative evaluation of the E test, agar dilution, and broth microdilution for testing susceptibilities of *Helicobacter pylori* strains to 20 antimicrobial agents. *Journal of clinical microbiology*. 1997; 35 (7):1842–1846. <https://doi.org/10.1128/jcm.35.7.1842-1846.1997> PMID: 9196205.
23. Joyce LF, Downes J, Stockman K, Andrew JH. Comparison of five methods, including the PDM Epsilon-meter test (E test), for antimicrobial susceptibility testing of *Pseudomonas aeruginosa*. *Journal of clinical microbiology*. 1992; 30 (10):2709–2713. <https://doi.org/10.1128/jcm.30.10.2709-2713.1992> PMID: 1400972.
24. Jacobson RK, Notaro MJ, Carr GJ. Comparison of *Neisseria gonorrhoeae* minimum inhibitory concentrations obtained using agar dilution versus microbroth dilution methods. *Journal of microbiological methods*. 2019; 157:93–99. <https://doi.org/10.1016/j.mimet.2019.01.001> PMID: 30629964.
25. Hachem CY, Clarridge JE, Reddy R, Flamm R, Evans DG, Tanaka SK, et al. Antimicrobial susceptibility testing of *Helicobacter pylori*. Comparison of E-test, broth microdilution, and disk diffusion for ampicillin, clarithromycin, and metronidazole. *Diagnostic microbiology and infectious disease*. 1996; 24 (1):37–41. [https://doi.org/10.1016/0732-8893\(95\)00252-9](https://doi.org/10.1016/0732-8893(95)00252-9) PMID: 8988762.
26. Osato MS, Reddy R, Reddy SG, Penland RL, Graham DY. Comparison of the Etest and the NCCLS-approved agar dilution method to detect metronidazole and clarithromycin resistant *Helicobacter pylori*. *International journal of antimicrobial agents*. 2001; 17 (1):39–44. [https://doi.org/10.1016/s0924-8579\(00\)00320-4](https://doi.org/10.1016/s0924-8579(00)00320-4) PMID: 11137647.
27. Gao Wen 1, Hong Cheng, Fulian Hu, Jiang Li, Lihui Wang, Guibin Yang, et al. The evolution of *Helicobacter pylori* antibiotics resistance over 10 years in Beijing, China. *Helicobacter*. 2010; 15 (5):460–466. <https://doi.org/10.1111/j.1523-5378.2010.00788.x> PMID: 21083752.
28. Jung Won Lee Nayoung Kim, Jung Mogg Kim Ryoung Hee Nam, Chang Hyun, Jae Yeon Kim, et al. Prevalence of primary and secondary antimicrobial resistance of *Helicobacter pylori* in Korea from 2003 through 2012. *Helicobacter*. 2013; 18 (3):206–214. <https://doi.org/10.1111/hel.12031> PMID: 23241101.
29. Megraud F, Coenen S, Versporten A, Goossens H, Glupczynski Y, Alexander MH, et al. *Helicobacter pylori* resistance to antibiotics in Europe and its relationship to antibiotic consumption. *Gut*. 2013; 62 (1):34–42. <https://doi.org/10.1136/gutjnl-2012-302254> PMID: 22580412.
30. Deyi V, Bontems P, Vanderpas J, Koster ED, Ntounda R, Van BC, et al. Multicenter survey of routine determinations of resistance of *Helicobacter pylori* to antimicrobials over the last 20 years (1990 to 2009) in Belgium. *Journal of clinical microbiology*. 2010; 49 (6):2200–2209. <https://doi.org/10.1128/JCM.02642-10> PMID: 21450969.
31. Alarcón T, Urruzuno P, Martínez MJ, Domingo D, Llorca L, Correa A, et al. Antimicrobial susceptibility of 6 antimicrobial agents in *Helicobacter pylori* clinical isolates by using EUCAST breakpoints compared with previously used breakpoints. *Enfermedades infecciosas y microbiología clínica*. 2017; 35 (5):278–282. <https://doi.org/10.1016/j.eimc.2016.02.010> PMID: 27017059.
32. Azzaya D, Gantuya B, Oyuntsetseg K, Davaadorj D, Matsumoto T, Akada J, et al. High Antibiotic Resistance of *Helicobacter pylori* and Its Associated Novel Gene Mutations among the Mongolian Population. *Microorganisms*. 2020; 8(7), 1062. <https://doi.org/10.3390/microorganisms8071062> PMID: 32708761.
33. Megraud F, Lehours P. *Helicobacter pylori* detection and antimicrobial susceptibility testing. *Clinical microbiology reviews*. 2007; 20 (2):280–322. <https://doi.org/10.1128/CMR.00033-06> PMID: 17428887.
34. Montes M, Villalon FN, Eizaguirre FJ, Maider D, Ignacio M, María FR. et al. *Helicobacter pylori* Infection in Children. Antimicrobial Resistance and Treatment Response. *Helicobacter*. 2015; 20 (3):169–175. <https://doi.org/10.1111/hel.12187> PMID: 25382231.

35. Liang CM, Cheng JW, Kuo CM, Chang KC, Chuah SK. Levofloxacin-containing second-line anti-*Helicobacter pylori* eradication in Taiwanese real-world practice. *Biomedical journal*. 2014; 37 (5):326–330. <https://doi.org/10.4103/2319-4170.125650> PMID: 25163495.
36. Morimoto Norihito, Takeuchi Hiroaki, Nishida Yoshie, et al. Clinical Application of the DiversiLab Microbial Typing System Using Repetitive Sequence-Based PCR for Characterization of *Helicobacter pylori* in Japan. *Journal of clinical laboratory analysis*. 2015; 29 (3):250–253. <https://doi.org/10.1002/jcla.21758> PMID: 24796534.
37. Liu DS, Wang YH, Zeng ZR, Zhang ZY, Lu H, Xu JM, et al. Primary antibiotic resistance of *Helicobacter pylori* in Chinese patients: a multiregion prospective 7-year study. *Clinical microbiology and infection*. 2018; 24 (7):780 e785–780 e788. <https://doi.org/10.1016/j.cmi.2017.11.010> PMID: 29138101.
38. Giulio MD, Campi ED, Bartolomeo SD, Cataldi V, Marzio L, Grossi L, et al. In vitro antimicrobial susceptibility of *Helicobacter pylori* to nine antibiotics currently used in Central Italy. *Scandinavian journal of gastroenterology*. 2016; 51 (3):263–269. <https://doi.org/10.3109/00365521.2015.1092577> PMID: 26554617.