

Acupoint Stimulation for Enhanced Recovery After Colon Surgery: A Prospective Multicenter Randomized Controlled Trial

Zhihong Lu^{1,*}, Ailin Luo^{2,*}, Su Min^{3,*}, Hailong Dong¹, Qiuju Xiong³, Xinhua Li², Qingzhu Deng², Tingting Liu¹, Xue Yang¹, Chen Li⁴, Qingchuan Zhao⁵, Lize Xiong^{1,6}

¹Department of Anaesthesiology and Perioperative Medicine, Xijing Hospital, Fourth Military Medical University, Xi'an, People's Republic of China; ²Department of Anaesthesiology, Tongji Hospital, Huazhong University of Science and Technology, Wuhan, People's Republic of China; ³Department of Anaesthesiology, First Affiliated Hospital of Chongqing Medical University, Chongqing, People's Republic of China; ⁴Department of Medical Statistics, Fourth Military Medical University, Xi'an, People's Republic of China; ⁵Xijing Hospital of Digestive Diseases, Fourth Military Medical University, Xi'an, People's Republic of China; ⁶Translational Research Institute of Brain and Brain-Like Intelligence and Department of Anesthesiology, Shanghai Fourth People's Hospital Affiliated to Tongji University School of Medicine, Shanghai, People's Republic of China

*These authors contributed equally to this work

Correspondence: Zhihong Lu; Lize Xiong, Department of Anesthesiology and Perioperative Medicine, Xijing Hospital, Fourth Military Medical University, Changle West Road 127, Xi'an, Shaanxi, 710032, People's Republic of China, Tel +86-29-84775337; Tel +86-29-84772126, Email deerlu23@163.com; xlmzk@163.com

Purpose: The aim of this study was to evaluate the efficacy of transcutaneous electrical acupoint stimulation (TEAS) in improving bowel function and thus shortening hospital stay after laparoscopic colon surgery within the ERAS pathway.

Patients and Methods: From November 2016 to March 2018, 100 patients who underwent elective colon surgery were enrolled and 94 finished study ($n = 47$ for each) in three university hospitals. Patients in the TEAS group received TEAS 30 min before surgery and once a day for 3 days after surgery, while those in the Control Group received no stimulation. Primary outcome was the time to discharge.

Results: Compared with standardized postoperative care, TEAS resulted in a shorter time to first flatus ($P=0.03$) and time to first defecation ($P=0.03$), as well as a reduction in the length of hospital stay ($P=0.02$). Median patient-controlled analgesia (PCA) deliveries and PCA attempts at 24h, 48h and 72h after surgery were less in the TEAS group ($P<0.01$). No evidence of significant advantages in postoperative pain intensity, nausea, vomiting, sleeping quality and expenses was found in the TEAS group.

Conclusion: Perioperative TEAS further shortens the time to meet discharge criteria after laparoscopic colon surgery in patients under ERAS strategy.

Keywords: transcutaneous electrical acupoint stimulation, colon surgery, laparoscopy, ERAS

Plain Language Summary

Recovery of bowel function is important for patients undergoing colorectal surgery. Enhanced recovery after surgery (ERAS) program makes great efforts on improving bowel function. In patients under ERAS program, transcutaneous electrical acupoint stimulation (TEAS) can further shorten the time to first flatus and to first defecation after colon surgery. The length of hospital stay and need of analgesics after surgery were decreased as well.

Introduction

Accumulating evidence supported that the enhanced recovery after surgery (ERAS) pathway and laparoscopic approach benefit patients undergoing colorectal resections by shortening the length of hospital stay and reducing morbidity after surgery.¹⁻⁵ For patients undergoing colon surgery, the return of bowel function is especially important for ERAS.⁶ Shortening fasting time, early mobilization and early oral intake have been used to facilitate the recovery of bowel

function.⁶ However, postoperative ileus still represents problems in the management.⁷ The time to first bowel motion was not satisfied in the laparoscopic arm in many of these trials, which is only about 1 day earlier than that in the open arm.^{7–9}

Acupuncture-related techniques have been used to treat bowel function disorders, including irritable bowel syndrome and chronic functional constipation^{10–12} Perioperative acupuncture-related techniques including transcutaneous electrical acupoint stimulation (TEAS) have shown benefits including narcotic-sparing effects and fewer complications.¹³ Besides, acupuncture has been used to treat gastrointestinal motility disorder.^{14–19} Evidence of acupuncture for postoperative gastrointestinal function is accumulating. In patients undergoing major abdominal surgery, Li et al¹⁹ observed shorter time to flatus and to ambulation after TEAS. In patients undergoing laparoscopic surgery, the time to defecation was shortened as well.²⁰ However, whether the potential benefit of TEAS on gastrointestinal function could shorten the length of stay after surgery is not clear. Therefore, the aim of this multicenter randomized trial was to evaluate whether perioperative TEAS could further improve the gastrointestinal function and shorten the time to meet the discharge criteria in patients undergoing laparoscopic colon surgery with current ERAS approaches.

Materials and Methods

Patients and Randomization

The randomized, prospective, sham controlled study was undertaken in 3 university hospitals in China (Xijing Hospital, Tongji Hospital, and First Affiliated Hospital of Chongqing Medical University). The study was approved by the Institutional Ethics Committee of each participating hospital and has been registered in ClinicalTrials.gov (NCT02921529). Written informed consent was obtained from all patients. Patients scheduled for laparoscopic colon surgery under general anaesthesia were screened. Inclusion criteria included age between 18 and 75 years, a body mass index between 18 and 30, and ASA status of I–III. Patients with contraindications for electrical stimulation, difficulty in communicating, confirmed or suspected drug abuse/addiction or alcohol abuse/addiction, or severe hepatic or renal dysfunction were excluded. Patients were randomly assigned to the TEAS or Control group in a ratio of 1:1 using a computer-generated random allocation sequence. The randomization code for each patient was put in sealed envelope and not opened until allocation.

Intervention and Blinding

For all patients, electrodes were placed at the bilateral PC6/LI4 and ST36/SP6 and connected to the stimulator (Model No. SDZ-II; Hwato electronic stimulator; Suzhou Medical Appliances Co., Ltd, Suzhou, China) (See [Supplemental Digital Content 1](#)). Patients in the TEAS group received stimulation at the bilateral PC6 and ST36 points. The device provided “disperse-dense” waves of alternating frequencies of 2 Hz and 10 Hz for 2 cycles. Increasing electrical stimulation intensity (4 mA to 11 mA) was applied to identify the threshold intensity. Stimulation began 30 min before induction and lasted for 30 min. Then the same stimulation for 30 min was given in the morning once a day for 3 days after surgery. Patients in the Control group received no stimulation.

For logistic reasons, blinding the patients was hard to perform. However, the threshold intensity was identified in all the patients, and they were told that they may or may not feel the electrical stimulation. Interventions were performed by a designated investigator who was not involved in the anaesthesia or the follow-up. The stimulator was placed in an opaque box. Investigators involved in the follow-up were blinded to the group allocation.

Procedures

Anaesthesia was induced with midazolam, propofol and fentanyl. Rocuronium was given to facilitate tracheal intubation. Intraoperative anaesthesia was maintained with a remifentanyl infusion and sevoflurane. Patients were managed according to ERAS methods previously reported^{21,22} (see [Supplemental Digital Content 2](#)) and followed up until discharge.

The primary outcome was the time to meet discharge criteria after surgery that we previously reported^{21,22} ([Table 1](#)). The secondary outcomes included time to flatus, time to first defecation, and time to first oral intake; pain intensity assessed by visual analogue scale (VAS), attempts and deliveries of patient-controlled analgesia (PCA), postoperative nausea and vomiting (PONV), quality of sleeping (QoS) and quality of recovery (QoR) evaluated at 24h, 48h and 72h after surgery. QoS was assessed using visual sleeping scale and the Richard-Campbell Sleep Questionnaire (RCSQ),²³ and QoR was assessed using

Table 1 Criteria for Discharge

	Content
Criteria 1	Normal body temperature, normal mobilization (off-bed mobilization >6h in 24 h), normal oral diet (oral intake of liquid diet >1000mL in 24h), normal gastrointestinal function (flatus and defecation)
Criteria 2	Pain could be controlled with oral analgesics
Criteria 3	No discomfort complaint, no parenteral nutrition

QoR-15 questionnaire.²⁴ Higher scores indicate better quality ([Supplemental Digital Contents 3 and 4](#)). Postoperative major complications ([Supplemental Digital Content 5](#)) and expenses were recorded as well.

Statistical Analysis

We performed all the analyses in a modified intention-to-treat population, which included all the patients who had undergone randomization and all the interventions. All the patients were followed for the duration of the trial.

Based on a previous study, the average time to meet discharge criteria was 5 days for patients undergoing colorectal surgery.²¹ A minimum of 43 patients in each group would give 90% power to detect a decrease of 1 day for the time to discharge, with a standard deviation of 1.3 days, at an overall 2-sided $P < 0.05$. The sample size was inflated to 94 patients ($n=47$ for each) to account for a rate of withdrawals and loss to follow-up by 10%.

Data were analyzed with SAS 9.2 (USA: SAS Institute Inc., 2010). Descriptive statistics were applied to present subjects' characters. Comparison of qualitative data between two groups was done using Chi-square test and Fisher's exact test. And comparison of quantitative data was evaluated by t -test. Also, Satterthwaite t' test was applied if the variances were unequal between two groups. Univariate analysis of survival data was estimated using the Log rank test. A two-tailed P -value of 0.05 was considered as statistically significant with a 95% confidence interval (95% CI).

Results

From November 2016 through March 2018, a total of 100 patients were enrolled at 3 hospitals. We randomly assigned 50 patients to each group ([Figure 1](#)). After randomization, 5 patients were excluded due to procedure converting from laparoscopic to open surgery and 1 patient was excluded due to rejection to TEAS ([Figure 1](#)). There was no significant difference between the groups with respect to demographic, anesthetic and surgical characteristics ([Table 2](#)), pre-existing diseases and pre-operative laboratory test results including nutritional status ([Supplemental Digital Contents 6 and 7](#)). Case numbers in each center were shown in [Supplemental Digital Content 8](#).

Primary Outcome

The median follow-up time was 7 days in each group. The median time to discharge was $6.0d \pm 1.8d$ (95% CI, 5.5–6.5) in the TEAS group compared with $7.0d \pm 2.4d$ (95% CI, 6.3–7.7, $P = 0.02$) in the Control group ([Table 3](#)). Log rank test showed a P value of 0.02 between the two groups ([Figure 2](#)). Subgroup analysis based on type of surgery showed same results concerning time to discharge ([Supplemental Digital Content 9](#)).

Secondary Outcomes

The mean time to flatus and mean time defecation were $34.5 \text{ h} \pm 16.7\text{h}$ (95% CI, 29.6–39.4) and $42.0\text{h} \pm 25.0\text{h}$ (95% CI, 34.7–49.3) respectively in the TEAS group, compared with $42.4\text{h} \pm 22.9\text{h}$ (95% CI, 35.6–49.1, $P = 0.03$) and $55.4\text{h} \pm 29.7\text{h}$ (95% CI, 46.7–64.1, $P = 0.03$) respectively in the Control group ([Table 3](#)). Log rank test of the two outcomes showed P values of 0.03 and 0.03 respectively between the two groups ([Supplemental Digital Contents 10 and 11](#)).

Incidences of PONV, the VAS at rest and at cough, and score of sleeping quality in 3 days after surgery were not different between the groups (all $P > 0.05$). However, median PCA deliveries and PCA attempts at 24h, 48h and 72h after surgery were less in the TEAS group than in the Control ($P < 0.01$). Median QoR-15 scores²² in the TEAS group were not different to the Control group at 48 h and 72h after surgery, but lower at 24h after surgery 50.5 (IQR 42.3–61.8) vs 44.5 (IQR 35.0–57.0), $P = 0.04$. Postoperative morbidity of infection and major complications were not different between the




Figure 1 Trial profile.

two groups. Mean postoperative expense was 62962 ± 28001 CNY in the TEAS group and $84,218 \pm 115,027$ CNY in the Control group, with no significant difference ($P=0.22$, Table 3).

Discussion

This multicenter prospective randomized trial suggests that in patients being managed by the current ERAS protocol, perioperative TEAS (30 min before anaesthetic induction, and 30 min once a day for 3 days after surgery) provided additional benefit in enhanced recovery after laparoscopic colon surgery. Patients needed less opioid after surgery, and the time to meet discharge criteria and recovery of bowel function were shortened significantly.

Our study was the first multicenter randomized controlled trial on the role of TEAS for postoperative hospital stay and bowel function in laparoscopic colon surgery. There has been limited randomized controlled trials published in the English literature that examined the role of acupuncture in preventing postoperative ileus after laparoscopic colorectal surgery. In a single-center trial, Ng et al²⁵ observed that electroacupuncture once a day for 4 days postoperatively decreased the time to flatus by 0.6 d and shortened the length of hospital stay from 8.5 to 6d. In our study, with TEAS, a non-invasive technique, we achieved enhanced recovery as well time to flatus decreased by 7.9h and time to discharge decreased by 1d. Additionally, we did TEAS both preoperatively (30min before anaesthesia induction) and postoperatively (once a day for 3 days). Previous studies have found that preoperative acupuncture-related technique potentially provided additional benefits including alleviating anxiety and decreasing anaesthetic consumption.¹³

The shortening of time to discharge or to flatus varied in different trials. In a trial in open colorectal surgery, Zhang et al²⁶ found that electroacupuncture decreased the time to flatus by 9 hours, close to the 7.9 hours in our study. But Ng's study and Li's study found stronger effect of bowel function acceleration.^{19,25} And the time to first oral intake was not shortened in our study, while in Ng's study the time to normal diet decreased by 0.8d. A possible explanation is that in our study patient were managed with ERAS protocol which encourages early oral intake and mobilization. The benefits of TEAS maybe overlapped by ERAS strategy. Ng et al acknowledged that one of the limitations of their study was that

Table 2 Demographic and Surgical Characteristics of the Patients

	TEAS (n=47)	Control (n=47)	P value
Age, years	56.2±9.0	55.6±9.9	0.75
Gender (male/female), n	26/21	23/24	0.54
BMI (kg/m ²)	22.5±2.6	22.8±2.8	0.63
ASA (I/II/III)	0/45/2	1/43/3	0.68
NRS 2002 score	2(0, 5)	2(0, 4)	0.62
Type of surgery (left/right/sigmoid)	6/18/23	5/18/24	0.95
Duration of surgery, min	163.7±49.2	160.1±50.3	0.73
Duration of anaesthesia, min	200.2±48.6	199.2±52.3	0.92
Fluid input, mL	1930.9±459.9	1996.8±529.3	0.52
Crystalloid, mL	1462.8±428.8	1437.2±543.2	0.80
Colloid, mL	446.8±214.5	553.2±238.5	0.05
Output, mL	434.8±273.5	430.0±236.0	0.93
Blood loss, mL	72.8±135.9	62.9±44.0	0.65
Use of vasoactive agents, n (%)	12 (25.5)	13 (27.7)	0.82

Note: Data are presented as mean±SD, n or median (min, max).

Abbreviations: BMI, body mass index; ASA, American Society of Anaesthesiologists; NRS, nutritional risk screening.

Table 3 Recovery Parameters of the Patients

	TEAS (n=47)	Control (n=47)	P value	Difference, 95% CI
Time to discharge, d	6.0±1.8 ^a	7.0±2.4	0.02	-0.98(-1.82,-0.14)
Time to flatus, h	34.5±16.7 ^a	42.4±22.9	0.03	-7.90(-12.35,-3.66)
Time to first defecation, h	42.0±25.0 ^a	55.4±29.7	0.03	-13.42(-23.62, -6.80)
Time to first oral intake of water, h	33.5±22.3	28.9±17.5	0.28	/
PONV, n (%)				
24 h after surgery	30(63.83)	22(46.81)	0.17	/
48h after surgery	10(21.28)	9(19.15)	0.76	/
72h after surgery	8(17.02)	4(8.7)	0.23	/
VAS at rest/at cough, median (min, max)				
24 h after surgery	2.2(0.0, 10.0)/5.8(0.4, 10.0)	1.2(0.0-8.1)/ 5.5(1.4-10.0)	0.41/0.94	/
48h after surgery	1.2(0.0, 6.7)/ 5.0(0.3, 10.0)	1.6(0.0-6.0)/ 5(0.7, 10.0)	0.80/0.98	/
72h after surgery	0.6(0.0, 5.0)/ 3.5(0.1, 10.0)	1.0(0.0, 6.2)/ 4.0(0.3, 10.0)	0.05/0.33	/
PCA attempts/deliveries, median (min, max)				
24 h after surgery	8.0(3.0, 18.0)/5.0.0(2.0, 12.0)	16.5(8.0, 38.0)/12(6, 22.0)	0.001/0.001	/
48h after surgery	13.0(5.0, 22.0)/7.0(3.0, 14.0)	26.0(11.5, 45.5)/16.5(8.0, 35.0)	0.002/0.001	/
72h after surgery	13.0(5.0, 24.0)/7.0(3.0, 20.0)	24.5(13.0, 46.0)/16.5(10.0, 36.0)	0.002/0.003	/

(Continued)


Table 3 (Continued).

	TEAS (n=47)	Control (n=47)	P value	Difference, 95% CI
QoR –15 score, median (min, max)				
24 h after surgery	54.5(15.0, 92.0)	59.0(27.0, 136.0)	0.01	/
48h after surgery	59.0(35.0, 104.0)	62.0(0.0, 130.0)	0.35	/
72h after surgery	67.0(31.0, 113.0)	67.0(11.0, 136.0)	0.75	/
VSS score, median (min, max)				
24 h after surgery	69.0(0.0, 100.0)	60.0(0.0, 96.0)	0.29	/
48h after surgery	44.5(0.0, 96.0)	52.5(1.0, 89.0)	0.27	/
72h after surgery	30.5(2.0, 82.0)	40.5(1.0, 93.0)	0.48	/
RCSQ score, median (min, max)				
24 h after surgery	61.4(18.7, 100.0)	57.0(19.0, 85.0)	0.26	/
48h after surgery	38.5(6.3, 91.0)	45.0(12.9, 85.1)	0.39	/
72h after surgery	34.0(5.0, 81.6)	39.9(8.0, 77.4)	0.44	/
Major complications, n (%)	0(0)	0(0)	1.00	/
Post-surgery expense, CNY	62962±28,001	84,218±115,027	0.21	/

Note: Data are presented as mean±SD, n (%) or mean (min, max).

Abbreviations: PONV, postoperative nausea and vomiting; VAS, visual analogue scale; PCA, patient-controlled analgesia; QoR, quality of recovery; VSS, visual sleeping scale; RCSQ, Richard–Campbell Sleep Questionnaire; CNY, Chinese Yuan.

they did not use a fast-track perioperative program. They also suggested that possible combined effects of EA and the fast-track program on the clinical outcomes after laparoscopic colorectal surgery will be an important area for further research.

**Figure 2** Kaplan–Meier estimation of incidence of meeting discharge criteria.

In our study, the cost after surgery was not different between the two groups ($P=0.22$). However, it should be prudent to conclude that TEAS has no cost-effective benefit. In our study, the time to meet discharge criteria was reduced by 1 day in patients receiving TEAS. It is estimated that a reduction of the length of hospital stay by 1 day may reduce the annual health care system costs in the United States by approximately US\$1 billion.²⁷ The cost-effective effect of TEAS may be shown in the hospital and overall health care system.

Mechanisms of acupuncture-related techniques improving gut function have been studied in animal and clinical studies. In patients undergoing open colon surgery, Meng et al²⁸ failed to prove any benefit of electroacupuncture on time to first bowel movement when the patients received epidural anaesthesia, which blocked the afferent and efferent pathway. This highlights the role of neural pathway during action of acupuncture. It was also reported that vagal nerve system-induced anti-inflammatory effect may be the underlying mechanism of acupuncture improving gut function. Electroacupuncture may suppress intestinal manipulation-induced inflammation via activation of the cholinergic anti-inflammatory pathway in macrophages.^{29,30} Besides, in our study, the postoperative analgesics consumption was fewer during the first 3 days after surgery. The use of opioid-based analgesia is usually thought to exacerbate postoperative ileus.^{31–33} The decreased opioid use in our study may contribute to the better bowel function recovery.

In our study, frequency of stimulation was set at 2Hz/10Hz, which is a low frequency. Both low frequency and high frequency of electrical acupoint stimulation have been used in the prevention and treatment of postoperative ileus. In Li et al¹⁹ and Wang et al²⁰ trials, 2/100Hz was used. While Yang et al found that 10 Hz and 30Hz are more effective in increasing the gastrointestinal motility and attenuating peripheral inflammation.³⁴ Besides, our previous studies showed that TEAS at 2/10Hz induces opioid-sparing effect,³⁵ which may decrease opioid-related postoperative ileus. The optimal frequency for improving gut function still needs further investigation.

The present trial has some limitations. First, investigators could not administer TEAS in a patient-blinded manner. The patient may tell the investigator who did the follow-up that he/she felt a stimuli. We tried to minimize the potential bias by applying electrodes at acupoints and testified stimulation threshold for all patients. We told the patients that they may not feel the stimulation during treatment. The stimulators were put in opaque box to blind the medical staff in the OR and in the ward. Second, flatus is sometimes regarded as an insensitive index, and the time to resume oral intake can be influenced by the patient's perception and the clinician. We therefore adopted time to meet discharge criteria as primary endpoint and time to defecation as one secondary endpoint, because they are more objective and can be recorded by the assessor without bias. Third, though we followed ERAS approaches and encouraged the patients to resume oral intake as early as 6h after surgery, the time to oral intake was longer than 24h in both groups. Adherence to ERAS protocols should be paid attention in future studies.

Conclusion

In conclusion, this multicenter randomized controlled study suggests that TEAS could further enhance short-term recovery and shorten hospital stay after laparoscopic colon surgery in patients being managed by ERAS program. Further studies are warranted to generalize our findings.

Data Sharing Statement

Data related to the specific manuscript will be made available upon reasonable request in adherence with transparency conventions in medical research and through requests to the corresponding author (xlzmzk@163.com).

Acknowledgments

The authors are grateful to Min Chen and Genlin Ji for supporting the anesthesia for the patients, Jielai Xia for suggesting the methods of statistical analysis and critical reading of the manuscript.

Funding

This work was supported by the National Key Basic Research Program of China (No. 2014CB543200), the National Natural Science Foundation of China (No. 81871028), the National Cultivation and Development Program (2021ZY019) and the Aeromedical Flow Principal Investigator Program of Air Force Medical University (2022HYPI02).

Disclosure

The authors declare no conflicts of interest in this work.

References

- Ni X, Jia D, Chen Y, Wang L, Suo J. Is the enhanced recovery after surgery (ERAS) program effective and safe in laparoscopic colorectal cancer surgery? A meta-analysis of randomized controlled trials. *J Gastrointest Surg.* 2019;23(7):1502–1512. doi:10.1007/s11605-019-04170-8
- Ota H, Ikenaga M, Hasegawa J, et al. Safety and efficacy of an “enhanced recovery after surgery” protocol for patients undergoing colon cancer surgery: a multi-institutional controlled study. *Surg Today.* 2017;47(6):668–675. doi:10.1007/s00595-016-1423-4
- García-Erce JA, Camps-Cervantes Á, Ferrando-Ortolá C. Association between use of enhanced recovery after surgery protocol and postoperative complications in colorectal surgery: the postoperative outcomes within enhanced recovery after surgery protocol (POWER) Study. *JAMA Surg.* 2019;154(8):725–736. doi:10.1001/jamasurg.2019.0995
- Godellas CV, Vaince FT, Singer MA, Godellas CV, Vaince FT. Enhanced recovery after surgery: a clinical review of implementation across multiple surgical subspecialties. *Am J Surg.* 2020;219(3):530–534. doi:10.1016/j.amjsurg.2019.11.009
- Ban KA, Berian JR, Ko CY. Does implementation of enhanced recovery after surgery (ERAS) protocols in colorectal surgery improve patient outcomes? *Clin Colon Rectal Surg.* 2019;32(2):109–113. doi:10.1055/s-0038-1676475
- Lassen K, Soop M, Nygren J, et al. Consensus review of optimal perioperative care in colorectal surgery: Enhanced Recovery After Surgery (ERAS) Group recommendations. *Arch Surg.* 2009;144(10):961–969. doi:10.1001/archsurg.2009.170
- Thillainadesan J, Yumol MF, Suen M, Hilmer S, Naganathan V. Enhanced recovery after surgery in older adults undergoing colorectal surgery: a systematic review and meta-analysis of randomized controlled trials. *Dis Colon Rectum.* 2021;64(8):1020–1028. doi:10.1097/DCR.0000000000002128
- Guillou PJ, Quirke P, Thorpe H, et al. Short-term endpoints of conventional versus laparoscopic-assisted surgery in patients with colorectal cancer (MRC CLASICC trial): multicentre, randomised controlled trial. *Lancet.* 2005;365(9472):1718–1726. doi:10.1016/S0140-6736(05)66545-2
- Huang Z, Li T, Zhang G, et al. Comparison of open, laparoscopic, and robotic left colectomy for radical treatment of colon cancer: a retrospective analysis in a consecutive series of 211 patients. *World J Surg Oncol.* 2022;20(1):345. doi:10.1186/s12957-022-02796-8
- Wang XY, Wang H, Guan YY, Cai RL, Shen GM. Acupuncture for functional gastrointestinal disorders: a systematic review and meta-analysis. *J Gastroenterol Hepatol.* 2021;36(11):3015–3026. doi:10.1111/jgh.15645
- Santucci NR, Chogle A, Leiby A, et al. Non-pharmacologic approach to pediatric constipation. *Complement Ther Med.* 2021;59:102711. doi:10.1016/j.ctim.2021.102711
- Lee HY, Kwon OJ, Kim JE, et al. Efficacy and safety of acupuncture for functional constipation: a randomised, sham-controlled pilot trial. *BMC Complement Altern Med.* 2018;18(1):186. doi:10.1186/s12906-018-2243-4
- Lu Z, Dong H, Wang Q, Xiong L. Perioperative acupuncture modulation: more than anaesthesia. *Br J Anaesth.* 2015;115(2):183–193. doi:10.1093/bja/aev227
- Liu YH, Dong GT, Ye Y, et al. Effectiveness of acupuncture for early recovery of bowel function in cancer: a systematic review and meta-analysis. *Evid Based Complement Alternat Med.* 2017;2017:2504021. doi:10.1155/2017/2504021
- Zhu YJ, Wu XY, Wang W, et al. Acupuncture for quality of life in gastric cancer patients undergoing adjuvant chemotherapy. *J Pain Symptom Manage.* 2022;63(2):210–220. doi:10.1016/j.jpainsymman.2021.09.009
- Kanza Gül D, Şolt Kirca A. Effects of acupressure, gum chewing and coffee consumption on the gastrointestinal system after caesarean section under spinal anaesthesia. *J Obstet Gynaecol.* 2021;41(4):573–580. doi:10.1080/01443615.2020.1787363
- Soylu D, Tekinsoy Kartın P. The effect on gastrointestinal system functions, pain and anxiety of acupressure applied following laparoscopic cholecystectomy operation: a randomised, placebo-controlled study. *Complement Ther Clin Pract.* 2021;43:101304. doi:10.1016/j.ctcp.2021.101304
- Zhou X, Cao SG, Tan XJ, et al. Effects of Transcutaneous Electrical Acupoint Stimulation (TEAS) on postoperative recovery in patients with gastric cancer: a randomized controlled trial. *Cancer Manag Res.* 2021;13:1449–1458. doi:10.2147/CMAR.S292325
- Li WJ, Gao C, An LX, Ji YW, Xue FS, Du Y. Perioperative transcutaneous electrical acupoint stimulation for improving postoperative gastrointestinal function: a randomized controlled trial. *J Integr Med.* 2021;19(3):211–218. doi:10.1016/j.joim.2021.01.005
- Wang Y, Yang JW, Yan SY, et al. Electroacupuncture vs sham electroacupuncture in the treatment of postoperative ileus after laparoscopic surgery for colorectal cancer: a multicenter, randomized clinical trial. *JAMA Surg.* 2022;2022:1.
- Feng F, Ji G, Li JP, et al. Fast-track surgery could improve postoperative recovery in radical total gastrectomy patients. *World J Gastroenterol.* 2013;19(23):3642–3648. doi:10.3748/wjg.v19.i23.3642
- Feng F, Li XH, Shi H, et al. Fast-track surgery combined with laparoscopy could improve postoperative recovery of low-risk rectal cancer patients: a randomized controlled clinical trial. *J Dig Dis.* 2014;15(6):306–313. doi:10.1111/1751-2980.12142
- Menear A, Elliott R, Lal LMA, Lal S, McKinley S. Repeated sleep-quality assessment and use of sleep-promoting interventions in ICU. *Nurs Crit Care.* 2017;22(6):348–354. doi:10.1111/nicc.12315
- Stark PA, Myles PS, Burke JA. Development and psychometric evaluation of a postoperative quality of recovery score: the QoR-15. *Anesthesiology.* 2013;118(6):1332–1340. doi:10.1097/ALN.0b013e318289b84b
- Ng SS, Leung WW, Mak TW, et al. Electroacupuncture reduces duration of postoperative ileus after laparoscopic surgery for colorectal cancer. *Gastroenterology.* 2013;144(2):307–313 e1. doi:10.1053/j.gastro.2012.10.050
- Zhang Z, Wang C, Li Q, et al. Electroacupuncture at ST36 accelerates the recovery of gastrointestinal motility after colorectal surgery: a randomised controlled trial. *Acupunct Med.* 2014;32(3):223–226. doi:10.1136/acupmed-2013-010490
- Sommer NP, Schneider R, Wehner S, Kalff JC, Vilz TO. State-of-the-art colorectal disease: postoperative ileus. *Int J Colorectal Dis.* 2021;36(9):2017–2025. doi:10.1007/s00384-021-03939-1
- Meng ZQ, Garcia MK, Chiang JS, et al. Electro-acupuncture to prevent prolonged postoperative ileus: a randomized clinical trial. *World J Gastroenterol.* 2010;16(1):104–111. doi:10.3748/wjg.v16.i1.104

29. Luo D, Liu S, Xie X, Hou X. Electroacupuncture at acupoint ST-36 promotes contractility of distal colon via a cholinergic pathway in conscious rats. *Dig Dis Sci*. 2008;53(3):689–693. doi:10.1007/s10620-007-9929-7
30. Yang NN, Yang JW, Ye Y, et al. Electroacupuncture ameliorates intestinal inflammation by activating $\alpha 7$ nAChR-mediated JAK2/STAT3 signaling pathway in postoperative ileus. *Theranostics*. 2021;11(9):4078–4089. doi:10.7150/thno.52574
31. Farmer AD, Holt CB, Downes TJ, Ruggeri E, Del Vecchio S, De Giorgio R. Pathophysiology, diagnosis, and management of opioid-induced constipation. *Lancet Gastroenterol Hepatol*. 2018;3(3):203–212. doi:10.1016/S2468-1253(18)30008-6
32. Sheng RY, Yan Y, Linh Dang HH. Acupuncture for hot flashes: a literature review of randomized controlled trials conducted in the last 10 years. *World J Tradit Chin Med*. 2021;7:397–407. doi:10.4103/wjtc.wjtc_27_21
33. De Giorgio R, Zucco FM, Chiarioni G, et al. Management of opioid-induced constipation and bowel dysfunction: expert opinion of an Italian multidisciplinary panel. *Adv Ther*. 2021;38(7):3589–3621. doi:10.1007/s12325-021-01766-y
34. Yang NN, Ye Y, Tian ZX, et al. Effects of electroacupuncture on the intestinal motility and local inflammation are modulated by acupoint selection and stimulation frequency in postoperative ileus mice. *Neurogastroenterol Motil*. 2020;32(5):e13808. doi:10.1111/nmo.13808
35. Wang H, Xie Y, Zhang Q, et al. Transcutaneous electric acupoint stimulation reduces intra-operative remifentanyl consumption and alleviates postoperative side-effects in patients undergoing sinusotomy: a prospective, randomized, placebo-controlled trial. *Br J Anaesth*. 2014;112(6):1075–1082. doi:10.1093/bja/aeu001

Journal of Multidisciplinary Healthcare

Dovepress

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>