

Article

Respiratory Syncytial Virus Persistence in Macrophages Upregulates Fcγ Receptors Expression

Jorge Gaona ¹, Carlos Santiago-Olivares ¹, Enrique Ortega ² and Beatriz Gómez ^{1,*}

¹ Department of Microbiology and Parasitology, Faculty of Medicine, National Autonomous University of Mexico (UNAM), Ciudad Universitaria, D.F. Mexico C.P. 04510, Mexico; E-Mails: jgaber2007@gmail.com (J.G.); carlosantiagolivares@yahoo.com.mx (C.S.-O.)

² Department of Immunology, Biomedical Research Institute, National Autonomous University of Mexico (UNAM), Ciudad Universitaria, D.F. México C.P. 04510, Mexico; E-Mail: ortsoto@unam.mx

* Author to whom correspondence should be addressed; E-Mail: begomez@unam.mx; Tel.: +52-55-5623-2469; Fax: +52-55-5623-2382.

Received: 14 October 2013; in revised form: 29 November 2013 / Accepted: 15 January 2014 / Published: 6 February 2014

Abstract: Viruses can persist in differentiated cells (*i.e.*, macrophages) over long periods of time, altering host cells functions but not inducing their death. We had previously reported that, in early passages (14–40) of a murine macrophage-like cell line persistently infected with respiratory syncytial virus (RSV) (MφP), FcγR-mediated phagocytosis and expression of FcγRIIB/RIII on the cell membrane were increased with respect to mock-infected macrophages (MφN). In this work, we explored the mechanism underlying such effects. Increases in FcγR expression and FcγR-mediated phagocytosis are preserved after more than 87 passages of the persistently infected culture. We analyzed the expression of FcγR isoforms at both mRNA and protein levels, and found out that RSV persistence distinctly affects the expression of FcγR isoforms. We also observed that the increase in FcγRs expression results neither from soluble factors (cytokines) or viral products released by the infected cells, nor from an increase in the rate of FcγR internalization. Our results suggest that RSV persistence in macrophages induce intracellular effects that have an impact on FcγRs gene expression at both mRNA and protein levels, and that the characteristics of RSV persistence were preserved for over 87 passages.

Keywords: RSV persistence; macrophages; FcγRs expression

1. Introduction

Viruses persisting in differentiated cells—*i.e.*, macrophages (Mφs)—can regulate the expression of both their own genes and those of the host cell in order to achieve residence in a non-lytic state, besides selectively affecting functions of the infected cell without destroying it [1]. Mφs play very important roles in innate and adaptive immune responses since they are involved in various processes, such as phagocytosis, antigen presentation, and cytokine production [2,3]. Mφs express on their membrane receptors for the Fc region of IgG antibodies (FcγRs). Alterations of FcγRs-mediated phagocytosis *in vitro* and *in vivo* have been reported in Mφs persistently infected with an RNA virus (HIV-1). This effect is caused by a decreased synthesis of the gamma chain of receptors, suggesting that viral persistence alters gene expression in the host cell [4,5]. Changes in host cell-gene expression resulting from persistence of respiratory syncytial virus (RSV), another RNA virus, have also been reported: in the human epithelial cell line HEp-2, viral persistence alters the expression of host genes involved in cell survival and chemokine production [6]. Also, we had previously shown that, in a macrophage-like murine cell line persistently infected with RSV (MφP) [7], FcγRs expression, FcγRs-mediated phagocytosis, and cytokines production are altered [8].

Cell-surface FcγRs comprise a family of integral membrane glycoproteins expressed by most leukocytes. Murine Mφs express four different FcγRs: while FcγRI, FcγRIII, and FcγRIV are activatory, FcγRIIB mediates the inhibition of FcγR-mediated signals [9,10]. Aggregation of FcγRs by antigen-antibody complexes or IgG-opsonized particles triggers several effects or functions, including phagocytosis [11]. FcγRs have also been shown to participate in both the afferent phase of immune responses and immune system homeostasis [12,13].

RSV, a paramyxovirus of the genus *Pneumovirus*, is prevalent and highly infectious [14]. Worldwide, it is a very important pathogen that causes frequent acute upper and lower respiratory tract infections, especially among infants and young children [15,16]. RSV is also a paramount cause of morbidity and mortality in the elderly and in immunocompromised patients [17], constituting the second leading cause of death due to viral infections in elderly individuals [18]. While seasonal RSV outbreaks occur each year globally, the RSV virus can be detected only during the winter epidemic season [14,19]. Young children who have recovered from severe RSV bronchiolitis often develop chronic and recurrent respiratory problems [20,21]. The link between RSV infection, the development of sequelae (wheezing, asthma) [22,23] and chronic obstructive pulmonary disease has been clearly established in several well-controlled prospective epidemiological studies [24,25], suggesting that RSV persists in individuals with this condition [26]. The delayed effects of severe RSV disease might be partially explained by viral persistence, which may cause chronic inflammation [26,27] and/or change cell genome expression patterns.

Persistent infection with RSV occurs in patients with T-cell immunodeficiencies, and establishing persistent infections in tissue culture of either epithelial or hematopoietic cells (e.g., HEp-2 [6] or Mφ [7]) is easy. RSV persistence in animal models has been reported in infected nude mice [28–30], which develop chronic airway function abnormalities [31–33]. Persistence of bovine respiratory syncytial virus, which is closely related to RSV, has also been demonstrated in B cells from naturally infected cows [34]. Although viral persistence in humans has not been clearly proven, circumstantial evidence suggests its occurrence. Immunohistological observations indicate the presence of RSV antigens in

osteoclasts and in multinucleated cells formed in bone marrow cultures from patients with Paget's disease [35]. In addition, RSV nucleic acid was detected in archival postmortem lung tissue from infants who had died during the summertime, without apparent clinical disease having been reported, suggesting that the virus might persist in lungs after an acute infection [36]. Since no animal reservoir of RSV has been found, persistent human infections by RSV may be implicated, at least partially, in preserving the virus during inter-epidemic periods. Despite the evidence that RSV is able to establish persistent infections in various cell types, and the possible relationship between RSV persistence and human disease [23–26,37], few studies have investigated the effects that persistent infection by this virus can have on the functions of infected cells.

We have previously reported that persistent infection by RSV in a murine macrophage-like cell line (M ϕ P) alters the expression of Fc γ RIIB/RIII on the cell membrane and Fc γ Rs-mediated phagocytosis [8]. In this work, we explored the mechanism underlying such effects, studying the effect of RSV persistence on the expression of the different Fc γ R isoforms at both mRNA and protein level, and determining whether the effect is caused by soluble factors (cytokines) or viral products released by infected cells.

2. Results and Discussion

2.1. Results

2.1.1. Persistent Infection by RSV Increases Fc γ R-mediated Phagocytosis and Fc γ RIIB/RIII Expression even at High Passage Numbers

Throughout this paper, we refer to M ϕ P as cells of the murine macrophage-like cell line (P388D1) persistently infected with RSV, and to M ϕ N as mock-infected P388D1 cells. Persistent infection in the M ϕ P cultures used in these experiments was confirmed by the presence of mRNA for the N gene and the expression of viral antigens in 92%–96% of cells (Figure 1). Viral persistence parameters (infective units in culture supernatants and % of RSV antigen-positive cells) did not differ significantly from earlier reported values [7].

Previously, we had reported that in M ϕ P early passages (14–40), Fc γ Rs-mediated phagocytosis and plasma membrane expression of Fc γ RIIB/RIII increase in comparison with non-infected M ϕ N [8]. In this work, we evaluated whether the same characteristics were present in M ϕ P later culture passages (72 to 87). Phagocytosis and expression of Fc γ RIIB/RIII were determined as previously described [8]. We found out that, after more than 70–80 passages, M ϕ P still showed higher levels of phagocytic activity and cell surface Fc γ RIIB/RIII expression than M ϕ N cells (Figure 2). As expected, phagocytic activity was dependent on the IgG concentration used for opsonization, and the phagocytosis of IgG-opsonized sheep red blood cells (SRBCs) by M ϕ P was always significantly higher (usually by a factor of 2) than that of M ϕ N. It is noteworthy that no differences were observed in the phagocytosis of non-opsonized erythrocytes, which suggests that the effect is characteristic of Fc γ R-mediated phagocytosis (Figure 2A). Increases in the expression of cell surface Fc γ RIIB/RIII in M ϕ P (50%–72%) (Figure 2B) was similar to those previously reported for early culture passages [8].

Figure 1. RSV persistence in MφP. (A) Presence of mRNA of the N gene in MφP. Agarose/EtBr gel (2%) electrophoresis of RT-PCR products. Total RNA from three different passages of MφP (72, 83 and 87) or MφN cells was harvested and converted to cDNA. DNA primers for the N gene were used to amplify a segment of 1,187 bp by PCR. As control, a segment of RNA for GAPDH was amplified; (B) Expression of RSV antigens on MφP membrane. MφN (empty histogram) or MφP (gray histogram) were stained with FITC-labeled anti RSV antibodies and analyzed by flow cytometry. Histogram is representative of several experiments with different passages of MφP and MφN during the course of these experiments.






Figure 2. FcγRIIB/RIII mediated phagocytic activity and expression of FcγRIIB/RIII receptors in MφP and MφN. (A) Phagocytosis of IgG-opsonized and non-opsonized SRBC by MφN and MφP. Cells were incubated with 2,4,6-trinitrophenyl-labeled SRBC opsonized with the indicated concentrations of the anti-2,4-dinitrophenol (anti-DNP) IgG antibody 4F8. Results are expressed as mean $\pm$ 1 SD from three independent experiments passages of MφN and passages 72, 83 and 87 of MφP (* $p < 0.05$); (B) Expression of FcγRIIB/RIII in the above mentioned passages of MφN or MφP cells. Cell-membrane FcγRIIB/RIII were stained with specific monoclonal antibody 2.4G2 and secondary FITC-labeled F(ab')₂ fragments of anti-rat antibodies, and analyzed by flow cytometry. Each individual bar represents the MFI of 10,000 cells. Statistical analysis of the average of the three different passages of MφN and MφP (38.67 ± 2.4 vs. 60.33 ± 4.9 , respectively) indicates a significant difference in FcγRIIB/RIII expression ($p = 0.016$).



2.1.2. Increase in FcγRIIB/RIII Cell Membrane Expression Is Not Mediated by Soluble Factors

Persistence of RSV in macrophages could induce the release of extracellular factors (e.g., cytokines, viral particles, viral products), which would act in a paracrine-like way in order to induce the increase in FcγRIIB/RIII expression. Seeking to investigate whether the increased expression of FcγRIIB/RIII was produced by factors released by persistently infected MφP, we treated MφN cells with a conditioned medium obtained from MφN or MφP cultures after an incubation period of 12 h or 24 h. Expression levels of FcγRIIB/RIII were determined by flow cytometry. Mean fluorescence intensity (MFI) of FcγRIIB/RIII expression by MφN was not significantly altered when MφN were incubated for 24 h with supernatants from MφN or MφP (Figure 3). So as to verify that MφN are able to increase the expression of FcγRIIB/RIII in response to a stimulus already known to increase the expression of these receptors [38], MφN were incubated with heat-killed NHTi, which induced a significant increase in FcγR expression as compared to cells treated with conditioned medium (Figure 3). This suggests that the MφN cell line is able to respond to an activating stimulus. These results show that the increase in the expression of FcγRIIB/RIII induced by RSV persistence is not mediated by extracellular factors released by persistently infected cells.


Figure 3. FcγRIIB/RIII expression in MφN after treatment with MφN or MφP conditioned medium (CM). MφN were treated with either 12 h- or 24 h-CM from MφN or MφP, or with heat-killed NHTi as control (see Materials and Methods for details). After 24 h, FcγRIIB/RIII was analyzed by flow cytometry. Results are expressed as mean ± 1 SD of mean fluorescence intensity in three independent experiments.



2.1.3. RSV Persistence does not Affect FcγRIIB/RIII Endocytosis

In order to determine whether the increase in membrane FcγRIIB/RIII was due to impaired receptor endocytosis, we measured the rate of FcγRIIB/RIII internalization as described in Materials and Methods. We found similar FcγRs internalization kinetics in both MφN and MφP. Average decreases in MFI at 120 min were 14.82 and 15.79 units for MφN and MφP, respectively, suggesting that FcγRIIB/RIII receptors endocytosis is not significantly altered by RSV persistence (Figure 4).


Figure 4. Internalization of FcγRIIB/RIII in MφN and MφP. Internalization kinetics of mAb 2.4G2-FcγRIIB/RIII complexes was monitored by flow cytometry during 120 min, as described in Materials and Methods. Results are expressed as the mean ± 1 SD from three independent experiments. No significant difference was observed between MφN and MφP in the net amount of internalized mAb 2.4G2-FcγRIIB/RIII complexes.



2.1.4. Intracellular Levels of FcγRIIB/RIII Proteins Are Increased in MφP

In order to investigate whether the increase of FcγRIIB/RIII expression on the membrane of MφPs was associated with increased receptor synthesis, we determined the total amount of receptor protein (membrane and intracellular) in the cells using flow cytometry. We found out that MφPs have more FcγRIIB/RIII protein than MφNs, both on the cell surface and intracellularly (Figure 5). These results suggest that viral persistence induces the upregulation of FcγRIIB/RIII synthesis.

Figure 5. Total FcγRIIB/RIII protein in MφN and MφP. Cell-membrane and total FcγRIIB/RIII protein content were determined in non-permeabilized or permeabilized MφN and MφP cells. FcγRIIB/RIII expression was evaluated with 2.4G2 monoclonal antibody and analyzed by flow cytometry. Results are expressed as mean ± 1 SD from three independent experiments, using different passages of MφN and MφP.* indicates $p < 0.05$.




2.1.5. Membrane Expression of FcγRIIB and FcγRIII, but Not of FcγRI, Is Increased in RSV Persistently Infected Cells

Four different types of murine Fcγ receptors have been described (RI to RIV). In our initial work we measured the expression of FcγRs with mAb 2.4G2, which recognizes both FcγRIIB and FcγRIII [39]. In order to determine the effect of RSV persistence on the expression levels of FcγRI, FcγRIIB and FcγRIII separately, we compared the amount of each protein in MφPs and MφNs lysates by Western blot using antibodies specific for each isoform. Lack of a suitable commercial antibody prevented us from determining FcγRIV.

The protein content of FcγRIIB and FcγRIII was observed to be higher in MφP than in MφN: fold increases were 1.9 and 0.6, respectively. In contrast, no significant difference was found in FcγRI levels between in MφP and MφN (Figure 6). It is interesting that the highest effect of RSV persistence was the increase in FcγRIIB, an inhibitory receptor, and yet phagocytosis is higher in MφPs cells. This discrepancy might be related to the IgG isotype used for opsonization (IgG2b), which is very weakly recognized by FcγRIIB [40].

Figure 6. Total FcγRI, RIIB, and RIII protein in MφN and MφP cell extracts. (A) Cells from three different passages of MφN and cells from passages 72, 83 and 87 of MφP were lysed and FcγRI, FcγRIIB, FcγRIII and GAPDH proteins in cell extracts were determined by Western blot with specific antibodies; (B) The relative amount of each FcγR isoform was determined as the ratio of the densitometric intensity of the FcγR band to the intensity of the GAPDH band in the correspondent cell extract. Results are expressed as mean ± 1 SD (* indicates $p < 0.05$).



2.1.6. mRNA Expression of FcγRs Is Distinctively Affected by Viral Persistence


Since we could not evaluate the effect of RSV persistence on the expression of FcγRIV by Western blot, and we wanted to have a better understanding of the effect of RSV persistence on the expression of FcγRs, we resorted to RT-PCR so as to determine mRNA levels for all four FcγR types in MφPs and MφNs. Results are shown in Figure 7. Levels of mRNA for FcγRIV were similar in MφNs and MφPs, suggesting that RSV persistence has no effect in the expression levels of this receptor. As expected from the results of Western blots, mRNAs for FcγRIIB and FcγRIII were higher in MφPs than in MφNs, with increases of 0.4 and 2.2 fold, respectively. Unexpectedly, we found that the FcγRI mRNA was also increased (0.6 fold) in MφPs with respect to MφNs, which suggests that this receptor's expression level is also regulated at the post-transcriptional level.

2.2. Discussion

Viral persistence in cell lines is a well established model that can be used to study alterations in the expression profile of the host genome, which are caused by the constant expression of viral genes [1]. We and other researchers have shown that RSV persistence in cell cultures alters the profile of host genome expression. RSV persistence has been reported in various cell lines, and in two of them, HEp-2, (human epithelial cell line) [6] and P338D1 (murine macrophage-like cell line) [7], the characteristics resulting from persistent viral infection have been extensively studied. Although these two cell models of RSV persistence share several characteristics, they differ in some others. Thus, for instance, expression

of viral antigens in RSV-infected HEp-2 cells ranges from nil to highly positive, a high titer of extracellular virus is produced, and syncytia are formed [6]. In contrast, we have found that the characteristics of viral persistence in an RSV-infected murine macrophage-like cell line, M ϕ P (derived from P338D1), are different: 92%–96% of the cells express viral antigens—as determined by flow cytometry—low titers of extracellular infective virus are produced, and no syncytia are observed. These characteristics are still present after 87 passages. Also, RSV persistence produces an increase in the expression of Fc γ Rs and phagocytic activity in M ϕ P as compared to M ϕ N. RSV persistence induced no cell activation, as evaluated by nitrite production. However, it is clear that RSV persistence induces alterations in the expression of host genes, as demonstrated by the downregulation of ICAM-1 [41] and upregulation of membrane receptors (Fc γ Rs [8]), cytokines [8,42] and chemokines [43]. In addition, expression of the anti-apoptotic proteins Bcl-2, Bcl-X, and XIAP was enhanced, while Bcl-X and XIAP were regulated post-transcriptionally [44].

Figure 7. Fc γ Rs mRNA expression in M ϕ N and M ϕ P. Total RNA from M ϕ N and M ϕ P cells was harvested and converted to cDNA, and mRNAs for Fc γ RI, RIIB, RIII and RIV were measured by real-time PCR. Transcript levels were normalized to GAPDH mRNA expression. Data are expressed as $2^{-\Delta CT}$. Results are shown as mean $\pm$ 1 SD from three independent experiments using cells from passages 72, 83 and 87, * indicates $p < 0.05$.



In this work, we focused on exploring the mechanisms possibly responsible for the increase in Fc γ R-mediated phagocytosis that we had previously reported to occur in early passages (14–40) of the persistently infected macrophage-like cell line M ϕ P [8]. This increase is still evident after 87 passages (Figure 2). The likelihood that the observed effect on Fc γ R expression resulted from an autocrine/paracrine-like action of factors (*i.e.*, cytokines, viral particles, viral proteins) released by infected cells was ruled out, because extracellular components in conditioned medium from M ϕ P did not alter Fc γ RIIB/RIII expression in M ϕ N (Figure 3). This suggests that the higher expression of Fc γ RIIB/RIII seen in M ϕ P is not due to stimulation either by cytokines (IL-1 β , IL-6, [8]) or chemokines (RANTES, MIP-1 α , MIP-1 β , MIP-2; Tirado R. personal communication), which have been detected in M ϕ P supernatants. We also analyzed the possibility that the observed increase in membrane expression of Fc γ RIIB/RIII resulted from a reduced rate of Fc γ R internalization induced by RSV persistence. However, we found no difference between M ϕ N and M ϕ P in the kinetics of Fc γ RIIB /RIII internalization (Figure 4).

Since neither endocytic activity nor components of the conditioned medium (viral particles, viral proteins, cytokines, chemokines, *etc.*) seemed to participate in the increased Fc γ R expression, we hypothesized that the observed effect of viral persistence was caused by RSV inducing an increase in

the synthesis of FcγRs. Results obtained by flow cytometry assays of cell-membrane and intracellular protein expression (Figure 5) were consistent with our hypothesis. Since mouse cells express four different FcγR isoforms, and because up to this point we had only evaluated the expression of FcγRIIB and FcγRIII isoforms with a monoclonal antibody which recognizes both, we set out to determine the effect of RSV persistence in the expression of each of the four FcγR isoforms expressed by mouse macrophages (FcγRI, FcγRIIB, FcγRIII and FcγRIV) by Western blot with isoform-specific antibodies and by quantitative RT-PCR. We found out that RSV persistence distinctively affects expression of FcγR isoforms. Protein levels of FcγRIIB and FcγRIII are increased in MφP as compared to MφN, while FcγRI levels are not affected. FcγRIV levels could not be determined because no suitable commercial antibody is available at present. However, since the levels of mRNA for FcγRIV were not altered, it is reasonable to expect that the membrane expression of this isoform is not altered either. Higher protein levels of FcγRIIB and FcγRIII are consistent with the increased levels of mRNA for these two isoforms.

The effect of the observed changes in the expression of FcγR isoforms on the activation of effector functions of macrophages, is expected to depend on the IgG isotype involved [40]. IgG1 isotype has a higher affinity for the inhibitory FcγRIIB isoform than for the activatory FcγRIII and FcγRIV isoforms. Thus, the increases in FcγRIIB and FcγRIII expression would be expected to result in a lower cell response to immune complexes or opsonized particles containing IgG1. In contrast, IgG2a and IgG2b show higher affinity for the activatory FcγRIII and FcγRIV isoforms than for FcγRIIB. Therefore, effector functions mediated by these isotypes are less sensitive to changes in FcγRIIB. In fact, this is consistent with our observation that FcγR-mediated phagocytosis of IgG2b-opsonized erythrocytes is increased in MφP. However, it should be kept in mind that phagocytosis is a complex cellular process, involving many proteins and signaling molecules—thus, it is not expected to depend solely on the expression of the receptor involved. For example, FcγRs-mediated phagocytosis in human monocyte-derived macrophages (MDM) increases after IL-10 treatment and decreases after IFN-γ treatment, but these opposite effects do not correlate with the changes in FcγR expression induced by these cytokines [45]. The effect of RSV persistence on the expression of FcγRs mRNA suggests that RSV persistence is able to affect transcription factors, such as PU.1, which participates in FcγRs transcription [46]. Further work is underway in our laboratory in order to examine the participation of PU.1 in FcγRs transcription in RSV infected cells.

It is interesting that, while RSV-induced changes in protein levels of FcγRIIB and FcγRIII correlated with changes in the respective mRNA levels, FcγRI protein expression levels were not significantly affected by RSV persistence, although its mRNA was found to be increased. This suggests that expression of cellular genes in MφP can be regulated at either the transcriptional and/or post-transcriptional levels. Consistent with this notion are studies reporting that the expression of transmembrane protein ICAM-1 is transcriptionally downregulated [41].

Changes in FcγRs expression in macrophages persistently infected with RSV, besides its effect on phagocytosis, could affect other effector functions mediated by these receptors, such as ADCC. However, as previously mentioned, effects of the observed changes are also expected to be highly influenced by the antibody isotypes involved. RSV persistence could also have an impact on the release of inflammatory mediators, thus affecting the functions of macrophages and other cells involved in host defense, although RSV persistence does not seem to alter endogenous or exogenous

viral antigen presentation [42]. Additional studies are needed to further characterize the effects of RSV persistence in macrophages. Finally, it is important to study the effects of RSV persistence in human cells in order to evaluate the possible significance of these results to human disease, since differences have been reported in immune cells between mice and human [47].

3. Experimental Section

3.1. Virus and Cell Lines

The RSV Long strain wild type (wt RSV) has been the prototype virus used in our laboratory. Both the virus origin and the procedures for propagating, purifying and measuring its infectivity in Vero cells have been previously reported [7]. The murine macrophage-like cell line P388D1 was originally obtained from ATCC (TIB 63; Rockville, MD, USA). A sub-line of P388D1 cells persistently infected with wt RSV (MφP) was obtained in our laboratory by infecting the original cell line with RSV at a multiplicity of infection (moi) of 1, and then culturing the surviving cells [7]. As control, P388D1 cells were mock-infected and subcultured in parallel conditions (MφN). Both cell lines are maintained by subculturing. MφP passages from 72 to 87 were used in this study. Throughout the passages, the presence of viral genome was monitored by detecting the mRNA of the gene N by conventional RT-PCR, using a primer pair to amplify a segment of 1,187 bp between nucleotides 1,140–2,327: forward 5'-ATGGCTCTTAGCAAAGTC-3', Reverse 5'-TTTTTTGTTAAGCTTCAAGCTCTACATC-3'. A segment of 260 bp of GAPDH was amplified as control. RSV antigen was detected by flow cytometry on the surface of 92%–97% of the MφP using FITC-labeled polyclonal anti-RSV (Oxoid, Hampshire, UK), diluted 1:10 in PBS containing 0.1% (w/v) BSA (PBSA) (Figure 1). Extracellular viral infectivity titer was $1\text{--}2 \times 10^2$ TCID₅₀/mL per $1\text{--}2 \times 10^6$ cells, and no syncytia were observed [7].

In order to verify lack of activation, the production of nitrates by MφN and MφP was routinely evaluated by the Griess reaction. Nitrate concentrations in supernatants usually ranged from: 7.1–8.0 μmol/mL per $5\text{--}6 \times 10^6$ cells for MφN and 8.3–10.2 μmol/mL per $5\text{--}6 \times 10^6$ cells for MφP.

Both cultures were propagated in RPMI 1640 medium (GIBCO/BRL, Grand Island, NY, USA) supplemented with 0.2% NaHCO₃, 10 mM HEPES, 1 μM 2-mercaptoethanol, and 10% heat-inactivated (56 °C; 30 min) fetal bovine serum (FBS) (Biowest, Veracruz, Mexico).

3.2. Phagocytosis Assays

Phagocytosis of non-opsonized and IgG-opsonized sheep red blood cells (IgG-SRBC) was performed as previously described [8,48], but with minor modifications. In brief, SRBC were labeled with trinitrobenzene-sulfonic acid (TNBS, Sigma Aldrich Corp., St. Louis, MO, USA) 1 mg/mL for 30 min at room temperature. After washing, SRBCs were opsonized by incubation with one out of two different sub-hemagglutinating concentrations (1.52 or 2.28 μg/mL) of affinity-purified murine monoclonal anti-DNP IgG (4F8; IgG2b, produced in our laboratory). Concentrations higher than 2.28 μg/mL could not be used due to agglutination of erythrocytes.

For phagocytosis assays, MφN or MφP were plated in 96-well plates (1×10^5 cells/200 μL of RPMI per well). Non-opsonized or opsonized SRBC (25 μL/well of a 2% suspension of SRBC) were added, and the plates were incubated for 60 min at 37 °C. Non-ingested SRBC were lysed with distilled water,

and the cells were exhaustively washed with PBS. Cells in the wells were lysed with sodium dodecyl sulfate (1% in PBS), and the pseudoperoxidase activity of the ingested SRBC hemoglobin was determined by a colorimetric assay using 3,3'-diaminobenzidine as substrate. Optical density was read at 492 nm. Results are expressed as phagocytic index = (number of cells $\times$ Optical Density/100). In all experiments, phagocytosis of opsonized and non-opsonized SRBC was determined in sextuplicate wells. Controls for endogenous enzymatic activity consisted in lysates of six additional wells to which no SRBC had been added, treated under the same conditions.

3.3. Flow Cytometry

Cell surface expression of Fc γ Rs was determined by flow cytometry using anti-Fc γ RIIB/III (2.4G2, a rat monoclonal antibody that recognizes Fc γ RIIB and RIII epitopes) as unconjugated primary antibody and goat anti-rat F(ab')₂-FITC as secondary antibody, both from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Cells (5×10^5) from different M ϕ P or M ϕ N passages were washed with PBSA and fixed with 4% paraformaldehyde (30 min; room temperature). Fixed cells were washed with PBSA and incubated in PBS + 1% BSA containing 15 μ g of 2.4G2 antibody (2 h; room temperature). After incubation, cells were washed twice with PBSA, and then incubated with goat anti-rat F(ab')₂-FITC (1 h; at room temperature and protected from light). Cells were resuspended in PBS (300 μ L), and 10,000 cells were analyzed by flow cytometry (FACScan, Becton Dickinson, Mountain View, CA, USA). So as to determine intracellular proteins, cells were fixed and permeabilized with cold methanol:acetone, 1:1 (1 min; room temperature). Afterwards, cells were washed with PBSA and incubated in PBS + 1% BSA containing 10 μ g of 2.4G2 antibody (2 h; room temperature). Following that incubation, cells were washed twice with PBSA and then incubated with goat anti-rat F(ab')₂-FITC (1 h; at room temperature and protected from light). Thereafter, cells were resuspended in 300 μ L of PBS and analyzed by flow cytometry.

3.4. Conditioned Medium

Confluent M ϕ N or M ϕ P cultures (approx. 5×10^6 cells) were washed with PBS, and then seven mL of supplemented RPMI were added. After 12 or 24 h of incubation at 37 °C, supernatants were collected and cellular debris was removed by centrifugation (352 $\times$ g). (We know that cytokine concentrations in supernatants reach a plateau at 24 h [43]). Two mL of this conditioned medium were added to M ϕ N cultures (1×10^6 cells) and the cultures were incubated for 24 h at 37 °C, 5% CO₂ atmosphere. As positive stimulation control, M ϕ N were incubated with heat-killed non-typeable *Haemophilus influenzae* (NTHi) strain 2019 (moi of 100), kindly provided by Dr. Michael Apicella (Department of Microbiology, College of Medicine, University of Iowa, Iowa City, IA, USA). NTHi had been heat-killed (60 min; 70 °C water bath), washed twice by centrifugation (20 min; 900 $\times$ g; 4 °C), and suspended in PBS at a final concentration of 1×10^{10} bacteria/mL. The expression of Fc γ RIIB/RIII was determined by flow cytometry, as described above.

3.5. Endocytosis Assay

The endocytosis of cell-membrane FcγRIIB/RIII receptors was monitored with the monoclonal antibody 2.4G2. Briefly, 2.4G2 antibody was added to MφN or MφP (5×10^5) cells at 20 μg/mL (30 min, 4 °C). After this period, cells were washed thoroughly with ice-cold medium without FBS. Endocytosis was started by adding medium at 37 °C and then incubating at 37 °C for different periods of time (0, 30, 60, and 120 min). Thereafter, cells were fixed with 4% paraformaldehyde (30 min; room temperature) and washed. Afterwards, F(ab')₂-FITC against the primary antibody 2.4G2 was added. Fluorescence intensity was determined by flow cytometry as described above.

3.6. Western Blot

Whole-cell extracts were prepared from MφN or MφP (3×10^6 cells) with ice-cold lysing buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, and 0.1% SDS) containing 1× protease inhibitors cocktail (Sigma Aldrich Corp., St. Louis, MO, USA). After incubation (10 min on ice), lysates were collected and detergent-insoluble material was removed by centrifugation (20 min, 4 °C; 8,800 ×g). Protein concentration was determined using the DC protein assay kit (Bio-Rad, Hercules, CA, USA). Samples containing 20 μg of protein each were then boiled in 4× SDS sample buffer and resolved on NovexBis-Tris Mini Gels NuPAGE (Invitrogen, Carlsbad, CA, USA). Subsequently, proteins were transferred to PVDF membranes (Amersham, Piscataway, NJ, USA) and nonspecific binding sites were blocked by immersing the membranes in blocking solution (PBS, 0.1% Tween-20, and 5% low-fat milk) (1 h, room temperature). After washing the membranes with PBS-0.1% Tween-20, they were incubated with the primary antibody diluted 1:500 in blocking solution (4 °C, overnight), followed by a peroxidase-conjugated appropriate secondary antibody (Santa Cruz Biotechnology, CA, USA), diluted 1:2,500 in blocking solution (1 h, room temperature). The primary antibodies used for Western blot were anti-FcγRI goat polyclonal antibody, anti-FcγRIIB rabbit polyclonal antibody, anti-FcγRIII mouse monoclonal antibody (clone ASH1975), and anti-GAPDH, all from Santa Cruz Biotechnology. Proteins were detected using a chemiluminescent substrate, Super Signal West Dura Extended Duration substrate (Pierce Thermo Scientific, Rockford, IL, USA), and their intensities were normalized to that of GAPDH. Densitometric analysis of results was performed in images obtained with the Chemidox XRS (BioRad, Hercules, CA, USA) and analyzed with the Quantity one software (Bio-Rad).

3.7. Real Time RT-PCR

Total RNA was extracted from MφN or MφP (2×10^6 cells) with TRIzol, following the manufacturer's instructions. RNA (2 μg) was reverse transcribed with RNA transcriptase Superscript II (Invitrogen, Carlsbad, CA, USA). TaqMan real-time PCR was performed with primers and probes (assay on demand 20× mix) for FcγRs genes and GAPDH as control gene, using TaqMan assay reagent master mix (Applied Biosystems, Foster City, CA, USA). Real-time PCR was performed with 80 ng of cDNA for both target genes and endogenous control. Cycling parameters were established according to the manufacturer's protocol. Triplicate CT values were analyzed in Microsoft Excel, using the comparative CT (ΔT) method as described by the manufacturer (Applied Biosystems, Branchburg, NJ,

USA). The amount of mRNA from each target gene ($2^{-\Delta CT}$) was obtained by normalizing it to the endogenous reference (GAPDH) sample.

3.8. Statistics

Data are expressed as mean $\pm$ 1 SD from the indicated number of experiments. Differences between groups were determined by Student's *t*-test, using GraphPad Prism 4.0 software (GraphPad, San Diego CA, USA), and considered statistically significant at $p < 0.05$.

4. Conclusions

RSV persistent infection in a murine macrophage-like cell line alters FcγR expression and induces an increase in FcγR-mediated phagocytosis. Increases in FcγR expression and FcγR-mediated phagocytosis are preserved after more than 87 passages of the persistently infected culture. The increase in FcγRs expression results neither from soluble factors (cytokines) or viral products released by the infected cells, nor from an increase in the rate of FcγR internalization. Persistence of RSV genome in infected cells distinctly affects the expression of FcγR isoforms at both the mRNA and protein levels. These results indicate that RSV persistence, a phenomenon possibly more common than usually assumed, might affect the expression of cellular genes and, consequently, normal cell functions as well.

Acknowledgments

The authors thank Yvonne Rosenstein Azoulay, Carlos Rosales Ledezma and Evelyn Rivera-Toledo for helpful discussions and constructive comments during the course of this work, Olivia Reynoso Ducoing for technical assistance with Western blot assays, Rafael Palma for technical assistance with conditioned medium assays, Veronica Yakoleff for helpful comments, and Ma Pilar Ortega for English editing. The research was supported by grants from the Consejo Nacional de Ciencia y Tecnología (CONACYT), México (Grant 78862); Dirección General de Asuntos del Personal Académico, Universidad Nacional Autónoma de México (Grant PAPIIT IN23610-3); and Facultad de Medicina and PDCB, UNAM. Jorge Gaona was recipient of a scholarship from CONACYT (Scholarship 20313). This work is part of the PhD dissertation submitted by Jorge Gaona in partial fulfillment of the degree requirements.

Author Contributions

Beatriz Gómez, Enrique Ortega, and Jorge Gaona designed the experiments; Carlos Santiago-Olivares and Jorge Gaona performed the experiments; Beatriz Gómez, Enrique Ortega and Jorge Gaona prepared the manuscript. All authors read and approved the final manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

References and Notes

1. Oldstone, M.B. Anatomy of viral persistence. *PLoS Pathog.* **2009**, *5*, e1000523.
2. Anderson, C.F.; Gerber, J.S.; Mosser, D.M. Modulating macrophage function with IgG immune complexes. *J. Endotoxin Res.* **2002**, *8*, 477–481.
3. Biswas, S.K.; Chittechath, M.; Shalova, I.N.; Lim, J.Y. Macrophage polarization and plasticity in health and disease. *Immunol. Res.* **2012**, *53*, 11–24.
4. Leeansyah, E.; Wines, B.D.; Crowe, S.M.; Jaworowski, A. The mechanism underlying defective Fcγ receptor-mediated phagocytosis by HIV-1-infected human monocyte-derived macrophages. *J. Immunol.* **2007**, *178*, 1096–1104.
5. Kedzierska, K.; Azzam, R.; Ellery, P.; Mak, J.; Jaworowski, A.; Crowe, S.M. Defective phagocytosis by human monocyte/macrophages following HIV-1 infection: Underlying mechanisms and modulation by adjunctive cytokine therapy. *J. Clin. Virol.* **2003**, *26*, 247–263.
6. Martinez, I.; Lombardia, L.; Herranz, C.; Garcia-Barreno, B.; Dominguez, O.; Melero, J.A. Cultures of HEp-2 cells persistently infected by human respiratory syncytial virus differ in chemokine expression and resistance to apoptosis as compared to lytic infections of the same cell type. *Virology* **2009**, *388*, 31–41.
7. Sarmiento, R.E.; Tirado, R.; Gomez, B. Characteristics of a respiratory syncytial virus persistently infected macrophage-like culture. *Virus Res.* **2002**, *84*, 45–58.
8. Guerrero-Plata, A.; Ortega, E.; Gomez, B. Persistence of respiratory syncytial virus in macrophages alters phagocytosis and pro-inflammatory cytokine production. *Viral Immunol.* **2001**, *14*, 19–30.
9. Ravetch, J.V.; Bolland, S. IgG Fc receptors. *Annu. Rev. Immunol.* **2001**, *19*, 275–290.
10. Nimmerjahn, F.; Ravetch, J.V. Divergent immunoglobulin g subclass activity through selective Fc receptor binding. *Science* **2005**, *310*, 1510–1512.
11. Nimmerjahn, F.; Ravetch, J.V. Analyzing antibody-Fc-receptor interactions. *Methods Mol. Biol.* **2008**, *415*, 151–162.
12. Nimmerjahn, F.; Ravetch, J.V. Fcγ receptors as regulators of immune responses. *Nat. Rev. Immunol.* **2008**, *8*, 34–47.
13. Nimmerjahn, F.; Ravetch, J.V. Antibody-mediated modulation of immune responses. *Immunol. Rev.* **2010**, *236*, 265–275.
14. Collins, P.L.; Melero, J.A. Progress in understanding and controlling respiratory syncytial virus: Still crazy after all these years. *Virus Res.* **2011**, *162*, 80–99.
15. Bont, L.; Houben, M.L. The epidemiology of respiratory syncytial virus lower respiratory tract infection on young children. *Pediatr. Infect. Dis. J.* **2012**, *30*, 778–784.
16. Van Drunen Littel-van den Hurk, S.; Watkiss, E.R. Pathogenesis of respiratory syncytial virus. *Curr. Opin. Virol.* **2012**, *2*, 300–305.
17. Habibi, M.S.; Openshaw, P.J. Benefit and harm from immunity to respiratory syncytial virus: Implications for treatment. *Curr. Opin. Infect. Dis.* **2012**, *25*, 687–694.
18. Walsh, E.E. Respiratory syncytial virus infection in adults. *Semin. Respir. Crit. Care Med.* **2011**, *32*, 423–432.
19. Welliver, R.C. Review of epidemiology and clinical risk factors for severe respiratory syncytial virus (RSV) infection. *J. Pediatr.* **2003**, *143*, S112–S117.

20. Zeng, R.; Li, C.; Li, N.; Wei, L.; Cui, Y. The role of cytokines and chemokines in severe respiratory syncytial virus infection and subsequent asthma. *Cytokine* **2011**, *53*, 1–7.
21. Hall, C.B.; Douglas, R.G.; Geiman, J.M. Respiratory syncytial virus infections in infants: Quantitation and duration of shedding. *J. Pediatr.* **1976**, *89*, 11–15.
22. Szabo, S.M.; Levy, A.R.; Gooch, K.L.; Bradt, P.; Wijaya, H.; Mitchell, I. Elevated risk of asthma after hospitalization for respiratory syncytial virus infection in infancy. *Paediatr. Respir. Rev.* **2013**, *13*, S9–S15.
23. Piedimonte, G. Respiratory syncytial virus and asthma: Speed-dating or long-term relationship? *Curr. Opin. Pediatr.* **2013**, *25*, 344–349.
24. Openshaw, P.J.; Dean, G.S.; Culley, F.J. Links between respiratory syncytial virus bronchiolitis and childhood asthma: Clinical and research approaches. *Pediatr. Infect. Dis. J.* **2003**, *22*, S58–S64.
25. Sikkil, M.B.; Quint, J.K.; Mallia, P.; Wedzicha, J.A.; Johnston, S.L. Respiratory syncytial virus persistence in chronic obstructive pulmonary disease. *Pediatr. Infect. Dis. J.* **2008**, *27*, S63–S70.
26. Tripp, R.A. The brume surrounding respiratory syncytial virus persistence. *Am. J. Respir. Crit. Care Med.* **2004**, *169*, 778–779.
27. McNamara, P.S.; Smyth, R.L. The pathogenesis of respiratory syncytial virus disease in childhood. *Br. Med. Bull.* **2002**, *61*, 13–28.
28. Mejias, A.; Chavez-Bueno, S.; Gomez, A.M.; Somers, C.; Estripeaut, D.; Torres, J.P.; Jafri, H.S.; Ramilo, O. Respiratory syncytial virus persistence: Evidence in the mouse model. *Pediatr. infect. Dis. J.* **2008**, *27*, S60–S62.
29. Sutton, T.C.; Tayyari, F.; Khan, M.A.; Manson, H.E.; Hegele, R.G. T helper 1 background protects against airway hyperresponsiveness and inflammation in guinea pigs with persistent respiratory syncytial virus infection. *Pediatr. Res.* **2007**, *61*, 525–529.
30. Schwarze, J.; O'Donnell, D.R.; Rohwedder, A.; Openshaw, P.J. Latency and persistence of respiratory syncytial virus despite T cell immunity. *Am. J. Respir. Crit. Care Med.* **2004**, *169*, 801–805.
31. Bramley, A.M.; Vitalis, T.Z.; Wiggs, B.R.; Hegele, R.G. Effects of respiratory syncytial virus persistence on airway responsiveness and inflammation in guinea-pigs. *Eur. Respir. J.* **1999**, *14*, 1061–1067.
32. Hegele, R.G.; Hayashi, S.; Bramley, A.M.; Hogg, J.C. Persistence of respiratory syncytial virus genome and protein after acute bronchiolitis in guinea pigs. *Chest* **1994**, *105*, 1848–1854.
33. Riedel, F.; Oberdieck, B.; Streckert, H.J.; Philippou, S.; Krusat, T.; Marek, W. Persistence of airway hyperresponsiveness and viral antigen following respiratory syncytial virus bronchiolitis in young guinea-pigs. *Eur. Respir. J.* **1997**, *10*, 639–645.
34. Valarcher, J.F.; Bourhy, H.; Lavenue, A.; Bourges-Abella, N.; Roth, M.; Andreoletti, O.; Ave, P.; Schelcher, F. Persistent infection of b lymphocytes by bovine respiratory syncytial virus. *Virology* **2001**, *291*, 55–67.
35. Mills, B.G.; Frausto, A.; Singer, F.R.; Ohsaki, Y.; Demulder, A.; Roodman, G.D. Multinucleated cells formed *in vitro* from paget's bone marrow express viral antigens. *Bone* **1994**, *15*, 443–448.
36. Cubie, H.A.; Duncan, L.A.; Marshall, L.A.; Smith, N.M. Detection of respiratory syncytial virus nucleic acid in archival postmortem tissue from infants. *Pediatr. Pathol. Lab. Med.* **1997**, *17*, 927–938.

37. Mohapatra, S.S.; Boyapalle, S. Epidemiologic, experimental, and clinical links between respiratory syncytial virus infection and asthma. *Clin. Microbiol. Rev.* **2008**, *21*, 495–504.
38. Herra, C.M.; Keane, C.T.; Whelan, A. Increased expression of Fc gamma receptors on neutrophils and monocytes may reflect ongoing bacterial infection. *J. Med. Microbiol.* **1996**, *44*, 135–140.
39. Unkeless, J.C. Characterization of a monoclonal antibody directed against mouse macrophage and lymphocyte fc receptors. *J. Exp. Med.* **1979**, *150*, 580–596.
40. Nimmerjahn, F.; Ravetch, J.V. Fcgamma receptors: Old friends and new family members. *Immunity* **2006**, *24*, 19–28.
41. Arrevillaga, G.; Gaona, J.; Sanchez, C.; Rosales, V.; Gomez, B. Respiratory syncytial virus persistence in macrophages downregulates intercellular adhesion molecule-1 expression and reduces adhesion of non-typeable haemophilus influenzae. *Intervirology* **2012**, *55*, 442–450.
42. Guerrero-Plata, A.; Ortega, E.; Ortiz-Navarrete, V.; Gomez, B. Antigen presentation by a macrophage-like cell line persistently infected with respiratory syncytial virus. *Virus Res.* **2004**, *99*, 95–100.
43. Tirado, R.; Gómez, B. National Autonomous University of Mexico (UNAM), Ciudad Universitaria, D.F., México. Unpublished work, 2013.
44. Nakamura-Lopez, Y.; Villegas-Sepulveda, N.; Sarmiento-Silva, R.E.; Gomez, B. Intrinsic apoptotic pathway is subverted in mouse macrophages persistently infected by RSV. *Virus Res.* **2011**, *158*, 98–107.
45. Frausto-Del-Río, D.; Soto-Cruz, I.; Garay-Canales, C.; Ambriz, X.; Soldevila, G.; Carretero-Ortega, J.; Vázquez-Prado, J.; Ortega, E. Interferon gamma induces actin polymerization, rac1 activation and down regulates phagocytosis in human monocytic cells. *Cytokine* **2012**, *57*, 158–168.
46. Berclaz, P.Y.; Shibata, Y.; Whitsett, J.A.; Trapnell, B.C. GM-CSF, via PU.1, regulates alveolar macrophage Fcgamma R-mediated phagocytosis and the IL-18/IFN-gamma -mediated molecular connection between innate and adaptive immunity in the lung. *Blood* **2002**, *100*, 4193–4200.
47. Mestas, J.; Hughes, C.C. Of mice and not men: Differences between mouse and human immunology. *J. Immunol.* **2004**, *172*, 2731–2738.
48. Jungi, T.W. A rapid and sensitive method allowing photometric determination of erythrophagocytosis by mononuclear phagocytes. *J. Immunol. Methods* **1985**, *82*, 141–153.