



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to the editor

Depression, anxiety, and insomnia symptoms among patients with COVID-19: It might be too soon for conclusions

ARTICLE INFO

Keywords

Prevalence
 Depression
 Anxiety
 Insomnia
 COVID-19

We read with great interest the recently published paper from Chaomeng et al. [1] on the “Prevalence of depression, anxiety, and insomnia symptoms among patients with COVID-19: A meta-analysis of quality effects model”. This study reinforces beliefs from previous studies suggesting that patients with COVID-19 may later develop insomnia [2]. More information about this condition is very important, considering it is the most common sleep disorder in the world [3], being responsible for over 5 million consultations a year in the United States of America (USA) alone [4]. Insomnia is a pathology that is related to many other medical conditions such as hypertension [5], diabetes [5], and neurological [6], and sleep disorders [6] [7]. However, as insomnia often occur in comorbidity, we should be aware of confounders which may hamper correct interpretations of different scenarios. For instance, in Lichstein’s [8] study the sample had insomnia, but there was no clear clinical indicator of obstructive sleep apnea (OSA). After objectively studied regarding sleep related respiratory function, 43% had an apnea-hypopnea index (AHI) indicative of mild OSA, and 29% had an AHI indicative of moderate OSA [8]. Because insomnia and OSA frequently co-occur [13], it would have been interesting to explore the presence of sleep-disordered breathing in the Chaomeng et al. [1] review. On the other hand, the studies included in the above-mentioned meta-analysis relied in insomnia diagnosis by Insomnia Severity Index (ISI). Wallace & Wohlgenuth [9] did a study in which the population with obstructive sleep apnea filled the ISI questionnaire being observed that 74% of them presented with profiles compatible with clinically significant insomnia. Therefore, it is plausible that, at least some part of the individuals who were diagnosed with insomnia had also sleep apnea (Comorbid Insomnia and Sleep Apnea - COMISA). Further, if SARS-Cov-2 infection could be confirmed, additional respiratory stress could play a role in this relevant comorbidity increasing its severity. However, the symptoms that Chaomeng et al. [1] studied may not have been caused by COVID-19. Brito et al. [10] concluded for an association between psychiatric disorders and social isolation [10]; likewise with the fact that a large portion of the general population had insomnia even before the SARS-CoV2 pandemic, the symptoms identified would not necessarily be a consequence of COVID-19 infection. Moreover, unemployment has already been shown to be an external source of insomnia [11], a scenario

that got worse in 2020. For example, in Brazil between December 2020–February 2021, 14.4 million people was unemployed, setting a historical record [12]. In this context, we believe that a direct relationship between COVID-19 and primary insomnia should not be inferred at this point, even if data from Chaomeng et al. [1] is inspiring and very welcome.

Declarations of interest: none.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

The work was carried out at the Bahiana School of Medicine and Public Health.

References

- [1] Chaomeng Liu Weigang Pan, Li Li, Yanping Ren, Y.R. Bing Li, X. Ma, Prevalence of depression, anxiety, and insomnia among patients with COVID-19: a meta-analysis of quality effects model, *J. Psychosom. Res.* 147 (2021).
- [2] Lin L. Yu, J. Wang, Ou-Yang X. Yong, Q. Miao, R. Chen, Liang F. Xia, et al., The immediate impact of the 2019 novel coronavirus (COVID-19) outbreak on subjective sleep status, *Sleep Med.* 77 (xxxx) (2021) 348–354 [Internet]. Available from: <https://doi.org/10.1016/j.sleep.2020.05.018>.
- [3] J.C. Souza, R. Reimao, *Epidemiology of the insomnia, Psicol Estud.* 9 (1) (2004) 3–7.
- [4] E.S. Ford, A.G. Wheaton, T.J. Cunningham, W.H. Giles, D.P. Chapman, J.B. Croft, Trends in outpatient visits for insomnia, sleep apnea, and prescriptions for sleep medications among US adults: findings from the national ambulatory medical care survey 1999–2010, *Sleep* 37 (8) (2014) 1283–1293.
- [5] D.J. Taylor, L.J. Mallory, K.L. Lichstein, H.H. Durrence, B.W. Riedel, A.J. Bush, Comorbidity of chronic insomnia with medical problems, *Sleep* 30 (2) (2007) 213–218.
- [6] K. Banno, M.H. Kryger, Comorbid Insomnia, *Sleep Med. Clin.* 1 (3) (2006) 367–374.
- [7] S.B. Krell, V.K. Kapur, Insomnia complaints in patients evaluated for obstructive sleep apnea, *Sleep Breath.* 9 (3) (2005) 104–110.
- [8] K.L. Lichstein, K.W. Lester, R.N. Aguillard, B.W. Riedel, Occult sleep apnea in a recruited sample of older adults with insomnia, *J. Consult. Clin. Psychol.* 67 (3) (1999) 405–410.
- [9] D.M. Wallace, W.K. Wohlgenuth, Predictors of insomnia severity index profiles in United States veterans with obstructive sleep apnea, *J. Clin. Sleep Med.* 15 (12) (2019) 1827–1837.
- [10] H.K.M. Brito, G.T. Lima, L.B.C. de Oliveira, M.F. Rocha, M.V.G. de Carvalho, N. S. Costa, et al., Aggravation of psychiatric diseases during the period of social isolation: a brief literary review, *Braz. J. Heal. Rev.* 4 (2) (2021) 4678–4691.

<https://doi.org/10.1016/j.jpsychores.2021.110567>

Received 8 June 2021; Accepted 30 June 2021

Available online 7 July 2021

0022-3999/© 2021 Elsevier Inc. All rights reserved.

- [11] M.M. Ohayon, Epidemiology of insomnia: what we know and what we still need to learn, *Sleep Med. Rev.* 6 (2) (2002) 97–111.
- [12] After a year of pandemic, Brazil has record unemployment. The State of Minas Gerais [Internet]. Estado de Minas. 2021. Available from: https://www.em.com.br/app/noticia/internacional/2021/04/30/interna_internacional,1262081/apos-um-ano-de-pandemia-brasil-tem-recorde-de-desempregados.shtml (accessed 25 May 2023).
- [13] M. Meira e Cruz, M. Kryger, C. Morin, L. Palombini, C. Salles, D. Gozal, Comorbid Insomnia and Sleep Apnea: mechanisms and implications of an underrecognized and misinterpreted sleep disorder, *Sleep Medicine* (2021), <https://doi.org/10.1016/j.sleep.2021.05.043>. In press.

Lucas Gonzaga^a, Miguel Meira e Cruz^{a,b,c,d,e,*}, Cristina Salles^a

^a *International Center on Clinical Sleep Medicine and Research, Bahiana School of Medicine and Public Health, Salvador, Bahia, Brazil*

^b *European Sleep Center, Portugal*

^c *Sleep Unit, Centro Cardiovascular da Universidade de Lisboa, Lisbon School of Medicine, Portugal*

^d *Neuroimmune Pain Interface Lab, Faculdade São Leopoldo Mandic, Campinas, Brazil*

^e *European Sleep Center, Brazil*

* Corresponding author at: Sleep Unit, Centro Cardiovascular da Universidade de Lisboa, Lisbon School of Medicine, Av. Professor Egas Moniz, 1649-028 Lisbon, Portugal.
E-mail address: mcruz@medicina.ulisboa.pt (M. Meira e Cruz).