

RESEARCH ARTICLE

Open Access

Evolution of the osteoblast: skeletogenesis in gar and zebrafish

B Frank Eames^{1,2*}, Angel Amores¹, Yi-Lin Yan¹ and John H Postlethwait¹

Abstract

Background: Although the vertebrate skeleton arose in the sea 500 million years ago, our understanding of the molecular fingerprints of chondrocytes and osteoblasts may be biased because it is informed mainly by research on land animals. In fact, the molecular fingerprint of teleost osteoblasts differs in key ways from that of tetrapods, but we do not know the origin of these novel gene functions. They either arose as neofunctionalization events after the teleost genome duplication (TGD), or they represent preserved ancestral functions that pre-date the TGD. Here, we provide evolutionary perspective to the molecular fingerprints of skeletal cells and assess the role of genome duplication in generating novel gene functions. We compared the molecular fingerprints of skeletogenic cells in two ray-finned fish: zebrafish (*Danio rerio*)—a teleost—and the spotted gar (*Lepisosteus oculatus*)—a “living fossil” representative of a lineage that diverged from the teleost lineage prior to the TGD (i.e., the teleost sister group). We analyzed developing embryos for expression of the structural collagen genes *col1a2*, *col2a1*, *col10a1*, and *col11a2* in well-formed cartilage and bone, and studied expression of skeletal regulators, including the transcription factor genes *sox9* and *runx2*, during mesenchymal condensation.

Results: Results provided no evidence for the evolution of novel functions among gene duplicates in zebrafish compared to the gar outgroup, but our findings shed light on the evolution of the osteoblast. Zebrafish and gar chondrocytes both expressed *col10a1* as they matured, but both species’ osteoblasts also expressed *col10a1*, which tetrapod osteoblasts do not express. This novel finding, along with *sox9* and *col2a1* expression in developing osteoblasts of both zebrafish and gar, demonstrates that osteoblasts of both a teleost and a basally diverging ray-finned fish express components of the supposed chondrocyte molecular fingerprint.

Conclusions: Our surprising finding that the “chondrogenic” transcription factor *sox9* is expressed in developing osteoblasts of both zebrafish and gar can help explain the expression of chondrocyte genes in osteoblasts of ray-finned fish. More broadly, our data suggest that the molecular fingerprint of the osteoblast, which largely is constrained among land animals, was not fixed during early vertebrate evolution.

Background

Skeletal tissues provide invaluable traits to document vertebrate evolution and to reveal the mechanistic basis for evolutionary change. Two main processes underlie skeletal development: histogenesis—histological differentiation of skeletal tissues—and morphogenesis—acquisition of skeletal element location, shape, and size. Skeletal histogenesis involves overt differentiation of cells that secrete the extracellular matrix of cartilage and bone (chondrocytes and osteoblasts, respectively),

and follows the mesenchymal condensation of skeletogenic cells. Skeletal morphogenesis directs such differentiation events in space and time. While many studies propose a molecular genetic basis for evolutionary changes to skeletal morphogenesis [1-5], the evolution of skeletal histogenesis among vertebrates is fertile ground for additional molecular analyses [6,7].

Each cell type achieves and performs its function by employing a specific set of genes, recently termed its molecular fingerprint [6,8,9]. Are molecular fingerprints free to evolve in each vertebrate clade in response to selective pressures? Or were molecular fingerprints fixed shortly after tissues first evolved and since then have remained rather constant? For example, skeletons of

* Correspondence: b.frank@usask.ca

¹Institute of Neuroscience, University of Oregon, Eugene, OR 97403-1254, USA

Full list of author information is available at the end of the article

land animals evolved under greater gravitational stress than skeletons of aquatic animals [10]. As a result, one might hypothesize that the molecular fingerprints of skeletal cells in tetrapods, for instance, would differ from those of teleosts. Alternatively, molecular fingerprints of skeletal cells may have been fixed at the evolutionary origin of the skeleton, and other mechanisms may adapt animals to lineage-specific selective pressures. Comparative studies can reveal the extent to which molecular fingerprints of skeletal cells evolved among vertebrate clades, although the few clades examined thus far are biased towards crown-groups of both ray-finned (actinopterygian) and lobe-finned (sarcopterygian) fish (see Figure 1). Therefore, analyses of more basally-diverging lineages, such as gar or lungfish, will provide evolutionary perspective to these studies.

Comparisons among human, mouse, and chick skeletal tissues suggest that the molecular fingerprints of chondrocytes or osteoblasts do not vary greatly among tetrapods (Figure 1A). Tetrapods exhibit one type of bone tissue, but they have three types of cartilage: elastic cartilage, hyaline cartilage, and fibrocartilage [6,23,24]. Here, we focus on the predominant type of cartilage in vertebrates, hyaline cartilage, which serves as the template for bone during endochondral ossification. Collagens are the most abundant proteinaceous skeletal matrix components, and tetrapod chondrocytes and osteoblasts typically express different fibrillar collagens. In tetrapods, Collagen type 1 alpha 2 (Col1a2) is expressed abundantly in bone and is absent from cartilage, while Collagen type 2 alpha 1 (Col2a1) typifies cartilage and is not expressed in tetrapod bone [23,24]. Hyaline cartilage chondrocytes undergo a maturation process during development, when they express Collagen type 10 alpha 1 (Col10a1), but tetrapod bone does not express Col10a1 [15,25]. Despite these fundamental differences in collagen expression of tetrapod osteoblasts and chondrocytes, their molecular fingerprints also overlap; for example, both cell types express Collagen type 11 alpha 2 (Col11a2; [26]).

Skeletogenic transcription factors control molecular fingerprints of chondrocytes and osteoblasts; Sox9 is required for chondrocyte differentiation, while Runx2 is necessary for osteoblast differentiation [15]. Sox9 and Runx2 dictate skeletal cell differentiation by binding to and promoting the transcription of genes that impart identity to skeletal tissues. For instance, Sox9 directly regulates *Col2a1* expression, while Runx2 activates *Col1a2* transcription [27,28]. Much of the overlap in molecular fingerprints of tetrapod chondrocytes and osteoblasts can be attributed to Runx2 activity, which is required for chondrocyte maturation in addition to its role in osteoblast differentiation [29,30]. Perhaps such overlap is not surprising, considering that tetrapod

chondrocytes and osteoblasts differentiate from a bipotential progenitor cell, the osteochondroprogenitor, during both embryonic and adult stem cell development [31,32]. Indicative of the delicate balance required for these transcription factors to direct discrete cell lineages, Sox9 can repress Runx2 activity [31,32].

Although hyaline cartilage chondrocytes show conserved molecular fingerprints among vertebrate clades, a few studies in fish suggest that the molecular fingerprint of osteoblasts varies among vertebrates (Figure 1A). In contrast to tetrapod osteoblasts, zebrafish osteoblasts express *col10a1*, and various teleosts show evidence of Col2 in their bone matrix [11,12,33,34]. A lineage-specific genome duplication event, the teleost genome duplication (TGD, or R3), occurred at the base of the teleost radiation, and genome duplications have been thought to facilitate the origin of new gene capabilities [35-40]. For example, Sox9 does not have a direct effect on osteoblast differentiation in tetrapods, but a Sox9 duplicate in teleosts (*sox9b*) has been reported to affect bone development [41]. Therefore, one of at least three evolutionary scenarios might explain differences in the molecular fingerprint of tetrapod and teleost osteoblasts (Figure 1B). Hypothesis 1 proposes that a new function for these collagens in osteoblasts evolved in the teleost lineage, perhaps facilitated by the TGD (i.e., neofunctionalization). Hypothesis 2, like Hypothesis 1, proposes that collagen gene neofunctionalization occurred, but that this event happened in the ancestral actinopterygian osteoblast, and hence ruling out the hypothesis that the TGD facilitated this novel gene function. Hypothesis 3 proposes that the last common ancestor to both tetrapods and teleosts had osteoblast expression patterns found in today's teleosts, but that these patterns were lost secondarily in the tetrapod lineage (Figure 1B). To distinguish among these possibilities, we compared molecular fingerprints of skeletogenic cells in the teleost *Danio rerio* with the spotted gar *Lepisosteus oculatus*, a member of a teleost sister group that diverged before the TGD. Currently, the molecular fingerprints of chondrocytes and osteoblasts are completely unknown for any non-teleost actinopterygian.

Results demonstrated that gar and zebrafish share molecular fingerprints of both chondrocytes and osteoblasts. As an indication of skeletal cell molecular fingerprints, we used *in situ* hybridization on developing gar and zebrafish embryos. Specifically, we analyzed expression of the structural collagen genes *col1a2*, *col2a1*, *col10a1*, and *col11a2* in well-developed cartilage and bone, and also revealed expression of the transcription factor genes *sox9* and *runx2* during mesenchymal condensation. We found that, like osteoblasts in the teleost *Danio rerio*, gar osteoblasts expressed *col2a1* and *col10a1*. Therefore, these data refute by parsimony the




Figure 1 Comparing molecular fingerprints of skeletal cells suggests evolution of the osteoblast among vertebrate clades. A. Published collagen and transcription factor gene expression in various vertebrate clades reveals variation in the molecular fingerprint of the osteoblast (red text), whereas the chondrocyte shows a conserved molecular fingerprint. References and species cited: Teleosts = *A. semicinctus*, *B. horae*, *C. aceratus*, *C. aeneus*, *D. rerio*, *G. aculeatus*, *G. aymonieri*, *J. floridae*, *M. sanctaefilomenae*, *O. latipes*, *P. antarcticum*, *P. pangasius*, *P. pungitius*, *P. reticulata*, *R. trilineata*, *T. ladigesii* [11-14]; Mammals = *H. sapiens*, *M. musculus*, *R. norvegicus* [15-18]; Birds = *A. platyrhynchos*, *C. coturnix japonica*, *G. gallus* [15,19-22]. Almost every reference cited in this table focused on cranial skeletal tissues, but published work suggests that the summarized molecular fingerprints can be applied to skeletal cells throughout the body [19]. **B.** Three possible evolutionary scenarios explain the expression of Col2 and Col10 in osteoblasts of Teleosts, but not Tetrapods (Mammals + Birds). Hypothesis 1 suggests neofunctionalization (ectopic expression of Col2 and Col10) in osteoblasts appeared after the teleost-specific genome duplication, or TGD. Hypothesis 2 suggests neofunctionalization in osteoblasts appeared in the ancestral Actinopterygian, or ray-finned fish. Hypothesis 3 suggests Col2 and Col10 expression was present in the common ancestor of Actinopterygians and Sarcopterygians (i.e. ancestral Osteichthyan), and was subsequently lost in the Sarcopterygian (lobe-finned fish) lineage. Abbreviations: Col1 = Collagen type 1a2; Col2 = Collagen type 2a1; Col10 = Collagen type 10a1; Col11 = Collagen type 11a2.

role of the TGD in the origin of lineage-specific skeletal molecular fingerprints (Hypothesis 1) and furthermore argue that the expression of “chondrocyte” genes in osteoblasts is a shared feature of actinopterygians. More experiments will be required to distinguish between Hypotheses 2 and 3. In efforts to explain the actinopterygian expression patterns reported here, we found, surprisingly, that the “chondrogenic” transcription factor *sox9* was expressed in developing gar and zebrafish osteoblasts. We discuss these findings in a phylogenetic context and suggest that the molecular fingerprint of the primitive vertebrate osteoblast was less fixed than previously expected from studies of tetrapods.

Methods

Fish

All fish and embryos were maintained with IACUC approval, according to established protocols [42,43]. Wild-type zebrafish were of the AB strain; gar originated from animals collected in Lafourche Parish, Louisiana (courtesy of Drs. A. Ferrara and Q. Fontenot).

Histology and confocal microscopy

Embryos and larvae were processed for Alcian blue/Alizarin red staining and sectioned for histology as described previously [44]. For zebrafish, we used transgenic lines to help visualize the location and organization of specific populations of cells. *Tg(foxp2.A:EGFP)zc42* fish produce chondrocyte fluorescence [45], *Tg(sp7:EGFP)b1212* fish make fluorescent osteoblasts [46], and *Tg(fli1a:EGFP)y1* fish have fluorescence broadly among neural crest cells of the head [47]. Animals were imaged live under a confocal microscope while stained with the vital dye Alizarin red, as described previously [44].

Molecular cloning and section in situ hybridization

Whole-mount and section RNA *in situ* hybridization were carried out as described [41,48]. Zebrafish probes used were *runx2a*, *runx2b*, *sox9a*, *sox9b*, *col1a2*, *col2a1a*, *col10a1*, and *col11a2* [13,49].

Results

Early and late stages of ceratohyal and dentary development

To elucidate molecular fingerprints of chondrocytes and osteoblasts in gar and zebrafish, we focused on two cranial skeletal elements: the ceratohyal, which in mammals forms the anterior horn of the hyoid bone and whose chondrocytes form directly from mesenchyme during endochondral ossification; and the dentary, which in mammals forms the mandible bone and whose osteoblasts differentiate directly from mesenchyme during intramembranous ossification. To define equivalent developmental stages of cranial skeletogenesis between

gar and zebrafish larvae, we performed whole-mount staining with Alcian blue and Alizarin red on a variety of stages for both species (Figure 2A, B). Our histological analyses of ceratohyal development demonstrated a well-defined, Alcian blue-stained cartilage rod in gar larvae by 14 days post-fertilization (dpf) and in zebrafish larvae by 6 dpf (Figure 2A-D). In the same region earlier in development, a faint Alcian blue-stained condensation of mesenchyme was apparent in gar (8 dpf) and zebrafish (54 hours post-fertilization; hpf) (Figure 2E, F). Dentary bone formation was evidenced by substantial Alizarin red staining adjacent to the anterolateral aspects of Meckel's cartilage at 14 dpf for gar and at 6 dpf for zebrafish (Figure 2A, B, G, H). No Alizarin red staining was found in the region of the dentary in gar at 11 dpf or in zebrafish at 3 dpf (Figure 2I, J). These results established the location and timing of equivalent stages of cartilage and bone development in gar and zebrafish.

Chondrocyte molecular fingerprint

To analyze cellular and molecular features of developing chondrocytes in gar and zebrafish, we examined histologically stained sections, imaged transgenic zebrafish by confocal microscopy, and studied gene expression patterns in tissue sections. At 14 dpf, the gar ceratohyal contained hundreds of chondrocytes, none of which showed evidence of hypertrophy (Figure 3A, C). By 28 dpf, however, the gar ceratohyal showed clearly hypertrophic chondrocytes in the mid-diaphyseal region (Figure 3D). At both 14 dpf and 28 dpf, Aniline blue-stained bone matrix was apparent in an extremely thin layer of the perichondrium. The 6 dpf zebrafish ceratohyal had dozens of chondrocytes, most of which already displayed evidence of hypertrophy (Figure 3B, E; [44]). Alizarin red staining of bone matrix in the perichondrium was also evident. Therefore, well-developed ceratohyals of both the gar and zebrafish displayed equivalent cellular and histological features of chondrocyte (and perichondral bone) development.

We next sought to explore the molecular fingerprints of well-developed chondrocytes in gar and zebrafish. In both gar and zebrafish, *col1a2* expression was absent or very low in chondrocytes of the well-formed ceratohyal, although it was expressed clearly in cells of the perichondrium of both species (Figure 3F, H). Ceratohyals in both gar and zebrafish had high levels of *col2a1* transcripts (Figure 3G, I), although more mature, mid-diaphyseal chondrocytes appeared to have down-regulated transcript levels (data not shown), which is consistent with similar findings in tetrapods [50]. In addition, *col2a1* expression was detected in the perichondrium of the ceratohyal in both gar and zebrafish. Expression of *col10a1* was high in mature chondrocytes and in





Figure 2 Temporal series of skeletal preparations reveal early and late stages of cartilage and bone development in gar and zebrafish.


A-J, Alcian blue (cartilage)/Alizarin red (mineralized bone) stained fish larvae. Lateral images of larval head skeletons show that many cartilage and a few bone elements were well-formed by 14 dpf in gar (A) and by 6 dpf in zebrafish (B). Dissected and flat-mounted ceratohyals of 14 dpf gar (C) and 6 dpf zebrafish (D) showed strong Alcian blue staining and distinct boundaries of the skeletal element, which are two features of well-developed cartilage. At earlier stages, more faint and diffuse Alcian blue staining was apparent in developing ceratohyal condensations of 8 dpf gar (E) and 54 hpf zebrafish (F). Ventral views show obvious Alizarin red staining of the dentary along anterolateral aspects of Meckel's cartilage in the lower jaws of 14 dpf gar (G) and 6 dpf zebrafish (H). At earlier timepoints, no Alizarin red staining was visible in regions of the dentary in 11 dpf gar (I) and 3 dpf zebrafish (J). Scale bars: A-J = 0.25 mm. Abbreviations: ch = ceratohyal; d = dentary; dpf = days post-fertilization; hpf = hours post-fertilization; hs = hyosymplectic; Mk = Meckel's.



perichondral cells of both gar and zebrafish ceratohyals, whereas surrounding, less mature chondrocytes did not express *col10a1* (Figure 3J, L). Transcripts for *col11a2* were evident in both chondrocytes and perichondral cells of gar and zebrafish ceratohyals, although the levels of expression in mature chondrocytes appeared to be reduced relative to adjacent chondrocytes (Figure 3K,

M, and data not shown), which again is consistent with published reports in tetrapods [51].

To help explain the collagen gene expression patterns observed in cells of the well-developed ceratohyal cartilage, we analyzed expression of genes encoding Sox9 and Runx2, two transcription factors known to regulate these collagen genes in tetrapods [15,27,28,52,53].



Because specification of cell types occurs prior to their overt differentiation and transcription factor expression at this timepoint predicts skeletal cell fates [19,31], we focused on the mesenchymal condensation phase of cartilage development. Progenitor cells of the gar ceratohyal had undergone mesenchymal condensation by 7 dpf, while chondrogenic condensation of the zebrafish ceratohyal had occurred by 53 hpf (Figure 4A-C). In gar, *sox9* transcripts were abundant in chondrogenic cells, whereas levels of *runx2* expression were only slightly above background (Figure 4D, G). Due to the TGD, zebrafish has two co-orthologs of both *sox9* and *runx2* [37,41,54]. Expression patterns of *sox9* and *runx2* genes in cells of the zebrafish ceratohyal condensation at 53 hpf were similar to those seen in gar, although levels of *runx2* gene expression were much higher (Figure 4E, F, H, I). In total, these data demonstrate that gar and zebrafish share molecular fingerprints of developing chondrocytes (Table 1).

Osteoblast molecular fingerprint

To analyze the cellular and molecular features of developing osteoblasts in gar and zebrafish, we performed histological stains, confocal imaging of transgenic zebrafish, and *in situ* hybridization on tissue sections. The

dentaries of 14 dpf gar and 6 dpf zebrafish showed abundant bone matrix adjacent to Meckel's cartilage (Figure 5A, B). In both gar and zebrafish, *col1a2* expression was high in osteoblasts of the well-developed dentary (Figure 5C, D). Transcripts for *col2a1* were apparent in both gar and zebrafish dentaries, although levels detected in the zebrafish were relatively lower (Figure 5E, F). Osteoblasts of both gar and zebrafish dentaries demonstrated abundant *col10a1* expression, as well as high levels of *col11a2* transcripts (Figure 5G-J).

Table 1 Phylogenetic comparison of chondrocyte molecular fingerprint

Chondrocyte	gar	zebrafish	chick	mouse
<i>collagen type 1a2</i>	-	-	-	-
<i>collagen type 2a1</i>	+	+	+	+
<i>collagen type 10a1</i>	+	+	+	+
<i>collagen type 11a2</i>	+	+	+	+
<i>sox9</i>	+	+	+	+
<i>runx2</i>	+/-	+	+	+

The detection of particular gene expression by *in situ* hybridization is indicated as follows: '+' = high detectable expression; '-' = undetectable expression; '+/-' = low detectable expression. Gar and zebrafish data are presented here, and chick and mouse data come from references cited in Figure 1




Figure 5 Collagen expression in osteoblasts of gar and zebrafish is equivalent. **A**, Trichrome-stained coronal gar section. **B**, confocal slice of *sp7:EGFP; foxp2.A:EGFP* zebrafish, which have GFP expression in both developing osteoblasts and developing chondrocytes, respectively [45,46] and also are stained with Alizarin red to visualize mineralized bone matrix. **C-J**, *in situ* hybridization on coronal sections. Aniline blue staining in the 14 dpf gar (**A**) and Alizarin red staining in the 6 dpf zebrafish (**B**) reveals bone matrix of the dentary. Osteoblasts of the zebrafish dentary are labeled with the *sp7:EGFP* transgene. Osteoblasts of both the gar and zebrafish dentaries expressed *col1a2* (**C**, **D**), *col2a1* (**E**, **F**), *col10a1* (**G**, **H**), and *col11a2* (**I**, **J**), although expression of *col2a1a* in zebrafish osteoblasts was relatively lower than seen in gar osteoblasts. Scale bars: A-J = 15 μ m. Abbreviations: b = bone; c = cartilage; dpf = days post-fertilization; ob = osteoblast; Tri = Trichrome.




Figure 6 Transcription factor expression in osteogenic condensations of gar and zebrafish is equivalent. **A**, Trichrome-stained coronal gar section. **B**, Differential interference contrast image of zebrafish. **C**, confocal slice of *fli1a:EGFP* zebrafish. **D-I**, *in situ* hybridization on coronal sections. Osteogenic cells of the 10 dpf gar dentary have begun to secrete Aniline blue-stained bone matrix, but do not stain with Alizarin red, as this new matrix is uncalcified (data not shown). Osteogenic cells of the 72 hpf zebrafish dentary are located in an equivalent position. Similar to expression of *sox9* in the gar dentary (**D**), the zebrafish dentary expressed both *sox9a* (**E**) and *sox9b* (**F**) co-orthologs. Osteogenic cells of the gar dentary expressed transcripts for *runx2* (**G**), and osteogenic cells of the zebrafish dentary expressed both *runx2a* (**H**) and *runx2b* (**I**) co-orthologs. Scale bars: A-I = 20 μ m. Abbreviations: DIC = differential interference contrast; dpf = days post-fertilization; hpf = hours post-fertilization; Mk = Meckel's; og = osteogenic cells; Tri = Trichrome.

To help understand the expression patterns of collagen genes observed in osteoblasts of the well-developed dentary bone, we again analyzed *sox9* and *runx2* expression during mesenchymal condensation. At 10 dpf, osteogenic cells of the gar dentary had undergone mesenchymal condensation, and were beginning to secrete Aniline blue-stained bone matrix (Figure 6A). Osteogenic cells of the zebrafish dentary condensation were visible at 72 hpf (Figure 6B, C). Both *sox9* and *runx2* transcripts were apparent in osteogenic cells of the gar dentary (Figure 6D, G). The expression of *sox9* in osteogenic cells of the gar dentary was dynamic during development. While *runx2* transcript levels were high in presumptive pre-osteoblasts lateral to Meckel's cartilage at 7 dpf, *sox9* expression was not detected in these cells (Additional file 1: Figure S1). These data argue that *sox9* expression increased in osteoblasts as they began to differentiate. Similar to the single *sox9* gene in gar, both *sox9a* and *sox9b* were expressed in osteogenic cells of the developing zebrafish dentary at 72 hpf (Figure 6D-F). Also overlapping with the single *runx2* expression in the gar dentary, *runx2a* and *runx2b* transcripts were abundant in the zebrafish dentary at 72 hpf (Figure 6G-I). In summary, these data demonstrate that gar and zebrafish share molecular fingerprints of developing osteoblasts (Table 2).

Discussion

Due to their preservation in the fossil record, cartilage and bone serve as invaluable traits in understanding vertebrate evolution. Evolutionary inferences, however, often assume that the histogenesis of skeletal tissues themselves remains constant among vertebrate lineages. To be fair, cells that produce cartilage and bone (i.e., chondrocytes and osteoblasts, respectively) may have been free to evolve since their appearance roughly 500 million years ago [55,56]. Here, we ask explicitly: To what extent do vertebrate clades share expression of the

Table 2 Phylogenetic comparison of osteoblast molecular fingerprint

Osteoblast	gar	zebrafish	chick	mouse
<i>collagen type 1a2</i>	+	+	+	+
<i>collagen type 2a1</i>	+	+/-	-	-
<i>collagen type 10a1</i>	+	+	-	-
<i>collagen type 11a2</i>	+	+	+	+
<i>sox9</i>	+	+	-	-
<i>runx2</i>	+	+	+	+

The detection of particular gene expression by *in situ* hybridization is indicated as follows: '+' = high detectable expression; '-' = undetectable expression; '+/-' = low detectable expression. Gar and zebrafish data are presented here, and chick and mouse data come from references cited in Figure 1

sets of genes that characterize skeletogenic cell types (i.e., molecular fingerprints; [6,8,9])?

A molecular fingerprint that is shared among vertebrate clades would suggest evolutionary constraints on that skeletal cell type (i.e., skeletal cell types are not free to vary). For instance, cells in cranial and appendicular skeletal tissues have different embryologic origins, and so developmental constraints may limit the molecular fingerprint of a skeletal cell that appears in both regions. While future experiments can test this hypothesis more extensively, skeletogenic cells in different embryonic regions (i.e., cranial vs. appendicular) of a given individual have been shown to exhibit a conserved molecular fingerprint [19]. Another interesting potential embryonic constraint is the fact that osteoblasts have two evolutionary and developmental origins within vertebrates. During vertebrate phylogeny, bone originated in the dermis (i.e., exoskeleton), and then later appeared in the perichondrium surrounding cartilage templates (i.e., endoskeleton) [57]. While not a focus of this study, we did not find differences between molecular fingerprints of osteoblasts from the exoskeleton (e.g., those in the dentary) and endoskeleton (e.g., those surrounding the ceratohyal). Therefore, our results do not support the notion that the exoskeleton and endoskeleton have separate embryonic constraints on the molecular fingerprints of osteoblasts, but testing this hypothesis could be a fruitful avenue of future research.

A molecular fingerprint that varies among clades suggests relaxed constraints on the evolution of that cell type. One might expect variation in molecular fingerprints of skeletogenic cells among various vertebrate lineages, especially given the different selective pressures to which each vertebrate clade has been exposed. For example, the skeletons of land animals withstand a stronger effective gravitational force than do the skeletons of water-borne animals [10]. Some aquatic lineages, including sharks and other “cartilaginous” fish, and some Antarctic fish, have even lost the majority of their bony skeleton at some point during phylogeny [13,58]. Are signatures of the embryonic response to these varied selective pressures seen in the molecular fingerprints of skeletogenic cells across vertebrates?

Spurred by the reported and unexpected expression of *col10a1* and *Col2*, two markers of tetrapod chondrocytes, in osteoblasts of teleosts (Figure 1, Tables 1, 2; [11,12]), we pursued the hypothesis that molecular fingerprints of skeletogenic cells vary among vertebrate clades. Experiments revealed collagen and transcription factor gene expression in skeletal cells of hyaline cartilage and bone in the zebrafish—a teleost—and gar, which diverged in the actinopterygian lineage prior to the teleost-specific genome duplication (TGD; Figure

1; [43]). Specifically, our data distinguish among competing hypotheses to explain why osteoblasts of teleosts express *col10a1* and *Col2*, which are not expressed in osteoblasts of tetrapods. Osteoblast expression of these collagens either represents a neofunctionalization event that was specific either to the teleost lineage subsequent to the TGD (Hypothesis 1) or to the actinopterygian lineage (Hypothesis 2), or they were expressed in osteoblasts of the common ancestor of tetrapods and teleosts and subsequently lost in tetrapods (Hypothesis 3, Figure 1). Admittedly, evaluation of molecular fingerprints based upon expression of a few genes is a limited approach, but our findings on gene expression in chondrocytes and osteoblasts of the gar and zebrafish suggest evolutionary trends that could be embellished by massively parallel transcriptomics (e.g., RNA-seq; [59]).

We demonstrate that gar and zebrafish share molecular fingerprints of both chondrocytes and osteoblasts (Tables 1, 2). Despite evidence that genome duplication can facilitate the origin of new gene functions [35,36,39,40], our data reject the proposed teleost neofunctionalization hypothesis for osteoblast evolution (Hypothesis 1, Figure 1). Because both gar and zebrafish express *col2a1* and *col10a1* in their osteoblasts, the most parsimonious explanation is that these markers were present in the molecular fingerprint of the ancestral actinopterygian osteoblast. Therefore, parsimony favors Hypothesis 2, although our results do not reject Hypothesis 3, and more experiments are required to distinguish between these two possibilities. The notion that osteoblasts achieved collagen neofunctionalization somewhere in the actinopterygian lineage (Hypothesis 2, Figure 1) could be tested further by revealing the molecular fingerprint of osteoblasts in bichir, an actinopterygian diverging more basally than the gar lineage [60]. Similar studies of the lungfish or coelocanth, basally-diverging sarcopterygians, would test the possibility that *col2a1* and *col10a1* expression was present in osteoblasts of the ancestral bony fish and subsequently was lost somewhere in the sarcopterygian lineage leading to tetrapods (Hypothesis 3, Figure 1).

Our studies of skeletogenic transcription factors suggest a functional framework to explain why *col2a1* and *col10a1* are expressed in osteoblasts of actinopterygians, but not in osteoblasts of sarcopterygians (Table 2). In addition to *runx2* expression, gar and zebrafish osteoblasts express *sox9* during mesenchymal condensation of dermal bones. Developing osteoblasts of tetrapods typically express *Runx2* but not *Sox9* during mesenchymal condensation of dermal bones [19,31,61]. We propose that the expression of *sox9* in gar and zebrafish osteoblasts may explain the presence of *col2a1* and *col10a1* transcripts, given two assumptions. First, actinopterygian

osteoblasts would have to translate the *sox9* transcript we observed into Sox9 protein. Second, Sox9-responsive cis-acting regulatory elements that drive *Col2a1* expression in tetrapods [27] would have to operate similarly in actinopterygian lineages. In support of this latter notion, *col2a1* gene expression is extinguished in *sox9* mutant zebrafish [41]. Currently, Sox9 has not been shown to bind to the *Col10a1* promoter, but mis-expression of Sox9 in developing avian osteoblasts causes ectopic *Col10a1* expression, and loss of Sox9 can abrogate *Col10a1* expression in mouse [31,62], showing that *Col10a1* is downstream of Sox9 control. Deciphering the molecular mechanism by which *sox9* expression in developing osteoblasts can vary among vertebrate clades will shed light on the evolution of cell type-specific molecular fingerprints.

If the primitive condition for osteoblasts in the common ancestor of actinopterygians and sarcopterygians included expression of *sox9*, *col2a1*, and *col10a1*, then tetrapod osteoblasts would have lost expression of these genes secondarily, as outlined in Hypothesis 3. This possibility would give an entirely fresh phylogenetic context to reports of a transient chondrogenic phase during tetrapod dermal bone development [61,63]. Interestingly, sub-populations of chondrocytes in the zebrafish may lose *sox9* and *col2a1* expression as they transition to osteocytes in response to Hh signaling [64], so a developmental precedence may exist for the transitions in molecular fingerprints that Hypothesis 3 proposes during evolution. More broadly, we reveal fundamental differences between the molecular fingerprints of osteoblasts in actinopterygian and sarcopterygian clades, a finding consistent with the hypothesis that the primitive osteoblast-like cell was under reduced constraint (i.e., free to vary) during early vertebrate phylogeny.

Comparison of our data with published data for tetrapods further argues that, while the osteoblast has evolved differently between actinopterygian and sarcopterygian lineages, the molecular fingerprint of the chondrocyte appears to be conserved among vertebrates (Tables 1, 2). Although sampling of vertebrate lineages in this manner is as yet too restricted to be confident of making generalizations, limited studies on chondrocytes of hyaline cartilage in amphibians and reptiles do support this conclusion [65-67].

What mechanisms allow the osteoblast to vary among extant vertebrates, then, but constrain the chondrocyte? We argue that cell types that appear earlier in phylogeny and ontogeny are less free to vary during subsequent evolution. Cartilage appeared in the fossil record in the primitive chordate *Haikouella* 530 million years ago, and hyaline cartilage is a shared trait among chordates,

hemichordates, and even some disparate protostome taxa [6,56,68]. Apart from two clades of diverged agnathans (i.e., hagfish and lamprey), all vertebrate lineages develop bone, which appeared in fossilized conodonts from 515 million years ago [55,69]. Therefore, cartilage appeared before bone during phylogeny. In addition, cartilage appears before bone during ontogeny. Taken together, we suggest that because the chondrocyte appears before the osteoblast during both phylogeny and ontogeny, the molecular fingerprint of the chondrocyte is more constrained than that of the osteoblast. As such, our interpretation is consistent with the notion of phyletic constraint [70] and may provide a novel system by which to analyze molecular details of a developmental constraint.

Conclusions

While the molecular genetic basis for evolutionary changes to skeletal morphology has received much attention, similar studies on the evolution of skeletal cell types is limited. The set of genes, or molecular fingerprint, expressed by a cartilage- or bone-forming cell (chondrocyte or osteoblast, respectively) has been determined largely from human, mouse, and chick, thus providing an extremely limited sampling among vertebrate clades. A couple of studies demonstrated that teleost osteoblasts express collagens that normally are expressed only in chondrocytes of tetrapods, allowing us to generate specific hypotheses on the evolution of the osteoblast among vertebrates (Figure 1). Here, we test the hypothesis that the molecular fingerprint of the osteoblast underwent neofunctionalization in the teleost lineage specifically, perhaps as a result of the teleost-specific genome duplication (TGD). We compare expression of collagen and transcription factor genes during embryonic development of cartilage and bone in the teleost zebrafish *Danio rerio* and the spotted gar *Lepisosteus oculatus*, which diverged in the actinopterygian lineage prior to TGD. We find equivalent expression patterns of these genes in chondrocytes and osteoblasts of zebrafish and gar, thus refuting by parsimony the hypothesis. In addition, we show expression of the “chondrocyte” transcription factor *sox9* in developing osteoblasts of zebrafish and gar, providing a molecular explanation for the expression of “chondrocyte” genes in fish osteoblasts. Finally, we argue from comparing our results to those of tetrapods that the molecular fingerprint of the osteoblast was not fixed during early vertebrate evolution, which supports previous work on bone and dentine tissues in the fossil record [57,71], whereas the molecular fingerprint of the hyaline chondrocyte is constrained among vertebrate clades.

Additional material

Additional file 1: Figure S1. Transcription factor expression in pre-osteoblasts of gar. **A**, Trichrome-stained coronal section. **B, C**, *in situ* hybridization on coronal sections. Trichrome staining of 7 dpf gar (A) shows mesenchymal cells lateral to the condensation of Meckel's cartilage. These pre-osteoblasts do not express *sox9* (B), but express high levels of *runx2* (C). Scale bars: A-C = 50 μ m. Abbreviations: dpf = days post-fertilization; Mk = Meckel's; po = pre-osteoblasts; Tri = Trichrome.

Acknowledgements

We wish to thank Allyse Ferrara and Quentin Fontenot of Nicholls State University for gar embryos, Poh Kheng Loi, Adam Burch, Charles Kimmel, and Ruth BreMiller for help with frozen sectioning, *in situ* hybridization, and section histology. We also recognize John Dowd, Amanda Rapp, and the Fish Facility at the University of Oregon for tremendous fish housing and husbandry. This work was supported in part by grant numbers F32 DE016778-03 (B.F.E.) and P01 HD22486 and R01 RR020833 (J.H.P.) from the National Institutes of Health.

Author details

¹Institute of Neuroscience, University of Oregon, Eugene, OR 97403-1254, USA. ²Department Anatomy and Cell Biology, University of Saskatchewan, 107 Wiggins Road, Saskatoon, SK S7N 5E5, Canada.

Authors' contributions

BFE, AA, and JHP contributed to the conception of this study. BFE, AA, YLY, and JHP designed the experiments. BFE stained bone and cartilage of staged specimens. AA cloned gar genes. YLY performed *in situ* hybridizations. BFE generated all images and wrote the manuscript. All authors read, revised, and approved the final manuscript.

Received: 25 October 2011 Accepted: 5 March 2012

Published: 5 March 2012

References

1. Cresko WA, Amores A, Wilson C, Murphy J, Currey M, Phillips P, Bell MA, Kimmel CB, Postlethwait JH: **Parallel genetic basis for repeated evolution of armor loss in Alaskan threespine stickleback populations.** *Proc Natl Acad Sci USA* 2004, **101**(16):6050-6055.
2. Depew MJ, Lufkin T, Rubenstein JL: **Specification of jaw subdivisions by *Dlx* genes.** *Science* 2002, **298**(5592):381-385.
3. Harris MP, Rohner N, Schwarz H, Perathoner S, Konstantinidis P, Nusslein-Volhard C: **Zebrafish *eda* and *edar* mutants reveal conserved and ancestral roles of ectodysplasin signaling in vertebrates.** *PLoS Genet* 2008, **4**(10):e1000206.
4. Kimmel CB, Walker MB, Miller CT: **Morphing the hyomandibular skeleton in development and evolution.** *J Exp Zool B Mol Dev Evol* 2007, **308**(5):609-624.
5. Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G Jr, Dickson M, Grimwood J, Schmutz J, Myers RM, Schluter D, Kingsley DM: **Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles.** *Science* 2005, **307**(5717):1928-1933.
6. Cole AG: **A review of diversity in the evolution and development of cartilage: the search for the origin of the chondrocyte.** *Eur Cell Mater* 2011, **21**:122-129.
7. Hecht J, Stricker S, Wiecha U, Stiege A, Panopoulou G, Podsiadlowski L, Poustka AJ, Dieterich C, Ehrlich S, Suvorova J, et al: **Evolution of a core gene network for skeletogenesis in chordates.** *PLoS Genet* 2008, **4**(3):e1000025.
8. Arendt D: **Genes and homology in nervous system evolution: comparing gene functions, expression patterns, and cell type molecular fingerprints.** *Theory Biosci* 2005, **124**(2):185-197.
9. Arendt D: **The evolution of cell types in animals: emerging principles from molecular studies.** *Nat Rev Genet* 2008, **9**(11):868-882.
10. Volkmann D, Baluska F: **Gravity: one of the driving forces for evolution.** *Protoplasm* 2006, **229**(2-4):143-148.
11. Eames BF, Singer A, Smith GA, Wood ZA, Yan YL, He X, Polizzi SJ, Catchen JM, Rodriguez-Mari A, Linbo T, et al: **UDP xylose synthase 1 is required for morphogenesis and histogenesis of the craniofacial skeleton.** *Dev Biol* 2010, **341**(2):400-415.
12. Benjamin M, Ralphs JR: **Extracellular matrix of connective tissues in the heads of teleosts.** *J Anat* 1991, **179**:137-148.
13. Albertson RC, Yan YL, Titus TA, Pisano E, Vacchi M, Yelick PC, Detrich HW, Postlethwait JH: **Molecular pedomorphism underlies craniofacial skeletal evolution in Antarctic notothenioid fishes.** *BMC Evol Biol* 2010, **10**:4.
14. Kluver N, Kondo M, Herpin A, Mitani H, Schartl M: **Divergent expression patterns of *Sox9* duplicates in teleosts indicate a lineage specific subfunctionalization.** *Dev Genes Evol* 2005, **215**(6):297-305.
15. Eames BF, de la Fuente L, Helms JA: **Molecular Ontogeny of the Skeleton.** *Birth Defects Research (Part C)* 2003, **69**:93-101.
16. Marks SC Jr, Lundmark C, Christersson C, Wurtz T, Odgren PR, Seifert MF, Mackay CA, Mason-Savas A, Popoff SN: **Endochondral bone formation in toothless (osteopetrotic) rats: failures of chondrocyte patterning and type X collagen expression.** *Int J Dev Biol* 2000, **44**(3):309-316.
17. Bronckers AL, Sasaguri K, Engelse MA: **Transcription and immunolocalization of *Runx2/Cbfa1/Pebp2alphaA* in developing rodent and human craniofacial tissues: further evidence suggesting osteoclast phagocytosis osteocytes.** *Microsc Res Tech* 2003, **61**(6):540-548.
18. de Crombrughe B, Lefebvre V, Behringer RR, Bi W, Murakami S, Huang W: **Transcriptional mechanisms of chondrocyte differentiation.** *Matrix Biol* 2000, **19**(5):389-394.
19. Eames BF, Helms JA: **Conserved molecular program regulating cranial and appendicular skeletogenesis.** *Dev Dyn* 2004, **231**(1):4-13.
20. Eames BF, Schneider RA: **The genesis of cartilage size and shape during development and evolution.** *Development* 2008, **135**(23):3947-3958.
21. Merrill AE, Eames BF, Weston SJ, Heath T, Schneider RA: **Mesenchyme-dependent BMP signaling directs the timing of mandibular osteogenesis.** *Development* 2008, **135**(7):1223-1234.
22. Walchli C, Koch M, Chiquet M, Odermatt BF, Trueb B: **Tissue-specific expression of the fibril-associated collagens XII and XIV.** *J Cell Sci* 1994, **107**(Pt 2):669-681.
23. Gray H, Williams PL: *Gray's anatomy* Edinburgh: C. Livingstone; 1989.
24. Ham AW, Cormack DH: *Ham's histology.* 9 edition. Philadelphia: Lippincott; 1987.
25. Linsenmayer TF, Eavey RD, Schmid TM: **Type X collagen: a hypertrophic cartilage-specific molecule.** *Pathol Immunopathol Res* 1988, **7**(1-2):14-19.
26. Li SW, Arita M, Kopen GC, Phinney DG, Prockop DJ: **A 1,064 bp fragment from the promoter region of the *Col11a2* gene drives *lacZ* expression not only in cartilage but also in osteoblasts adjacent to regions undergoing both endochondral and intramembranous ossification in mouse embryos.** *Matrix Biol* 1998, **17**(3):213-221.
27. Lefebvre V, Huang W, Harley VR, Goodfellow PN, de Crombrughe B: **SOX9 is a potent activator of the chondrocyte-specific enhancer of the pro alpha1(I) collagen gene.** *Mol Cell Biol* 1997, **17**(4):2336-2346.
28. Kern B, Shen J, Starbuck M, Karsenty G: ***Cbfa1* contributes to the osteoblast-specific expression of type I collagen genes.** *J Biol Chem* 2001, **276**(10):7101-7107.
29. Hoshi K, Komori T, Ozawa H: **Morphological characterization of skeletal cells in *Cbfa1*-deficient mice.** *Bone* 1999, **25**(6):639-651.
30. Kim IS, Otto F, Zabel B, Mundlos S: **Regulation of chondrocyte differentiation by *Cbfa1*.** *Mech Dev* 1999, **80**(2):159-170.
31. Eames BF, Sharpe PT, Helms JA: **Hierarchy revealed in the specification of three skeletal fates by *Sox9* and *Runx2*.** *Dev Biol* 2004, **274**(1):188-200.
32. Zhou G, Zheng Q, Engin F, Munivez E, Chen Y, Sebald E, Krakow D, Lee B: **Dominance of SOX9 function over RUNX2 during skeletogenesis.** *Proc Natl Acad Sci USA* 2006, **103**(50):19004-19009.
33. Avaron F, Hoffman L, Guay D, Akimenko MA: **Characterization of two new zebrafish members of the hedgehog family: atypical expression of a zebrafish indian hedgehog gene in skeletal elements of both endochondral and dermal origins.** *Dev Dyn* 2006, **235**(2):478-489.
34. Li N, Felber K, Elks P, Croucher P, Roehl HH: **Tracking gene expression during zebrafish osteoblast differentiation.** *Dev Dyn* 2009, **238**(2):459-466.
35. Brodie ED: **How an ancient genome duplication electrified modern fish.** *Proc Natl Acad Sci USA* 2010, **107**(51):21953-21954.
36. Wagner A: **Gene duplications, robustness and evolutionary innovations.** *Bioessays* 2008, **30**(4):367-373.

37. Amores A, Force A, Yan YL, Joly L, Amemiya C, Fritz A, Ho RK, Langeland J, Prince V, Wang YL, et al: **Zebrafish hox clusters and vertebrate genome evolution.** *Science* 1998, **282**(5394):1711-1714.
38. Postlethwait J, Amores A, Cresko W, Singer A, Yan YL: **Subfunction partitioning, the teleost radiation and the annotation of the human genome.** *Trends Genet* 2004, **20**(10):481-490.
39. Ohno S: **Gene duplication and the uniqueness of vertebrate genomes circa 1970-1999.** *Semin Cell Dev Biol* 1999, **10**(5):517-522.
40. Taylor JS, Raes J: **Duplication and divergence: the evolution of new genes and old ideas.** *Annu Rev Genet* 2004, **38**:615-643.
41. Yan YL, Willoughby J, Liu D, Crump JG, Wilson C, Miller CT, Singer A, Kimmel C, Westerfield M, Postlethwait JH: **A pair of Sox: distinct and overlapping functions of zebrafish sox9 co-orthologs in craniofacial and pectoral fin development.** *Development* 2005, **132**(5):1069-1083.
42. Westerfield M: **The zebrafish book. A guide for the laboratory use of zebrafish (*Danio rerio*)** Eugene: Univ. of Oregon Press; 2007.
43. Amores A, Catchen J, Ferrara A, Fontenot Q, Postlethwait JH: **Genome evolution and meiotic maps by massively parallel DNA sequencing: spotted gar, an outgroup for the teleost genome duplication.** *Genetics* 2011, **188**(4):799-808.
44. Eames B, Yan Y, Swartz M, Levic D, Knapik E, Postlethwait J, Kimmel C: **Mutations in fam20b and xylosyltransferase1 reveal that cartilage matrix controls timing of endochondral ossification through inhibition of chondrocyte maturation.** *PLoS Genet* 2011, **7**(8):e1002246.
45. Bonkowski JL, Wang X, Fujimoto E, Lee JE, Chien CB, Dorsky RI: **Domain-specific regulation of foxP2 CNS expression by lef1.** *BMC Dev Biol* 2008, **8**:103.
46. DeLaurier A, Eames BF, Blanco-Sanchez B, Peng G, He X, Swartz ME, Ullmann B, Westerfield M, Kimmel CB: **Zebrafish sp7:EGFP: a transgenic for studying otic vesicle formation, skeletogenesis, and bone regeneration.** *Genesis* 2010, **48**(8):505-511.
47. Crump JG, Swartz ME, Eberhart JK, Kimmel CB: **Moz-dependent Hox expression controls segment-specific fate maps of skeletal precursors in the face.** *Development* 2006, **133**(14):2661-2669.
48. Rodriguez-Mari A, Yan YL, Bremiller RA, Wilson C, Canestro C, Postlethwait JH: **Characterization and expression pattern of zebrafish Anti-Mullerian hormone (Amh) relative to sox9a, sox9b, and cyp19a1a, during gonad development.** *Gene Expr Patterns* 2005, **5**(5):655-667.
49. Yokoi H, Yan YL, Miller MR, BreMiller RA, Catchen JM, Johnson EA, Postlethwait JH: **Expression profiling of zebrafish sox9 mutants reveals that Sox9 is required for retinal differentiation.** *Dev Biol* 2009, **329**(1):1-15.
50. Zhao Q, Eberspaecher H, Lefebvre V, De Crombrughe B: **Parallel expression of Sox9 and Col2a1 in cells undergoing chondrogenesis.** *Dev Dyn* 1997, **209**(4):377-386.
51. Wai AW, Ng LJ, Watanabe H, Yamada Y, Tam PP, Cheah KS: **Disrupted expression of matrix genes in the growth plate of the mouse cartilage matrix deficiency (cmd) mutant.** *Dev Genet* 1998, **22**(4):349-358.
52. Ducey P, Zhang R, Geoffroy V, Ridall AL, Karsenty G: **Osf2/Cbfa1: a transcriptional activator of osteoblast differentiation.** *Cell* 1997, **89**(5):747-754.
53. Ng LJ, Wheatley S, Muscat GE, Conway-Campbell J, Bowles J, Wright E, Bell DM, Tam PP, Cheah KS, Koopman P: **SOX9 binds DNA, activates transcription, and coexpresses with type II collagen during chondrogenesis in the mouse.** *Dev Biol* 1997, **183**(1):108-121.
54. Flores MV, Tsang VW, Hu W, Kaley-Zylinska M, Postlethwait J, Crosier P, Crosier K, Fisher S: **Duplicate zebrafish runx2 orthologues are expressed in developing skeletal elements.** *Gene Expr Patterns* 2004, **4**(5):573-581.
55. Sansom IJ, Smith MP, Armstrong HA, Smith MM: **Presence of the earliest vertebrate hard tissue in conodonts.** *Science* 1992, **256**(5061):1308-1311.
56. Chen J-Y, Huang D-Y, Li C-W: **An early Cambrian craniate-like chordate.** *Nature* 1999, **402**(6761):518-522.
57. Smith MM, Hall BK: **Development and evolutionary origins of vertebrate skeletogenic and odontogenic tissues.** *Biol Rev Camb Philos Soc* 1990, **65**(3):277-373.
58. Eames BF, Allen N, Young J, Kaplan A, Helms JA, Schneider RA: **Skeletogenesis in the swell shark *Cephaloscyllium ventriosum*.** *J Anat* 2007, **210**(5):542-554.
59. Tariq MA, Kim HJ, Jejelowo O, Pourmand N: **Whole-transcriptome RNAseq analysis from minute amount of total RNA.** *Nucleic Acids Res* 2011, **39**(18):e120.
60. Inoue JG, Miya M, Tsukamoto K, Nishida M: **Basal actinopterygian relationships: a mitogenomic perspective on the phylogeny of the "ancient fish".** *Mol Phylogenet Evol* 2003, **26**(1):110-120.
61. Aberg T, Rice R, Rice D, Thesleff I, Waltimo-Siren J: **Chondrogenic potential of mouse calvarial mesenchyme.** *J Histochem Cytochem* 2005, **53**(5):653-663.
62. Ikegami D, Akiyama H, Suzuki A, Nakamura T, Nakano T, Yoshikawa H, Tsumaki N: **Sox9 sustains chondrocyte survival and hypertrophy in part through Pik3ca-Akt pathways.** *Development* 2011, **138**(8):1507-1519.
63. Abzhanov A, Rodda SJ, McMahon AP, Tabin CJ: **Regulation of skeletogenic differentiation in cranial dermal bone.** *Development* 2007, **134**(17):3133-3144.
64. Hammond CL, Schulte-Merker S: **Two populations of endochondral osteoblasts with differential sensitivity to Hedgehog signalling.** *Development* 2009, **136**(23):3991-4000.
65. Morishi T, Shibata Y, Tsukazaki T, Yamaguchi A: **Expression profile of *Xenopus* banded hedgehog, a homolog of mouse Indian hedgehog, is related to the late development of endochondral ossification in *Xenopus laevis*.** *Biochem Biophys Res Commun* 2005, **328**(4):867-873.
66. Kerney R, Hanken J: **Gene expression reveals unique skeletal patterning in the limb of the direct-developing frog, *Eleutherodactylus coqui*.** *Evol Dev* 2008, **10**(4):439-448.
67. Lopez D, Duran AC, de Andres AV, Guerrero A, Blasco M, Sans-Coma V: **Formation of cartilage in the heart of the Spanish terrapin, *Mauremys leprosa* (Reptilia, Chelonia).** *J Morphol* 2003, **258**(1):97-105.
68. Rychel AL, Smith SE, Shimamoto HT, Swalla BJ: **Evolution and development of the chordates: collagen and pharyngeal cartilage.** *Mol Biol Evol* 2006, **23**(3):541-549.
69. Janvier P: *Early vertebrates* Oxford: Oxford University Press; 1996.
70. Gilbert SF: *Developmental Biology*. 2 edition. Sunderland: Sinauer Associates, Inc; 1988.
71. Donoghue PC, Sansom IJ, Downs JP: **Early evolution of vertebrate skeletal tissues and cellular interactions, and the canalization of skeletal development.** *J Exp Zool B Mol Dev Evol* 2006, **306**(3):278-294.

doi:10.1186/1471-2148-12-27

Cite this article as: Eames et al.: Evolution of the osteoblast: skeletogenesis in gar and zebrafish. *BMC Evolutionary Biology* 2012 12:27.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

