

# SEACing the GAP that nEGOCiates TORC1 activation

## Evolutionary conservation of Rag GTPase regulation

Nicolas Panchaud<sup>†</sup>, Marie-Pierre Péli-Gullit<sup>†</sup>, and Claudio De Virgilio\*

Department of Biology; Division of Biochemistry; University of Fribourg; Fribourg, Switzerland

<sup>†</sup>These authors contributed equally to this work.

**T**he target of rapamycin complex 1 (TORC1) regulates eukaryotic cell growth in response to a variety of input signals. In *S. cerevisiae*, amino acids activate TORC1 through the Rag guanosine triphosphatase (GTPase) heterodimer composed of Gtr1 and Gtr2 found together with Ego1 and Ego3 in the EGO complex (EGOC). The GTPase activity of Gtr1 is regulated by the SEA complex (SEAC). Specifically, SEACIT, a SEAC subcomplex containing Iml1, Npr2, and Npr3 functions as a GTPase activator (GAP) for Gtr1 to decrease the activity of TORC1 and, consequently, growth, after amino acid deprivation. Here, we present genetic epistasis data, which show that SEACAT, the other SEAC subcomplex, containing Seh1, Sea2–4, and Sec13, antagonizes the GAP function of SEACIT. Orthologs of EGOC (Ragulator), SEACIT (GATOR1), and SEACAT (GATOR2) are present in higher eukaryotes, highlighting the remarkable conservation, from yeast to man, of Rag GTPase and TORC1 regulation.

**Keywords:** Rag GTPases, TOR complex 1, EGO complex, SEA complex, amino acid signaling, Iml1-Npr2-Npr3 Rag GTPase GAP complex, GATOR1, GATOR2, Seh1, Sec13

**Abbreviations:** TORC1, TOR complex 1; EGOC, EGO complex; SEAC, SEA complex

Submitted: 07/12/2013

Accepted: 08/01/2013

<http://dx.doi.org/10.4161/cc.26000>

\*Correspondence to: Claudio De Virgilio;  
Email: Claudio.DeVirgilio@unifr.ch

### Introduction

The target of rapamycin complex 1 (TORC1) is a structurally and functionally conserved regulator of eukaryotic cell growth that adapts anabolic and catabolic processes in response to a variety of inputs, such as growth factors, cellular stress, energy, and nutrients.<sup>1–4</sup> Amino acids, especially branched-chain amino acids like leucine, represent essential stimuli for TORC1 activation.<sup>5–7</sup> Members of the conserved Rag family of guanosine

triphosphatases (GTPases) mediate amino acid signaling to TORC1: in higher eukaryotes, RagA or RagB forms a heterodimer with RagC or RagD, whereas in *S. cerevisiae*, Gtr1 dimerizes with Gtr2. When RagA, RagB, or Gtr1 is bound to GTP, and RagC, RagD, or Gtr2 to GDP, the respective heterodimer is in its active, TORC1-stimulating conformation.<sup>8–11</sup> In mammalian cells, Rag GTPases do not directly activate TORC1, but trigger TORC1 relocalization from the cytoplasm to the limiting membrane of the lysosome, where it can be activated by the GTPase Rheb.<sup>5,10,12</sup> In *S. cerevisiae*, TORC1 remains associated with the limiting membrane of the vacuole (the yeast equivalent to the lysosome) irrespective of the presence or absence of leucine. Moreover, the yeast Rheb ortholog, Rhb1, is likely not required for the regulation of TORC1.<sup>4,9</sup> Thus, the mechanisms by which the Gtr1–Gtr2 heterodimer controls TORC1 function in *S. cerevisiae* remains mysterious.

Gtr and Rag heterodimers are core switches that fulfill their function as part of larger protein complexes. In *S. cerevisiae*, Gtr1–Gtr2 associates with Ego1 and Ego3 to form the EGO complex (EGOC). Ego1 is N-terminally myristoylated and palmitoylated and thus tethers the EGOC to the vacuolar membrane.<sup>9,13–17</sup> Ego3, the precise function of which remains unknown, forms homodimers that, like the C-terminal domains of Gtr1 and Gtr2, are structurally similar to members of the Roadblock/LC7 superfamily of proteins.<sup>15,17</sup> In mammals, Rag GTPase heterodimers associate with the Ego1 equivalent p18 (LAMTOR1), the


Ego3-Ego3-related heterodimer p14-MP1 (LAMTOR2-LAMTOR3),<sup>17</sup> C7orf59 (LAMTOR4), and HBXIP (LAMTOR5), which form the Ragulator complex.<sup>12</sup> Like the EGO, the Ragulator complex sits on the limiting membrane of the lysosome by virtue of lipidation of p18, which is the only Rag–Ragulator subunit lacking structural resemblance with Roadblock domains (RDs).<sup>5,6,12</sup> The entire pentameric Ragulator complex is proposed to act as the guanine nucleotide exchange factor (GEF) for RagA and RagB.<sup>18</sup> Whether the EGO possesses similar GEF activity remains questionable, because *S. cerevisiae* cells lack apparent orthologs of HBXIP and C7orf59, and guanine nucleotide exchange on Gtr1 is rather proposed to be stimulated by a Vam6-dependent mechanism.<sup>9</sup> A GTPase-activating protein

(GAP) that regulates Rag/Gtr proteins has, until recently, remained elusive.


Recently, subunits of the octameric vacuolar Seh1-associated complex (SEAC) were implicated in negative regulation of TORC1 in yeast.<sup>19–22</sup> In an effort to clarify the relationship between SEAC and TORC1, we discovered in genetic epistasis analyses that the Iml1–Npr2–Npr3 SEAC subcomplex, which we now name SEACIT (for SEAC subcomplex Inhibiting TORC1 signaling), negatively regulates TORC1 through Gtr1 within the EGO.<sup>23</sup> Moreover, in line with our genetic data, we found that leucine deprivation triggered Iml1 to transiently interact with Gtr1 (in a Npr2- and Npr3-dependent manner) to stimulate its intrinsic GTPase activity. Of note, both Npr2 and Npr3 contain a N-terminal longin domain, the structure of which is closely related to RDs and may serve as platform for Rag GTPases.<sup>24</sup> The GAP activity of SEACIT is conserved, as the orthologous complex in *Drosophila* and human cells (i.e., DEPDC5–Npr2–Npr13), coined GATOR1, also acts as a GAP toward RagA and RagB.<sup>25</sup> Intriguingly, various glioblastomas and ovarian cancers contain nonsense or frameshift

mutations or truncating deletions in GATOR1-encoding genes, and a number of cancer cell lines with homozygous deletions in *DEPDC5*, *NPRL2*, or *NPRL3* exhibit hyperactive mTORC1 that is insensitive to amino acid deprivation.<sup>25</sup> Since these GATOR1-inactivating mutations also cause hypersensitivity to the TORC1 inhibitor rapamycin in mammalian cells, they may help to predict the therapeutic benefit of clinically approved TORC1 inhibitors in cancer treatments.<sup>25</sup>

In addition to Iml1, Npr2, and Npr3 (SEACIT), the octameric SEAC also contains Sea2, Sea3, Sea4, Seh1, and Sec13, orthologs of the mammalian and *Drosophila* GATOR2 subcomplex proteins WDR24, WDR59, Mios, Seh1L, and Sec13, respectively. These proteins form the other SEAC-subcomplex, which we now name SEACAT (for SEAC subcomplex Activating TORC1 signaling). Except for Sec13, all of the GATOR2 components have been implicated in negative regulation of GATOR1 in higher eukaryotes.<sup>25</sup> Similarly, yeast Sea2, Sea3, and Sea4 antagonize, although redundantly, the SEACIT-mediated TORC1 inhibition.<sup>23</sup> However, roles for yeast Seh1, or either



**Figure 1.** Loss of Iml1 suppresses the TORC1 activation defect in *sec13*Δ (A) and *seh1*Δ (B) cells. Indicated (prototrophic) strains expressing a plasmid-based copy of Sch9<sup>T570A</sup>-HA<sub>5</sub> were grown exponentially at 25 °C (A) or 30 °C (B). Immunoblots detecting the level of phosphorylation within the C terminus of Sch9 were used to quantify in vivo TORC1 activity as previously described.<sup>50</sup> Bar graphs refer to the mean ratio (± S.D.) of hyperphosphorylated/hypophosphorylated Sch9 from 3 independent experiments, normalized to the values for wild-type cells.



**Figure 2.** Conserved regulators of the Rag-family GTPases. The yeast SEAC is composed of 2 subcomplexes, SEACIT and SEACAT. SEACAT antagonizes the GAP-function of SEACIT. Vam6 is thought to be the GEF for Gtr1, which resides in the EGO on the vacuolar membrane. Similarly, the mammalian (and *Drosophila*) GATOR complex is composed of the 2 subcomplexes GATOR1 and GATOR2. GATOR2 antagonizes the GAP-function of GATOR1. Whether or not mammalian Vam6 orthologs (i.e., the TGF-β receptor-associated protein 1 [TRAP1 or TGFBRAP1] and the TRAP1-like protein [TLP], aka hVPS39)<sup>45,46,52–54</sup> act as a RagA/B GEF is unclear, rather the pentameric Ragulator complex, acting downstream of the vacuolar ATPase, is reported to serve this function. For details, please see text.

yeast or metazoan Sec13 upstream of the Rag GTPases are currently not reported.

## Results and Discussion

To determine if Sec13, like other SEACAT components, controls TORC1 activity via SEACIT, we assayed TORC1 activity in a temperature-sensitive *sec13<sup>ts</sup>* (*sec13-1*)<sup>26</sup> mutant. As is shown in **Figure 1A**, the *sec13<sup>ts</sup>* mutant exhibited significantly reduced TORC1 activity when grown at the permissive temperature of 25 °C. This reduced TORC1 activity matches well with the observation that *sec13-1* is synthetic lethal when combined with a hypomorphic allele of LST8 (i.e., *lst8-1* for lethal with sec-thirteen), which encodes a stimulatory component in TOR-containing complexes.<sup>27,28</sup> Importantly, loss of Iml1 strongly activated TORC1 in both wild-type and *sec13<sup>ts</sup>* mutant cells. Similarly, we also observed that loss of Seh1 resulted in a significant reduction of TORC1 activity, which was fully suppressed in the absence of Iml1 (**Fig. 1B**). These genetic data therefore support a model in which Sec13 and Seh1, together with the other SEACAT components, promote TORC1 activity through inhibition of the GAP function of SEACIT. These results extend the remarkable evolutionary conservation of TORC1 regulation by Rag GTPases and delineate an inhibitory role for the pentameric SEACAT/GATOR2 subcomplex upstream of the SEACIT/GATOR1 subcomplex (**Fig. 2**).

Curiously, both Sec13 and Seh1 not only function within the SEAC, but also within the nuclear pore complex (NPC) as part of the conserved heptameric Nup84 subcomplex that is essential for the overall architecture of the NPC and consequently the transport of mRNAs and macromolecules (e.g., pre-ribosomes) across the nuclear membrane.<sup>29</sup> Moreover, Sec13 also associates with Sec31 into a heterotetramer, which forms the outer shell of coatmer complex II (COPII) coated vesicles of the secretory pathway that bud off from the endoplasmic reticulum (ER).<sup>30,31</sup> The occurrence of Sec13 and Seh1 in functionally different protein complexes suggests that their 3-dimensional structure, which is characterized, like those of all other SEACAT subunits, by the


presence of WD-40 repeats that form  $\beta$ -propellers,<sup>19,21</sup> renders them particularly well suited to serve as building and/or scaffolding blocks within larger protein complexes. Given these observations, it is tempting to speculate that Sec13/Seh1 serve to couple nuclear-to-cytoplasmic mRNA/protein transport or protein secretion to TORC1 control. For instance, compromised nuclear pore function or secretion may tie up or jam Seh1 and/or Sec13, thereby causing reduced SEAC assembly and, consequently, downregulation of TORC1. Interestingly, a genome-scale RNA interference screen by dsRNA reverse-transfection on living *Drosophila* cell microarrays identified nuclear pore components as TORC1 regulators.<sup>32</sup> In a similar vein, alterations in the yeast secretory pathway have also been found to converge on TORC1 regulation. For instance, loss of the Golgi  $\text{Ca}^{2+}/\text{Mn}^{2+}$  ATPase Pmr1 strongly increased the secretion of (heterologous) proteins that transit through the secretory pathway and, based on genetic experiments, also caused TORC1 activation (e.g., *pmr1* $\Delta$  suppressed the rapamycin-sensitive phenotype of the *lst8-1* mutation).<sup>33,34</sup> Conversely, addition of the secretory pathway inhibitor tunicamycin and inactivation of the Rab escort protein Mrs6 both strongly inhibited TORC1-dependent phosphorylation of Sch9.<sup>28,35</sup> In sum, these observations lend support to a model in which both NPC function

and secretory pathway flux are part of an increasing number of physiological cues (including v-ATPase activity, leucyl-tRNA synthetase function, glutaminolysis-driven production of  $\alpha$ -ketoglutarate, glucose and amino acid levels, vesicle trafficking, or actin polarization),<sup>9,36-46</sup> which may converge on Rag GTPase-mediated control of TORC1 (**Fig. 3**). Future studies should therefore aim at deciphering whether any of these cues may fine-tune TORC1 by regulating the GTP loading status of Rag GTPases through the SEACIT/GATOR1 and/or SEACAT/GATOR2 complexes.

## Materials and Methods

### Growth conditions, strains, and plasmids

Unless stated otherwise, prototrophic strains were pre-grown overnight in synthetic defined dropout (SD; 0.17% yeast nitrogen base, 0.5% ammonium sulfate, 0.2% [-adenine/-histidine/-leucine/-uracil/-tryptophan] dropout mix, and 2% glucose). For TORC1 activity assays, cells were diluted to an OD<sub>600</sub> of 0.2 and further grown at 30 °C until they reached an OD<sub>600</sub> of 0.8. The following isogenic *S. cerevisiae* strains (all wild-type for *LYS2* and *MET15* in the BY4741/2 background)<sup>47</sup> were used in this study: MAT $\alpha$  *his3* $\Delta$ 1, *leu2* $\Delta$ 0, *ura3* $\Delta$ 0 (YL515; WT)<sup>9</sup>; MAT $\alpha$  *iml1* $\Delta$ ::*kanMX*, *his3* $\Delta$ 1, *leu2* $\Delta$ 0, *ura3* $\Delta$ 0 (NP04-4C)<sup>23</sup>; MAT $\alpha$  *seh1* $\Delta$ ::*kanMX*,



**Figure 3.** Physiological cues, which may regulate TORC1 through the Rag GTPase module. Red check marks indicate the existence of experimental data supporting (in yeast or mammalian cells) a model in which the respective cue impinges on Rag GTPase regulation (please see text for corresponding references). Currently speculative processes are denoted with a question mark.



*his3Δ1, leu2Δ0, ura3Δ0* (MP308-7A); MAT $\alpha$  *seh1Δ::kanMX, iml1Δ::kanMX, his3Δ1, leu2Δ0, ura3Δ0* (MP308-8B); MAT $\alpha$  *sec13<sup>ts</sup>-kanMX, his3Δ1, leu2Δ0, ura3Δ0* (MP309-5D); and MAT $\alpha$  *sec13<sup>ts</sup>-kanMX, iml1Δ::kanMX, his3Δ1, leu2Δ0, ura3Δ0* (MP309-9A). The original *sec13<sup>ts</sup>* (MAT $\alpha$  *sec13-1-kanMX, his3Δ1, leu2Δ0, ura3Δ0, met15Δ0*)<sup>48</sup> and *seh1Δ* (MAT $\alpha$  *seh1Δ::kanMX, his3Δ1, leu2Δ0, ura3Δ0, met15Δ0*)<sup>49</sup> mutants were rendered wild-type for *MET15* by backcrossing with YL515. Sequencing of the *sec13<sup>ts</sup>* ORF revealed that this allele carries 2 mutations, which change Lys<sup>44</sup> and Ser<sup>224</sup> in Sec13 to Glu<sup>44</sup> and Asp<sup>224</sup>, respectively. All strains carried the following plasmids: pRS413-Sch9<sup>T570A</sup>-HA<sub>5</sub>,<sup>50</sup> pRS415,<sup>51</sup> and pRS416.<sup>51</sup>

#### TORC1 activity assays

TORC1 activity was determined by quantification of the phosphorylation of the C-terminal part of HA<sub>5</sub>-tagged Sch9<sup>T570A</sup>, which contains 5 TORC1 phosphorylation sites, as described previously.<sup>9,50</sup> Briefly, following chemical cleavage with NTCB, extracts were separated by 7.5% SDS-PAGE, and membranes were probed with anti-HA antibodies (12CA5) and anti-mouse IgG antibodies coupled to HRP (Biorad).

#### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

#### Acknowledgments

We thank Michael Costanzo for strains, Floriane Jaquier for technical assistance, and R Loewith for plasmids and critical and valuable comments on the manuscript. This research was supported by the Canton of Fribourg and the Swiss National Science Foundation (CDV).

#### References

- Soulard A, Cohen A, Hall MN. TOR signaling in invertebrates. *Curr Opin Cell Biol* 2009; 21:825-36; PMID:19767189; <http://dx.doi.org/10.1016/j.ceb.2009.08.007>
- Zoncu R, Efeyan A, Sabatini DM. mTOR: from growth signal integration to cancer, diabetes and ageing. *Nat Rev Mol Cell Biol* 2011; 12:21-35; PMID:21157483; <http://dx.doi.org/10.1038/nrm3025>
- De Virgilio C, Loewith R. The TOR signalling network from yeast to man. *Int J Biochem Cell Biol* 2006; 38:1476-81; PMID:16647875; <http://dx.doi.org/10.1016/j.biocel.2006.02.013>

- De Virgilio C, Loewith R. Cell growth control: little eukaryotes make big contributions. *Oncogene* 2006; 25:6392-415; PMID:17041625; <http://dx.doi.org/10.1038/sj.onc.1209884>
- Efeyan A, Zoncu R, Sabatini DM. Amino acids and mTORC1: from lysosomes to disease. *Trends Mol Med* 2012; 18:524-33; PMID:22749019; <http://dx.doi.org/10.1016/j.molmed.2012.05.007>
- Jewell JL, Russell RC, Guan KL. Amino acid signalling upstream of mTOR. *Nat Rev Mol Cell Biol* 2013; 14:133-9; PMID:23361334; <http://dx.doi.org/10.1038/nrm3522>
- Dodd KM, Tee AR. Leucine and mTORC1: a complex relationship. *Am J Physiol Endocrinol Metab* 2012; 302:E1329-42; PMID:22354780; <http://dx.doi.org/10.1152/ajpendo.00525.2011>
- Kim E, Goraksha-Hicks P, Li L, Neufeld TP, Guan KL. Regulation of TORC1 by Rag GTPases in nutrient response. *Nat Cell Biol* 2008; 10:935-45; PMID:18604198; <http://dx.doi.org/10.1038/ncb1753>
- Binda M, Péli-Gulli MP, Bonfils G, Panchaud N, Urban J, Sturgill TW, Loewith R, De Virgilio C. The Vam6 GEF controls TORC1 by activating the EGO complex. *Mol Cell* 2009; 35:563-73; PMID:19748353; <http://dx.doi.org/10.1016/j.molcel.2009.06.033>
- Sancak Y, Peterson TR, Shaul YD, Lindquist RA, Thoreen CC, Bar-Peled L, Sabatini DM. The Rag GTPases bind raptor and mediate amino acid signaling to mTORC1. *Science* 2008; 320:1496-501; PMID:18497260; <http://dx.doi.org/10.1126/science.1157535>
- Kim E, Guan KL. RAG GTPases in nutrient-mediated TOR signaling pathway. *Cell Cycle* 2009; 8:1014-8; PMID:19270521; <http://dx.doi.org/10.4161/cc.8.7.8124>
- Sancak Y, Bar-Peled L, Zoncu R, Markhard AL, Nada S, Sabatini DM. Ragulator-Rag complex targets mTORC1 to the lysosomal surface and is necessary for its activation by amino acids. *Cell* 2010; 141:290-303; PMID:20381137; <http://dx.doi.org/10.1016/j.cell.2010.02.024>
- Dubouloz F, Deloche O, Wanke V, Cameroni E, De Virgilio C. The TOR and EGO protein complexes orchestrate microautophagy in yeast. *Mol Cell* 2005; 19:15-26; PMID:15989961; <http://dx.doi.org/10.1016/j.molcel.2005.05.020>
- Binda M, Bonfils G, Panchaud N, Péli-Gulli MP, De Virgilio C. An EGOcentric view of TORC1 signaling. *Cell Cycle* 2010; 9:221-2; PMID:20023374; <http://dx.doi.org/10.4161/cc.9.2.10585>
- Kogan K, Spear ED, Kaiser CA, Fass D. Structural conservation of components in the amino acid sensing branch of the TOR pathway in yeast and mammals. *J Mol Biol* 2010; 402:388-98; PMID:20655927; <http://dx.doi.org/10.1016/j.jmb.2010.07.034>
- De Virgilio C. The essence of yeast quiescence. *FEMS Microbiol Rev* 2012; 36:306-39; PMID:21658086; <http://dx.doi.org/10.1111/j.1574-6976.2011.00287.x>
- Zhang T, Péli-Gulli MP, Yang H, De Virgilio C, Ding J. Ego3 functions as a homodimer to mediate the interaction between Gtr1-Gtr2 and Ego1 in the EGO complex to activate TORC1. *Structure* 2012; 20:2151-60; PMID:23123112; <http://dx.doi.org/10.1016/j.str.2012.09.019>
- Bar-Peled L, Schweitzer LD, Zoncu R, Sabatini DM. Ragulator is a GEF for the rag GTPases that signal amino acid levels to mTORC1. *Cell* 2012; 150:1196-208; PMID:22980980; <http://dx.doi.org/10.1016/j.cell.2012.07.032>
- Dokudovskaya S, Waharte F, Schlessinger A, Pieper U, Devos DP, Cristea IM, Williams R, Salamero J, Chait BT, Salvi A, et al. A conserved coatomer-related complex containing Sec13 and Seh1 dynamically associates with the vacuole in *Saccharomyces cerevisiae*. *Mol Cell Proteomics* 2011; 10:006478; PMID:21454883; <http://dx.doi.org/10.1074/mcp.M110.006478>
- Neklesa TK, Davis RW. A genome-wide screen for regulators of TORC1 in response to amino acid starvation reveals a conserved Npr2/3 complex. *PLoS Genet* 2009; 5:e1000515; PMID:19521502; <http://dx.doi.org/10.1371/journal.pgen.1000515>
- Dokudovskaya S, Rout MP. A novel coatomer-related SEA complex dynamically associates with the vacuole in yeast and is implicated in the response to nitrogen starvation. *Autophagy* 2011; 7:1392-3; PMID:21804352; <http://dx.doi.org/10.4161/autophagy.7.11.17347>
- Wu X, Tu BP. Selective regulation of autophagy by the Iml1-Npr2-Npr3 complex in the absence of nitrogen starvation. *Mol Biol Cell* 2011; 22:4124-33; PMID:21900499; <http://dx.doi.org/10.1091/mbc.E11-06-0525>
- Panchaud N, Péli-Gulli MP, De Virgilio C. Amino acid deprivation inhibits TORC1 through a GTPase-activating protein complex for the Rag family GTPase Gtr1. *Sci Signal* 2013; 6:ra42; PMID:23716719; <http://dx.doi.org/10.1126/scisignal.2004112>
- Levine TP, Daniels RD, Wong LH, Gatta AT, Gerondopoulos A, Barr FA. Discovery of new Longin and Roadblock domains that form platforms for small GTPases in Ragulator and TRAPP-II. *Small GTPases* 2013; 4:62-9; PMID:23511850; <http://dx.doi.org/10.4161/sgtp.24262>
- Bar-Peled L, Chantranupong L, Cherniack AD, Chen WW, Ottina KA, Grabiner BC, Spear ED, Carter SL, Meyerson M, Sabatini DM. A Tumor suppressor complex with GAP activity for the Rag GTPases that signal amino acid sufficiency to mTORC1. *Science* 2013; 340:1100-6; PMID:23723238; <http://dx.doi.org/10.1126/science.1232044>
- Novick P, Field C, Schekman R. Identification of 23 complementation groups required for post-translational events in the yeast secretory pathway. *Cell* 1980; 21:205-15; PMID:6996832; [http://dx.doi.org/10.1016/0092-8674\(80\)90128-2](http://dx.doi.org/10.1016/0092-8674(80)90128-2)
- Roberg KJ, Bickel S, Rowley N, Kaiser CA. Control of amino acid permease sorting in the late secretory pathway of *Saccharomyces cerevisiae* by SEC13, LST4, LST7 and LST8. *Genetics* 1997; 147:1569-84; PMID:9409822
- Loewith R, Jacinto E, Wullschlegel S, Lorberg A, Crespo JL, Bonenfant D, Oppliger W, Jenoe P, Hall MN. Two TOR complexes, only one of which is rapamycin sensitive, have distinct roles in cell growth control. *Mol Cell* 2002; 10:457-68; PMID:12408816; [http://dx.doi.org/10.1016/S1097-2765\(02\)00636-6](http://dx.doi.org/10.1016/S1097-2765(02)00636-6)
- Hoelz A, Deblor EW, Blobel G. The structure of the nuclear pore complex. *Annu Rev Biochem* 2011; 80:613-43; PMID:21495847; <http://dx.doi.org/10.1146/annurev-biochem-060109-151030>
- Zanetti G, Pahuja KB, Studer S, Shim S, Schekman R. COPII and the regulation of protein sorting in mammals. *Nat Cell Biol* 2011; 14:20-8; PMID:22193160; <http://dx.doi.org/10.1038/ncb2390>
- Gürkan C, Stagg SM, Lapointe P, Balch WE. The COPII cage: unifying principles of vesicle coat assembly. *Nat Rev Mol Cell Biol* 2006; 7:727-38; PMID:16990852; <http://dx.doi.org/10.1038/nrm2025>

32. Lindquist RA, Ottina KA, Wheeler DB, Hsu PP, Thoreen CC, Guertin DA, Ali SM, Sengupta S, Shaul YD, Lamprocht MR, et al. Genome-scale RNAi on living-cell microarrays identifies novel regulators of *Drosophila melanogaster* TORC1-S6K pathway signaling. *Genome Res* 2011; 21:433-46; PMID:21239477; <http://dx.doi.org/10.1101/gr.111492.110>
33. Devasahayam G, Ritz D, Helliwell SB, Burke DJ, Sturgill TW. Pmr1, a Golgi Ca<sup>2+</sup>/Mn<sup>2+</sup>-ATPase, is a regulator of the target of rapamycin (TOR) signaling pathway in yeast. *Proc Natl Acad Sci U S A* 2006; 103:17840-5; PMID:17095607; <http://dx.doi.org/10.1073/pnas.0604303103>
34. Rudolph HK, Antebi A, Fink GR, Buckley CM, Dorman TE, LeVitre J, Davidow LS, Mao JI, Moir DT. The yeast secretory pathway is perturbed by mutations in PMR1, a member of a Ca<sup>2+</sup> ATPase family. *Cell* 1989; 58:133-45; PMID:2526682; [http://dx.doi.org/10.1016/0092-8674\(89\)90410-8](http://dx.doi.org/10.1016/0092-8674(89)90410-8)
35. Lempiäinen H, Uotila A, Urban J, Dohnal I, Ammerer G, Loewith R, Shore D. Sfp1 interaction with TORC1 and Msr6 reveals feedback regulation on TOR signaling. *Mol Cell* 2009; 33:704-16; PMID:19328065; <http://dx.doi.org/10.1016/j.molcel.2009.01.034>
36. Zoncu R, Bar-Peled L, Efeyan A, Wang S, Sancak Y, Sabatini DM. mTORC1 senses lysosomal amino acids through an inside-out mechanism that requires the vacuolar H<sup>+</sup>-ATPase. *Science* 2011; 334:678-83; PMID:22053050; <http://dx.doi.org/10.1126/science.1207056>
37. Bonfils G, Jaquenoud M, Bontron S, Ostrowicz C, Ungermann C, De Virgilio C. Leucyl-tRNA synthetase controls TORC1 via the EGO complex. *Mol Cell* 2012; 46:105-10; PMID:22424774; <http://dx.doi.org/10.1016/j.molcel.2012.02.009>
38. Han JM, Jeong SJ, Park MC, Kim G, Kwon NH, Kim HK, Ha SH, Ryu SH, Kim S. Leucyl-tRNA synthetase is an intracellular leucine sensor for the mTORC1-signaling pathway. *Cell* 2012; 149:410-24; PMID:22424946; <http://dx.doi.org/10.1016/j.cell.2012.02.044>
39. Durán RV, Opliger W, Robitaille AM, Heiserich L, Skendaj R, Gottlieb E, Hall MN. Glutaminolysis activates Rag-mTORC1 signaling. *Mol Cell* 2012; 47:349-58; PMID:22749528; <http://dx.doi.org/10.1016/j.molcel.2012.05.043>
40. Goranov AI, Gulati A, Dephore N, Takahara T, Maeda T, Gygi SP, Manalis S, Amon A. Changes in cell morphology are coordinated with cell growth through the TORC1 pathway. *Curr Biol* 2013; 23:1269-79; PMID:23810534; <http://dx.doi.org/10.1016/j.cub.2013.05.035>
41. Zurita-Martinez SA, Puria R, Pan X, Boeke JD, Cardenas ME. Efficient Tor signaling requires a functional class C Vps protein complex in *Saccharomyces cerevisiae*. *Genetics* 2007; 176:2139-50; PMID:17565946; <http://dx.doi.org/10.1534/genetics.107.072835>
42. Efeyan A, Zoncu R, Chang S, Gumper I, Snitkin H, Wolfson RL, Kirak O, Sabatini DD, Sabatini DM. Regulation of mTORC1 by the Rag GTPases is necessary for neonatal autophagy and survival. *Nature* 2013; 493:679-83; PMID:23263183; <http://dx.doi.org/10.1038/nature11745>
43. Li L, Kim E, Yuan H, Inoki K, Goraksha-Hicks P, Schiesher RL, Neufeld TP, Guan KL. Regulation of mTORC1 by the Rab and Arf GTPases. *J Biol Chem* 2010; 285:19705-9; PMID:20457610; <http://dx.doi.org/10.1074/jbc.C110.102483>
44. Durán RV, Hall MN. Glutaminolysis feeds mTORC1. *Cell Cycle* 2012; 11:4107-8; PMID:23095634; <http://dx.doi.org/10.4161/cc.22632>
45. Flinn RJ, Backer JM. mTORC1 signals from late endosomes: taking a TOR of the endocytic system. *Cell Cycle* 2010; 9:1869-70; PMID:20436274; <http://dx.doi.org/10.4161/cc.9.10.11679>
46. Flinn RJ, Yan Y, Goswami S, Parker PJ, Backer JM. The late endosome is essential for mTORC1 signaling. *Mol Biol Cell* 2010; 21:833-41; PMID:20053679; <http://dx.doi.org/10.1091/mbc.E09-09-0756>
47. Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, Boeke JD. Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* 1998; 14:115-32; PMID:9483801; [http://dx.doi.org/10.1002/\(SICI\)1097-0061\(19980130\)14:2<115::AID-YEA204>3.0.CO;2-2](http://dx.doi.org/10.1002/(SICI)1097-0061(19980130)14:2<115::AID-YEA204>3.0.CO;2-2)
48. Li Z, Vizeacoumar FJ, Bahr S, Li J, Warringer J, Vizeacoumar FS, Min R, Vandersluis B, Bellay J, Devit M, et al. Systematic exploration of essential yeast gene function with temperature-sensitive mutants. *Nat Biotechnol* 2011; 29:361-7; PMID:21441928; <http://dx.doi.org/10.1038/nbt.1832>
49. Winzeler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, Andre B, Bangham R, Benito R, Boeke JD, Bussey H, et al. Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* 1999; 285:901-6; PMID:10436161; <http://dx.doi.org/10.1126/science.285.5429.901>
50. Urban J, Souillard A, Huber A, Lippman S, Mukhopadhyay D, Deloche O, Wanke V, Anrather D, Ammerer G, Riezman H, et al. Sch9 is a major target of TORC1 in *Saccharomyces cerevisiae*. *Mol Cell* 2007; 26:663-74; PMID:17560372; <http://dx.doi.org/10.1016/j.molcel.2007.04.020>
51. Sikorski RS, Hieter P. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* 1989; 122:19-27; PMID:2659436
52. Caplan S, Hartnell LM, Aguilar RC, Naslavsky N, Bonifacino JS. Human Vam6p promotes lysosome clustering and fusion in vivo. *J Cell Biol* 2001; 154:109-22; PMID:11448994; <http://dx.doi.org/10.1083/jcb.200102142>
53. Felici A, Würthner JU, Parks WT, Giam LR, Reiss M, Karpova TS, McNally JG, Roberts AB. TLP, a novel modulator of TGF- $\beta$  signaling, has opposite effects on Smad2- and Smad3-dependent signaling. *EMBO J* 2003; 22:4465-77; PMID:12941698; <http://dx.doi.org/10.1093/emboj/cdg428>
54. Messler S, Kropp S, Episkopou V, Felici A, Würthner J, Lemke R, Jerabek-Willemsen M, Willecke R, Scheu S, Pfeffer K, et al. The TGF- $\beta$  signaling modulators TRAP1/TGFBRAP1 and VPS39/Vam6/TLP are essential for early embryonic development. *Immunobiology* 2011; 216:343-50; PMID:20961651; <http://dx.doi.org/10.1016/j.imbio.2010.07.006>