

SCIENTIFIC DATA

OPEN Data Descriptor: A collection of European sweet cherry phenology data for assessing climate change

Bénédicte Wenden¹, José Antonio Campoy¹, Julien Lecourt², Gregorio López Ortega³, Michael Blanke⁴, Sanja Radičević⁵, Elisabeth Schüller⁶, Andreas Spornberger⁶, Danilo Christen⁷, Hugo Magein⁸, Daniela Giovannini⁹, Carlos Campillo¹⁰, Svetoslav Malchev¹¹, José Miguel Peris¹², Mekjell Meland¹³, Rolf Stehr¹⁴, Gérard Charlot¹⁵ & José Quero-García¹

Received: 20 July 2016

Accepted: 18 October 2016

Published: 6 December 2016

Professional and scientific networks built around the production of sweet cherry (*Prunus avium* L.) led to the collection of phenology data for a wide range of cultivars grown in experimental sites characterized by highly contrasted climatic conditions. We present a dataset of flowering and maturity dates, recorded each year for one tree when available, or the average of several trees for each cultivar, over a period of 37 years (1978–2015). Such a dataset is extremely valuable for characterizing the phenological response to climate change, and the plasticity of the different cultivars' behaviour under different environmental conditions. In addition, this dataset will support the development of predictive models for sweet cherry phenology exploitable at the continental scale, and will help anticipate breeding strategies in order to maintain and improve sweet cherry production in Europe.

Design Type(s)	observation design • data integration objective
Measurement Type(s)	phenology
Technology Type(s)	visual observation method
Factor Type(s)	selectively maintained organism
Sample Characteristic(s)	Prunus avium • French Republic • Germany • Serbia • Kingdom of Spain • Switzerland • United Kingdom • Italy • Belgium • Bulgaria • Kingdom of Norway • Austria

¹UMR 1332 BFP, INRA, Univ. Bordeaux, Villenave d'Ornon 33140, France. ²NIAB EMR, East Malling, ME19 6BJ, UK. ³IMIDA, Murcia Institute of Agri-Food Research and Development, Murcia 30150, Spain. ⁴INRES—Horticultural Science, University of Bonn, 53121 Bonn, Germany. ⁵Fruit Research Institute (FRI), Čačak 32000, Republic of Serbia. ⁶University of Natural Resources and Life Sciences, Vienna, Department of Crop Sciences, Division of Viticulture and Pomology, Vienna 1180, Austria. ⁷Agroscope Changins—Wädenswil Research Station ACW, 1964 Conthey, Switzerland. ⁸Walloon Agricultural Research Center (CRA-W), Gembloux 5030, Belgium. ⁹CREA, Council for Agricultural Research and Economics, Fruit Tree Research Unit of Forlì, Forlì 47121, Italy. ¹⁰Scientific and Technological Research Center of Extremadura (CICYTEX), Guadajira (Badajoz) 06187, Spain. ¹¹Fruit Growing Institute—Plovdiv, Plovdiv 4004, Bulgaria. ¹²IRTA, Experimental Station of Lleida, Lleida 25003, Spain. ¹³Norwegian Institute of Bioeconomy Research—NIBIO Ullensvang, Lofthus N-5781, Norway. ¹⁴Fruit Research Center Jork, Jork D-21635, Germany. ¹⁵Ctifl, 751 chemin de Balandran, Bellegarde 30127, France. Correspondence and requests for materials should be addressed to B.W. (email: benedicte.wenden@bordeaux.inra.fr).

Background & Summary

The impact of climate change on plant phenology has been described in recent decades, highlighting a hastening of flowering phenology in response to increasing winter and spring temperatures^{1–7}. Long records of flowering dates have been proven extremely valuable to reconstruct past phenology and to predict phenology under future climatic scenarios. In Japan for example, flowering dates of cherry trees (*Prunus jamasakura*) have been recorded for centuries and analyses of these records have revealed that cherry flowering is currently occurring earlier than at any time in the previous seven to 12 centuries, due to the impact of warming and urbanization on phenology^{5,8,9}. This earlier flowering trend has been observed for other fruit trees^{10–14}, which are particularly vulnerable to temperature changes due to their long life span. Peach (*Prunus persica*), apricot (*Prunus armeniaca*), almond (*Prunus dulcis*), plum (*Prunus salicina* and *domestica*) and sweet cherry (*Prunus avium* L.) are amongst the most commercially important *Prunus* fruit tree species planted in temperate climate zones. In Europe, cherry tree blossom of early maturing cultivars showed an advance up to 4.7 days/°C in Germany^{10,14}, and warmer winters have dramatically affected the sweet cherry production in South-Western France, with a 30% yield in 2007. Warmer winters can as well be associated with delayed spring phenology for some species and, occasionally resulting in abnormal flowering phenology and reduced productivity^{15–18}. Models for chill availability predict an increase in the delaying effect of mild winters as temperature increase becomes more pronounced^{6,19,20}, especially in warmer locations²¹. In the context of substantial changes to environmental conditions induced by climate change, it will be essential that plant cultivars are well adapted to warmer winter and spring temperatures and to more extreme climatic events such as erratic spring frosts and summer heat waves. This is especially true for perennial fruit crops, which require more than a decade before a new cultivar is released.

Large phenological datasets are key for the development of phenological models (e.g., refs 22–25), which are valuable tools to support breeding strategies. Although recent studies have shown the value of a wide range of data^{26,27}, most analyses for fruit tree crops are based on phenological data for a very limited number of experimental sites, and rarely include more than two cultivars within a species (e.g., refs 28,29).

Sweet cherry trees are particularly interesting for phenology studies, their long orchard life providing the potential for long-term datasets. Reference cultivars have been planted and observed for decades for phenology and productivity traits in trials dedicated to new hybrids characterization. For example, at the Fruit Experimental Station (Toulence, INRA Bordeaux, France), phenological data have been recorded for ‘Burlat’ cultivar for 35 years. Consequently, large phenological datasets are available for reference cultivars in many European orchards involved in breeding programmes. Despite this, long historical datasets of fruit tree phenology are rarely analysed together or made available to the scientific community. A few analyses on sweet cherry phenology in Europe were published using the phenological observations of fruit trees by the German Weather Service (DWD)^{10,12}, from non-publicly available datasets^{30,31} or from the PEP725 data¹⁴. In addition, published studies often focus on specific location^{10,12,30–33}.

In this study, we describe a unique dataset of sweet cherry flowering and maturity records for 25 sites in Europe (Fig. 1) with highly contrasted climates. Past studies showed that phenology data spanning




Figure 1. Experimental orchards location. Flowering and maturity dates were recorded in 25 sites from 11 European countries, indicated by the red circles. Size of the circle is proportional to the number of cultivars recorded in each site.

Country	Institute	Location	Altitude (m)	Lat	Long	Nb of cultivars	Years of observation	1980	1985	1990	1995	2000	2005	2010
Germany	Bonn University	Bonn	180	50.62	6.98	1	1978 - 2015							
United Kingdom	EMR	East Malling	38	51.28	0.45	4	1978 - 2014							
Switzerland	Agroscope	Conthey	504	46.22	7.33	1	1979 - 2007							
France	INRA	Toulenne	8	44.58	-0.28	191	1981 - 2015							
Austria	BOKU	Vienna	162	48.29	16.43	8	1982 - 2014							
France	CTIFL	Balandran	52	43.76	4.46	170	1982 - 2015							
France	Sefra	Etoile	134	44.82	4.89	154	1985 - 2014							
Belgium	CRA-W	Gembloux	170	50.55	4.66	1	1989 - 2014							
Italy	CREA-FRF	Forli	34	44.22	12.02	3	1991 - 2015							
Serbia	FRI	Čačak	260	43.88	20.35	16	1992 - 2014							
Germany	OVA	Jork	-1	53.51	9.74	79	1993 - 2011							
France	La Tapy	Carpentras	84	44.09	5.05	57	1994 - 2014							
Norway	NIBIO	Ullensvang	27	60.32	6.66	10	1995 - 1998							
Spain	IRTA	Corbins	233	41.70	0.69	16	1996 - 2006							
France	Serfel	St Gilles	38	43.69	4.43	37	2004 - 2013							
France	Verexal	Obernai	179	48.47	7.51	21	2006 - 2014							
Spain	IMIDA	Cieza	244	38.29	-1.5	3	2006 - 2010							
Bulgaria	Fruitgrowing Institute	Plovdiv	130	42.11	24.7	8	2006 - 2011							
France	Sefra	Bozas	536	45.06	4.65	24	2007 - 2011							
France	La Morinière	St Epain	95	47.16	0.60	25	2007 - 2014							
France	Sefra	St Laurent d'Agnay	388	45.63	4.68	37	2007 - 2014							
France	Cefel	Montauban	90	44.04	1.31	29	2004 - 2012							
France	Centrex	Torreilles	10	42.76	2.98	20	2008 - 2014							
Spain	CAEM	El Torno	463	40.14	-5.95	2	1994, 2010 - 2013							
Spain	IMIDA	Jumilla	395	38.04	-1.98	36	2010 - 2015							

Beginning of flowering (■) – Full flowering (■) – End of flowering (■) – Maturity (■)

Figure 2. Description of data source, time-span, number of observed cultivars, and location within Europe.

20 or 30 years were valuable for climate change related analyses^{7,34}. Thus the dataset presented here, with an overall 37 year-period (1978–2015, Fig. 2) will be valuable for phenology and climate change studies. This dataset covers a wide range of European latitudes and longitudes and is unique in its collection of cultivars (between 1 and 191 cultivars per site), each cultivar being represented by clones of the same original tree in each country, which supports robust analyses of plasticity and response to climatic conditions.

Since data were collected from various experimental stations, the dataset is not homogeneous regarding the number of cultivars (Figs 1 and 2, Table 1) or the record length (Fig. 2). Past research have shown the value of using heterogeneous records combined from different sites and cultivars for the evaluation of climate change response and phenology modelling approaches (e.g., refs 21,35–38). In particular, phenology models have been successfully tested and optimized using data sourced from different sites^{27,38}. Therefore, we want to highlight the value of this dataset, combining data from various geographical sites and contrasted cultivars, for potential multi-environment analyses yet to be implemented. 13 out of 25 sites meet the criteria of more than 15 recorded years that was shown to be useful for climate change analysis⁷ (Fig. 3; Table 1). Single site analyses can thus be applied to track phenological climate shifts in a given environment. Subsequently, phenology models can be further evaluated by pooling data across geographically and climatically varied sites, leading to a better

Site	Number of observed cultivars	Number of data points				
		Beginning of flowering	Full flowering	End of flowering	Maturity	Total
Balandran	170	2530	2714	2192	2555	9991
Bonn	1	30	6			36
Bozas	24	72	100	87	32	291
Čačak	16	338	336	286	49	1009
Carpentras	57		552	8	282	842
Cieza	3	15	15	15		45
Conthey	1		29			29
Corbins	16	157	156	147	10	470
East Malling	4	167	167	167		501
El Torno	2	10	10	9	10	39
Etoile	154	1675	1686	1409	1229	5999
Forli	3	38	38	31	7	114
Gembloux	1	27	27	27	22	103
Jork	79	685	685	682		2052
Jumilla	36	234	234	205	234	907
Montauban	29	9	79			88
Obernai	21		154		95	249
Plovdiv	8	45	45	23	14	127
St Epain	25		221		216	437
St Gilles	37	325	401	2	255	983
St Laurent d'Agny	37	67	236	53	9	365
Torreilles	20	128	128	85	121	462
Toulonne	191	4218	3518	3603	3476	14815
Ullensvang	10	231	29	23		283
Vienna	8	153	147	146	42	488

Table 1. Overall description of the data records.




Figure 3. Information on the number of observed cultivars and the overall record length for each site. Some site names were omitted the ease the comprehension. The numbers between brackets indicate the number of cultivars with more than 15 years of observation. The dash line indicate the 15 years limit for the record length, criteria mentioned in (Fu *et al.*⁷) as sufficient to perform climate change response analysis.








BBCH code	Description	
60	First flowers open	
61	Beginning of flowering: about 10% of flowers open	
65	Full flowering: at least 50% of flowers open, first petals falling	
67	Flowers fading: majority of petals fallen	
69	End of flowering: all petals fallen	
87	Fruit ripe for picking	
89	Fruit ripe for consumption: fruit have typical taste and firmness	

Figure 4. BBCH descriptors (Meier⁴⁰) corresponding to the flowering and maturity stages observed in the dataset. Pictures @INRA.

knowledge of climate-phenology relations. Extreme European climate, e.g., South of Spain, can be used to investigate climate analogues for projected climatic scenarios³⁹. In addition, for eight sites, two or more cultivars were observed for at least 15 years (Fig. 3). Between cultivar differences can be assessed at a single site and across sites with common cultivars. These analyses can reveal whether some cultivars are more susceptible to evolution in climatic conditions. The dataset offers the possibility to study different flowering phase data, namely beginning, full and end of flowering, together with maturity dates for some cultivars (Table 1), and to perform sequential phenology phase assessments.

Overall, despite the fact that they are heterogeneous, data from the different sites can be combined in integrated analysis to study their responses to environment. Such dataset is extremely valuable for characterizing the phenology response to climate change and the plasticity of the different cultivar behaviour under various environmental conditions. In addition, these records can support the development of predictive models exploitable at the continental scale that can be used by growers and breeders, and to anticipate breeding strategies in order to maintain and improve sweet cherry production in Europe.

Methods

Sweet cherry phenological data were collated from French and European networks. Established in 1952, CTIFL is a non-profit organization involved in the French fruit and vegetable industry. It developed a private database dedicated to information on cultivars planted in experimental orchards. Flowering and maturity dates for up to 191 reference cultivars grown in French experimental stations were extracted from the database. At the European scale, in the context of the COST Action 1104 (2012–2016; <https://www.bordeaux.inra.fr/cherry/>), which aimed at creating a dynamic network of scientists and other professionals conducting research to improve sweet or sour cherry production in Europe, we established a working group (WG) for phenology studies. Flowering and maturity dates together with the protocol details for the observations were collected. Although a standardisation of the recorded stages is on-going within the group, past observation standards for the different flowering stages are not homogeneous and

Country	Location	Beginning of flowering		Full flowering		End of flowering		Maturity	
Belgium	Gembloux	10% open flowers	BBCH 61	50% open flowers	BBCH 65	petal fall	BBCH 67	ripe fruits, harvest	BBCH 89
Bulgaria	Plovdiv	5% open flowers	BBCH 61	25–75% open flowers	BBCH 65	Petal falls for 75% of flowers	BBCH 67		
France	All experimental sites	10% open flowers	BBCH 61	80% open flowers	BBCH 65	10–20% petal fall	BBCH 67	30% ripe fruits (color, taste)	BBCH 89
Germany	Bonn	5–10% open flowers	BBCH 61	50–75% open flowers to beginning of petal fall	BBCH 65		BBCH 67		
Germany	Jork	1% open flowers	BBCH 60	80% open flowers	BBCH 65	Petal falls for 90% of flowers	BBCH 67		
Italy	Forli	5–10% open flowers	BBCH 61	80% open flowers	BBCH 65	No more pollen is shed by the open flowers	BBCH 67		
Norway	Ullensvang	1% open flowers	BBCH 60		BBCH 65		BBCH 67		
Serbia	Čačak	10–20% open flowers	BBCH 61	90–100% open flowers	BBCH 65	90% petal fall	BBCH 67		
Spain	Cieza	5–10% open flowers	BBCH 61	50–75% open flowers to beginning of petal fall	BBCH 65	75% open flower to the most petals have fallen	BBCH 67		
Spain	Jumilla	5–10% open flowers	BBCH 61	50–75% open flowers to beginning of petal fall	BBCH 65	75% open flower to the most petals have fallen	BBCH 67		
Spain	Lleida	1% open flowers	BBCH 60	80% open flowers	BBCH 65	100% open flowers	BBCH 67		
Switzerland	Changins			80% open flowers	BBCH 65				
United Kingdom	East Malling	1% open flowers	BBCH 60	50% open flowers	BBCH 65	90% petal fall	BBCH 67		

Table 2. Stages recorded in the experimental sites, when available, associated with the corresponding BBCH stage.

Header	Description
Country	
Institute	Research or experimental institute attached to the experimental station
Station	Name of station
Latitude	Latitude in decimal degrees
Longitude	Longitude in decimal degrees
Altitude	Altitude in meters
Plantation	Year of plantation of the tree
Year	Year of observation
Cultivar	Registered name of the cultivar
Clone	Number of clone
Rootstock	Name of the rootstock
Beginning of flowering (date)	Date observed for the BBCH stage corresponding to beginning of flowering
Full flowering (date)	Date observed for the BBCH stage corresponding to full flowering
End of flowering (date)	Date observed for the BBCH stage corresponding to end of flowering
Beginning of maturity (date)	Date observed for the BBCH stage corresponding to maturity
Beginning of flowering	Number of days in year [1–365/366] for the BBCH stage corresponding to beginning of flowering
Full flowering	Number of days in year [1–365/366] for the BBCH stage corresponding to full flowering
End of flowering	Number of days in year [1–365/366] for the BBCH stage corresponding to end of flowering
Beginning of maturity	Number of days in year [1–365/366] for the BBCH stage corresponding to maturity
Flowering duration	Number of days between beginning and end of flowering

Table 3. Metadata.

Compared sites		Distance (km)	Cultivar	Number of common years	Spearman correlation	Average date for beginning of flowering
Balandran	Etoile	123	Burlat	22	0.93	89
Balandran	Etoile	123	Ferdouce	15	0.67	82
Balandran	Etoile	123	Summit	26	0.84	93
Balandran	Etoile	123	Sweetheart Sumtare	21	0.87	85
Gembloux	Bonn	164	Burlat	20	0.85	104
Balandran	Toulence	390	Arcina Fercer	18	0.95	87
Balandran	Toulence	390	Burlat	24	0.88	87
Balandran	Toulence	390	Coralise Gardel	15	0.57	81
Balandran	Toulence	390	Duroni 3	15	0.77	94
Balandran	Toulence	390	Earlise Rivedel	19	0.76	81
Balandran	Toulence	390	Ferdouce	17	0.88	80
Balandran	Toulence	390	Ferobri	18	0.84	85
Balandran	Toulence	390	Fertard	16	0.68	92
Balandran	Toulence	390	Folfer	16	0.93	81
Balandran	Toulence	390	Hedelfingen	21	0.88	90
Balandran	Toulence	390	Kordia	17	0.72	88
Balandran	Toulence	390	Lapins	20	0.78	84
Balandran	Toulence	390	Napoleon	18	0.90	87
Balandran	Toulence	390	Rainier	27	0.88	85
Balandran	Toulence	390	Regina	17	0.79	92
Balandran	Toulence	390	Stark Hardy Giant	24	0.90	85
Balandran	Toulence	390	Summit	29	0.84	93
Balandran	Toulence	390	Sweetheart Sumtare	19	0.87	85
Balandran	Toulence	390	Van	17	0.90	83
Toulence	Etoile	411	Burlat	28	0.93	89
Toulence	Etoile	411	Duroni 3	16	0.87	96
Toulence	Etoile	411	Summit	26	0.91	93
Toulence	Etoile	411	Sweetheart Sumtare	15	0.86	85

Table 4. Consistency of data between sites. Spearman correlation and average date for beginning of flowering (BBCH 61) were calculated for cultivars grown in the different sites.

are described in Fig. 4 and Table 2. They correspond to a percentage of open flowers or fallen petals. Since the development of the BBCH scale (*Biologische Bundesanstalt, Bundessortenamt und Chemische Industrie*), this standard has been applied as a coding system for the characterization of the entire developmental cycle of annual and perennials plants^{40,41}. Here, where possible, we associated the recorded stages, as defined in each experimental location for observations, with the corresponding BBCH stage (Table 2). In every location, one or two observers were in charge of recording the phenology dates. At the end of the season, records were added to the dataset. We calculated the length of the flowering season, which is the number of days between beginning and end of flowering, where these dates were available.

Data Records

Flowering and maturity dates from all sites can be found in the dataset file stored in the Dryad Digital Repository (Data Citation 1). The spread sheet consists of a table with the description of all phenological data (Table 3). The experimental sites are described by name, latitude, longitude and altitude. Each row corresponds to the dates (beginning of flowering, full flowering, end of flowering, beginning of maturity) documented each year for one tree when available, or the average of several trees for each cultivar. For registration reasons, one cultivar can be registered and observed under different clone numbers, ranging from 1 to 7 clone accession numbers, so the clone number was indicated when available. The cultivar name was always indicated and when available the rootstock information was provided. Dates recorded were also provided as day of year (starting with 1 for January 1st) and the duration of flowering was calculated (days).

Technical Validation

All data were checked for consistency and anomalous values were corrected or removed (Supplementary Table 1). Some old cultivars, that can be found in many countries, have names that differ slightly between




Figure 5. Comparison of flowering dates in days (from 1st of January) for 6 cultivars from our dataset and, unless otherwise stated, for BBCH60 stage for the ‘Early cultivar’ and ‘Late cultivar’ collected from PEP725 (<http://www.pep725.eu>) for stations geographically close to our experimental sites. For the Bonn site, PEP725 data were obtained from the station 1360, located in Zülpich (Germany; longitude: 6.67; latitude: 50.68; distance from Bonn: 23 km). Gembloux data were compared to flowering dates from the station 1498, located in Übach-Palenberg (Germany, longitude: 6.12; latitude: 50.92; distance: 110 km). For Jork, PEP725 data were obtained from the station 346, located in Hamburg-Altengamme (Germany; longitude: 10.28; latitude: 53.43; distance: 37 km). Conthey data were compared to full flowering (BBCH stage 65) dates from station 3260 (Müllheim, Germany; longitude: 7.63; latitude: 47.82, distance: 178 km). Spearman’s correlation coefficient was calculated for each site between our data and corresponding PEP725 data.

the sites so we arbitrarily chose one common name: ‘Badacsony’ was selected as the common name for ‘Badacsony’, ‘Badacsoner’ and ‘Badacsonyi’; ‘Francesca’ and ‘Francesca’ were regrouped as ‘Francesca’. In total, 51 records were corrected (Supplementary Table 1).

For records of more than 15 years, we checked the consistency of collected data between sites. This cross-checking showed data for a given cultivar were highly correlated, even for sites as far as 400 km from each other (Table 4), confirming that the collected data are consistent. In addition, when possible, we chose to compare our data to similar phenological observational records from the European phenology database PEP725 (<http://www.pep725.eu>). Data for ‘early cultivar’ and ‘late cultivar’ were retrieved from the database and correlated with close-by sites when at least 15 common years of data were available. We identified PEP stations located within a range of 200 km for Bonn, Conthey, Gembloux and

Jork. Strong correlations and minimal differences were found between our data and the flowering dates recorded and validated in PEP725 (Fig. 5). Flowering dates records for six cultivars met the criteria of 15 year-records and the Spearman correlations were all higher than 0.76, regardless of the flowering precocity of the cultivar or the site (Fig. 5).

Usage Notes

The sweet cherry phenology dataset was collected with the objective to support climate change and phenological analyses for varied European environments. These data can be associated with other *Prunus avium* flowering data provided by the European phenology database PEP725 (<http://www.pep725.eu>) to perform a wide evaluation of phenology in early and late cultivars.

References

- Chmielewski, F.-M. & Rötzer, T. Response of tree phenology to climate change across Europe. *Agric. For. Meteorol.* **108**, 101–112 (2001).
- Cook, B. I. *et al.* Sensitivity of Spring Phenology to Warming Across Temporal and Spatial Climate Gradients in Two Independent Databases. *Ecosystems* **15**, 1283–1294 (2012).
- Gordo, O. & Sanz, J. J. Phenology and climate change: a long-term study in a Mediterranean locality. *Oecologia* **146**, 484–495 (2005).
- Menzel, A. *et al.* European phenological response to climate change matches the warming pattern. *Glob. Chang. Biol.* **12**, 1969–1976 (2006).
- Primack, R. B., Higuchi, H. & Miller-Rushing, A. J. The impact of climate change on cherry trees and other species in Japan. *Biol. Conserv.* **142**, 1943–1949 (2009).
- Vitasse, Y. *et al.* Assessing the effects of climate change on the phenology of European temperate trees. *Agric. For. Meteorol.* **151**, 969–980 (2011).
- Fu, Y. H. *et al.* Declining global warming effects on the phenology of spring leaf unfolding. *Nature* **526**, 104–107 (2015).
- Aono, Y. & Omoto, Y. Variation in the March mean temperature deduced from cherry blossom in Kyoto since the 14th century. *J. Agr. Met.* **48**, 635–638 (1993).
- Aono, Y. & Kazui, K. Phenological data series of cherry tree flowering in Kyoto, Japan, and its application to reconstruction of springtime temperatures since the 9th century. *Int. J. Biometeorol.* **28**, 905–914 (2008).
- Chmielewski, F.-M., Müller, A. & Bruns, E. Climate changes and trends in phenology of fruit trees and field crops in Germany, 1961–2000. *Agric. For. Meteorol.* **121**, 69–78 (2004).
- Dose, V. & Menzel, A. Bayesian analysis of climate change impacts in phenology. *Glob. Chang. Biol.* **10**, 259–272 (2004).
- Estrella, N., Sparks, T. H. & Menzel, A. Trends and temperature response in the phenology of crops in Germany. *Glob. Chang. Biol.* **13**, 1737–1747 (2007).
- Legave, J.-M., Guédon, Y., Malagi, G., El Yaacoubi, A. & Bonhomme, M. Differentiated Responses of Apple Tree Floral Phenology to Global Warming in Contrasting Climatic Regions. *Front. Plant Sci.* **6**, 1054 (2015).
- Jochner, S., Sparks, T. H., Laube, J. & Menzel, A. Can we detect a nonlinear response to temperature in European plant phenology? *Int. J. Biometeorol.* **60**, 1551–1561 (2016).
- Doi, H., Gordo, O. & Katano, I. Heterogeneous intra-annual climatic changes drive different phenological responses at two trophic levels. *Clim. Res.* **36**, 181–190 (2008).
- Yu, H., Luedeling, E. & Xu, J. Winter and spring warming result in delayed spring phenology on the Tibetan Plateau. *Proc. Natl Acad. Sci.* **107**, 22151–22156 (2010).
- Sherry, R. A. *et al.* Divergence of reproductive phenology under climate warming. *Proc. Natl. Acad. Sci. USA* **104**, 198–202 (2007).
- Cook, B. I., Wolkovich, E. M. & Parmesan, C. Divergent responses to spring and winter warming drive community level flowering trends. *Proc. Natl. Acad. Sci. USA* **109**, 9000–9005 (2012).
- Luedeling, E. Climate change impacts on winter chill for temperate fruit and nut production: A review. *Sci. Hortic.* **144**, 218–229 (2012).
- Pope, K. S. *et al.* Detecting nonlinear response of spring phenology to climate change by Bayesian analysis. *Glob. Chang. Biol.* **19**, 1518–1525 (2013).
- Guo, L., Dai, J., Wang, M., Xu, J. & Luedeling, E. Responses of spring phenology in temperate zone trees to climate warming: A case study of apricot flowering in China. *Agric. For. Meteorol.* **201**, 1–7 (2015).
- Chung, U., Mack, L., Yun, J. I. & Kim, S. H. Predicting the timing of cherry blossoms in Washington, DC and Mid-Atlantic States in response to climate change. *PLoS ONE* **6**, e27439 (2011).
- Ruml, M., Milatovic, D., Vulic, T. & Vukovic, A. Predicting apricot phenology using meteorological data. *Int. J. Biometeorol.* **55**, 723–732 (2011).
- Legave, J. M. *et al.* A comprehensive overview of the spatial and temporal variability of apple bud dormancy release and blooming phenology in Western Europe. *Int. J. Biometeorol.* **57**, 317–331 (2013).
- Kalvāns, A., Bitāne, M. & Kalvāne, G. Forecasting plant phenology: evaluating the phenological models for *Betula pendula* and *Padus racemosa* spring phases, Latvia. *Int. J. Biometeorol.* **59**, 165–179 (2015).
- Legave, J. M. *et al.* A comprehensive overview of the spatial and temporal variability of apple bud dormancy release and blooming phenology in Western Europe. *Int. J. Biometeorol.* **57**, 333–335 (2013).
- Andreini, L. *et al.* Understanding dormancy release in apricot flower buds (*Prunus armeniaca* L.) using several process-based phenological models. *Agric. For. Meteorol.* **184**, 210–219 (2014).
- Luedeling, E., Zhang, M., McGranahan, G. & Leslie, C. Validation of winter chill models using historic records of walnut phenology. *Agric. For. Meteorol.* **149**, 1854–1864 (2009).
- Matzneller, P., Blümel, K. & Chmielewski, F.-M. Models for the beginning of sour cherry blossom. *Int. J. Biometeorol.* **58**, 703–715 (2014).
- de Vries, P. Phenological stages in sweet cherry with regard to pre-selection. *Euphytica* **16**, 177–182 (1967).
- Luedeling, E., Kunz, A. & Blanke, M. M. Identification of chilling and heat requirements of cherry trees—a statistical approach. *Int. J. Biometeorol.* **57**, 679–689 (2013).
- Gratacós, E. & Cortés, A. Phenology and production of sweet cherry cultivars in a low chilling area of central Chile. *Acta. Hortic.* **795**, 239–244 (2008).
- Albuquerque, N., García-Montiel, F., Carrillo, A. & Burgos, L. Chilling and heat requirements of sweet cherry cultivars and the relationship between altitude and the probability of satisfying the chill requirements. *Environ. Exp. Bot.* **64**, 162–170 (2008).
- Sparks, T. H. & Menzel, A. Observed changes in seasons: An overview. *Int. J. Climatol.* **22**, 1715–1725 (2002).
- Webb, L. B. *et al.* Earlier wine-grape ripening driven by climatic warming and drying and management practices. *Nat. Clim. Chang* **2**, 259–264 (2012).

36. Fu, Y. H., Campioli, M., Van Oijen, M., Deckmyn, G. & Janssens, I. a. Bayesian comparison of six different temperature-based budburst models for four temperate tree species. *Ecol. Modell* **230**, 92–100 (2012).
37. Pope, K. S., Dose, V., Da Silva, D., Brown, P. H. & DeJong, T. M. Nut crop yield records show that budbreak-based chilling requirements may not reflect yield decline chill thresholds. *Int. J. Biometeorol.* **59**, 707–715 (2014).
38. Darbyshire, R., Pope, K. & Goodwin, I. An evaluation of the chill overlap model to predict flowering time in apple tree. *Sci. Hortic* **198**, 142–149 (2016).
39. Webb, L. B., Watterson, I., Bhend, J., Whetton, P. H. & Barlow, E. W. R. Global climate analogues for winegrowing regions in future periods: Projections of temperature and precipitation. *Aust. J. Grape Wine Res.* **19**, 331–341 (2013).
40. Meier, U. *Growth stages of mono-and dicotyledonous plants BBCH Monograph. Federal Biological Research Centre for Agriculture and Forestry* (2001).
41. Fadón, E., Herrero, M. & Rodrigo, J. Flower development in sweet cherry framed in the BBCH scale. *Sci. Hortic.* **192**, 141–147 (2015).

Data Citation

1. Wenden, B. *et al.* *Dryad Digital Repository* <http://dx.doi.org/10.5061/dryad.1d28m> (2016).

Acknowledgements

The authors would like to thank the EU COST FA1104 action for enabling interaction and data sharing between sweet cherry collaborators. We warmly thank Dr Rebecca Darbyshire for her remarks and suggestions on the manuscript. The authors are grateful to the organisations that contributed to the French data (Koala database, administrated by Ctifl): Ctifl (Balandran), Sefra (Bozas, Etoile, Saint Laurent d'Agny), Serfel (Saint Gilles), Centrex (Torreilles), Cefel (Montauban), Verexal (Obernai), La Morinière (Saint Epain), La Tapy (Carpentras), Fruit Experimental Unit (INRA Bordeaux, Toulence). Data from the Republic of Serbia were obtained within the scope of the Research Project TR-31064, supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia. We thank INRA, Aquitaine Region and CEP Innovation (AQUIPRU project 2014-1R20102-2971) for financing postdoctoral fellowship to JAC. The authors would like to thank NIAB EMR technical team who collected the data for decades, and more recently Mike Davies, Clare Hobson, Carlos Angulo Costa and Helen Longbottom.

Author Contributions

B.W. collected, checked the data, performed the technical validation and wrote the paper. J.A.C. and J.L. contributed to the technical validation and corrected the paper. J.L., G.L.O., M.B., S.R., E.S., A.S., D.C., H.M., D.G., C.C., S.M., J.M.P., M.M., R.S. and G.C. provided data. J.Q.G. organized the phenology network and provided data.

Additional Information

Supplementary Information accompanies this paper at <http://www.nature.com/sdata>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Wenden, B. *et al.* A collection of European sweet cherry phenology data for assessing climate change. *Sci. Data* **3**:160108 doi: 10.1038/sdata.2016.108 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0>

Metadata associated with this Data Descriptor is available at <http://www.nature.com/sdata/> and is released under the CC0 waiver to maximize reuse.

© The Author(s) 2016