


A LuxR Homolog in a Cottonwood Tree Endophyte That Activates Gene Expression in Response to a Plant Signal or Specific Peptides

Amy L. Schaefer,^a Yasuhiro Oda,^a Bruna Goncalves Coutinho,^a Dale A. Pelletier,^b Justin Weiburg,^a Vittorio Venturi,^c E. Peter Greenberg,^a  Caroline S. Harwood^a

University of Washington, Seattle, Washington, USA^a; Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA^b; International Centre for Genetic Engineering and Biotechnology, Trieste, Italy^c

A.L.S., Y.O., and B.G.C. contributed equally to this work.

ABSTRACT Homologs of the LuxR acyl-homoserine lactone (AHL) quorum-sensing signal receptor are prevalent in *Proteobacteria* isolated from roots of the Eastern cottonwood tree, *Populus deltoides*. Many of these isolates possess an orphan LuxR homolog, closely related to OryR from the rice pathogen *Xanthomonas oryzae*. OryR does not respond to AHL signals but, instead, responds to an unknown plant compound. We discovered an OryR homolog, PipR, in the cottonwood endophyte *Pseudomonas* sp. strain GM79. The genes adjacent to *pipR* encode a predicted ATP-binding cassette (ABC) peptide transporter and peptidases. We purified the putative peptidases, PipA and AapA, and confirmed their predicted activities. A transcriptional *pipA-gfp* reporter was responsive to PipR in the presence of plant leaf macerates, but it was not influenced by AHLs, similar to findings with OryR. We found that PipR also responded to protein hydrolysates to activate *pipA-gfp* expression. Among many peptides tested, the tripeptide Ser-His-Ser showed inducer activity but at relatively high concentrations. An ABC peptide transporter mutant failed to respond to leaf macerates, peptone, or Ser-His-Ser, while peptidase mutants expressed higher-than-wild-type levels of *pipA-gfp* in response to any of these signals. Our studies are consistent with a model where active transport of a peptidic signal is required for the signal to interact with PipR, which then activates peptidase gene expression. The identification of a peptide ligand for PipR sets the stage to identify plant-derived signals for the OryR family of orphan LuxR proteins.

IMPORTANCE We describe the transcription factor PipR from a *Pseudomonas* strain isolated as a cottonwood tree endophyte. PipR is a member of the LuxR family of transcriptional factors. LuxR family members are generally thought of as quorum-sensing signal receptors, but PipR is one of an emerging subfamily of LuxR family members that respond to compounds produced by plants. We found that PipR responds to a peptidic compound, and we present a model for Pip system signal transduction. A better understanding of plant-responsive LuxR homologs and the compounds to which they respond is of general importance, as they occur in dozens of bacterial species that are associated with economically important plants and, as we report here, they also occur in members of certain root endophyte communities.

Received 21 June 2016 Accepted 27 June 2016 Published 2 August 2016

Citation Schaefer AL, Oda Y, Coutinho BG, Pelletier D, Weiburg J, Venturi V, Greenberg EP, Harwood CS. 2016. A LuxR homolog in a cottonwood tree endophyte that activates gene expression in response to a plant signal or specific peptides. *mBio* 7(4):e01101-16. doi:10.1128/mBio.01101-16.

Editor Edward G. Ruby, University of Hawaii—Manoa

Copyright © 2016 Schaefer et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Caroline S. Harwood, csh5@uw.edu.

This article is a direct contribution from a Fellow of the American Academy of Microbiology. External solicited reviewers: Steven Lindow, University of California, Berkeley; Juan Gonzalez, University of Texas at Dallas.

The fast-growing Eastern cottonwood tree, *Populus deltoides*, possesses a distinct microbiota of endophytic (dominated by *Gamma*- and *Alphaproteobacteria*) and rhizosphere-associated (dominated by *Acidobacteria* and *Alphaproteobacteria*) bacteria (1). We have shown that acyl-homoserine lactone (AHL)-type quorum-sensing (QS) genes are prevalent in the genomes of *Proteobacteria* isolated from *Populus* roots (2). Quorum sensing is a cell-to-cell signaling system that allows bacteria to control the expression of genes in a cell density-dependent manner. The AHL QS regulatory circuits include both signal synthases (encoded by *luxI*-type genes) and signal receptors (encoded by *luxR*-type genes) (3, 4). Often the AHL synthase and its coevolved receptor genes are linked on the chromosome, but some *luxR* homologs are not linked to a *luxI* gene. Such *luxR* genes are termed orphans or

solos (2, 5) and are abundant in genomes of bacteria isolated from *P. deltoides* (2). Some of the better-studied orphan LuxRs respond to AHLs made by another paired LuxI-LuxR system present in the same cell (6) or by AHLs exogenously provided from neighboring bacteria (7, 8), while the recently described orphan LuxRs from *Photorhabdus* species have been shown to detect endogenous, non-AHL metabolites (9, 10).

Interestingly, many of the *Populus* root isolates encode members of a particular subfamily of LuxR orphan receptors (2) that are responsive to plant-derived chemical elicitors rather than AHLs (reviewed in references 5, 11, and 12). Apparently these LuxR homologs sense their plant host, rather than a QS signal (12, 13). Compared with the AHL-responsive LuxRs, little is known about how these plant-responsive homologs function, and the




FIG 1 *Pseudomonas* sp. GM79 genomic region surrounding the *oryR* homolog *pipR* (red, *PMI36_04623*). The region includes genes predicted to encode peptidases (yellow, *PMI36_04622* and *PMI36_04624*) and an ABC-type peptide transporter (blue, *PMI36_04617-04621*). There are five peptide transporter genes coding for one periplasmic binding protein, two nucleotide-binding domain (NBD) proteins, and two transmembrane domain (TMD) proteins. The positive numbers below the genes indicate the number of bases in the intergenic region separating the genes; a negative number indicates there is overlap of the two genes.

plant-associated compounds that serve as their ligands have yet to be identified. The best-studied examples are from plant-pathogenic members of the genus *Xanthomonas* (14–17), but similar systems are found in other plant-associated bacteria (11–13), including plant symbionts (18) and biocontrol agents (12). LuxR homologs from several of these bacteria have been shown to activate the transcription of adjacent genes annotated as encoding proline iminopeptidases (*pip* genes). The *pip* genes have been implicated as virulence factors in some bacteria (14, 15). To distinguish the plant-responsive LuxR homologs from the AHL-responsive LuxR homologs, we refer to this subfamily of regulators as OryR regulators, because *X. oryzae* OryR was one of the earliest described plant-responsive LuxR homologs (16).

Here, we describe an OryR regulator that we name PipR, encoded in the *Populus* root endophyte *Pseudomonas* sp. strain GM79 (2), a member of the *Pseudomonas fluorescens* subfamily (19, 20). The genes flanking *pipR* are predicted to encode peptidases and an ATP-binding cassette (ABC) peptide transporter. We show that, similar to *X. oryzae* OryR, PipR activates the transcription of a flanking peptidase gene in response to plant leaf macerates but not in response to AHLs. PipR also responded to protein hydrolysates and a specific peptide (Ser-His-Ser) to activate the expression of the flanking peptidase gene. We show that the PipR response requires the ABC transporter and is modulated by the adjacent peptidase enzymes, perhaps forming a feedback loop. We propose that because we have identified a specific signal molecule, the *Pseudomonas* sp. GM79 PipR system can serve as a model for molecular analyses of the plant-responsive OryR family of signal-

ing systems, which are found in a large number of diverse, plant-associated bacteria.

RESULTS

GM79 possesses an *oryR* homolog, which is flanked by peptidase genes. The genome of *Pseudomonas* sp. GM79 (21) contains two orphan *luxR* homologs (2), *PMI36_01833* and *PMI36_04623*. The polypeptide encoded by *PMI36_01833* is a homolog of the PpoR orphan from *Pseudomonas putida*, which responds to the AHL signal, 3-oxo-hexanoyl-L-homoserine lactone (3-oxo-C6-HSL) (2, 22). The other *luxR* homolog, *PMI36_04623*, is predicted to be a member of the OryR subfamily of plant-responsive LuxR homologs, based on its amino acid sequence and the context of neighboring *pip* genes (2, 12). Like other OryR-type polypeptides, *PMI36_04623* has a tryptophan in place of a tyrosine that is conserved in the AHL-responsive LuxR homologs, but unlike the *Xanthomonas* and *Ensifer* OryR homologs, a conserved tryptophan residue remains unchanged (see Fig. S1 in the supplemental material) (reviewed in reference 12).

All known *oryR* homologs are flanked by at least one gene annotated as a proline iminopeptidase gene (*pip*) (15). In GM79, the *oryR* homolog is flanked by two genes predicted to encode proline iminopeptidases (<http://img.jgi.doe.gov>) (21) (Fig. 1), in a genomic arrangement similar to that of the *oryR* homolog (*nesR*) in *Ensifer meliloti* (18). To confirm whether the genes flanking the GM79 *oryR* homolog actually code for peptidases, both enzymes were purified as hexahistidine-tagged fusion proteins and assayed for their ability to cleave N-terminal amino acid residues from a

TABLE 1 Substrate specificities of purified His₆-PipA and His₆-AapA enzymes

Substrate	Mean activity ± SD ^a	
	His ₆ -PipA	His ₆ -AapA
L-Proline-β-naphthylamide	100.0 ± 13.4	9.7 ± 0.8
L-Alanine-β-naphthylamide	79.4 ± 12.5	331.1 ± 39.9
L-Hydroxy-proline-β-naphthylamide	30.4 ± 1.3	23.5 ± 2.6
L-Serine-β-naphthylamide	21.5 ± 2.6	12.0 ± 0.5
L-Leucine-β-naphthylamide	7.7 ± 1.8	2.1 ± 1.0
L-Histidine-β-naphthylamide	ND	2.9 ± 0.8
L-Glutamic acid-β-naphthylamide	ND	ND
L-Proline- <i>p</i> -nitroanilide	0.72 ± 0.01	0.006 ± 0.001
L-Methionine- <i>p</i> -nitroanilide	0.17 ± 0.02	0.160 ± 0.021
L-Lysine- <i>p</i> -nitroanilide	ND	ND

^a Enzyme (PipA [*PMI36_04624*] and AapA [*PMI36_04622*]) purification and assay conditions are described in Materials and Methods; the results are the mean activities from 4 to 8 assays. Naphthylamide substrate results were measured as relative fluorescence units (RFU) per min per mg of protein and normalized to the activity exhibited by His₆-PipA with L-proline-β-naphthylamide as the substrate. Nitroanilide substrate results are reported as millimoles cleaved per min per mg of protein. ND, not detected (not above the background of the no-added-enzyme control).

variety of fluorescent (β -naphthylamide) and chromogenic (*p*-nitroanilide) substrates (Table 1). The PMI36_04622 enzyme was most active in cleaving an N-terminal alanine, while the PMI36_04624 enzyme exhibited good activity in cleaving N-terminal proline and, to a slightly smaller degree, alanine. Both enzymes had moderate activity with hydroxy-proline-, serine-, and methionine-linked substrates, while little-to-no peptidase activity was observed with histidine-, glutamic acid-, and lysine-linked substrates (Table 1). Based on the substrate specificities exhibited by the purified GM79 enzymes, we propose naming PMI36_04622 and PMI36_04624 *aapA* for alanine aminopeptidase and *pipA* for proline iminopeptidase, respectively.

A bioassay for the plant-derived signal. To aid in the identification of the predicted plant-derived signal for *Pseudomonas* sp. GM79, we required a promoter that uses the PMI36_04623 OryR homolog for activation. In other systems, the *pip* gene adjacent to the *oryR*-type gene is often under OryR control (14–16). In the presence of the plant-derived ligands, the OryR homologs are believed to bind inverted repeat DNA elements (23) and activate gene transcription. The gene encoding the *Pseudomonas* sp. GM79 OryR homolog is also upstream from a proline iminopeptidase gene (*pipA*), and thus, we have named it *pipR* (Fig. 1). Previously, we reported that an inverted repeat sequence centered -71.5 bp upstream from the translational start site of the GM79 *pipA* gene matched the published DNA-binding site for *X. oryzae* OryR in 13 of 20 bases (2). We created the reporter plasmid pP_{*pipA*}-*gfp* (see Materials and Methods; see also Table S1 in the supplemental material), which contains a transcriptional fusion of the GM79 *pipA* promoter with the green fluorescent protein gene (*gfp*) (Fig. 2a). We hypothesized that the GM79 *pipA* promoter would be active when GM79 (pP_{*pipA*}-*gfp*) was grown in the presence of plant macerates but not when grown with AHLs (16). For these experiments, we grew the GM79 (pP_{*pipA*}-*gfp*) strain in minimal medium (see Materials and Methods) to avoid the potential activation of the PipR system, as has been reported for OryR when *X. oryzae* is grown in rich medium even in the absence of rice macerates (24). We tested six AHL signals (see Materials and Methods) with various side-chain lengths and substitutions and found that, even at relatively high concentrations (1 μ M), pP_{*pipA*}-*gfp* expression was not higher than in the controls with only water added. Our initial experiments using *Populus* leaf macerates were unsuccessful, as the growth of our reporter strain was inhibited. *Populus* leaves are known to contain high concentrations of phenolics (25), which can be toxic to bacteria. Therefore, we utilized a protocol to remove the growth inhibition activity from the *Populus* leaf macerates (see Materials and Methods). The partially purified leaf macerates, referred to hereinafter as leaf macerates, induced pP_{*pipA*}-*gfp* activity by a modest but reproducible twofold (Fig. 2b). These results are quantitatively similar to those observed with *X. oryzae* (24).

PipR can respond to protein hydrolysates and specific tripeptides. Because the genes flanking *pipR* are involved in peptide metabolism, we hypothesized that the plant signal may be peptide-like. We tested a variety of peptide-rich protein hydrolysates and found several that could activate the expression of the pP_{*pipA*}-*gfp* gene fusion (Fig. 3a). Enzymatic digests of animal tissue (Bacto-peptone), soybean meal (Bacto-soytone), and pancreatic digest of casein (Bacto-tryptone) each activated pP_{*pipA*}-*gfp* expression.

Because protein hydrolysates are rich in small peptides (26), we




FIG 2 Activities of *pipA* and *aapA* promoters in cells grown in the presence of leaf macerates or peptone. (a) DNA sequences of the *pipA* and *aapA* promoter regions cloned into HindIII-BamHI sites of the promoter-*gfp* transcriptional fusion plasmid pPROBE-NT (see Materials and Methods; see also Table S1 in the supplemental material). Blue letters indicate the first three codons of the *pipA* (top) or *aapA* (bottom) ORF, black letters indicate the intergenic, non-coding sequences, and red letters show the *pipR* DNA sequence (top, 3' end of *pipR*; bottom, noncoding strand of the 5' end of *pipR*). The 20-bp DNA sequence below both promoter sequences is the *Xanthomonas oryzae* OryR-binding sequence (24); bases identical to those in the *pipA* or *aapA* (overlapping the *pipR* ORF) promoter regions are indicated by black dots. Translation start codons (or their complements) are underlined, and the *pipR* stop codon is boxed. The two mutations in the predicted PipR-binding site of pP_{*pipAmut*}-*gfp* (Materials and Methods) are indicated by the black arrows (top, CT changed to TA). (b) Activity of the indicated promoter-*gfp* probe in GM79 wild type (WT) or the *pipR* mutant (PipR⁻) grown in the presence of water control (white bars), 0.25% leaf macerates (green bars), or 0.5% peptone (orange bars). The data are the mean relative fluorescence units (RFU) per optical density (OD) unit from six replicates, and the error bars represent the standard deviations.

screened a small library of compounds (268 dipeptides and 14 tripeptides) that are available as part of the Biolog phenotype microarrays for microbial cells for the ability to activate pP_{*pipA*}-*gfp*. Five dipeptides induced GFP above background levels: Gly-Cys, His-Gly, His-Pro, His-Ser, and Ser-Pro. Small amounts (1 mg) of His-Ser, His-Pro, and Ser-Pro are available for purchase (AnaSpec), so we retested these dipeptides using known concentrations, but only His-Ser had appreciable pP_{*pipA*}-*gfp* reporter activity (data not shown). We purchased a larger amount (100 mg) of His-Ser from another vendor (Sigma-Aldrich) but were surprised to find that this material failed to activate our reporter. Mass spectrometry analysis confirmed that the primary species (100% relative abundance) found in both samples was His-Ser ($M + H = 243.1090$, 0 ppm); however, a minor species ($\sim 5\%$ relative abundance) with a mass consistent with a tripeptide compound con-




FIG 3 The $pP_{pipA-gfp}$ reporter is activated by the addition of *Populus* leaf macerates, protein hydrolysates, and the SHS tripeptide. (a) Activity of the $pP_{pipA-gfp}$ reporter in wild-type cells grown in the presence of the following: water control, 0.5% leaf macerates (leaf), 1% Bacto-peptone (pep), 1% Bacto-soy (soy), and 1% Bacto-tryptone (tryp). (b) Dose-response for $pP_{pipA-gfp}$ activation by peptone (orange squares), leaf macerates (green squares), or SHS (red circles), HSS (blue circles), or SSH (black circles) tripeptide. The leaf macerate and peptone concentrations indicated were calculated by using the original concentrations prior to the cleanup protocol (Materials and Methods). The data are the mean RFU per OD unit from six replicates, and the error bars represent the standard deviations.

taining one histidine and two serine residues ($M + H = 330.1407$, 0 ppm) was found only in the active sample (AnaSpec). To test the hypothesis that this minor tripeptide species was responsible for the $pP_{pipA-gfp}$ reporter activation, we tested all three possible tripeptide variations (SSH, SHS, and HSS) (Fig. 3b). Two of the tripeptides, SSH and HSS, had little to no activity (Fig. 3b, black and blue circles) even at the highest concentration tested (16.5 mg/ml or 50 mM). However, the SHS tripeptide showed a moderate level of $pP_{pipA-gfp}$ reporter expression (Fig. 3b, red circles), but only at relatively high concentrations (≥ 0.33 mg/ml or 1 mM). We suspect that the signal(s) present in the leaf macerate is not the SHS tripeptide, as LuxR homologs usually respond to nM (or lower) levels of their ligand (27): at 1 mM concentrations, SHS would be easily detected by mass spectrometry of plant macerates, and we cannot find it there. However, the $pP_{pipA-gfp}$ reporter expression with the specific SHS tripeptide is further evidence that the native ligand may be peptidellike.

The PipR protein is the receptor for the response to plant macerates and the transcription activator of *pipA* expression. Leaf macerate, peptone, and the SHS tripeptide all failed to activate the expression of the $pP_{pipA-gfp}$ reporter in a *pipR* deletion mutant, thus implicating the PipR protein as the signal receptor




FIG 4 Influence of mutations in the *pipR*-flanking genes on $pP_{pipA-gfp}$ activity. In all panels, the strains are wild type (WT), *pipR* (*PMI36_04623*) mutant (PipR⁻), *aapB* (*PMI36_04621*) TMD transporter mutant (AapB⁻), *pipA* (*PMI36_04624*) mutant (PipA⁻), *aapA* (*PMI36_04622*) mutant (AapA⁻), and *pipA aapA* (*PMI36_04624* and *PMI36_04622*) double mutant (PipA⁻AapA⁻). Data are the activities of the $pP_{pipA-gfp}$ reporter grown in the water control (white bars) or in the presence of the following additions: 0.25% leaf macerates (green bars) (a), 0.5% peptone (orange bars) (b), and 1 mM (0.03%) SHS tripeptide (red bars) (c). The data are the RFU per OD unit from six replicates, and the error bars represent the standard deviations.

(Fig. 4). To confirm whether the DNA region of dyad symmetry predicted to bind the PipR protein was required for $pP_{pipA-gfp}$ activation, we mutated two conserved bases known to be important for binding of LuxR homologs (28) to create $pP_{pipAmut-gfp}$ (see Table S1 and Fig. S2a in the supplemental material) and found that PipR protein-dependent transcription from the *pipA* pro-




FIG 5 A model for PipR activation of *pipA* in GM79. The unknown signal(s) from plant macerates or peptone (stars) are taken up via the ABC-type transporter (4-component blue complex; the periplasmic-binding protein is not pictured). Once inside the cell, the signal can bind PipR, converting it to a form capable of binding the *pipA* promoter region and activating *pipA* and, possibly, *aapA*, resulting in high expression levels of peptidases (yellow lightning bolts). We hypothesize that these two peptidases act on the signal(s) or a bacterium-derived version of the signal(s) to reduce activity, thus creating a negative-feedback control loop.

motor was abolished (Fig. 2b). The *pipR* mutation was complemented by expressing *pipR* from a plasmid—although overexpression of *pipR* on a multicopy plasmid resulted in high GFP expression levels even in the absence of signal (see Fig. S2a).

There is also a potential PipR-binding site centered -91.5 bases upstream from the ATG start of the *aapA* gene (2), although this sequence overlaps the 5' coding region of the *pipR* gene (Fig. 2a). To test whether the *aapA* gene was also under control of PipR, we created an *aapA* promoter reporter plasmid, pP_{*aapA*}-*gfp* (see Table S1 and Fig. S2a in the supplemental material). The basal *gfp* expression levels of pP_{*aapA*}-*gfp* were about five times higher than those of pP_{*pipA*}-*gfp* in wild-type cells. The addition of leaf macerates had a very small effect, but peptone stimulated pP_{*aapA*}-*gfp* expression by about 1.5-fold (Fig. 2b). The expression of pP_{*aapA*}-*gfp* in a PipR deletion strain was reduced in cells grown in the presence of peptone (Fig. 2b). These results indicate that PipR strongly controls downstream *pipA* expression and has a small but measurable effect on *aapA* expression.

A mutation in the putative ABC transporter gene *aapB* abolishes induction of pP_{*pipA*}-*gfp* by plant macerates, peptone, and SHS tripeptide. The *aapA* gene and the downstream ABC-type transporter genes, now named *aapB*, *-C*, *-D*, *-E*, and *-F*, are likely cotranscribed as an operon (the *aapA*-*F* operon), as there is little intergenic sequence between them (Fig. 1). The transmembrane domain (TMD) polypeptides (encoded by *PMI36_04621* and *_04620*; *aapBC*) are predicted to have six transmembrane domains each (<http://www.cbs.dtu.dk/services/TMHMM-2.0/>), placing this transporter in the type 1 family of ABC importers (29, 30). Because a similarly annotated ABC-type peptide transporter is adjacent to the *pipR* homolog in *E. meliloti* (18) (as well as several bacterial isolates from *Populus* roots [2, 21, 31]) and because PipR responds to the tripeptide SHS, we wondered whether the putative transporter was required for the PipR signal(s) to enter the cell. To assess the role of *aapB*-*F* in *pipA* activation, we created an in-frame deletion mutation in *aapB*. This AapB mutant did not respond to leaf macerates, peptone, or the SHS tripeptide (Fig. 4). The *aapB* mutation could be complemented with an *aapB* expression plasmid (see Fig. S2b in the supplemental material). These data are consistent with the idea that the PipR signal is taken up by cells via the *aap* operon-encoded ABC-type transporter.

Peptidase mutants exhibit an enhanced *pipA*-*gfp* response. We showed as described above that *aapA* and *pipA* encode peptidases capable of cleaving several different N-terminal amino acid

residues (Table 1). We investigated whether peptidase gene inactivation had an effect on PipR signaling and found that pP_{*pipA*}-*gfp* expression was much higher in the peptidase mutants than in the wild-type GM79 when grown with leaf macerate or peptone. When grown with leaf macerates, pP_{*pipA*}-*gfp* expression in the peptidase single mutants and the *pipA aapA* double mutant was about twofold and sixfold higher, respectively, than in the wild type (Fig. 4a). These levels were even higher when cells were grown with peptone (2- to 5-fold higher for the single peptidase mutants and 14-fold higher in the *pipA aapA* double mutant relative to the levels in the wild type) (Fig. 4b). The higher pP_{*pipA*}-*gfp* activities in the single *aapA* and *pipA* peptidase mutants were complemented to nearly wild-type levels by the expression of the respective peptidase gene (see Fig. S2c and d in the supplemental material). The AapA and PipA enzymes of GM79 are both predicted to localize to the cytoplasm (32). Our results are consistent with a model where the transported plant or peptone signals are degraded by the enzymatic activities of AapA and/or PipA (Fig. 5). However, we cannot exclude the possibility that the imported signal is modified by GM79 and that this modified form of the signal is a substrate for the peptidases or that the peptidases target other components of the PipR system.

DISCUSSION

We show here that, as in several plant-associated bacteria (14–16, 18, 33), the *Populus* tree endophyte *Pseudomonas* sp. GM79 possesses a LuxR homolog that does not respond to AHL signals but instead recognizes an unknown compound in *Populus* leaf macerates. We call this LuxR homolog PipR. Our work demonstrates that PipR binds to a specific DNA sequence to activate the expression of its downstream proline iminopeptidase gene (*pipA*) in response to an unknown plant signal (Fig. 2b and 3). These results are similar to those found previously in *X. oryzae* (16, 24).

To extend our work in *Pseudomonas* sp. GM79 beyond what is known about the homologous *Xanthomonas* systems (14–17), we examined whether the genes surrounding *pipR* contribute to its activity. These flanking genes are annotated as being involved in peptide degradation and transport, leading us to hypothesize that PipR could respond to peptidic compounds. Indeed, we found that a variety of peptide-rich peptones (including Bacto-peptone) and a specific tripeptide (SHS) could activate a PipR-dependent reporter.

A strain with a mutation in a transmembrane domain (TMD)

protein gene (*aapB*) of the ABC transporter near *pipR* (Fig. 1) did not respond to plant leaf macerates, peptone, or the SHS tripeptide (Fig. 4), suggesting that these signal(s) enter cells by active transport. Transporters are not required for entry of AHL signals into cells, as AHLs can diffuse into and out of bacterial cells (34, 35). However, ABC-type transporters are used in many of the Gram-positive quorum-sensing systems for the import of peptide pheromone signals (reviewed in Cook and Federle [36]). There are no ABC-type transporters genetically linked to the *oryR* homologs in *Xanthomonas* species (<http://img.jgi.doe.gov>); however, upstream from the *oryR*-type genes is a gene annotated as a member of the amino acid/polyamine/organocation (APC) transporter superfamily (TC 2.A.3); interestingly this transporter gene is highly expressed (12-fold higher than in the wild type) in an *X. axonopodis* strain overexpressing an OryR (XagR) homolog (14). One could imagine that this APC transporter may play a role in *Xanthomonas* species similar to that of the GM79 ABC transporter: import of the OryR-responsive plant signal(s).

Strains with mutations in the flanking peptidase genes showed elevated expression of pP_{*pipA*}-*gfp* compared to the level in the wild type when grown in the presence of leaf macerates and peptone (Fig. 4a and b). A similar result, increased *pip* expression compared to the level in the wild type, was reported for an *X. campestris* Pip⁻ mutant (15).

One interpretation of these results is that the peptidases enzymatically degrade the PipR signal(s) and in the peptidase mutants, less signal degradation occurs, resulting in higher PipR-dependent gene activation. A model of the PipR system consistent with these data is depicted in Fig. 5. Signal(s) enter the cell via the ABC-type transporter and activate PipR-dependent transcription of *pipA*. Although the Pip activity from *X. campestris* has been reported as localized to the periplasm (15), both AapA and PipA of *Pseudomonas* sp. GM79 are predicted to be cytoplasmic (32). For *Pseudomonas* sp. GM79, our data suggest that AapA and PipA can utilize a transported PipR ligand as a substrate, although we cannot exclude the possibility that they act on a compound derived from the ligand or on some other component of the PipR signaling system. This arrangement constitutes a negative-feedback loop for the system, which would ensure a rapid inactivation of *pipA* transcription when the signal becomes limited.

There is increasing evidence that not all orphan LuxR homologs sense AHLs. In addition to the plant-responsive OryR-type transcription factors discussed here, the LuxR homologs CarR (*Serratia* sp. strain 39006) (37) and MalR (*Burkholderia thailandensis*) (38), which both retain all of the conserved amino acid residues in the AHL-binding domain of LuxR homologs, do not require an AHL for activity. There are also examples of orphan LuxR homologs that utilize endogenous non-AHL compounds as signal ligands, including PluR (*Photorhabdus luminescens*) (9) and PauR (*Photorhabdus asymbiotica*) (10), which respond to α -pyrones and dialkylresorcinols, respectively. In addition, activators of AHL-responsive LuxR homologs have been identified which bear little resemblance to the native AHL signal ligand (39). Our work suggests that the GM79 PipR ligand is peptidic. It will be interesting to purify and elucidate the structures of the PipR signals from both the plant macerate and peptone material. We predict that the plant and peptone signals will be structurally similar but not necessarily identical.

We are curious to test whether the PipR system mutants created here are also impaired in *Populus* host interactions, as is the

case with PipR homologs in several plant pathogens (14–17) and mutualists (12, 18). We are also interested to know which GM79 genes, other than the peptidase genes, are under the control of PipR. In other bacteria, PipR homologs regulate not only proline iminopeptidase gene expression but additional traits, including those important for colonization of and movement through the plant host (motility [40] and biosurfactant and adhesin production [14]), accumulation of osmoprotectants (14), and synthesis of antifungal compounds (12).

PipR homologs are encoded in the genomes of several plant-associated bacterial genera, including *Xanthomonas*, *Dickeya*, *Agrobacterium*, *Rhizobium*, *Ensifer*, and *Pseudomonas* (reviewed in references 5, 11, and 12), and whether or not all these transcription factors respond to the same plant signal or different but related compounds is not known. The plant-responsive OryRs are of general importance, as they appear to play a role in the health of economically important plants (14–17). We believe *Pseudomonas* GM79 is a useful model to begin to understand the chemistry of what may prove to be a new family of interkingdom signals, or cues, involved in plant-bacterium interactions.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The bacterial strains and plasmids used are described in Table S1 in the supplemental material. *Pseudomonas* sp. GM79 and its derived strains were grown in R2A or M9 minimal medium (41) with 10 mM succinate (M9-suc) at 30°C. *E. coli* strains were grown in LB broth (42) and incubated at 37°C with shaking. Antibiotics were used when required at the following concentrations: 50 μ g/ml (*Escherichia coli*) or 25 μ g/ml (GM79) kanamycin, 100 μ g/ml ampicillin, 20 μ g/ml (*E. coli*) or 50 μ g/ml (GM79) gentamicin, and 10 μ g/ml tetracycline.

Chemicals. AHL signals were tested at 1 μ M concentrations and included *N*-butanoyl-L-homoserine lactone (C4-HSL); *N*-3-oxo-hexanoyl-L-HSL (3-oxo-C6-HSL), *N*-3-oxo-octanoyl-L-HSL (3-oxo-C8-HSL), *N*-3-hydroxyoctanoyl-L-HSL (3-hydroxy-C8-HSL), *N*-3-oxododecanoyl-L-HSL (3-oxo-C12-HSL), and *N*-(*p*-coumaroyl)-L-HSL (*p*-coumaroyl-HSL) (purchased from Sigma-Aldrich, St. Louis, MO, or the University of Nottingham, Nottingham, United Kingdom). The β -naphthylamide and *p*-nitroanilide amino acid substrates were purchased from Sigma-Aldrich. Bacto-peptone, Bacto-soytone, and Bactotryptone were purchased from Becton, Dickinson, and Company (Franklin Lakes, NJ). The HS dipeptide was purchased from both AnaSpec (Fremont, CA) and Sigma-Aldrich. The tripeptides HSS, SHS, and SSH were custom synthesized by Peptide 2.0 (Chantilly, VA).

Reporters, mutants, and plasmids. All plasmids and primer sequences are described in Tables S1 and S2, respectively, in the supplemental material. We created the reporter plasmids pP_{*pipA*}-*gfp* and pP_{*aapA*}-*gfp* by PCR amplifying 263-bp DNA fragments containing the intergenic promoter regions, using GM79 genomic DNA as the template, and cloning the PCR products into HindIII-BamHI-digested pPROBE-NT (43). To create pP_{*pipA*}^{mut}-*gfp*, we ordered a gBlock gene fragment (Integrated DNA Technologies, Coralville, IA) containing the exact promoter sequence that was cloned into pP_{*pipA*}-*gfp*, except that the CT nucleotides present in the predicted PipR-binding site were changed to TA. Mutant constructions were performed similarly: DNA sequences of about 500 bp from both up- and downstream of the desired in-frame deletion locations were either created by two-step overlap extension PCR amplification (Δ *pipA* mutation) or synthesized as a single DNA fragment of about 1 kb (Eurofins Genomics, Huntsville, AL) and cloned into EcoRI-BamHI-digested suicide vector pEX19-Gm (44). The knockout suicide vector was introduced into *Pseudomonas* GM79 strains by conjugal mating, and single-crossover mutants were selected by plating on M9-suc agar containing gentamicin. Double-crossover mutants were selected by streaking onto R2A agar containing 5% sucrose and screened for loss of Gm^r.

For complementation of the *pipR* mutant, we PCR amplified a DNA fragment containing 250 bp of the *pipR* promoter sequence, the *pipR* gene, and the intergenic region between *pipR* and *pipA* and cloned the PCR product into HindIII-BamHI-digested pPROBE-NT (43). For complementation of the *pipA* mutant, the *pipA* gene and 254 bp of its promoter sequence were PCR amplified by using GM79 genomic DNA as the template, and the product was cloned into the BamHI-HindIII sites of pMMB67EH-TetRA. The plasmid for *aapA* complementation was constructed similarly except that only 190 bp of its promoter sequence was included. Because the *aapB* gene likely shares a promoter with the upstream *aapA* gene, we used the same forward primer as was used for complementation of the *aapA* mutant (Aap-CompFOR) plus a reverse primer for the 3' end of the TMD gene (TsptCompREV) and used genomic DNA from the *aapA* mutant (79ΔAapA strain; see Table S1 in the supplemental material) as a PCR template. The PCR product was cloned into BamHI-HindIII-digested pMMB67EH-TetRA. Complementation plasmids (or pMMB67EH-TetRA vector controls) were introduced into the appropriate mutant strains harboring the pP_{*pipA*}-*gfp* reporter by conjugal mating. All mutant and plasmid constructs were confirmed by DNA sequencing.

Purification of His₆-tagged proteins. To obtain purified PipA and AapA, the genes were cloned into the His₆-tagged protein expression vector pQE-30, creating plasmids pQEpipA and pQEaapA, respectively (see Tables S1 and S2 in the supplemental material). *E. coli* M15 pRep4 containing either pQEpipA or pQEaapA was grown at 30°C in 500 ml of LB plus antibiotics to an optical density at 600 nm of 0.6 (OD₆₀₀). The production of His-tagged protein was then induced by the addition of 1 mM isopropyl-β-D-thiogalactopyranoside (IPTG) and incubation was continued at 16°C overnight, after which cells were pelleted, resuspended in buffer (50 mM NaH₂PO₄, 300 mM NaCl, 10 mM imidazole, pH 8), broken by French pressure cell, and centrifuged for 20 min at 14,000 × g. The His₆-tagged proteins were purified from clarified cell extracts by cobalt resin column chromatography (Qiagen, Valencia, CA).

Peptidase assays. Enzyme assays were performed in 0.1-ml volumes containing 50 mM Tris, pH 7.4, 10 mM MnCl₂, 0.75 mM amino acid substrate, and 0.6 μg His-tagged protein. Reaction mixtures were incubated for 20 min at 30°C and stopped by equivalent addition of 0.1 M acetic acid. Substrate cleavage was assessed by measuring either fluorescence (excitation at 355 nm and emission at 415 nm) for the β-naphthylamine-linked substrates or color [410 nm, molar extinction coefficient (M⁻¹ cm⁻¹) = 8,000] for the *p*-nitroanilide-linked substrates.

Reporter assays. Bioassays were performed in M9-suc for two reasons. (i) OryR accumulated in *X. oryzae* when grown in rich medium (peptone-yeast extract-salts) in the absence of plant macerates (24), suggesting that something in complex medium can induce the system. Therefore, we decided to use a minimal medium so as not to confound our results. (ii) Succinate was chosen as the carbon and energy source in the minimal medium because there were no significant growth rate differences between the wild-type and *pipR* mutant strains in this medium. Strains containing pP_{*pipA*}-*gfp* were incubated overnight (24 h) in M9-suc plus kanamycin at 30°C with shaking. Cells were diluted 1:100 into fresh medium, 150-μl aliquots were added to individual wells of a 96-well microtiter dish containing 7.5 μl (except as indicated in Fig. 3) of material to be tested (leaf macerates, peptone, peptides, or AHLs), and the plates were sealed with Breathe-Easy sealing membrane (Research Products International, Mount Prospect, IL) and incubated at room temperature for ~24 h. GFP fluorescence (excitation at 485 nm and emission at 535 nm) and growth (OD₅₉₅) were assessed using a Tecan Genios pro plate reader, and data were plotted as relative fluorescence units (RFU) per OD unit.

Preparation of partially purified *Populus* leaf macerates and peptone material. Because various additions to the bioassay strain culture showed both inhibitory (leaf macerates) and stimulatory (Bacto-peptone) growth effects, we developed a two-step cleanup protocol to produce the partially purified material used in all of our experiments. For leaf macerates, 5 g of *P. deltoides* WV94 leaves (greenhouse grown) were frozen in liquid nitrogen, macerated with a mortar and pestle, added to 100 ml of

Milli-Q water (5% weight/vol), sterilized by autoclaving, and then filtered to remove plant tissue (as described in reference 24). Peptone was prepared in Milli-Q water at a concentration of 10 g/100 ml (10% wt/vol). Both leaf and peptone material were then passed over a C₁₈ reverse-phase (RP) solid-phase extraction (SPE) cartridge (Waters Corp., Milford, MA). The C₁₈-RP cartridge did not bind the active material but did retain a large amount of nonactive material (including the bacterial-growth-inhibiting activity in the leaf macerates). The flowthrough fraction was passed through an Amicon ultra-15 filter with a nominal molecular weight limit of 3,000 (Merck Millipore, Cork, Ireland) to remove any higher-mass, nonactive compounds. Partially purified material was concentrated, resuspended in Milli-Q water to its original concentration, and filter sterilized with a 0.2-μm syringe filter.

Peptide screening with Biolog plates. Biolog phenotype microarray plates for nitrogen utilization assays (PM6, PM7, and PM8) were used (Biolog, Inc., Hayward, CA). GM79 (pP_{*pipA*}-*gfp*) cells in M9-suc medium were incubated in the Biolog plates for 18 h, and then GFP fluorescence (excitation at 485 nm and emission at 535 nm) and growth (OD₅₉₅) were determined. As a control for PipR activity, 1% peptone was added to the L-glutamine positive control present on every Biolog plate.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <http://mbio.asm.org/lookup/suppl/doi:10.1128/mBio.01101-16/-/DCSupplemental>.

Figure S1, EPS file, 1.1 MB.

Figure S2, EPS file, 0.5 MB.

Table S1, DOCX file, 0.1 MB.

Table S2, DOCX file, 0.1 MB.

ACKNOWLEDGMENTS

We thank Dave Weston (ORNL) for providing *Populus* leaf material; Hemantha Don Kulasekara and Sam Miller (University of Washington) for sharing the pMMB67EH-TetRA vector; and Colin Manoil (University of Washington) for the gift of Biolog plates.

FUNDING INFORMATION

This work, including the efforts of Amy L. Schaefer, Yasuhiro Oda, Bruna Goncalves Coutinho, Dale A. Pelletier, Justin Weiburg, Everett Peter Greenberg, and Caroline S. Harwood, was funded by Department of Energy (BER) Genomic Science Program (DE-AC05-00OR22725).

This research was sponsored by the Genomic Science Program, U.S. Department of Energy, Office of Science, Biological and Environmental Research, as part of the Plant Microbe Interfaces Scientific Focus Area (<http://pmi.ornl.gov>). Oak Ridge National Laboratory is managed by UT-Battelle LLC, for the U.S. Department of Energy under contract DE-AC05-00OR22725.

REFERENCES

- Gottel NR, Castro HF, Kerley M, Yang Z, Pelletier DA, Podar M, Karpinets T, Uberbacher E, Tuskan GA, Vilgalys R, Doktycz MJ, Schadt CW. 2011. Distinct microbial communities within the endosphere and rhizosphere of *Populus deltoides* roots across contrasting soil types. *Appl Environ Microbiol* 77:5934–5944. <http://dx.doi.org/10.1128/AEM.05255-11>.
- Schaefer AL, Lappala CR, Morlen RP, Pelletier DA, Lu TY, Lankford PK, Harwood CS, Greenberg EP. 2013. LuxR- and LuxI-type quorum-sensing circuits are prevalent in members of the *Populus deltoides* microbiome. *Appl Environ Microbiol* 79:5745–5752. <http://dx.doi.org/10.1128/AEM.01417-13>.
- Whitehead NA, Barnard AM, Slater H, Simpson NJ, Salmond GP. 2001. Quorum-sensing in gram-negative bacteria. *FEMS Microbiol Rev* 25: 365–404. <http://dx.doi.org/10.1111/j.1574-6976.2001.tb00583.x>.
- Waters CM, Bassler BL. 2005. Quorum sensing: cell-to-cell communication. *Annu Rev Cell Dev Biol* 21:319–346. <http://dx.doi.org/10.1146/annurev.cellbio.21.012704.131001>.
- Patankar AV, González JE. 2009. Orphan LuxR regulators of quorum sensing. *FEMS Microbiol Rev* 33:739–756. <http://dx.doi.org/10.1111/j.1574-6976.2009.00163.x>.

6. Lee JH, Lequette Y, Greenberg EP. 2006. Activity of purified QscR, a *Pseudomonas aeruginosa* orphan quorum-sensing transcription factor. *Mol Microbiol* 59:602–609. <http://dx.doi.org/10.1111/j.1365-2958.2005.04960.x>.
7. Sperandio V. 2010. SdiA sensing of acyl-homoserine lactones by enterohemorrhagic *E. coli* (EHEC) serotype O157:H7 in the bovine rumen. *Gut Microbes* 1:432–435. <http://dx.doi.org/10.4161/gmic.1.6.14177>.
8. Dyszel JL, Soares JA, Swearingen MC, Lindsay A, Smith JN, Ahmer BM. 2010. *E. coli* K-12 and EHEC genes regulated by SdiA. *PLoS One* 5:e8946. <http://dx.doi.org/10.1371/journal.pone.0008946>.
9. Brachmann AO, Brameyer S, Kresovic D, Hitkova I, Kopp Y, Manske C, Schubert K, Bode HB, Heermann R. 2013. Pyrones as bacterial signaling molecules. *Nat Chem Biol* 9:573–578. <http://dx.doi.org/10.1038/nchembio.1295>.
10. Brameyer S, Kresovic D, Bode HB, Heermann R. 2015. Dialkylresorcinols as bacterial signaling molecules. *Proc Natl Acad Sci U S A* 112:572–577. <http://dx.doi.org/10.1073/pnas.1417685112>.
11. González JF, Venturi V. 2013. A novel widespread interkingdom signaling circuit. *Trends Plant Sci* 18:167–174. <http://dx.doi.org/10.1016/j.tplants.2012.09.007>.
12. Subramoni S, Gonzalez JF, Johnson A, Péchy-Tarr M, Rochat L, Paulsen I, Loper JE, Keel C, Venturi V. 2011. Bacterial subfamily of LuxR regulators that respond to plant compounds. *Appl Environ Microbiol* 77:4579–4588. <http://dx.doi.org/10.1128/AEM.00183-11>.
13. Patel HK, Suárez-Moreno ZR, Degraasi G, Subramoni S, González JF, Venturi V. 2013. Bacterial LuxR solos have evolved to respond to different molecules including signals from plants. *Front Plant Sci* 4:447. <http://dx.doi.org/10.3389/fpls.2013.00447>.
14. Chatnaparat T, Prathuangwong S, Ionescu M, Lindow SE. 2012. XagR, a LuxR homolog, contributes to the virulence of *Xanthomonas axonopodis* pv. *glycines* to soybean. *Mol Plant Microbe Interact* 25:1104–1117. <http://dx.doi.org/10.1094/MPMI-01-12-0008-R>.
15. Zhang L, Jia Y, Wang L, Fang R. 2007. A proline iminopeptidase gene upregulated in plant by a LuxR homolog is essential for pathogenicity of *Xanthomonas campestris* pv. *campestris*. *Mol Microbiol* 65:121–136. <http://dx.doi.org/10.1111/j.1365-2958.2007.05775.x>.
16. Ferluga S, Bigirimana J, Höfte M, Venturi V, Lux A. 2007. R homolog of *Xanthomonas oryzae* pv. *oryzae* is required for optimal rice virulence. *Mol Plant Pathol* 8:529–538. <http://dx.doi.org/10.1111/j.1364-3703.2007.00415.x>.
17. Xu H, Zhao Y, Qian G, Liu F. 2015. XocR, a LuxR solo requested for virulence in *Xanthomonas oryzae* pv. *oryzicola*. *Front Cell Infect Microbiol* 5:37. <http://dx.doi.org/10.3389/fcimb.2015.00037>.
18. Patankar AV, González JE. 2009. An orphan LuxR homolog of *Sinorhizobium meliloti* affects stress adaptation and competition for nodulation. *Appl Environ Microbiol* 75:946–955. <http://dx.doi.org/10.1128/AEM.01692-08>.
19. Timm CM, Campbell AG, Utturkar SM, Jun SR, Paraless RE, Tan WA, Robeson MS, Lu TY, Jawdy S, Brown SD, Ussery DW, Schadt CW, Tuskan GA, Doktycz MJ, Weston DJ, Pelletier DA. 2015. Metabolic functions of *Pseudomonas fluorescens* strains from *Populus deltoides* depend on rhizosphere or endosphere isolation compartment. *Front Microbiol* 6:1118. <http://dx.doi.org/10.3389/fmicb.2015.01118>.
20. Jun SR, Wassenaar TM, Nookaew I, Hauser L, Wanchai V, Land M, Timm CM, Lu TY, Schadt CW, Doktycz MJ, Pelletier DA, Ussery DW. 2016. Diversity of *Pseudomonas* genomes, including *Populus*-associated isolates, as revealed by comparative genome analysis. *Appl Environ Microbiol* 82:375–383. <http://dx.doi.org/10.1128/AEM.02612-15>.
21. Brown SD, Utturkar SM, Klingeman DM, Johnson CM, Martin SL, Land ML, Lu TY, Schadt CW, Doktycz MJ, Pelletier DA. 2012. Twenty-one *Pseudomonas* genomes and nineteen genomes from diverse bacteria isolated from the rhizosphere and endosphere of *Populus deltoides*. *J Bacteriol* 194:5991–5993. <http://dx.doi.org/10.1128/JB.01243-12>.
22. Subramoni S, Venturi V. 2009. PpoR is a conserved unpaired LuxR solo of *Pseudomonas putida* which binds N-acyl homoserine lactones. *BMC Microbiol* 9:125. <http://dx.doi.org/10.1186/1471-2180-9-125>.
23. Devine JH, Shadel GS, Baldwin TO. 1989. Identification of the operator of the lux regulon from the *Vibrio fischeri* strain ATCC7744. *Proc Natl Acad Sci U S A* 86:5688–5692. <http://dx.doi.org/10.1073/pnas.86.15.5688>.
24. Ferluga S, Venturi V. 2009. OryR is a LuxR-family protein involved in interkingdom signaling between pathogenic *Xanthomonas oryzae* pv. *oryzae* and rice. *J Bacteriol* 191:890–897. <http://dx.doi.org/10.1128/JB.01507-08>.
25. Tsai CJ, Harding SA, Tschaplinski TJ, Lindroth RL, Yuan Y. 2006. Genome-wise analysis of the structural genes regulating defense phenylpropanoid metabolism in *Populus*. *New Phytol* 172:47–62. <http://dx.doi.org/10.1111/j.1469-8137.2006.01798.x>.
26. Becton, Dickinson and Company. 2006. BD Bionutrients technical manual: advanced bioprocessing, 3rd ed, revised. BD, Sparks, MD.
27. Ahlgren NA, Harwood CS, Schaefer AL, Giraud E, Greenberg EP. 2011. Aryl-homoserine lactone quorum sensing in stem-nodulating photosynthetic bradyrhizobia. *Proc Natl Acad Sci U S A* 108:7183–7188. <http://dx.doi.org/10.1073/pnas.1103821108>.
28. Hirakawa H, Oda Y, Phattarasukol S, Armour CD, Castle JC, Raymond CK, Lappala CR, Schaefer AL, Harwood CS, Greenberg EP. 2011. Activity of the *Rhodospseudomonas palustris* p-coumaroyl-homoserine lactone-responsive transcription factor RpaR. *J Bacteriol* 193:2598–2607. <http://dx.doi.org/10.1128/JB.01479-10>.
29. Locher KP. 2009. Structure and mechanism of ATP-binding cassette transporters. *Philos Trans R Soc Lond B Biol Sci* 364:239–245. <http://dx.doi.org/10.1098/rstb.2008.0125>.
30. Davidson AL, Chen J. 2004. ATP-binding cassette transporters in bacteria. *Annu Rev Biochem* 73:241–268. <http://dx.doi.org/10.1146/annurev.biochem.73.011303.073626>.
31. Brown SD, Klingeman DM, Lu TY, Johnson CM, Utturkar SM, Land ML, Schadt CW, Doktycz MJ, Pelletier DA. 2012. Draft genome sequence of *Rhizobium* sp. strain PD01-076, a bacterium isolated from *Populus deltoides*. *J Bacteriol* 194:2383–2384. <http://dx.doi.org/10.1128/JB.00198-12>.
32. Yu NY, Wagner JR, Laird MR, Melli G, Rey S, Lo R, Dao P, Sahinalp SC, Ester M, Foster LJ, Brinkman FS. 2010. PSORTb 3.0: improved protein subcellular localization prediction with refined localization sub-categories and predictive capabilities for all prokaryotes. *Bioinformatics* 26:1608–1615. <http://dx.doi.org/10.1093/bioinformatics/btq249>.
33. Patel HK, Ferrante P, Covaceuszach S, Lamba D, Scortichini M, Venturi V. 2014. The kiwifruit emerging pathogen *Pseudomonas syringae* pv. *actinidiae* does not produce AHLs but possesses three LuxR solos. *PLoS One* 9:e87862. <http://dx.doi.org/10.1371/journal.pone.0087862>.
34. Kaplan HB, Greenberg EP. 1985. Diffusion of autoinducer is involved in regulation of the *Vibrio fischeri* luminescence system. *J Bacteriol* 163:1210–1214.
35. Pearson JP, Van Delden C, Iglewski BH. 1999. Active efflux and diffusion are involved in transport of *Pseudomonas aeruginosa* cell-to-cell signals. *J Bacteriol* 181:1203–1210.
36. Cook LC, Federle MJ. 2014. Peptide pheromone signaling in *Streptococcus* and *Enterococcus*. *FEMS Microbiol Rev* 38:473–492. <http://dx.doi.org/10.1111/1574-6976.12046>.
37. Poulter S, Carlton TM, Spring DR, Salmond GP. 2011. The *Serratia* LuxR family regulator CarR 39006 activates transcription independently of cognate quorum sensing signals. *Mol Microbiol* 80:1120–1131. <http://dx.doi.org/10.1111/j.1365-2958.2011.07634.x>.
38. Truong TT, Seyedsayamdot M, Greenberg EP, Chandler JR. 2015. A *Burkholderia thailandensis* acyl-homoserine lactone-independent orphan LuxR homolog that activates production of the cytotoxin malleilactone. *J Bacteriol* 197:3456–3462. <http://dx.doi.org/10.1128/JB.00425-15>.
39. Müh U, Hare BJ, Duerkop BA, Schuster M, Hanzelka BL, Heim R, Olson ER, Greenberg EP. 2006. A structurally unrelated mimic of a *Pseudomonas aeruginosa* acyl-homoserine lactone quorum-sensing signal. *Proc Natl Acad Sci U S A* 103:16948–16952. <http://dx.doi.org/10.1073/pnas.0608348103>.
40. González JF, Myers MP, Venturi V. 2013. The inter-kingdom solo OryR regulator of *Xanthomonas oryzae* is important for motility. *Mol Plant Pathol* 14:211–221. <http://dx.doi.org/10.1111/j.1364-3703.2012.00843.x>.
41. Miller JH. 1972. Experiments in Molecular Genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
42. Sambrook J, Russell DW. 2001. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
43. Miller WG, Leveau JH, Lindow SE. 2000. Improved *gfp* and *inaZ* broad-host-range promoter-probe vectors. *Mol Plant Microbe Interact* 13:1243–1250. <http://dx.doi.org/10.1094/MPMI.2000.13.11.1243>.
44. Hoang TT, Karkhoff-Schweizer RR, Kutchma AJ, Schweizer HP. 1998. A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* 212:77–86. [http://dx.doi.org/10.1016/S0378-1119\(98\)00130-9](http://dx.doi.org/10.1016/S0378-1119(98)00130-9).