

POSTER PRESENTATION

Open Access

Development of dynamical weather-disease models to project and forecast malaria in Africa

Volker Ermert^{1*}, Andreas H Fink¹, Andrew P Morse², Anne E Jones², Heiko Paeth³, Francesca Di Giuseppe⁴, Adrian M Tompkins⁵

From Challenges in malaria research
Basel, Switzerland. 10-12 October 2012

Background

Weather and climate play an important role in the spread of malaria. Suitable weather conditions for malaria are found in sub-Saharan Africa, where most of the worldwide malaria cases and deaths are found. For this reason, integrated weather-disease malaria models are useful tools to project the malaria future and to provide monthly-to-seasonal forecasts.

Methods

Malaria projections and forecasts are undertaken by two dynamical mathematical-biological malaria models: (i) the LMM (Liverpool Malaria Model) [1-3] and (ii) VECTRI (VEctor-borne disease community model of the International Centre for Theoretical Physics, TRIeste). Both models are driven by daily temperature and precipitation values. An improved version of the LMM was introduced by [2], which was calibrated by malaria field observations from West Africa [3]. Regarding the assessment of the impact of climate change on malaria [4], the LMM was driven by data from the REgional MOdel (REMO) including the effect of land surface changes.

For the QWeCI (Quantifying Weather and Climate Impacts on health in developing countries) project, a seamless weather prediction system has been developed at ECMWF by appending the first 25 days of the monthly forecasting system with the Seasonal Forecasting System 4 to provide a continuous 120 day lead time prediction. The forecast is calibrated to correct for displacement errors of West African monsoonal precipitation.

Results and outlook

The malaria projections up to 2050 [4] based on the integrated REMO-LMM reveal a southward shift of the epidemic malaria area in West Africa due to the precipitation decline. The increased temperatures lead to an increase of transmission in highland territories. Formerly, malaria free areas become epidemic, whereas the epidemic risk is decreased in lower-altitude regions.

Actual research within the EU Seventh Framework Programme (FP7) QWeCI and HEALTHY FUTURES projects is underway to exploit the feasibility of monthly-to-seasonal malaria forecasts. QWeCI is currently developing prototype seamless malaria forecasts for Malawi (<http://nwmstest.ecmwf.int/products/forecasts/d/inspect/catalog/research/qweci/>).

The LMM and VECTRI neglect various important malaria factors like immunity, malaria control activities, or different vector characteristics. Further development of VECTRI will be undertaken to include other relevant malaria factors. Note that VECTRI represents a community model meaning that the model and code is publicly available (<http://users.ictp.it/~tompkins/vectri/>). The LMM is included in the so-called Disease Model Cradle (DMC) that is downloadable (http://www.liv.ac.uk/qweci/project_outputs/). Open-access web versions of both models are applicable for point data (see <http://qweci.uni-koeln.de>).

Author details

¹Institute of Geophysics and Meteorology, University of Cologne, Cologne, Germany. ²School of Environmental Sciences, University of Liverpool, Liverpool, UK. ³Institute of Geography, University of Würzburg, Würzburg, Germany. ⁴European Centre for Medium-range Weather Forecasts (ECMWF), Reading, UK. ⁵Earth System Physics, Abdus Salam International Centre for Theoretical Physics (ICTP), Trieste, Italy.

Published: 9 November 2012

¹Institute of Geophysics and Meteorology, University of Cologne, Cologne, Germany

Full list of author information is available at the end of the article

References

1. Hoshen MB, Morse AP: A weather-driven model of malaria transmission. *Malar J* 2004, **3**:32.
2. Ermert V, Fink AH, Jones AE, Morse AP: Development of a new version of the Liverpool Malaria Model. I. Refining the parameter settings and mathematical formulation of basic processes based on a literature review. *Malar J* 2011, **10**:35.
3. Ermert V, Fink AH, Jones AE, Morse AP: Development of a new version of the Liverpool Malaria Model. II. Calibration and validation for West Africa. *Malar J* 2011, **10**:62.
4. Ermert V, Fink AH, Morse AP, Paeth H: The Impact of Regional Climate Change on Malaria Risk due to Greenhouse Forcing and Land-Use Changes in Tropical Africa. *EHP* 2012, **120**:77-84.

doi:10.1186/1475-2875-11-S1-P133

Cite this article as: Ermert *et al.*: Development of dynamical weather-disease models to project and forecast malaria in Africa. *Malaria Journal* 2012 **11**(Suppl 1):P133.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

