

# SARS-CoV-2 nonstructural protein 6 from Alpha to Omicron: evolution of a transmembrane protein

Shuchen Feng,<sup>1</sup> Amornrat O'Brien,<sup>1</sup> Da-Yuan Chen,<sup>2,3</sup> Mohsan Saeed,<sup>2,3</sup> Susan C. Baker<sup>1</sup>

**AUTHOR AFFILIATIONS** See affiliation list on p. 6.

**ABSTRACT** We recently reported that mutations in both the spike glycoprotein and nonstructural protein 6 (nsp6) were associated with attenuation of the SARS-CoV-2 Omicron BA.1 variant. While mutations in spike allow evasion of neutralizing antibodies and promote specific modes of viral entry, the role of nsp6 mutations in pathogenesis is less clear. Nsp6 is essential for modifying the endoplasmic reticulum and generating double-membrane vesicles, the site of viral RNA replication. To investigate the evolution of nsp6, we evaluated 91,596 high-confidence human SARS-CoV-2 whole-genome sequences across 19 variants and lineages. While nsp6 of early variants of concern, such as Alpha, Beta, and Gamma, carried a triple amino acid deletion (106–108, termed  $\Delta$ SGF), the Delta, Epsilon, and Mu lineages retained the ancestral nsp6 sequence. For nsp6 in the emerging Omicron variants, we report a transition from an amino acid 105–107  $\Delta$ LSG deletion in BA.1 to increased dominance of the  $\Delta$ SGF in BA.2 and subsequent lineages. Our findings indicate that deletion within nsp6 was independently selected in multiple lineages of SARS-CoV-2, both early and late in the pandemic. Analysis of SARS-CoV-2-related coronaviruses in bats and pangolins revealed nsp6 sequences similar to the ancestral SARS-CoV-2 virus, indicating that the deletion in nsp6 may be an adaptation to replication in humans. Analysis of nsp6 sequences from multiple coronaviruses predicts a multipass transmembrane protein with a conserved C-terminal domain. Monitoring and evaluating changes in nsp6 and other nonstructural proteins will contribute to our understanding of factors associated with the attenuation of pandemic coronaviruses.

**IMPORTANCE** There is an ongoing need to evaluate genetic changes in SARS-CoV-2 for effects on transmission and pathogenesis. We recently reported an unexpected role for replicase component nsp6, in addition to changes in spike, in the attenuation of Omicron BA.1. In this commentary, we document a triple-amino-acid deletion in a predicted luminal domain of nsp6 that was found in multiple, independent variants of SARS-CoV-2, including all recent Omicron lineages. Furthermore, we modeled the predicted structure of nsp6, implicating a multipass transmembrane architecture as conserved in members of the *Coronaviridae* family. This information can guide future studies investigating the role of nsp6 in the pathogenesis of existing and emerging coronaviruses.

**KEYWORDS** SARS-CoV-2, nonstructural protein 6, nsp6, SARS-CoV-2 evolution, nsp6 topology model, SARS-CoV-2 attenuation

One major challenge of the ongoing COVID-19 pandemic is that emerging variants of SARS-CoV-2 may escape neutralizing antibodies generated by vaccination and/or previous infection. To date, variants of concern (VOCs) designated Alpha to Omicron have been documented (1). Substitutions and deletions in the spike (S) glycoprotein facilitate escape from neutralizing antibodies and can result in alternative mechanisms of virus entry (2, 3). Additionally, changes in the nucleocapsid protein have been associated with increased virulence, likely due to enhanced packaging of viral RNA (4).

**Editor** J. S. Malik Peiris, University of Hong Kong, Pokfulam, Hong Kong

Address correspondence to Susan C. Baker, sbaker1@luc.edu.

The authors declare no conflict of interest.

See the funding table on p. 6.

**Received** 17 March 2023

**Accepted** 10 June 2023

**Published** 21 July 2023

Copyright © 2023 Feng et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).


Since its emergence in November 2021 in South Africa (5), the Omicron variant has been associated with increased transmission but attenuated virulence compared to the ancestral virus (Wuhan-Hu-1) or the Delta VOC in multiple studies of animal models (6, 7) and human outcomes (8, 9). Prior vaccination or infection may also mitigate the pathogenesis of emerging variants (10). However, the contributing mutations were unknown. Recently, we reported an unexpected finding that mutations in spike and nonstructural protein 6 (nsp6), one of the 16 non-structural proteins generated by the proteolytic processing of the viral pp1ab polyprotein, were associated with the attenuation of Omicron BA.1 in K18-hACE2 mice (11). We found that the Omicron spike alone was not sufficient to fully attenuate the virulence of SARS-CoV-2, yet the combination of both Omicron spike and Omicron nsp6 attenuated the virulence to the level of Omicron BA.1. These findings were corroborated by a recent study that employed chimeric combinations of Delta and Omicron variants, identifying the predominant role of spike and nsp6 in Omicron BA.1's attenuation (12). Omicron BA.1 nsp6 contains a triple amino acid deletion (105–107,  $\Delta$ LSG) and an amino acid substitution (I189V). Here, we provide an updated analysis on the evolution and predicted structure and function of SARS-CoV-2 nsp6.

### **Nsp6 deletion mutants found in early variants enhance ER membrane zippering**

Previous studies documented a critical role of nsp6 in coronavirus replication via the formation of an nsp3-nsp4-nsp6 complex, a pre-requisite for the biogenesis of double-membrane vesicles (DMVs) (13–16). DMVs are the site of viral RNA synthesis for coronaviruses (17, 18). SARS-CoV-2, like other coronaviruses, induces ER-derived DMVs as part of the viral replication and transcription mechanisms (18–20). An elegant study by Ricciardi and co-workers suggested that SARS-CoV-2 nsp6 is responsible for organizing and connecting the DMVs to the ER in a process termed ER zippering (21). Interestingly, they reported that nsp6 with a triple amino acid deletion (106–108,  $\Delta$ SGF) found in early VOCs (Alpha, Beta, and Gamma) conferred an increased ER-zipper activity. This  $\Delta$ SGF deletion partially overlaps with the 105–107  $\Delta$ LSG deletion in Omicron BA.1 nsp6 (Fig. 1A). We also noted that the ancestral sequence of nsp6 is maintained in the more pathogenic Delta VOC. These observations of different nsp6 sequences in VOCs motivated us to evaluate the evolution of nsp6 over the course of the pandemic.

### **Amino acid deletions in nsp6 are independently selected for human adaptation**

To understand SARS-CoV-2 nsp6's evolution, we examined 91,596 human SARS-CoV-2 whole-genome sequences across 19 variants and/or lineages obtained from the Global Initiative on Sharing Avian Influenza Data (GISAID) platform (Table S1). The nsp6  $\Delta$ SGF deletion is found in early variants, including Alpha, Beta, Gamma, and later Eta, Iota, and Lambda (21), reported in 95% of the sequences. In contrast, the triple amino acid deletion is not present in Delta, Epsilon, and Mu VOCs, where the ancestral nsp6 sequence is retained in 99.6% of these variants. The Omicron BA.1 lineage has the most diversity in the reported sequences, with 65%  $\Delta$ LSG, 8.5% ancestral nsp6, 0.1%  $\Delta$ SGF, and a surprisingly high number of sequences that would result in a frameshift in nsp6 (10.3% 5-nucleotide deletion; 4.3% 2-nucleotide deletion; 11.7% other nsp6 nucleotide changes,  $n = 4,976$ ). The 5- and 2-nucleotide deletions (~15% of BA.1 sequences) could be sequencing or assembly errors in the consensus genomes available on GISAID. Currently, it is unclear why these deletions are more prevalent in BA.1 genomes compared to other Omicron sublineages. Interestingly, the  $\Delta$ SGF nsp6 again becomes the dominant deletion type in all Omicron VOCs from BA.2 to XBB. This deletion is reported in 78% of BA.2, 83% in BA.2.75 sequences, and >97% in the more recent Omicron lineages. The accession numbers for the 91,596 genomes evaluated for this study are available upon request.



**FIG 1** Analysis of SARS-CoV-2 nsp6 reveals SGF/LSG deletions in 16 of 19 lineages. (A) Relative abundance of deletion of SGF and LSG sequences in the ancestral SARS-CoV-2 strain (Wuhan-Hu-1) and 19 subsequent lineages. The evaluated human SARS-CoV-2 whole-genome sequences are downloaded from the Global Initiative on Sharing Avian Influenza Data (GISAID) platform with a limit to a maximum of 5,000 sequences per variant except for Alpha and Delta, where more (Continued on next page)

**FIG 1 (Continued)**

sequences are included. Sequences are screened to be “high coverage” or “low coverage excluded,” “collection date complete,” and include those submitted from sites around the world. The prevalence of the mutation types in each variant/lineage is shown on the right, with colors from light blue to deep pink representing increased values. (B) Analysis using Outbreak.info documents the prevalence of mutations in the ORF1ab region across the 19 lineages (accessed on 6 January 2023). The nsp6 deletion region is highlighted in the red box. The mutation prevalence in each variant/lineage is indicated by the intensity of the purple color. (C) Analysis of the nucleotide sequence encoding LSGF in nsp6 from viruses isolated from 19 bats and 13 pangolins. The alignment of the LSGF region is shown in light blue. Synonymous mutations are highlighted in gray and non-synonymous in green (F to Y) and blue (F to L).

When extending our scope to evaluate all mutations in the ORF1ab region, we identified only one additional mutation retained consistently across all variants, ORF1b:P314L in nsp12 (Fig. 1B). This mutation was reported as a co-evolution with the spike D614G substitution (22, 23). Figure 1B also shows that Omicron variants encode multiple mutations in ORF1ab compared to earlier lineages. The relative contribution of individual or combined mutations in ORF1ab to the attenuation of Omicron variants remains to be determined.


Taken together, the dominance of the nsp6  $\Delta$ SGF/ $\Delta$ LSG deletion in multiple, independent VOCs (Fig. 1; Table S1) suggests that SARS-CoV-2 nsp6  $\Delta$ SGF/ $\Delta$ LSG deletions may have been selected for viral adaptation in humans.

**Nsp6 deletion mutants are not present in bat or pangolin coronaviruses**

To determine if the  $\Delta$ SGF/ $\Delta$ LSG deletions were present in nsp6 of other SARS-like viruses, we compared whole-genome sequences of bats and pangolins coronaviruses to the SARS-CoV-2 reference genome (NCBI GenBank accession number [NC\\_045512.2](#)), including the bat coronaviruses that share the highest nucleotide identity to human SARS-CoV-2 (Yunnan/RaTG13/2013 and Laos/BANAL/2020) (24–27) (Fig. 1C). We show that these bat and pangolin coronaviruses do not carry any deletion in this region of nsp6; actually, they are quite similar to the ancestral SARS-CoV-2, with the same codons for the 106S and 107G amino acids. Synonymous mutations at the 108<sup>th</sup> position (F) are observed (indicated in gray color, Fig. 1C), resulting in 11 of bat (58%) and 7 of pangolin (54%) CoV genomes with identical amino acids to that of ancestral SARS-CoV-2 nsp6. This consistency of nsp6 in bat and pangolin coronaviruses and SARS-CoV-2 further supports that the nsp6 deletion found in VOCs may represent an adaptation of SARS-CoV-2 to the human host.

**Predicting the topology of SARS-CoV-2 nsp6**

The deletion mutations in nsp6 and its association with the attenuated Omicron phenotype raise several questions. Exactly where is this deletion in the overall structure of nsp6 and what function(s) are altered in the mutant? To date, there is no structural information available for SARS-CoV-2 nsp6. Previous studies using the model murine coronavirus, mouse hepatitis virus (MHV), revealed that nsp6 had both its termini exposed on the cytoplasmic face of the ER membrane while spanning the lipid bilayer six times (14, 28). Here, we analyzed the structure of nsp6 in SARS-CoV-2 with comparison to other human coronaviruses using multiple programs: HMMTOP, SOSUI, Phobius, and TMHMM (29–32) (Fig. 2A and C). Our analysis agreed with the six transmembrane (TM) topologies and identified a likely hydrophobic domain (HD) as membrane-associated in SARS-CoV-2 nsp6. All programs revealed similar helical structures for the first four and the final TM domains. The deletion mutation ( $\Delta$ SGF/ $\Delta$ LSG) found in VOCs is located within the predicted second and longest luminal loop (Fig. 2B). Further comparison of SARS-CoV-2 nsp6 with six other human coronaviruses and MHV suggested that the number of predicted TM domains and the likely-HD is highly conserved (Fig. 2C). Interestingly, the non-SARS viruses had an overall high amino acid identity and conserved sequence length in the C-terminal domain, while the N-terminal and middle domains showed lower identity in amino acid sequences (Fig. 2D), suggesting that the C-terminal domain may play an important role in nsp6 function.



**FIG 2** Analysis of predicted hydrophobic and transmembrane domains of nsp6 from seven human coronaviruses and mouse hepatitis virus (MHV). (A) Sequences of SARS-CoV-2 nsp6 (NCBI accession number [YP\\_009725302.1](https://www.ncbi.nlm.nih.gov/nuclot/YP_009725302.1)) are analyzed for the prediction of the hydrophobic domain (HD) and transmembrane domain (TM) using HMMTOP, SOSUI, Phobius, and TMHMM programs shown in gray boxes. The residues predicted as likely HD or TM by three of the four programs are indicated in the yellow boxed areas. Red and green fonts represent hydrophobic and hydrophilic amino acids, respectively. (B) The diagram illustrates a proposed model of the topology of SARS-CoV-2 nsp6. The asterisk indicates the position of LSGF residues located in between TM3 and TM4. (C) Alignment of predicted topologies for six beta coronaviruses (SARS-CoV-2, SARS-CoV [[NP\\_828864.1](https://www.ncbi.nlm.nih.gov/nuclot/NP_828864.1)], MERS-CoV [[YP\\_009047218.1](https://www.ncbi.nlm.nih.gov/nuclot/YP_009047218.1)], HCoV-OC43 [[YP\\_009924324.1](https://www.ncbi.nlm.nih.gov/nuclot/YP_009924324.1)], HCoV-HKU1 [[YP\\_009944274.1](https://www.ncbi.nlm.nih.gov/nuclot/YP_009944274.1)], and MHV-A59 [[YP\\_009915678.1](https://www.ncbi.nlm.nih.gov/nuclot/YP_009915678.1)]) and two alpha coronaviruses (HCoV-NL63 [[YP\\_010229075.1](https://www.ncbi.nlm.nih.gov/nuclot/YP_010229075.1)] and HCoV-229E [[NP\\_073549.1](https://www.ncbi.nlm.nih.gov/nuclot/NP_073549.1)]). The topologies were predicted by three of the four programs as described previously. (D) Amino acid identity in the N-terminal, middle, and C-terminal domains of nsp6 reveals more conservation in the C-terminal domain.

Targeting nsp6 may be productive for limiting the replication of potentially emerging coronaviruses. Previous studies point to the essential role of nsp6 in the assembly and ER



tethering of viral replication complexes and in the recruitment of lipid droplets to DMVs, whose association is thought to be essential for the maintenance and proliferation of viral replication organelles (14, 16, 21). Further studies are needed to identify host factors interacting with nsp6 during membrane zippering and DMV formation as well as to elucidate the role of the protein C-terminal domain and luminal and cytoplasmic loops in viral replication.

## ACKNOWLEDGMENTS

This work is supported by the NIH R01 AI59945 (to S.C.B. and M.S.) and the Walder Foundation's Chicago Coronavirus Assessment Network (Chicago CAN) initiative (to S.C.B.).

S.F. and A.O. collected data, performed analysis, and drafted the manuscript. All authors critically revised the manuscript and provided intellectual contributions.

## AUTHOR AFFILIATIONS

<sup>1</sup>Department of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Chicago, Illinois, USA

<sup>2</sup>Department of Biochemistry and Cell Biology, Boston University Chobanian and Avedisian School of Medicine, Boston, Massachusetts, USA

<sup>3</sup>National Emerging Infectious Diseases Laboratories (NEIDL), Boston, Massachusetts, USA

## AUTHOR ORCIDS

Shuchen Feng  <http://orcid.org/0000-0002-8878-8036>

Mohsan Saeed  <http://orcid.org/0000-0001-8505-7054>

Susan C. Baker  <http://orcid.org/0000-0001-6485-8143>

## FUNDING

Funder	Grant(s)	Author(s)
HHS   National Institutes of Health (NIH)	AI59945	Susan C Baker Mohsan Saeed
The Walder Foundation	Chicago Coronavirus Assessment Network (Chicago CAN)	Susan C Baker

## AUTHOR CONTRIBUTIONS

Shuchen Feng, Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing – review and editing, Visualization, Writing – original draft, Writing – review and editing | Amornrat O'Brien, Data curation, Formal analysis, Methodology, Writing – review and editing, Visualization, Writing – original draft | Da-Yuan Chen, Writing – review and editing | Mohsan Saeed, Writing – original draft, Writing – review and editing | Susan C. Baker, Funding acquisition, Writing – review and editing

## ADDITIONAL FILES

The following material is available [online](#).

### Supplemental Material

**Table S1 (mBio00688-23-S0001.eps).** Sequence types of the nsp6 deletion region across 19 SARS-CoV-2 variants and/or lineages.

## REFERENCES

- Centers for Disease Control and Prevention. 2023. SARS-CoV-2 variant classifications and definitions. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-classifications.html>
- Harvey WT, Carabelli AM, Jackson B, Gupta RK, Thomson EC, Harrison EM, Ludden C, Reeve R, Rambaut A, Peacock SJ, Robertson DL. 2021.

- SARS-CoV-2 variants, spike mutations and immune escape. *Nat Rev Microbiol* 19:409–424. <https://doi.org/10.1038/s41579-021-00573-0>
3. Jackson CB, Farzan M, Chen B, Choe H. 2022. Mechanisms of SARS-CoV-2 entry into cells. *Nat Rev Mol Cell Biol* 23:3–20. <https://doi.org/10.1038/s41580-021-00418-x>
  4. Syed AM, Taha TY, Tabata T, Chen IP, Ciling A, Khalid MM, Sreekumar B, Chen P-Y, Hayashi JM, Soczek KM, Ott M, Doudna JA. 2021. Rapid assessment of SARS-CoV-2-evolved variants using virus-like particles. *Science* 374:1626–1632. <https://doi.org/10.1126/science.abc16184>
  5. World Health Organization (WHO). 2021 Classification of Omicron (B.1.1.529): SARS-CoV-2 variant of concern. Available from: [https://www.who.int/news/item/26-11-2021-classification-of-omicron-\(b.1.1.529\)-sars-cov-2-variant-of-concern](https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov-2-variant-of-concern)
  6. Halfmann PJ, Iida S, Iwatsuki-Horimoto K, Maemura T, Kiso M, Scheaffer SM, Darling TL, Joshi A, Loeber S, Singh G, Foster SL, Ying B, Case JB, Chong Z, Whitener B, Moliva J, Floyd K, Ujie M, Nakajima N, Ito M, Wright R, Uraki R, Warang P, Gagne M, Li R, Sakai-Tagawa Y, Liu Y, Larson D, Osorio JE, Hernandez-Ortiz JP, Henry AR, Ciuderis K, Florek KR, Patel M, Odle A, Wong L-Y, Bateman AC, Wang Z, Edara V-V, Chong Z, Franks J, Jeevan T, Fabrizio T, DeBeauchamp J, Kercher L, Seiler P, Gonzalez-Reiche AS, Sordillo EM, Chang LA, van Bakel H, Simon V, Consortium Mount Sinai Pathogen Surveillance (PSP) study group, Douek DC, Sullivan NJ, Thackray LB, Ueki H, Yamayoshi S, Imai M, Perlman S, Webby RJ, Seder RA, Suthar MS, García-Sastre A, Schotsaert M, Suzuki T, Boon ACM, Diamond MS, Kawaoka Y. 2022. SARS-CoV-2 Omicron virus causes attenuated disease in mice and hamsters. *Nature* 603:687–692. <https://doi.org/10.1038/s41586-022-04441-6>
  7. van Doremalen N, Singh M, Saturday TA, Yinda CK, Perez-Perez L, Bohler WF, Weishampel ZA, Lewis M, Schulz JE, Williamson BN, Meade-White K, Gallogly S, Okumura A, Feldmann F, Lovaglio J, Hanley PW, Shaia C, Feldmann H, de Wit E, Munster VJ, Rosenke K. 2022. SARS-CoV-2 Omicron BA.1 and BA.2 are attenuated in rhesus macaques as compared to Delta. *Sci Adv* 8:eade1860. <https://doi.org/10.1126/sciadv.ade1860>
  8. Nyberg T, Ferguson NM, Nash SG, Webster HH, Flaxman S, Andrews N, Hinsley W, Bernal JL, Kall M, Bhatt S, Blomquist P, Zaidi A, Volz E, Aziz NA, Harman K, Funk S, Abbott S, COVID-19 Genomics UK (COG-UK) consortium, Hope R, Charlett A, Chand M, Ghani AC, Seaman SR, Dabrera G, De Angelis D, Presanis AM, Thelwall S. 2022. Comparative analysis of the risks of hospitalisation and death associated with SARS-CoV-2 Omicron (B.1.1.529) and Delta (B.1.617.2) variants in England: a cohort study. *Lancet* 399:1303–1312. [https://doi.org/10.1016/S0140-6736\(22\)00462-7](https://doi.org/10.1016/S0140-6736(22)00462-7)
  9. Lewnard JA, Hong VX, Patel MM, Kahn R, Lipsitch M, Tartof SY. 2022. Clinical outcomes associated with SARS-CoV-2 Omicron (B.1.1.529) variant and BA.1/BA.1.1 or BA.2 subvariant infection in Southern California. *Nat Med* 28:1933–1943. <https://doi.org/10.1038/s41591-022-01887-z>
  10. Wong JY, Cheung JK, Lin Y, Bond HS, Lau EHY, Ip DKM, Cowling BJ, Wu P. 2023. Intrinsic and effective severity of COVID-19 cases infected with the ancestral strain and Omicron BA.2 variant in Hong Kong. *J Infect Dis*:jia236. <https://doi.org/10.1093/infdis/jia236>
  11. Chen D-Y, Chin CV, Kenney D, Tavares AH, Khan N, Conway HL, Liu G, Choudhary MC, Gertje HP, O'Connell AK, Adams S, Kotton DN, Herrmann A, Ensser A, Connor JH, Bosmann M, Li JZ, Gack MU, Baker SC, Kirchdoerfer RN, Kataria Y, Crossland NA, Douam F, Saeed M. 2023. Spike and Nsp6 are key determinants of SARS-CoV-2 Omicron BA.1 attenuation. *Nature* 615:143–150. <https://doi.org/10.1038/s41586-023-05697-2>
  12. Taha TY, Chen IP, Hayashi JM, Tabata T, Walcott K, Kimmerly GR, Syed AM, Ciling A, Suryawanshi RK, Martin HS, Bach BH, Tsou C-L, Montano M, Khalid MM, Sreekumar BK, Renuka Kumar G, Wyman S, Doudna JA, Ott M. 2023. Rapid assembly of SARS-CoV-2 genomes reveals attenuation of the Omicron BA.1 variant through NSP6. *Nat Commun* 14:2308. <https://doi.org/10.1038/s41467-023-37787-0>
  13. Knoop K, Kikkert M, Worm S van den, Zevenhoven-Dobbe JC, van der Meer Y, Koster AJ, Mommaas AM, Snijder EJ. 2008. SARS-Coronavirus replication is supported by a reticulovesicular network of modified endoplasmic reticulum. *PLoS Biol* 6:e226. <https://doi.org/10.1371/journal.pbio.0060226>
  14. Angelini MM, Akhlaghpour M, Neuman BW, Buchmeier MJ. 2013. Severe acute respiratory syndrome Coronavirus nonstructural proteins 3, 4, and 6 induce double-membrane vesicles. *mBio* 4:e00524-13. <https://doi.org/10.1128/mBio.00524-13>
  15. Mihelc EM, Baker SC, Lanman JK. 2021. Coronavirus infection induces progressive restructuring of the endoplasmic reticulum involving the formation and degradation of double membrane vesicles. *Virology* 556:9–22. <https://doi.org/10.1016/j.virol.2020.12.007>
  16. Oostra M, Hagemeijer MC, van Gent M, Bekker CPJ, te Lintelo EG, Rottier PJM, de Haan CAM. 2008. Topology and membrane anchoring of the Coronavirus replication complex: not all hydrophobic domains of Nsp3 and Nsp6 are membrane spanning. *J Virol* 82:12392–12405. <https://doi.org/10.1128/JVI.01219-08>
  17. Gosert R, Kanjanahaluethai A, Egger D, Bienz K, Baker SC. 2002. RNA replication of mouse hepatitis virus takes place at double-membrane vesicles. *J Virol* 76:3697–3708. <https://doi.org/10.1128/jvi.76.8.3697-3708.2002>
  18. Snijder EJ, Limpens R, de Wilde AH, de Jong AWM, Zevenhoven-Dobbe JC, Maier HJ, Faas F, Koster AJ, Bárcena M. 2020. A unifying structural and functional model of the Coronavirus replication organelle: tracking down RNA synthesis. *PLoS Biol* 18:e3000715. <https://doi.org/10.1371/journal.pbio.3000715>
  19. Cortese M, Lee J-Y, Cerikan B, Neufeldt CJ, Oorschot VMJ, Köhrer S, Hennies J, Schieber NL, Ronchi P, Mizzon G, Romero-Brey I, Santarella-Mellwig R, Schorb M, Boermeier M, Mocaer K, Beckwith MS, Templin RM, Gross V, Pape C, Tischer C, Frankish J, Horvat NK, Laketa V, Stanifer M, Boulant S, Ruggieri A, Chatel-Chaix L, Schwab Y, Bartschlag R. 2020. Integrative imaging reveals SARS-CoV-2-induced reshaping of subcellular morphologies. *Cell Host Microbe* 28:853–866. <https://doi.org/10.1016/j.chom.2020.11.003>
  20. Twu W-I, Lee J-Y, Kim H, Prasad V, Cerikan B, Haselmann U, Tabata K, Bartschlag R. 2021. Contribution of autophagy machinery factors to HCV and SARS-CoV-2 replication organelle formation. *Cell Rep* 37:110049. <https://doi.org/10.1016/j.celrep.2021.110049>
  21. Ricciardi S, Guarino AM, Giaquinto L, Polishchuk EV, Santoro M, Di Tullio G, Wilson C, Panariello F, Soares VC, Dias SSG, Santos JC, Souza TML, Fusco G, Viscardi M, Brandi S, Bozza PT, Polishchuk RS, Venditti R, De Matteis MA. 2022. The role of Nsp6 in the biogenesis of the SARS-CoV-2 replication organelle. *Nature* 606:761–768. <https://doi.org/10.1038/s41586-022-04835-6>
  22. Biswas SK, Mudi SR. 2020. Spike protein D614G and RdRp P323L: the SARS-CoV-2 mutations associated with severity of COVID-19. *Genomics Inform* 18:e44. <https://doi.org/10.5808/GI.2020.18.4.e44>
  23. Volz E, Hill V, McCrone JT, Price A, Jorgensen D, O'Toole Á, Southgate J, Johnson R, Jackson B, Nascimento FF, Rey SM, Nicholls SM, Colquhoun RM, da Silva Filipe A, Shepherd J, Pascall DJ, Shah R, Jesudason N, Li K, Jarrett R, Pacchiarini N, Bull M, Geidelberg L, Siveroni I, COG-UK Consortium, Goodfellow I, Loman NJ, Pybus OG, Robertson DL, Thomson EC, Rambaut A, Connor TR. 2021. Evaluating the effects of SARS-CoV-2 spike mutation D614G on transmissibility and pathogenicity. *Cell* 184:64–75. <https://doi.org/10.1016/j.cell.2020.11.020>
  24. Wacharapluesadee S, Tan CW, Maneerorn P, Duengkhae P, Zhu F, Joyinda Y, Kaewpom T, Chia WN, Ampoot W, Lim BL, Worachotsueptrakun K, Chen V-W, Sirichan N, Ruchisrisarod C, Rodpan A, Noradechanon K, Phaichana T, Jantarat N, Thongnumchaima B, Tu C, Cramer G, Stokes MM, Hemachudha T, Wang L-F. 2021. Evidence for SARS-CoV-2 related Coronaviruses circulating in bats and pangolins in Southeast Asia. *Nat Commun* 12:1430. <https://doi.org/10.1038/s41467-021-21768-2>
  25. Zhou H, Ji J, Chen X, Bi Y, Li J, Wang Q, Hu T, Song H, Zhao R, Chen Y, Cui M, Zhang Y, Hughes AC, Holmes EC, Shi W. 2021. Identification of novel bat coronaviruses sheds light on the evolutionary origins of SARS-CoV-2 and related viruses. *Cell* 184:4380–4391. <https://doi.org/10.1016/j.cell.2021.06.008>
  26. Temmam S, Vongphayloth K, Baquero E, Munier S, Bonomi M, Regnault B, Douangboubpha B, Karami Y, Chrétien D, Sanamxay D, Xayaphet V, Paphaphanh P, Lacoste V, Somlor S, Lakeomany K, Phommavanh N, Pérot P, Dehan O, Amara F, Donati F, Bigot T, Nilges M, Rey FA, van der Werf S, Brey PT, Eloit M. 2022. Bat Coronaviruses related to SARS-CoV-2 and infectious for human cells. *Nature* 607:E19. <https://doi.org/10.1038/s41586-022-05048-7>
  27. Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, Si H-R, Zhu Y, Li B, Huang C-L, Chen H-D, Chen J, Luo Y, Guo H, Jiang R-D, Liu M-Q, Chen Y, Shen X-R, Wang X, Zheng X-S, Zhao K, Chen Q-J, Deng F, Liu L-L, Yan B, Zhan F-X, Wang Y-Y, Xiao G-F, Shi Z-L. 2020. A pneumonia outbreak associated with a new Coronavirus of probable bat origin. *Nature* 579:270–273. <https://doi.org/10.1038/s41586-020-2012-7>

28. Baliji S, Cammer SA, Sobral B, Baker SC. 2009. Detection of nonstructural protein 6 in murine Coronavirus-infected cells and analysis of the transmembrane topology by using bioinformatics and molecular approaches. *J Virol* 83:6957–6962. <https://doi.org/10.1128/JVI.00254-09>
29. Tusnády GE, Simon I. 2001. The HMMTOP transmembrane topology prediction server. *Bioinformatics* 17:849–850. <https://doi.org/10.1093/bioinformatics/17.9.849>
30. Hirokawa T, Boon-Chieng S, Mitaku S. 1998. SOSUI: classification and secondary structure prediction system for membrane proteins. *Bioinformatics* 14:378–379. <https://doi.org/10.1093/bioinformatics/14.4.378>
31. Käll L, Krogh A, Sonnhammer ELL. 2004. A combined transmembrane topology and signal peptide prediction method. *J Mol Biol* 338:1027–1036. <https://doi.org/10.1016/j.jmb.2004.03.016>
32. Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* 305:567–580. <https://doi.org/10.1006/jmbi.2000.4315>