

Barriers and Facilitators to Delivering Inpatient Cardiac Rehabilitation: A Scoping Review

Marina Wasilewski^{1,3}, Abirami Vijayakumar¹, Zara Szigeti¹, Sahana Sathakaran², Kuan-Wen Wang², Adam Saporta¹, Sander L Hitzig¹

¹St. John's Rehab, Sunnybrook Research Institute, North York, Ontario, Canada; ²Department of Occupational Science & Occupational Therapy, University of Toronto, Toronto, Ontario, Canada; ³Rehabilitation Sciences Institute (RSI), University of Toronto, Toronto, Ontario, Canada

Correspondence: Marina Wasilewski, St. John's Rehab, Sunnybrook Research Institute, North York, Ontario, Canada, Email marina.wasilewski@sunnybrook.ca

Objective: The purpose of this scoping review was to summarize the literature on barriers and facilitators that influence the provision and uptake of inpatient cardiac rehabilitation (ICR).

Methods: A literature search was conducted using PsycINFO, MEDLINE, EMBASE, CINAHL and AgeLine. Studies were included if they were published in English after the year 2000 and focused on adults who were receiving some form of ICR (eg, exercise counselling and training, education for heart-healthy living). For studies meeting inclusion criteria, descriptive data on authors, year, study design, and intervention type were extracted.

Results: The literature search resulted in a total of 44,331 publications, of which 229 studies met inclusion criteria. ICR programs vary drastically and often focus on promoting physical exercises and patient education. Barriers and facilitators were categorized through patient, provider and system level factors. Individual characteristics and provider knowledge and efficacy were categorized as both barriers and facilitators to ICR delivery and uptake. Team functioning, lack of resources, program coordination, and inconsistencies in evaluation acted as key barriers to ICR delivery and uptake. Key facilitators that influence ICR implementation and engagement include accreditation and professional associations and patient and family-centred practices.

Conclusion: ICR programs can be highly effective at improving health outcomes for those living with CVDs. Our review identified several patient, provider, and system-level considerations that act as barriers and facilitators to ICR delivery and uptake. Future research should explore how to encourage health promotion knowledge amongst ICR staff and patients.

Keywords: inpatient, cardiovascular diseases, cardiac rehabilitation, early mobilization, scoping review

Introduction

Contributing to over 17.9 million deaths each year, cardiovascular disease (CVD) are the leading cause of death worldwide.¹ CVD, such as ischemic heart disease (IHD) and coronary artery disease (CAD), are a group of disorders of the heart and blood vessels resulting from an array of biological, behavioural, environmental, and social risk factors including unhealthy diets, physical inactivity, and substance abuse.¹ In the last thirty years, the global incidence of CVD has doubled, with nearly 528 million cases of CVD being reported across the world in 2019.² In the United States alone, the direct and indirect costs of CVD amounted to \$863 billion in 2010 and are estimated to rise by 22% to \$1044 billion by 2030.³ To reduce the incidence of CVD, primary and secondary prevention efforts should be intensified.

Cardiac rehabilitation (CR) is a comprehensive and evidence-based model of care for managing CVD that involves a combination of exercise, education on modifiable risk factors and lifestyle changes, and/or psychological and social support.^{4,5} CR programs are integral to limit the physiological and psychological effects of cardiac illness, reduce the risk of sudden death and re-hospitalization, control cardiac symptoms, contribute to secondary prevention, and enhance the psychosocial and vocational status of cardiac patients.^{4,5} Prior research has highlighted the importance of early intervention, such as inpatient CR (ICR) as it optimizes patient independence and autonomy through the promotion of regular physical movement, education, and medication adherence.⁶ ICR is specifically known to promote functional recovery,

psychosocial well-being, long-term survival, and improve overall quality of life in cardiac patients.⁶ ICR is particularly important as the knowledge and skills obtained in the hospital is known to help the patient reach the next level of activity as an outpatient and establish patterns of care at home.⁷

Although there is ample evidence supporting the effectiveness of ICR, these programs are still being underutilized worldwide with attendance rates of only 20%-50%.^{5,6} Barriers to enrollment in CR programs include but are not limited to distance, lack of time, accessibility, and prior work commitments.⁸⁻¹⁰ There are also home-based CR programs (HBCR) which are used to increase participation in patient populations. For instance, HBCR programs can enhance the accessibility of CR programs by removing travel-related barriers and accommodating those that are immunocompromised.¹¹

Further, little is known about ICR practice patterns and implementation considerations, such as what is working well and what can be improved within these programs.¹² To our knowledge, no review exists that summarizes the barriers and facilitators to delivering ICR programs, highlighting a need to explore and map the available evidence related to the implementation of ICRs. As such, we aimed to identify and categorize the various barriers and facilitators that influence the provision and uptake of ICR.

Methods

This scoping review follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews (PRISMA-ScR) checklist.¹³ The review was conducted according to the five stages described by the Arksey and O'Malley's methodological framework.¹⁴ This methodology was appropriate given our interest in the broad topic of ICR and the heterogeneous nature of the body of literature. This scoping reviews' protocol has been registered in the Open Science Framework (OSF) (10.17605/OSF.IO/XQE7R).

Stage I: Identifying the Research Question

The guiding research question for this scoping reviewer is: What are the barriers and facilitators that influence the use, provision, engagement and overall implementation, of ICR? We also aimed to describe the scope of ICR, including how these programs are developed, who implements these programs, and which outcome measures are used to quantify the success of ICR.

Stage II: Identifying Relevant Studies

A research librarian was consulted for the development of a comprehensive search strategy for the following databases: PsycINFO, Medline, Embase, CINAHL and AgeLine on October 28, 2020. The search strategy included terms such as "cardiovascular disease", "rehab", and "inpatient/in-hospital" (see [Appendix A](#) for Medline search strategy). There have been substantial changes in our understanding of cardiac conditions, structure, and organization of the healthcare system in the last two decades, leading to progression in ICR intervention strategies.⁶ Thus, this scoping review omitted literature published prior to the year 2000. Detailed inclusion and exclusion criteria are described below.

Stage III: Study Selection

Articles were included if they were in English and focused on adults who were receiving some form of ICR, such as physical exercise or nutritional counselling. Only primary research articles, regardless of study design, were included. We excluded review articles, unpublished dissertations, study protocols, and conference proceedings. Articles were also excluded if they focused on pediatric populations, primary prevention, or did not describe the hospital setting in sufficient detail.

Records identified in the database search were uploaded into Covidence systematic review software for screening. Three reviewers (K-W.W., S.S., and Z.S.) screened 10% of the titles and abstracts to establish inter-rater reliability ($k=0.837$, 96% agreement). Once reliability was established, the remaining titles and abstracts, as well as full-texts, were independently screened. If there was disagreement between the reviewers, an additional team member was consulted to reach consensus on the inclusion of the article.

Stage IV: Charting the Data

A data extraction chart was created to extract key information that would help us answer our research question, including study design and details (eg author, country), ICR program characteristics (eg interventions provided, staffing model), and relevant findings (eg benefits of ICR, barriers/facilitators to ICR implementation and utilization). Similar to full-text screening, the included articles were distributed across four reviewers (A.V, K-W.W., S.S., and Z.S) for independent extraction of the relevant information.

Stage V: Collating, Summarizing and Reporting Results

The present review employed a descriptive summary and a narrative analysis. Our descriptive summary highlighted details on the type and frequency of studies included, and the location where the study was conducted. We then categorized and narratively summarized findings into barriers and facilitators of ICR delivery and uptake.

Results

Our database search identified a total of 44,331 records in addition to one record identified through network sharing. We removed 8423 duplicates, leaving 35,909 unique records for title and abstract screening and 2099 were selected for full-text review. The most common reasons for exclusion were due to being an ineligible publication type (conference abstracts, letters, or editorials [$n = 766$]) or the CR program did not occur in an inpatient setting ($n = 732$). 48 of the full texts could not be located and were excluded. In total, 229 articles were included, as shown in [Figure 1](#).

Study Characteristics

Studies used a variety of designs and methodologies including randomized-case studies ($n=48$), observation-based methods ($n=32$), general pre- and post-test ($n=21$), surveys ($n=19$), case studies/reports ($n=10$), and qualitative methods ($n=8$). Most studies were conducted in Germany ($n=40$), the USA ($n=31$), Italy ($n=30$), and Japan ($n=27$). Studies focused on a variety of surgical conditions including coronary artery bypass graft surgery (CABG) ($n=34$) and valvular surgeries ($n=13$), as well as cardiac conditions including myocardial infarction (MI)/acute myocardial infarction (AMI) ($n=59$), CAD ($n=39$), and heart failure (HF) ($n=35$). Some studies also reported on comorbidities, with the most common being hypertension ($n=50$) and diabetes ($n=50$). Out of the 228 included articles, 157 included samples of males and females,^{15–169} 25 included males only,^{170–194} and four included a female-only sample.^{195–198} Studies were conducted in various acute settings including acute hospitals ($n=125$), designated inpatient rehabilitation hospitals ($n=51$), and rehabilitation centres ($n=23$). Full study details and characteristics can be found in [Appendix B](#). Below, we report specifically on our synthesis of the barriers and facilitators to ICR delivery and uptake from the included studies.

Barriers to ICR Delivery and Uptake

Patient Level

Individual Characteristics: Studies reported factors, behaviours, and attitudes that impaired patients' ability to participate in and receive ICR. These factors included a lack of knowledge, interest, and motivation to participate in ICR, and pain and anxiety when participating in ICR exercises and therapies.^{70,71,73,75,87,113,115,119,134,142,143,148,166–168,190,199–205} Additionally, sociodemographic conditions impacted patients' ability to participate, with some studies noting that older adults and those coming from lower socioeconomic status were less likely to participate in ICR.^{28,39,166,200,206}

Provider Level

Provider Knowledge and Efficacy: Studies indicate that providers lack expertise and interest in ICR^{131,152,166,206} and have limited knowledge about the types of ICR interventions that exist and their importance and benefit.^{39,58,70,80,84,99,108,113,130,142,164,166,168,199,201,206–211} Further, many of them have difficulty forming new concepts and skills to implement novel interventions⁸⁴ and are challenged by time/scheduling restrains as well as narrow scopes of practice when it comes to implementing ICR.^{51,78,108,124,142,201,207,209,210}




Figure 1 Flowchart.

Notes: Adapted from Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. doi: 10.1136/bmj.n71.

Team Functioning: The absence of strong social networks impacted program delivery. Some studies described multidisciplinary teams that lacked interdisciplinary communication, leadership, and administrative support.^{39,70,108,130,142,164,166,199,201,207,209–211}

System Level

Financial Costs and Resource Limitations: Broadly speaking, inadequate funding and resources both negatively influenced the ability to access and deliver ICR.^{39,70,80,99,108,130,142,164,166,199,201,207,209–211} Vincent et al¹⁴⁵ and Schweikert et al¹²⁸ reported that it cost their institutions \$4709 USD and €3388 EUR for total therapy charges per patient, respectively. Costs are significantly higher for those who require ICR in comparison to outpatient or community-based rehabilitation.^{128,145} Many of these charges are covered by insurance and national health coverage, however not all recipients of ICR are insured or have insurance that covers ICR.²¹⁰ Therefore, this creates a gap in access to ICR programs. Namely, those who have the financial means may have greater access to ICR and as a result, those who do not

have monetary access are forced to miss the opportunity to participate in such programs. Moreover, Weiser et al²¹² reported that it costs Swiss health insurance companies over \$462 million CHF each year to cover the costs of ICR. Out-of-pocket costs present significant barriers for the uninsured, especially for those who access ICR programs multiple times a week. Overall, costs impact participation in ICR programs.

Poorly Coordinated Processes: Referrals for ICR were inconsistent across the system and more likely to be provided when there was infrastructure that allowed for easy communication and collaboration (eg electronic medical record sharing between discharging and receiving facilities).¹⁸⁰ A lack of referrals and poor cross-regional collaborations (eg not having healthcare alliances across the region) acted as a barrier to patient participation in ICR. Acute care centres offering very early ICR did not always offer referrals to continue CR in an outpatient or tertiary setting. Moreover, not all facilities or clinicians offered referrals for patients to participate in ICR entirely. This often occurred when CR facilities and programs were limited and inaccessible, such as in rural regions.^{39,70,80,99,108,130,142,164,166,199,201,207,209–211}

Inconsistency in Evaluation: There was inconsistency in the literature related to how patient and provider feedback was obtained and incorporated to improve the quality of delivery and program design.^{84,96,166,207} Some studies described the use of validated and systemic processes to evaluate the ICRs, including patient surveys of changes in cardiac health outcomes, illness beliefs, and expectations.^{84,96} However, other research relied on qualitative interviews as a form of feedback to improve program use. For example, patients provided frontline clinicians with suggestions to improve the quality of care patient-centeredness of future programmes, particularly as it relates to consultation, accurate information, and equal access to treatment options.^{84,166} Other research reported using seminars or medical and nursing grand rounds as a means to gain feedback, information, and support from care providers to continuously improve patient outcomes.²⁰⁷

Facilitators of ICR Delivery and Uptake

Patient Level

Individual Characteristics: A facilitator for patient participation included recognition of their illness stage and the severity of their condition. This often occurred in tandem with a belief in the efficacy and importance of ICR, as those with more serious illnesses were more likely to endorse and participate in ICR.²⁰⁰ Prior knowledge and habits, including prior experiences with leading a healthy lifestyle, also facilitated patient participation and long-term benefits in ICR.^{51,52,58,70,71,73,77,78,92,115,124,142,143,153,157,158,165,167,168,174,191,200,201,203}

Provider Level

Provider Knowledge and Efficacy: Much like the absence of provider knowledge and efficacy can act as a barrier, its presence functions as a facilitator. When providers believed in the value and benefit of ICR programs, they were more likely to offer ICR and refer patients to continue with CR in an outpatient setting.^{51,78,108,124,142,201,206,207,209,210} ICR delivery was facilitated by providers who were confident in their capabilities and keenly aware of patients' needs.^{51,78,108,124,142,201,207,209,210} For example, physiotherapists (PTs) perceived themselves as trusted clinicians in the context of exercise-based CR, with patient relying on their support and motivation during rehabilitation.¹⁶⁵ Similarly, nurses acknowledged their role in supporting the day-to-day care of patients compared to other medical professionals and recognized their influence on promoting health-related behaviour changed.^{26,87} Notably, several studies indicated that provider knowledge and efficacy can be promoted by providing clinicians with appropriate supports such as education modules^{26,84,118,152,159,166,188,193} and well as support and communication from organizational leadership.²⁰⁷

Accreditation and Professional Associations: In one study, clinicians were more likely to offer and facilitate the use of ICR if they were affiliated with professional associations, such as the American Association of Cardiovascular and Pulmonary Rehabilitation (AACVPR).¹⁴²

Patient- and Family-Centered Practices: Building a strong rapport between clinicians, patients and families facilitated engagement in ICR and promoted its long-term benefits.^{49,92,167,168} When families are included in care and have the opportunity to gain knowledge of cardiac conditions, risk factors, and general healthy living habits, patients report more long-term positive health outcomes.^{96,114,145,158,162,163,179} When clinicians recognized patients' needs and resources, such as whether they were supported by family caregivers or whether they had healthy food options that were geographically accessible, patients were more likely to participate in ICR. Incorporating patients' needs and resources allowed for ICR

to be more tailored, relevant, and applicable and led to more long-term success, particularly with increasing independence at home.^{39,46,70,75,77,87,96,115,116,119,142,149,165–167,169,201}

System Level

Interestingly, no studies reported on facilitators at the system level.

Discussion

The goal of this scoping review was to provide a comprehensive synthesis of the factors that act as barriers and facilitators to ICR delivery and uptake. In total, we included 228 eligible articles. Our review identified several patient, provider, and system-level considerations that act as barriers and facilitators to ICR delivery and uptake.

Knowledge and Attitudes Towards ICR

Knowledge of ICR and a belief in the importance and necessity of the intervention were highlighted as key factors that influence the provision and uptake of ICR. For patients, knowledge and attitudes emerged as having the potential to both challenge *or* facilitate the uptake of ICR. Namely, when patients were aware of and convinced of the benefits of ICR, they were more likely to participate in the intervention. Conversely, when patients lacked knowledge, interest or motivation to participate in ICR, uptake was lower. Patients' lack of knowledge of what ICR entails may lead them to think they do not need the therapeutic benefits or attribute the content of ICR as being synonymous with physical exercise. Our review, as well as prior research, have highlighted that patients often have negative views towards physical exercise and hold a belief that they can undertake CR independently.^{164,213}

Similarly at the provider level, beliefs and attitudes towards ICR appeared to influence delivery, with poor knowledge and lack of expertise limiting providers' provision of ICR and strong knowledge and efficacy with ICR promoting its application. In a survey of clinicians' attitudes towards ICR, findings revealed that their belief in the efficacy of ICR was highly influenced by whether they value lifestyle modification as an effective means for behaviour change, rather than just as a mandatory way to change health behaviour.²¹⁴ In addition to personal values, a review of the literature suggests that a lack of clinician knowledge of ICR, such as not understanding who would most benefit from ICR or who should be referred, affects whether patients attend ICR.²¹⁵ Individual attitudes of clinicians can influence patients' own views on CR and can even promote healthy behaviours amongst patients.²¹⁶ In order to increase ICR participation, further efforts are needed to educate healthcare providers, healthcare systems, patients, and their families about the benefits of ICR, including efforts to change the perception that ICR is forced upon individuals or is less important than pharmacological or surgical therapy.²¹⁷ Future research should explore how to promote health promotion knowledge amongst ICR staff and patients and knowledge of behaviour change techniques amongst ICR staff.

Patient Characteristics Influence ICR Uptake

Our review begins to elucidate that patient characteristics have an important influence on ICR participation and uptake—particularly sex and gender. Out of 228 articles included in this scoping review, only 13 had more female participants than males,^{43,50,61,81,94,110,114,131,155,195–197,218} with some studies consisting solely of males (n=25). While this sex disparity might be due to the higher incidence of CVD among males,²¹⁹ it remains concerning that the female population is under-represented in these studies given that females have a higher mortality and hospitalization rates and experience poorer prognosis after a cardiac event, including future pregnancy complications.²¹⁹ Women have also been reported to have poorer participation rates in and referrals to CR programs,²²⁰ illustrating a need to understand how to optimize ICR programs to mitigate low attendance and compliance in this population. Ways to improve women's CR attendance may include exercise regimes (ie, yoga) and can be used to increase participation and engagement.¹⁹⁸ Future research that evaluates ICR programming should consider the factors associated with women's health, and future ICR programs should consider how to minimize non-adherence when catering interventions for female participants.

ICR is Highly Challenged by System-Level Issues

A notable finding from this review was that that system-level issues acted exclusively as barriers, with no system-level facilitators identified from the included studies. These issues included financial and resource limitations, inconsistent patient evaluations, and poorly coordinated processes across the continuum of care. A key issue was the lack of referrals to both ICR itself and referrals to continuing CR after ICR discharge. When patients receive long-term follow-up for at least 12 months following ICR discharge, they are more likely to sustain the benefits gained during their stay in ICR.^{116,130,148,153,190} Despite the recognized importance of long-term follow up, few ICR programs in the present review explicitly described a clear follow-up plan in the discharge planning process. This echoes the broader body of literature on CR which has identified a lack of referrals as a recurring issue that challenges the provision of CR.^{221,222} As has been demonstrated by Shanmugasagaram et al, these system-level issues can have a “trickle down” effect on patients in as far as a lack of physician referral for CR can generate patient perceptions that CR is not needed. As was discussed earlier in our paper, patients’ knowledge and attitudes towards ICR have an important impact on their participation in this intervention. Thus, addressing provider-level knowledge and attitudes as well as system-level challenges around integrated referrals can both have an ameliorative effect on ICR uptake at the patient level.

Need for Standardized Reporting of ICR Research

Our scoping review revealed that some confusion exists around the conceptualization of ICR. Many studies defined ICR as a part of acute care while others defined ICR as a rehabilitation period where patients stay in a dedicated institution, further illustrating the lack of uniformity in ICR delivery, goals, content, etc. These variable ways of classifying ICR makes summarizing the evidence challenging since the programs – though all referred to as ICR – entail different goals, staffing models, and resources depending on their settings. Moreover, the variability in classifying ICR means that the barriers and facilitators that influence ICR may differ depending on how it is contextualized. For example, while distance and transportation is often cited by patients as barriers to cardiac intervention participation,²²³ these barriers are not applicable to ICR as patients are staying in the institution during the rehabilitation period. To this end, standardization of design and reporting of ICR studies would better-enable evidence synthesis and comparison across existing and future studies.

Strengths and Limitations

Strengths of this scoping review include the rigorous development of search strategies tailored for each database and consultation with an academic librarian to identify proper keywords for each database. The search strategy was also validated with a test set which includes a list of relevant studies the reviewers identified a priori. The agreement rate among the reviewers was well above 80%, indicating a high level of accuracy in the study selection process. This review does have limitations, including the exclusion of non-English literature and studies that did not clearly state the care setting. Secondly, this scoping review did not undertake a bias and quality assessment. Given that this review’s focus was to synthesize the nature of ICR and the breadth of factors that influence implementation, critical appraisals and bias assessments were deemed “not applicable” by the review team, a distinction that is supported by current PRISMA-ScR reporting guidelines.

Conclusion

ICR programs can be extremely effective at improving health outcomes for those living with CVDs. This review identified several patient, provider, and system-level considerations that act as barriers and facilitators to ICR delivery and uptake. Future research should explore how to enhance health promotion knowledge amongst ICR staff and patients and knowledge of behaviour change techniques amongst ICR staff.

Acknowledgments

We wish to Mr. Henry Lam, academic librarian at Sunnybrook Health Sciences Centre, for his assistance with the literature search.

Funding

We wish to acknowledge funding support from the Malcolm Moffat Research Fund at St. John's Rehab Sunnybrook.

Disclosure

All authors declare no conflicts of interest in this work.

References

1. World Health Organization. Cardiovascular diseases. Available from: https://www.who.int/health-topics/cardiovascular-diseases/#tab=tab_1. Accessed July 27, 2023.
2. Roth GA, Mensah GA, Johnson CO, et al. Global burden of cardiovascular diseases and risk factors, 1990–2019: update from the GBD 2019 study. *J Am Coll Cardiol*. 2020;76(25):2982–3021. doi:10.1016/j.jacc.2020.11.010
3. Bloom DE, Cafiero ET, Jané-Llopis E, et al. *The Global Economic Burden of Non-Communicable Diseases*. Geneva: World Economic Forum; 2011.
4. Fernandes AC, McIntyre T, Coelho R, Prata J, Maciel MJ. Brief psychological intervention in Phase I of cardiac rehabilitation after acute coronary syndrome. *Port J Cardiol*. 2017;36(9):641–649. doi:10.1016/j.repc.2017.01.005
5. Kim JS, Kim GS, Kang SM, Chu SH. Symptom experience as a predictor of cardiac rehabilitation education programme attendance after percutaneous coronary intervention: a prospective questionnaire survey. *Eur J Cardiovasc Nurs*. 2021;20(3):183–191. doi:10.1177/1474515120940534
6. Mampuya WM. Cardiac rehabilitation past, present and future: an overview. *Cardiovasc Diagn Ther*. 2012;2(1):38–49. doi:10.3978/j.issn.2223-3652.2012.01.02
7. Rion JHK, Donald D. The walk to save: benefits of inpatient cardiac rehabilitation. *Medsurg Nurs*. 2016;25(3):159–162.
8. Winnige P, Filakova K, Hnatiak J, et al. Validity and reliability of the cardiac rehabilitation barriers scale in the Czech Republic (CRBS-CZE): determination of key barriers in east-central Europe. *Int J Environ Res Public Health*. 2021;18(24):13113. doi:10.3390/ijerph182413113
9. Antoniou V, Pasiak K, Loukidis N, et al. Translation, cross-cultural adaptation and psychometric validation of the Greek version of the cardiac rehabilitation barriers scale (CRBS-GR): what are the barriers in South-East Europe? *Int J Environ Res Public Health*. 2023;20(5):4064. doi:10.3390/ijerph20054064
10. Neubeck L, Freedman SB, Clark AM, Briffa T, Bauman A, Redfern J. Participating in cardiac rehabilitation: a systematic review and meta-synthesis of qualitative data. *Eur J Prev Cardiol*. 2012;19(3):494–503. doi:10.1177/1741826711409326
11. Nkonde-Price C, Reynolds K, Najem M, et al. Comparison of home-based vs center-based cardiac rehabilitation in hospitalization, medication adherence, and risk factor control among patients with cardiovascular disease. *JAMA Netw Open*. 2022;5(8):e2228720–e2228720. doi:10.1001/jamanetworkopen.2022.28720
12. Pack QR, Priya A, Lagu T, et al. Cardiac rehabilitation utilization during an acute cardiac hospitalization: a national sample. *J Cardiopulm Rehabil Prev*. 2019;39(1):19–26. doi:10.1097/hcr.0000000000000374
13. Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med*. 2018;169(7):467–473. doi:10.7326/M18-0850
14. Arksey H, O'malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol*. 2005;8(1):19–32. doi:10.1080/1364557032000119616
15. Albert NM, Buchsbaum R, Li J. Randomized study of the effect of video education on heart failure healthcare utilization, symptoms, and self-care behaviors. *Patient Educ Couns*. 2007;69(1–3):129–139. doi:10.1016/j.pec.2007.08.007
16. Altenhoner T, Baczkiewicz C, Weishaar H, Kutschmann M. Inequalities in therapeutic treatment during cardiac inpatient rehabilitation in Germany. *Int J Public Health*. 2012;57(1):175–184. doi:10.1007/s00038-011-0298-9
17. Ambrosetti M, Griffo R, Tramarin R, et al. Prevalence and 1-year prognosis of transient heart failure following coronary revascularization. *Intern Emerg Med*. 2014;9(6):641–647.
18. Ambrosetti M, Mariani P. Metabolic syndrome and related dietary intervention among patients with coronary and peripheral arterial disease attending cardiovascular rehabilitation programs. *Monaldi Arch Chest Dis*. 2007;68(4):227–230.
19. Ambrosetti M, Salerno M, Zambelli M, Mastropasqua F, Tramarin R, Pedretti RFE. Deep vein thrombosis among patients entering cardiac rehabilitation after coronary artery bypass surgery. *Chest*. 2004;125(1):191–196. doi:10.1378/chest.125.1.191
20. Anchah L, Hassali MA, Han Lim MS, et al. Health related quality of life assessment in acute coronary syndrome patients: the effectiveness of early phase I cardiac rehabilitation. *Health Qual Life Outcomes*. 2017;15:1–14. doi:10.1186/s12955-016-0583-7
21. Andjic M, Spiroski D, Ilic Stojanovic O, et al. Effect of short-term exercise training in patients following acute myocardial infarction treated with primary percutaneous coronary intervention. *Eur J Phys Rehabil Med*. 2016;52(3):364–369.
22. Ansari Z, Rafat S, Jorat MV, Ghanbari-Firoozabadi M, Mirzaei M, Sarebanhassanabadi M. Effect of inpatient cardiac rehabilitation on QT dispersion in patients with acute myocardial infarction. *Acta Med Iran*. 2013;51(9):604–610.
23. Arai Y, Kimura T, Takahashi Y, Hashimoto T, Arakawa M, Okamura H. Preoperative frailty is associated with progression of postoperative cardiac rehabilitation in patients undergoing cardiovascular surgery. *Gen Thorac Cardiovasc Surg*. 2019;67(11):917–924. doi:10.1007/s11748-019-01121-7
24. Arai Y, Kimura T, Takahashi Y, Hashimoto T, Arakawa M, Okamura H. Preoperative nutritional status is associated with progression of postoperative cardiac rehabilitation in patients undergoing cardiovascular surgery. *Gen Thorac Cardiovasc Surg*. 2018;66(11):632–640. doi:10.1007/s11748-018-0961-7
25. Asha Jyothi K, Madhavi K, Charan K, Thabita P. Study on physiological outcomes after phase I cardiac rehabilitation in mitral valve replacement individuals- an observational study. *Indian J Physiother Occup Therapy*. 2012;6(1):40–43.
26. Awoke MS, Baptiste DL, Davidson P, Roberts A, Dennison-Himmelfarb C. A quasi-experimental study examining a nurse-led education program to improve knowledge, self-care, and reduce readmission for individuals with heart failure. *Contemp Nurse*. 2019;55(1):15–26. doi:10.1080/10376178.2019.1568198

27. Aziz EF, Javed F, Pulimi S, et al. Implementing a pathway for the management of acute coronary syndrome leads to improved compliance with guidelines and a decrease in angina symptoms. *Int J Health Care Qual.* 2012;34(4):5–14. doi:10.1111/j.1945-1474.2011.00145.x
28. Babu AS, Maiya AG, George MM, Padmakumar R, Guddattu V. Effects of combined early in-patient cardiac rehabilitation and structured home-based program on function among patients with congestive heart failure: a randomized controlled trial. *Heart Views.* 2011;12(3):99–103. doi:10.4103/1995-705X.95064
29. Babu AS, Noone MS, Haneef M, Naryanan SM. Protocol-guided phase-I cardiac rehabilitation in patients with ST-elevation myocardial infarction in A rural hospital. *Heart Views.* 2010;11(2):52–56. doi:10.4103/1995-705X.73209
30. Baessler A, Hengstenberg C, Holmer S, et al. Long-term effects of in-hospital cardiac rehabilitation on the cardiac risk profile: a case-control study in pairs of siblings with myocardial infarction. *Eur Heart J.* 2001;22(13):1111–1118. doi:10.1053/euhj.2000.2444
31. Balestroni G, Panzeri A, Omarini P, et al. Psychophysical health of elderly inpatients in cardiac rehabilitation: a retrospective cohort study. *Eur J Phys Rehabil Med.* 2020;56(2):197–205. doi:10.23736/S1973-9087.20.05970-5
32. Bannerman A, Hamilton K, Isles C, et al. Myocardial infarction in men and women under 65 years of age: no evidence of gender bias. *Scott Med J.* 2001;46(3):73–78. doi:10.1177/003693300104600304
33. Barth J, Paul J, Harter M, Bengel J. Inpatient psychotherapeutic treatment for cardiac patients with depression in Germany: short-term results. *GMS Psycho Soc Med.* 2005;2:1–8.
34. Bauer BA, Cutshall SA, Anderson PG, et al. Effect of the combination of music and nature sounds on pain and anxiety in cardiac surgical patients: a randomized study. *Altern Ther Health Med.* 2011;17(4):16–23.
35. Bierbauer W, Scholz U, Bermudez T, et al. Improvements in exercise capacity of older adults during cardiac rehabilitation. *Eur J Prev Cardiol.* 2020;27:1747–1755. doi:10.1177/2047487320914736
36. Borowicz-Bienkowska S, Deskur-Smielecka E, Maleszka M, et al. The impact of short-term cardiac rehabilitation on changing dietary habits in patients after acute coronary syndrome. *J Cardiopulm Rehabil Prev.* 2013;33(4):234–238. doi:10.1097/HCR.0b013e318293b47b
37. Borzou SR, Amiri S, Salavati M, Soltanian AR, Safarpour G. Effects of the first phase of cardiac rehabilitation training on self-efficacy among patients undergoing coronary artery bypass graft surgery. *J Tehran Univ Heart Cent.* 2018;13(3):126–131.
38. Boyd J, Paratz J, Tronstad O, Caruana L, Walsh J. Exercise is feasible in patients receiving vasoactive medication in a cardiac surgical intensive care unit: a prospective observational study. *Aust Crit Care.* 2020;33(3):244–249. doi:10.1016/j.aucc.2020.02.004
39. Boyde M, Grenfell K, Brown R, et al. What have our patients learnt after being hospitalised for an acute myocardial infarction? *Aust Crit Care.* 2015;28(3):134–139. doi:10.1016/j.aucc.2014.05.003
40. Broadbent E, Ellis CJ, Thomas J, Gamble G, Petrie KJ. Further development of an illness perception intervention for myocardial infarction patients: a randomized controlled trial. *J Psychosom Res.* 2009;67(1):17–23. doi:10.1016/j.jpsychores.2008.12.001
41. Broadbent E, Leggat A, McLachlan A, Kerr A. Providing cardiovascular risk management information to acute coronary syndrome patients: a randomized trial. *Br J Health Psychol.* 2013;18(1):83–96. doi:10.1111/j.2044-8287.2012.02081.x
42. Busch C, Baumbach C, Willemsen D, et al. Supervised training with wireless monitoring of ECG, blood pressure and oxygen-saturation in cardiac patients. *J Telemed Telecare.* 2009;15(3):112–114. doi:10.1258/jtt.2009.003002
43. Busch JC, Lillou D, Wittig G, et al. Resistance and balance training improves functional capacity in very old participants attending cardiac rehabilitation after coronary bypass surgery. *J Am Geriatr Soc.* 2012;60(12):2270–2276. doi:10.1111/jgs.12030
44. Cacciatore F, Anello CB, Ferrara N, et al. Determinants of prolonged intensive care unit stay after cardiac surgery in the elderly. *Aging Clin Exp Res.* 2012;24(6):627–634. doi:10.3275/8521
45. Cai H, Zheng Y, Liu Z, et al. Effect of pre-discharge cardiopulmonary fitness on outcomes in patients with ST-elevation myocardial infarction after percutaneous coronary intervention. *BMC Cardiovasc Disord.* 2019;19(1):10–Jan. doi:10.1186/s12872-019-1189-x
46. Cameron LD, Petrie KJ, Ellis CJ, Buick D, Weinman JA. Trait negative affectivity and responses to a health education intervention for myocardial infarction patients. *Psychol Health.* 2005;20(1):1–18. doi:10.1080/08870440412331300011
47. Ranghi F, De Benedetti S, Caminiti G, et al. Cognitive impairment affects physical recovery of patients with heart failure undergoing exercise training. *Monaldi Arch Chest Dis.* 2012;78(2):113–114.
48. Cartledge S, Finn J, Bray JE, et al. Incorporating cardiopulmonary resuscitation training into a cardiac rehabilitation programme: a feasibility study. *Eur J Cardiovasc Nurs.* 2018;17(2):148–158. doi:10.1177/1474515117721010
49. Chu SK, McCormick Z, Hwang S, Sliwa JA, Rydberg L. Outcomes of acute inpatient rehabilitation of patients with left ventricular assist devices. *PMR.* 2014;6(11):1008–1012. doi:10.1016/j.pmrj.2014.05.004
50. Eichler S, Salzwedel A, Reibis R, et al. Multicomponent cardiac rehabilitation in patients after transcatheter aortic valve implantation: predictors of functional and psychocognitive recovery. *Eur J Prev Cardiol.* 2017;24(3):257–264. doi:10.1177/2047487316679527
51. Ekman I, Wolf A, Olsson LE, et al. Effects of person-centred care in patients with chronic heart failure: the PCC-HF study. *Eur Heart J.* 2012;33(9):1112–1119. doi:10.1093/eurheartj/ehr306
52. English ML, Speed J. Effectiveness of acute inpatient rehabilitation after left ventricular assist device placement. *Am J Phys Med Rehabil.* 2013;92(7):621–626. doi:10.1097/PHM.0b013e31827442a4
53. Eto Y, Koike A, Matsumoto A, et al. Early aerobic training increases end-tidal CO₂ pressure during exercise in patients after acute myocardial infarction. *Circ J.* 2004;68(8):778–783. doi:10.1253/circj.68.778
54. Eysenbach G, Schulz D, Schweier R, et al. A web-based peer-modeling intervention aimed at lifestyle changes in patients with coronary heart disease and chronic back pain: sequential controlled trial. *J Med Internet Res.* 2014;16(7):e177–e177. doi:10.2196/jmir.3434
55. Fernandes AC, McIntyre T, Coelho R, Prata J, Maciel MJ. Impact of a brief psychological intervention on lifestyle, risk factors and disease knowledge during phase I of cardiac rehabilitation after acute coronary syndrome. *Rev Port Cardiol.* 2019;38(5):361–368. doi:10.1016/j.repc.2018.09.009
56. Fiorina C, Vizzardi E, Lorusso R, et al. The 6-min walking test early after cardiac surgery. Reference values and the effects of rehabilitation programme. *Eur J Cardiothorac Surg.* 2007;32(5):724–729. doi:10.1016/j.ejcts.2007.08.013
57. Forestieri P, Guizilini S, Peres M, et al. A cycle ergometer exercise program improves exercise capacity and inspiratory muscle function in hospitalized patients awaiting heart transplantation: a pilot study. *Braz Cardiovasc Surg.* 2016;31(5):389–395. doi:10.5935/1678-9741.20160078

58. Forrest G, Radu G, Rifenburg K, Shields E, Clift S. Left Ventricular Assist Device: care on inpatient rehabilitation facility. *Rehabil Nurs*. 2015;40(6):378–383. doi:10.1002/rnj.192
59. Frengley JD, Sansone GR, Alba A, Uppal K, Kleinfeld J. Influence of age on rehabilitation outcomes and survival in post-acute inpatient cardiac rehabilitation. *J Cardiopulm Rehabil Prev*. 2011;31(4):230–238. doi:10.1097/hcr.0b013e318207d314
60. Frizzelli R, Tortelli O, Di Comite V, Ghirardi R, Pinzi C, Scarduelli C. Deep venous thrombosis of the neck and pulmonary embolism in patients with a central venous catheter admitted to cardiac rehabilitation after cardiac surgery: a prospective study of 815 patients. *Intern Emerg Med*. 2008;3(4):325–330. doi:10.1007/s11739-008-0142-2
61. Fukui S, Ogo T, Takaki H, et al. Efficacy of cardiac rehabilitation after balloon pulmonary angioplasty for chronic thromboembolic pulmonary hypertension. *Heart*. 2016;102(17):1403–1409. doi:10.1136/heartjnl-2015-309230
62. Galante A, Pietroliusti A, Cavazzini C, et al. Incidence and risk factors associated with cardiac arrhythmias during rehabilitation after coronary artery bypass surgery. *Arch Phys Med Rehabil*. 2000;81(7):947–952. doi:10.1053/apmr.2000.5587
63. Ghanbari-Firoozabadi M, Mirzaei M, Nasiriani K, et al. Cardiac specialists' perspectives on barriers to cardiac rehabilitation referral and participation in a low-resource setting. *Rehabil Proc Outcome*. 2020;9:1–7. doi:10.1177/1179572720936648
64. Gonçalves Mendes R, Polaquini Simões R, Melo Costa FS, et al. Short-term supervised inpatient physiotherapy exercise protocol improves cardiac autonomic function after coronary artery bypass graft surgery – a randomised controlled trial. *Disabil Rehabil*. 2010;32(16):1320–1327. doi:10.3109/09638280903483893
65. Griffo R, Ambrosetti M, Tramari R, et al. Effective secondary prevention through cardiac rehabilitation after coronary revascularization and predictors of poor adherence to lifestyle modification and medication. Results of the ICAROS Survey. *Int J Cardiol*. 2013;167(4):1390–1395. doi:10.1016/j.ijcard.2012.04.069
66. Hansen D, Eijnde BO, Roelants M, et al. Clinical benefits of the addition of lower extremity low-intensity resistance muscle training to early aerobic endurance training intervention in patients with coronary artery disease: a randomized controlled trial. *J Rehabil Med*. 2011;43(9):800–807. doi:10.2340/16501977-0853
67. Hanson LC, Taylor NF, McBurney H. The 10 m incremental shuttle walk test is a highly reliable field exercise test for patients referred to cardiac rehabilitation: a retest reliability study. *Physiotherapy*. 2016;102(3):243–248. doi:10.1016/j.physio.2015.08.004
68. Harada H, Kai H, Niiyama H, et al. Effectiveness of cardiac rehabilitation for prevention and treatment of sarcopenia in patients with cardiovascular disease - a retrospective cross-sectional analysis. *J Nutr Health Aging*. 2017;21(4):449–456. doi:10.1007/s12603-016-0743-9
69. Herdy AH, Marcchi PLB, Vila A, et al. Pre- and postoperative cardiopulmonary rehabilitation in hospitalized patients undergoing coronary artery bypass surgery a randomized controlled trial. *Am J Phys Med Rehabil*. 2008;87(9):714–719. doi:10.1097/PHM.0b013e3181839152
70. Hildebrandt A, Willemsen D, Reiss N, Bartsch P, Schmidt T, Bjarnason-Wehrens B. Characteristics, therapeutic needs, and scope of patients with a continuous-flow left ventricular device entering cardiac rehabilitation: a RETROSPECTIVE ANALYSIS. *J Cardiopulm Rehabil Prev*. 2019;39(2):91–96. doi:10.1097/HCR.0000000000000342
71. Hirschhorn AD, Richards D, Mungovan SF, Morris NR, Adams L. Supervised moderate intensity exercise improves distance walked at hospital discharge following coronary artery bypass graft surgery-A randomised controlled trial. *Heart Lung Circ*. 2008;17(2):129–138. doi:10.1016/j.hlc.2007.09.004
72. Hofer S, Kullich W, Graninger U, et al. Cardiac rehabilitation in Austria: long term health-related quality of life outcomes. *Health Qual Life Outcomes*. 2009;7:99. doi:10.1186/1477-7525-7-99
73. Hojskov IE, Thygesen LC, Moons P, Egerod I, Olsen PS, Berg SK. The challenge of non-adherence to early rehabilitation after coronary artery bypass surgery: secondary results from the SheppHeartCABG trial. *Eur J Cardiovasc Nurs*. 2020;19(3):238–247. doi:10.1177/1474515119883454
74. Hotta K, Kamiya K, Shimizu R, et al. Stretching exercises enhance vascular endothelial function and improve peripheral circulation in patients with acute myocardial infarction. *Int Heart J*. 2013;54(2):59–63. doi:10.1536/ihj.54.59
75. Huffman JC, Mastromauro CA, Sowden GL, Wittmann C, Rodman R, Januzzi JL. A collaborative care depression management program for cardiac inpatients: depression characteristics and in-hospital outcomes. *Psychosomatics*. 2011;52(1):26–33. doi:10.1016/j.psych.2010.11.021
76. Istvanovic N, Smalcelj A, Filakovic P, Cerovec D, Plecko D. Influence of in-hospital cardiac rehabilitation on psychological status after myocardial infarction in patients with D-type personality. *Coll Antropol*. 2011;35(3):797–807.
77. Kamke W, Dovifat C, Schranz M, Behrens S, Moesenthin J, Voller H. Cardiac rehabilitation in patients with implantable defibrillators: feasibility and complications. *Z Kardiol*. 2003;92(10):869–875. doi:10.1007/s00392-003-0997-1
78. Kanazawa N, Ueshima K, Tominari S, Nakayama T. Underuse of cardiac rehabilitation in workers with coronary artery disease: claims database survey in Japan. *Circ J*. 2017;81(10):1424–1431. doi:10.1253/circj.CJ-16-1260
79. Karakas M, Jaensch A, Breitling LP, Brenner H, Koenig W, Rothenbacher D. Prognostic value of midregional pro-A-type natriuretic peptide and N-terminal pro-B-type natriuretic peptide in patients with stable coronary heart disease followed over 8 years. *Clin Chem*. 2014;60(11):1441–1449. doi:10.1373/clinchem.2013.220202
80. Kato J, Koike A, Kuroki K, et al. Safety and efficacy of in-hospital cardiac rehabilitation following antiarrhythmic therapy for patients with electrical storm. *J Cardiol*. 2019;73(2):171–178. doi:10.1016/j.jjcc.2018.08.004
81. Katano S, Hashimoto A, Ohori K, et al. Nutritional status and energy intake as predictors of functional status after cardiac rehabilitation in elderly inpatients with heart failure - A retrospective cohort study. *Circ J*. 2018;82(6):1584–1591. doi:10.1253/circj.CJ-17-1202
82. Kim C, Choi HE. The effect and safety of aerobic interval training according to exercise intensity in acute coronary syndrome. *J Cardiopulm Rehabil Prev*. 2020;40(3):178–182. doi:10.1097/HCR.0000000000000455
83. Kozomara S, Krstic I. Effects of dynamic exercise and dynamic resistance training in rehabilitation of patients after surgical coronary revascularization. *Acta Fac Naissensis*. 2017;34(2):169–178. doi:10.1515/afmnai-2017-0019
84. Lau-Walker M, Landy A, Murrells T. Personalised discharge care planning for postmyocardial infarction patients through the use of the Personalised Patient Education Protocol-implementing theory into practice. *J Clin Nurs*. 2016;25(9–10):1292–1300. doi:10.1111/jocn.13177
85. Macchi C, Fattiroli F, Lova RM, et al. Early and late rehabilitation and physical training in elderly patients after cardiac surgery. *Am J Phys Med Rehabil*. 2007;86(10):826–834. doi:10.1097/PHM.0b013e318151fd86
86. Macchi C, Polcaro P, Cecchi F, et al. One-year adherence to exercise in elderly patients receiving postacute inpatient rehabilitation after cardiac surgery. *Am J Phys Med Rehabil*. 2009;88(9):727–734. doi:10.1097/PHM.0b013e3181b332a1

87. Machado B, Fernandes A, Cruzeiro S, Jesus R, Araújo N, Araújo I. Cardiac rehabilitation program and health literacy levels: a cross-sectional, descriptive study. *Nurs Health Sci*. 2019;21(3):406–412. doi:10.1111/nhs.12615
88. Manzoni GM, Villa V, Compare A, et al. Short-term effects of a multi-disciplinary cardiac rehabilitation programme on psychological well-being, exercise capacity and weight in a sample of obese in-patients with coronary heart disease: a practice-level study. *Psychol Health Med*. 2011;16(2):178–189. doi:10.1080/13548506.2010.542167
89. Masnaghetti SE, Braga SS, Vaninetti R, Baiardi P, Pedretti RFE. Altered fasting glycemia in cardiac patients during in-hospital rehabilitation: impact on short and long-term follow-up. *J Cardiovasc Med*. 2017;18(8):625–630. doi:10.2459/JCM.0000000000000478
90. Mayer-Berger W, Simic D, Mahmoodzad J, et al. Efficacy of a long-term secondary prevention programme following inpatient cardiovascular rehabilitation on risk and health-related quality of life in a low-education cohort: a randomized controlled study. *Eur J Prev Cardiol*. 2014;21(2):145–152. doi:10.1177/2047487312465526
91. Mendes RG, Simoes RP, de Souza Melo Costa F, et al. Left-ventricular function and autonomic cardiac adaptations after short-term inpatient cardiac rehabilitation: a prospective clinical trial. *J Rehabil Med*. 2011;43(8):720–727. doi:10.2340/16501977-0843
92. Meng K, Seekatz B, Haug G, et al. Evaluation of a standardized patient education program for inpatient cardiac rehabilitation: impact on illness knowledge and self-management behaviors up to 1 year. *Health Educ Res*. 2014;29(2):235–246. doi:10.1093/her/cyt107
93. Mohammed HG, Shabana AM. Effect of cardiac rehabilitation on cardiovascular risk factors in chronic heart failure patients. *Egypt Heart J*. 2018;70(2):77–82. doi:10.1016/j.ehj.2018.02.004
94. Motoki H, Nishimura M, Kanai M, et al. Impact of inpatient cardiac rehabilitation on Barthel Index score and prognosis in patients with acute decompensated heart failure. *Int J Cardiol*. 2019;293:125–130. doi:10.1016/j.ijcard.2019.06.071
95. Müller-Nordhorn J, Roll S, Willich N, Willich SN. Comparison of the short form (SF)-12 health status instrument with the SF-36 in patients with coronary heart disease. *Heart*. 2004;90(5):523–527. doi:10.1136/hrt.2003.013995
96. Murphy MC, George MV, Driscoll AL. Concordance with phase-one cardiac rehabilitation guidelines in the inpatient setting. *Aust J Adv Nurs*. 2007;25(1):31–37.
97. Musekamp G, Schuler M, Seekatz B, Bengel J, Faller H, Meng K. Does improvement in self-management skills predict improvement in quality of life and depressive symptoms? A prospective study in patients with heart failure up to one year after self-management education. *BMC Cardiovasc Disord*. 2017;17:51–51. doi:10.1186/s12872-017-0486-5
98. Nakamura K, Nakamura E, Niina K, Kojima K. Outcome after valve surgery in octogenarians and efficacy of early mobilization with early cardiac rehabilitation. *Gen Thorac Cardiovasc Surg*. 2010;58(12):606–611. doi:10.1007/s11748-010-0665-0
99. Nguyen TN, Abramson BL, Galluzzi A, Tan M, Yan AT, Goodman SG. Temporal trends and referral factors for cardiac rehabilitation post-acute coronary syndrome in Ontario: insights from the Canadian global registry of acute coronary events. *Can J Cardiol*. 2013;29(12):1604–1609. doi:10.1016/j.cjca.2013.10.002
100. Ogawa M, Satomi-Kobayashi S, Yoshida N, et al. Effects of acute-phase multidisciplinary rehabilitation on unplanned readmissions after cardiac surgery. *J Thorac Cardiovasc Surg*. 2019;161:1853–1860.e2. doi:10.1016/j.jtcvs.2019.11.069
101. Opasich C, De Feo S, Pinna GD, et al. Distance walked in the 6-minute test soon after cardiac surgery: toward an efficient use in the individual patient. *Chest*. 2004;126(6):1796–1801. doi:10.1378/chest.126.6.1796
102. Pantoni CBF, Di Thommazo-Luporini L, Mendes RG, et al. Continuous positive airway pressure during exercise improves walking time in patients undergoing inpatient cardiac rehabilitation after coronary artery bypass graft surgery: a randomized controlled trial. *J Cardiopulm Rehabil Prev*. 2016;36(1):20–27. doi:10.1097/HCR.0000000000000144
103. Pasquini G, Vannetti F, Molino-Lova R. Ability to work in anaerobic condition is associated with physical performance on the six-minute walk test in older patients receiving cardiac rehabilitation. *J Rehabil Med*. 2015;47(5):472–477. doi:10.2340/16501977-1956
104. Patman S, Sanderson D, Blackmore M. Physiotherapy following cardiac surgery: is it necessary during the intubation period? *Aust J Physiother*. 2001;47(1):7–16. doi:10.1016/S0004-9514(14)60294-4
105. Peschel T, Sixt S, Beitz F, et al. High, but not moderate frequency and duration of exercise training induces downregulation of the expression of inflammatory and atherogenic adhesion molecules. *Eur J Cardiovasc Prev Rehabil*. 2007;14(3):476–482. doi:10.1097/HJR.0b013e328167239d
106. Pizzorno M, Desilvestri M, Lippi L, et al. Early cardiac rehabilitation: could it improve functional outcomes and reduce length of stay and sanitary costs in patients aged 75 years or older? A retrospective case-control study. *Aging Clin Exp Res*. 2020;33:957–964. doi:10.1007/s40520-020-01589-x
107. Polcaro P, Lova RM, Guarducci L, et al. Left-ventricular function and physical performance on the 6-min walk test in older patients after inpatient cardiac rehabilitation. *Am J Phys Med Rehabil*. 2008;87(1):46–83. doi:10.1097/phm.0b013e31815e67d8
108. Racca V, Di Rienzo M, Mazzini P, et al. ICF-based approach to evaluating functionality in cardiac rehabilitation patients after heart surgery. *Eur J Phys Rehabil Med*. 2015;51(4):457–468.
109. Racca V, Bordonni B, Castiglioni P, Modica M, Ferratini M. Osteopathic manipulative treatment improves heart surgery outcomes: a randomized controlled trial. *Ann Thor Surg*. 2017;104(1):145–152. doi:10.1016/j.athoracsur.2016.09.110
110. Rahmani R, Niyazi S, Sobh-Rakhshankhah A, et al. Effects of a cardiac rehabilitation program versus usual care on cardiopulmonary function in patients with cardiac syndrome X. *J Cardiopulm Rehabil Prev*. 2020;40(1):41–47. doi:10.1097/HCR.0000000000000439
111. Ratajska M, Chochowska M, Kulik A, Bugajski P. Myofascial release in patients during the early postoperative period after revascularisation of coronary arteries. *Disabil Rehabil*. 2019;1–12. doi:10.1080/09638288.2019.1593518
112. Reibis R, Salzwedel A, Buhler H, Wegscheider K, Eichler S, Voller H. Impact of training methods and patient characteristics on exercise capacity in patients in cardiovascular rehabilitation. *Eur J Prev Cardiol*. 2016;23(5):452–459. doi:10.1177/2047487315600815
113. Riley H, Headley S, Lindenauer PK, et al. Patient perception of how smoking status influences cardiac rehabilitation attendance after an acute cardiac hospitalization. *J Cardiopulm Rehabil Prev*. 2019;39(3):181–186. doi:10.1097/HCR.0000000000000366
114. Rossi Ferrario S, Bacich D, Beltrame L, Balestroni G, Pistono M. Does a comprehensive inpatient rehabilitation program improve patients' and caregivers' emotional state in LVAD patients? *Artif Organs*. 2019;43(3):229–233. doi:10.1111/aor.13351
115. Salzwedel A, Heidler MD, Meng K, et al. Impact of cognitive performance on disease-related knowledge six months after multi-component rehabilitation in patients after an acute cardiac event. *Eur J Prev Cardiol*. 2019;26(1):46–55. doi:10.1177/2047487318791609

116. Salzwedel A, Wegscheider K, Schulz-Behrendt C, Dorr G, Reibis R, Voller H. No impact of an extensive social intervention program on return to work and quality of life after acute cardiac event: a cluster-randomized trial in patients with negative occupational prognosis. *Int Arch Occup Environ Health*. 2019;92(8):1109–1120. doi:10.1007/s00420-019-01450-3
117. Salzwedel A, Nosper M, Rohrig B, Linck-Eleftheriadis S, Strandt G, Voller H. Outcome quality of in-patient cardiac rehabilitation in elderly patients - Identification of relevant parameters. *Eur J Prev Cardiol*. 2014;21(2):172–180. doi:10.1177/2047487312469475
118. Salzwedel A, Koran I, Langheim E, et al. Patient-reported outcomes predict return to work and health-related quality of life six months after cardiac rehabilitation: results from a German multi-centre registry (OutCaRe). *PLoS One*. 2020;15(5):e0232752. doi:10.1371/journal.pone.0232752
119. Salzwedel A, Reibis R, Hadzic M, Buhler H, Voller H. Patients' expectations of returning to work, co-morbid disorders and work capacity at discharge from cardiac rehabilitation. *Vasc Health Risk Manag*. 2019;15:301–308. doi:10.2147/VHRM.S216039
120. Salzwedel A, Heidler MD, Haubold K, et al. Prevalence of mild cognitive impairment in employable patients after acute coronary event in cardiac rehabilitation. *Vasc Health Risk Manag*. 2017;13:55–60. doi:10.2147/VHRM.S121086
121. Sansone GR, Alba A, Frengley JD. Analysis of FIM instrument scores for patients admitted to an inpatient cardiac rehabilitation program. *Arch Phys Med Rehabil*. 2002;83(4):506–512. doi:10.1053/apmr.2002.31183
122. Santos-Hiss MD, Melo RC, Neves VR, et al. Effects of progressive exercise during phase I cardiac rehabilitation on the heart rate variability of patients with acute myocardial infarction. *Disabil Rehabil*. 2011;33(10):835–842. doi:10.3109/09638288.2010.514016
123. Scalvini S, Zanelli E, Comini L, et al. Home-based versus in-hospital cardiac rehabilitation after cardiac surgery: a nonrandomized controlled study. *Phys Ther*. 2013;93(8):1073–1083. doi:10.2522/ptj.20120212
124. Scalvini S, Grossetti F, Paganoni AM, La Rovere MT, Pedretti RF, Frigerio M. Impact of in-hospital cardiac rehabilitation on mortality and readmissions in heart failure: a population study in Lombardy, Italy, from 2005 to 2012. *Eur J Prev Cardiol*. 2019;26(8):808–817. doi:10.1177/2047487319833512
125. Schmidt T, Bjarnason-Wehrens B, Bartsch P, et al. Exercise capacity and functional performance in heart failure patients supported by a left ventricular assist device at discharge from inpatient rehabilitation. *Artif Organs*. 2018;42(1):22–30. doi:10.1111/aor.12936
126. Schnoll F, Laimer H, Altenberger J, et al. Reduction of coronary risk factors immediately and 1 year after inpatient rehabilitation in a highly motivated patient cohort. *Wien Med Wochenschr*. 2015;165(3–4):71–78. doi:10.1007/s10354-014-0339-0
127. Schwaab B, Waldmann A, Katalinic A, Sheikhzadeh A, Raspe H. In-patient cardiac rehabilitation versus medical care - A prospective multicentre controlled 12 months follow-up in patients with coronary heart disease. *Eur J Cardiovasc Prev Rehabil*. 2011;18(4):581–586. doi:10.1177/1741826710389392
128. Schweikert B, Hahmann H, Steinacker JM, et al. Intervention study shows outpatient cardiac rehabilitation to be economically at least as attractive as inpatient rehabilitation. *Clin Res Cardiol*. 2009;98(12):787–795. doi:10.1007/s00392-009-0081-6
129. Scrutinio D, Passantino A, Catanzaro R, et al. Inpatient cardiac rehabilitation soon after hospitalization for acute decompensated heart failure: a propensity score study. *J Cardiopulm Rehabil Prev*. 2012;32(2):71–77. doi:10.1097/hcr.0b013e31823be124
130. Seo YG, Park WH, Oh S, et al. Clinical outcomes of inpatient cardiac rehabilitation for patients with treated left ventricular assist device in Korea: 1-year follow-up. *J Exerc Rehabil*. 2019;15(3):481–487. doi:10.12965/jer.1938124.062
131. Sharp PB, Salyer J. Self-efficacy and barriers to healthy diet in cardiac rehabilitation participants and nonparticipants. *J Cardiovasc Nurs*. 2012;27(3):253–262. doi:10.1097/jcn.0b013e31821efdc2
132. Skobel EC, Dreher M, Knackstedt C. Effect of cardiac training therapy on minute ventilation/carbon dioxide production slope and exercise parameters in patients with severe chronic heart failure in short-time rehabilitation. *Res Cardiovasc Med*. 2017;6(4):20–23. doi:10.4103/rcm.rcm_30_17
133. Smialek J, Lelakowski J, Majewski J. Efficacy and safety of early comprehensive cardiac rehabilitation following the implantation of cardioverter-defibrillator. *Kardiologia Pol*. 2013;71(10):1021–1028. doi:10.5603/KP.2013.0256
134. Socha M, Wronecki K, Sobiech KA. Gender and age-dependent differences in body composition changes in response to cardiac rehabilitation exercise training in patients after coronary artery bypass grafting. *Ann Agric Environ Med*. 2017;24(3):517–521. doi:10.5604/12321966.1230731
135. Soleimannejad K, Nouzari Y, Ahsani A, Nejatian M, Sayehmiri K. Evaluation of the effect of cardiac rehabilitation on left ventricular diastolic and systolic function and cardiac chamber size in patients undergoing percutaneous coronary intervention. *J Tehran Univ Heart Cent*. 2014;9(2):54–58.
136. Song KJ. The effects of self-efficacy promoting cardiac rehabilitation program on self-efficacy, health behavior, and quality of life. *Taehan Kanho Hakhoe chi*. 2003;33(4):510–518. doi:10.4040/jkan.2003.33.4.510
137. Spiroski D, Andjic M, Stojanovic OI, et al. Very short/short-term benefit of inpatient/outpatient cardiac rehabilitation programs after coronary artery bypass grafting surgery. *Clin Cardiol*. 2017;40(5):281–286. doi:10.1002/clc.22656
138. Steinacker JM, Liu Y, Muche R, et al. Long term effects of comprehensive cardiac rehabilitation in an inpatient and outpatient setting. *Swiss Med Wkly*. 2011;141:w13141. doi:10.4414/sm.w.2011.13141
139. Tabet JY, Meurin P, Benzidi Y, et al. Greater prognostic value of peak VO₂ after exercise training program completion in heart failure patients. *Int J Cardiol*. 2013;168(4):4139–4144. doi:10.1016/j.ijcard.2013.07.076
140. Taya M, Amiya E, Hatano M, et al. High-intensity aerobic interval training can lead to improvement in skeletal muscle power among in-hospital patients with advanced heart failure. *Heart Vessels*. 2018;33(7):752–759. doi:10.1007/s00380-018-1120-x
141. Tsai YJ, Li MH, Chen CH, Tuan SH, Chen YJ, Lin KL. Improved oxygen uptake efficiency slope in acute myocardial infarction patients after early phase I cardiac rehabilitation. *Int J Rehabil Res*. 2017;40(3):215–219. doi:10.1097/MRR.0000000000000229
142. Turk-Adawi KI, Oldridge NB, Tarima SS, Stason WB, Shepard DS. Cardiac rehabilitation enrollment among referred patients: PATIENT AND ORGANIZATIONAL FACTORS. *J Cardiopulm Rehabil Prev*. 2014;34(2):114–122. doi:10.1097/HCR.000000000000017
143. Twardella D, Merx H, Hahmann H, Wusten B, Rothenbacher D, Brenner H. Long term adherence to dietary recommendations after inpatient rehabilitation: prospective follow up study of patients with coronary heart disease. *Heart*. 2006;92(5):635–640. doi:10.1136/hrt.2005.067611
144. Ul-Haq Z, Khan D, Hisam A, et al. Effectiveness of cardiac rehabilitation on health-related quality of life in patients with myocardial infarction in Pakistan. *J Coll Physicians Surg Pak*. 2019;29(9):803–809. doi:10.29271/jcpsp.2019.09.803

145. Vincent HK, Vincent KR. Functional and economic outcomes of cardiopulmonary patients: a preliminary comparison of the inpatient rehabilitation and skilled nursing facility environments. *Am J Phys Med Rehabil*. 2008;87(5):371–380. doi:10.1097/phm.0b013e31816dd251
146. Weberg M, Hjermstad MJ, Hilmarsen CW, Oldervoll L. Inpatient cardiac rehabilitation and changes in self-reported health related quality of life - a pilot study. *Ann Phys Rehabil Med*. 2013;56(5):342–355. doi:10.1016/j.rehab.2013.03.004
147. Weeks DL, Sprint GL, Stilwill V, Meisen-Vehrs AL, Cook DJ. Implementing wearable sensors for continuous assessment of daytime heart rate response in inpatient rehabilitation. *Telemed J E Health*. 2018;24:1014–1020. doi:10.1089/tmj.2017.0306
148. Willich SN, Muller-Nordhorn J, Kulig M, et al. Cardiac risk factors, medication, and recurrent clinical events after acute coronary disease: a prospective cohort study. *Eur Heart J*. 2001;22(4):307–313. doi:10.1053/euhj.2000.2294
149. Yildiz BS, Findikoglu G, Alihanoglu YI, Kilic ID, Evrengul H, Senol H. How do patients understand safety for cardiac implantable devices? Importance of postintervention education. *Rehabil Res Pract*. 2018;2018:1–9. doi:10.1155/2018/5689353
150. Yamamoto S, Matsunaga A, Wang G, et al. Effect of balance training on walking speed and cardiac events in elderly patients with ischemic heart disease. *Int Heart J*. 2014;55(5):397–403. doi:10.1536/ihj.14-017
151. Yamamoto S, Matsunaga A, Kamiya K, et al. Walking speed in patients with first acute myocardial infarction who participated in a supervised cardiac rehabilitation program after coronary intervention. *Int Heart J*. 2012;53(6):347–352. doi:10.1536/ihj.53.347
152. Wurst R, Kinkel S, Lin J, Goehner W, Fuchs R. Promoting physical activity through a psychological group intervention in cardiac rehabilitation: a randomized controlled trial. *J Behav Med*. 2019;42(6):1104–1116. doi:10.1007/s10865-019-00047-y
153. Wita K, Kulach A, Sikora J, et al. Managed care after acute myocardial infarction (Mc-AMI) reduces total mortality in 12-month follow-up results from a Poland's national health fund program of comprehensive post-mi care-a population-wide analysis. *J Clin Med*. 2020;9(10):1–10. doi:10.3390/jcm9103178
154. Windmöller P, Teixeira Bodnar E, Casagrande J, et al. Physical exercise combined with CPAP in subjects who underwent surgical myocardial revascularization: a randomized clinical trial. *Respir Care*. 2020;65(2):150–157. doi:10.4187/respcare.06919
155. Yokota J, Takahashi R, Matsukawa Y, Matsushima K. Examination of independent predictors of discharge disposition in acute phase hospitalized heart failure patients undergoing phase I cardiac rehabilitation. *Eur J Phys Rehabil Med*. 2020;56(6):780–786. doi:10.23736/S1973-9087.20.06347-9
156. Zolfaghari M, Mirhosseini SJ, Baghbehehti M, Bauer BA. The effect of classic chest physiotherapy on postoperative pain scores and hospital stay in patients undergoing off-pump coronary artery bypass graft surgery: a randomized clinical trial. *Galen Med J*. 2018;7(1):e838. doi:10.22086/gmj.v0i0.838
157. Zoch-Lesniak B, Dobberke J, Schlitt A, et al. Performance measures for short-term cardiac rehabilitation in patients of working age: results of the prospective observational multicenter registry OutCaRe. *Arch Rehabil Res Clin Transl*. 2020;2(2):100043. doi:10.1016/j.arct.2020.100043
158. Zhou Y, Cai X, Yu S, et al. Effect of comprehensive cardiac rehabilitation management on saq score, risk stratification in coronary artery disease, exercise tolerance and quality of life in patients with PCI. *Int J Clin Exp Med*. 2020;13(3):1620–1627.
159. Zanini M, Nery RM, De lima JB, Buhler RP, Da Silveira AD, Stein R. Effects of different rehabilitation protocols in inpatient cardiac rehabilitation after coronary artery bypass graft surgery: a RANDOMIZED CLINICAL TRIAL. *J Cardiopulm Rehabil Prev*. 2019;39(6):E19–E25. doi:10.1097/HCR.0000000000000431
160. Yu J, Wang M, Fang J, Lin Z. The effects of cardiac exercise rehabilitation training on the cardiopulmonary function and quality of life in patients with chronic stable heart failure. *Int J Clin Exp Med*. 2020;13(8):5983–5990.
161. Yu CM, Li LSW, Lam MF, Siu DCW, Miu RKM, Lau CP. Effect of a cardiac rehabilitation program on left ventricular diastolic function and its relationship to exercise capacity in patients with coronary heart disease: experience from a randomized, controlled study. *Am Heart J*. 2004;147(5):874–818. doi:10.1016/j.ahj.2003.12.004
162. Yost G, Coyle L, Milkevitch K, Adair R, Tatooles A, Bhat G. Efficacy of inpatient rehabilitation after left ventricular assist device implantation. *PMR*. 2017;9(1):40–45. doi:10.1016/j.pmrj.2016.05.013
163. Yoshida T, Yoshida K, Yamamoto C, et al. Effects of a two-week, hospitalized Phase II cardiac rehabilitation program on physical capacity, lipid profiles and psychological variables in patients with acute myocardial infarction. *Jpn Circ J*. 2001;65(2):87–93. doi:10.1253/jcj.65.87
164. Alavi M, Irajpour A, Giles T, Rabie K, Sarrafzadegan N. Barriers to education in cardiac rehabilitation within an Iranian society: a qualitative descriptive study. *Contemp Nurs*. 2013;44(2):204–214. doi:10.5172/conu.2013.44.2.204
165. Back M, Oberg B, Krevers B. Important aspects in relation to patients' attendance at exercise-based cardiac rehabilitation - facilitators, barriers and physiotherapist's role: a qualitative study. *BMC Cardiovasc Disord*. 2017;17(1):77. doi:10.1186/s12872-017-0512-7
166. Maddocks S, Cobbing SP. Experiences of and perspectives on phase I cardiac rehabilitation after coronary artery bypass graft surgery. *Physiother Can*. 2017;69(4):333–340. doi:10.3138/ptc.2016-39GH
167. Simonő CP, Pedersen BD, Dreyer P, Birkelund R. Dealing with existential anxiety in exercise-based cardiac rehabilitation: a phenomenological-hermeneutic study of patients' lived experiences. *J Clin Nurs*. 2015;24(17–18):2581–2590. doi:10.1111/jocn.12867
168. Svavarsdóttir MH, Sigurðardóttir ÁK, Steinsbekk A. Knowledge and skills needed for patient education for individuals with coronary heart disease: the perspective of health professionals. *Eur J Cardiovasc Nurs*. 2016;15(1):55–63. doi:10.1177/1474515114551123
169. White S, Bissell P, Anderson C. A qualitative study of cardiac rehabilitation patients' perspectives on making dietary changes. *J Hum Nutr Diet*. 2011;24(2):122–127. doi:10.1111/j.1365-277X.2010.01136.x
170. Amao R, Imamura T, Nakahara Y, et al. Reversible motor paralysis and early cardiac rehabilitation in patients with advanced heart failure receiving left ventricular assist device therapy. *Int Heart J*. 2016;57(6):766–768. doi:10.1536/ihj.16-153
171. Amaravathi E, Ramarao NH, Raghuram N, Pradhan B. Yoga-based postoperative cardiac rehabilitation program for improving quality of life and stress levels: fifth-year follow-up through a randomized controlled trial. *Int J Yoga*. 2018;11(1):44–52. doi:10.4103/ijoy.IJOY_57_16
172. Anderson JA, Petersen NJ, Kistner C, Soltero ER, Willson P. Determining predictors of delayed recovery and the need for transitional cardiac rehabilitation after cardiac surgery. *J Am Acad Nurse Pract*. 2006;18(8):386–392. doi:10.1111/j.1745-7599.2006.00152.x
173. Brown KD, Adams J, Meyer DM. Exercise training with cycle ergometry in the intensive care unit after total artificial heart implantation. *Bayl Univ Med Cent Proc*. 2020;33(4):674–676. doi:10.1080/08998280.2020.1789265
174. Brzek A, Nowak Z, Plewa M. Modified programme of in-patient (phase I) cardiac rehabilitation after acute myocardial infarction. *Int J Rehabil Res*. 2002;25(3):225–229. doi:10.1097/00004356-200209000-00008

175. Callahan MP, Pham T, Rashbaum I, Pineda H, Greenspan N. Cardiopulmonary rehabilitation in a patient with Noonan syndrome. *Arch Phys Med Rehabil*. 2000;81(2):230–232. doi:10.1016/S0003-9993(00)90147-3
176. Compostella L, Compostella C, Truong LVS, et al. History of erectile dysfunction as a predictor of poor physical performance after an acute myocardial infarction. *Eur J Prev Cardiol*. 2017;24(5):460–467. doi:10.1177/2047487316686434
177. Duru F, Candinas R, Dziekan G, Goebbels U, Myers J, Dubach P. Effect of exercise training on heart rate variability in patients with new-onset left ventricular dysfunction after myocardial infarction. *Am Heart J*. 2000;140(1):157–161. doi:10.1067/mhj.2000.106606
178. Froger-Bompas C, Laviolle B, Guillo P, et al. Sustained positive impact of a coronary rehabilitation programme on adherence to dietary recommendations. *Arch Cardiovasc Dis*. 2009;102(2):97–104. doi:10.1016/j.acvd.2008.10.020
179. Golabchi A, Basati F, Kargarfard M, Sadeghi M. Can cardiac rehabilitation programs improve functional capacity and left ventricular diastolic function in patients with mechanical reperfusion after ST elevation myocardial infarction?: a double-blind clinical trial. *ARYA Atheroscler*. 2012;8(3):2.
180. Goodwin-Esola MJ, Gatti G, Chokshi B. Rehabilitation barriers recognized in a geriatric patient after cardiac revascularization procedure. *Phys Occup Ther Geriatr*. 2014;32(3):183–187. doi:10.3109/02703181.2014.914619
181. Hambrecht R, Hilbrich L, Erbs S, et al. Correction of endothelial dysfunction in chronic heart failure: additional effects of exercise training and oral L-arginine supplementation. *J Am Coll Cardiol*. 2000;35(3):706–713. doi:10.1016/S0735-1097(99)00602-6
182. Helmy ZM, Mehani SHM, El-Refaey BH, Al-Salam EHA, Felaya ES. Low-level laser therapy versus trunk stabilization exercises on sternotomy healing after coronary artery bypass grafting: a randomized clinical trial. *Lasers Med Sci*. 2019;34(6):1115–1124. doi:10.1007/s10103-018-02701-4
183. Karaszewski D. Comparison of two models of hospital rehabilitation in patients after coronary artery bypass grafting. *Kardiochirurgia Torakochirurgia Pol*. 2014;11(1):86–89. doi:10.5114/kitp.2014.41940
184. Kim C, Choi HE, Lee BJ. Cardiac rehabilitation of a patient with an advanced dilated cardiomyopathy: a case report. *Ann Rehabil Med*. 2014;38(4):554–558. doi:10.5535/arm.2014.38.4.554
185. Kocur P, Deskur-Smielecka E, Wilk M, Dylewicz P. Effects of Nordic Walking training on exercise capacity and fitness in men participating in early, short-term inpatient cardiac rehabilitation after an acute coronary syndrome – a controlled trial. *Clin Rehabil*. 2009;23(11):995–1004. doi:10.1177/0269215509337464
186. Mlakar P, Salobir B, Cobo N, Prezelj M, Tercelj M, Sabovic M. Influence of short-term cardiac rehabilitation on oxidative stress in men after myocardial infarction depends upon smoking status. *J Cardiopulm Rehabil Prev*. 2013;33(6):401–405. doi:10.1097/hcr.000000000000016
187. Mohammed Mehani SH. Autonomic adaptation and functional capacity outcomes after hospital-based cardiac rehabilitation post coronary artery by pass graft. *Indian J Physiother Occup Therapy*. 2012;6(3):257–261.
188. Mourrot L, Teffaha D, Bouhaddi M, et al. Exercise rehabilitation restores physiological cardiovascular responses to short-term head-out water immersion in patients with chronic heart failure. *J Cardiopulm Rehabil Prev*. 2010;30(1):22–27. doi:10.1097/HCR.0b013e3181c8595c
189. Nissinoff J, Tian F, Therattil M, Salvarrey RM, Lee SW. Acute inpatient rehabilitation after left ventricular assist device implantation for congestive heart failure. *PMR*. 2011;3(6):586–589. doi:10.1016/j.pmrj.2010.11.010
190. Sarah S, Wolfgang MB, Claudia P. Effect of telerehabilitation on long-term adherence to yoga as an antihypertensive lifestyle intervention: results of a randomized controlled trial. *Complement Ther Clin Pract*. 2019;35:148–153. doi:10.1016/j.ctcp.2019.02.001
191. Sasanuma N, Takahashi K, Yamauchi S, et al. A five-year follow-up of a patient with fulminant myocarditis who underwent a stepwise and goal-oriented individualized comprehensive cardiac rehabilitation program. *J Cardiol Case*. 2015;11(6):160–163. doi:10.1016/j.jccase.2015.02.001
192. Sato S, Makita S, Majima M. Additional physical activity during cardiac rehabilitation leads to an improved heart rate recovery in male patients after coronary artery bypass grafting. *Circ J*. 2005;69(1):69–71. doi:10.1253/circj.69.69
193. Teffaha D, Mourrot L, Vernochet P, et al. Relevance of water gymnastics in rehabilitation programs in patients with chronic heart failure or coronary artery disease with normal left ventricular function. *J Card Fail*. 2011;17(8):676–683. doi:10.1016/j.cardfail.2011.04.008
194. Ugata Y, Wada H, Sakakura K, et al. High-intensity interval training for severe left ventricular dysfunction treated with left ventricular assist device. *Int Heart J*. 2018;59(1):216–219. doi:10.1536/ihj.17-090
195. Bahrami T, Rejeh N, Heravi-Karimooi M, Davood Tadrissi S, Vaismoradi M. The effect of foot reflexology on hospital anxiety and depression in female older adults: a randomized controlled trial. *Int J Ther Massage Bodywork*. 2019;12(3):16–21.
196. Bahrami T, Rejeh N, Heravi-Karimooi M, Vaismoradi M, Tadrissi SD, Sieloff C. Effect of aromatherapy massage on anxiety, depression, and physiologic parameters in older patients with the acute coronary syndrome: a randomized clinical trial. *Int J Nurs Pract*. 2017;23(6):1. doi:10.1111/ijn.12601
197. Bahrami T, Rejeh N, Heravi-Karimooi M, Vaismoradi M, Tadrissi SD, Sieloff CL. Aromatherapy massage versus reflexology on female elderly with acute coronary syndrome. *Nurs Crit Care*. 2018;23(5):229–236. doi:10.1111/nicc.12302
198. Murphy BA-O, Zaman S, Tucker K, et al. Enhancing the appeal of cardiac rehabilitation for women: development and pilot testing of a women-only yoga cardiac rehabilitation programme. *Eur J Cardiovasc Nurs*. 2021;20(7):633–640. doi:10.1093/eurjcn/zvab008
199. Westerdahl E, Möller M. Physiotherapy-supervised mobilization and exercise following cardiac surgery: a national questionnaire survey in Sweden. *J Cardiothorac Surg*. 2010;5:67–67. doi:10.1186/1749-8090-5-67
200. Kim JS, Kim GS, Kang SM, Chu SH. Symptom experience as a predictor of cardiac rehabilitation education programme attendance after percutaneous coronary intervention: a prospective questionnaire survey. *Eur J Cardiovasc Nurs*. 2020. doi:10.1177/1474515120940534
201. Jackson AC, Higgins RO, Murphy BM, Rogerson M, Le Grande MR. Cardiac rehabilitation in Australia: a brief survey of program characteristics. *Heart Lung Circ*. 2018;27(12):1415–1420. doi:10.1016/j.hlc.2017.08.024
202. Chockalingam P, Rajaram A, Maiya A, Contractor A. A multicentre retrospective study on quality and outcomes of cardiac rehabilitation programs in India. *Indian Heart J*. 2020;72(1):55–57. doi:10.1016/j.ihj.2020.03.002
203. Bestehorn K, Jannowitz C, Horack M, Karmann B, Halle M, Voller H. Current state of cardiac rehabilitation in Germany: patient characteristics, risk factor management and control status, by education level. *Vasc Health Risk Manag*. 2011;7:639–647. doi:10.2147/VHRM.S22971
204. Ghanbari-Firoozabadi M, Mirzaei M, Nasab MV, et al. Cross-cultural adaptation and psychometric validation of the Persian version of the cardiac rehabilitation barriers scale (CRBS-P). *BMJ Open*. 2020;10(6):e034552. doi:10.1136/bmjopen-2019-034552

205. Graversen CB, Johansen MB, Eichhorst R, et al. Influence of socioeconomic status on the referral process to cardiac rehabilitation following acute coronary syndrome: a cross-sectional study. *BMJ Open*. 2020;10(4):e036088. doi:10.1136/bmjopen-2019-036088
206. Kamiya K, Yamamoto T, Tsuchihashi-Makaya M, et al. Nationwide survey of multidisciplinary care and cardiac rehabilitation for patients with heart failure in Japan - An analysis of the AMED-CHF study. *Circ J*. 2019;83(7):1546–1552. doi:10.1253/circj.CJ-19-0241
207. Flynn FM, Cafarelli M, Petrakos K, Christophersen P. Improving outcomes for acute coronary syndrome patients in the hospital setting: successful implementation of the American heart association 'get with the guidelines' program by phase I cardiac rehabilitation nurses. *J Cardiovasc Nurs*. 2007;22(3):166–176. doi:10.1097/01.jcn.0000267824.27449.65
208. Goto Y, Saito M, Iwasaka T, et al. Poor implementation of cardiac rehabilitation despite broad dissemination of coronary interventions for acute myocardial infarction in Japan - A nationwide survey. *Circ J*. 2007;71(2):173–179. doi:10.1253/circj.71.173
209. Seo YG, Jang MJ, Park WH, Hong KP, Sung J. Inpatient cardiac rehabilitation programs' exercise therapy for patients undergoing cardiac surgery: national Korean Questionnaire Survey. *J Exerc Rehabil*. 2017;13(1):76–83. doi:10.12965/jer.1732806.403
210. De Gruyter E, Ford G, Stavreski B. Economic and social impact of increasing uptake of cardiac rehabilitation services--A cost benefit analysis. *Heart Lung Crit*. 2016;25(2):175–183. doi:10.1016/j.hlc.2015.08.007
211. Ben Gal T, Piepoli MF, Corra U, et al. Exercise programs for LVAD supported patients: a snapshot from the ESC affiliated countries. *Int J Cardiol*. 2015;201:215–219. doi:10.1016/j.ijcard.2015.08.081
212. Wieser S, Ruthemann I, De Boni S, et al. Cost of acute coronary syndrome in Switzerland in 2008. *Swiss Med Wkly*. 2012;142:w13655. doi:10.4414/SMW.2012.13655
213. Mildestvedt T, Meland E, Eide GE. How important are individual counselling, expectancy beliefs and autonomy for the maintenance of exercise after cardiac rehabilitation? *Scand J Public Health*. 2008;36(8):832–840. doi:10.1177/1403494808090633
214. Sarikaya S, Tur BS, Kurtas Y, et al. The awareness of physicians and allied health professionals about cardiopulmonary rehabilitation: a cross-sectional survey study. *Türkiye Fiziksel Tip Ve Rehabilitasyon Dergisi*. 2014;60(1):19–24. doi:10.5152/tftrd.2014.72677
215. Gallagher R, Neubeck L, Du H, et al. Facilitating or getting in the way? The effect of clinicians' knowledge, values and beliefs on referral and participation. *Eur J Prev Cardiol*. 2016;23(11):1141–1150. doi:10.1177/2047487316630085
216. Shanmugasagaram S, Oh P, Fau - Reid RD, et al. Cardiac rehabilitation barriers by rurality and socioeconomic status: a cross-sectional study. *Int J Equity Health*. 2013;12(1):1–81.
217. Arena R, Williams M, Forman DE, et al. Increasing referral and participation rates to outpatient cardiac rehabilitation: the valuable role of healthcare professionals in the inpatient and home health settings: a science advisory from the American Heart Association. *Circulation*. 2012;125(10):1321–1329. doi:10.1161/CIR.0b013e318246b1e5
218. Vitale G, Sarullo S, Vassallo L, et al. Prognostic value of the 6-min walk test after open-heart valve surgery: Experience of a Cardiovascular Rehabilitation Program. *J Cardiopulm Rehabil Prev*. 2018;38(5):304–308. doi:10.1097/HCR.0000000000000340
219. Gao Z, Chen Z, Sun A, Deng X. Gender differences in cardiovascular disease. *Med Nov Technol Devices*. 2019;4:100025. doi:10.1016/j.medntd.2019.100025
220. Taylor GH, Wilson SL, Sharp J. Medical, psychological, and sociodemographic factors associated with adherence to cardiac rehabilitation programs: a systematic review. *J Cardiovasc Nurs*. 2011;26(3):202–209. doi:10.1097/JCN.0b013e3181ef6b04
221. Gurewicz D, Prottas J, Bhalotra S, Suaya JA, Shepard DS. System-level factors and use of cardiac rehabilitation. *J Cardiopulm Rehabil Prev*. 2008;28(6):380–385. doi:10.1097/HCR.0b013e31818c3b5b
222. Shanmugasagaram S, Gagliese L, Oh P, et al. Psychometric validation of the cardiac rehabilitation barriers scale. *Clin Rehabil*. 2012;26(2):152–164. doi:10.1177/0269215511410579
223. Grace SL, Turk-Adawi K, Santiago de Araújo Pio C, Alter DA. Ensuring cardiac rehabilitation access for the majority of those in need: a call to action for Canada. *Can J Cardiol*. 2016;32(10 Suppl 2):S358–s364. doi:10.1016/j.cjca.2016.07.001
224. Orme MW, Clague-Baker NJ, Richardson M, Drewry S, Robinson TG, Singh SJ. Does cardiac rehabilitation for people with stroke in the sub-acute phase of recovery lead to physical behaviour change? Results from compositional analysis of accelerometry-derived data. *Physiotherapy*. 2020;107:234–242. doi:10.1016/j.physio.2019.10.003
225. Proudfoot C, Thow M, Rafferty D. A UK survey of phase I cardiac rehabilitation for patients with acute coronary syndrome. *Physiotherapy*. 2007;93(3):183–188. doi:10.1016/j.physio.2006.11.013
226. Kanazawa N, Iijima H, Fushimi K. In-hospital cardiac rehabilitation and clinical outcomes in patients with acute myocardial infarction after percutaneous coronary intervention: a retrospective cohort study. *BMJ open*. 2020;10(9):e039096. doi:10.1136/bmjopen-2020-039096
227. Altenhoener T, Leppin A, Grande G, Romppel M. Social inequality in patients' physical and psychological state and participation in rehabilitation after myocardial infarction in Germany. *Int J Rehabil Res*. 2005;28(3):251–257. doi:10.1097/00004356-200509000-00008
228. Reibis R, Jannowitz C, Halle M, Pittrow D, Gitt A, Voller H. Management and outcomes of patients with reduced ejection fraction after acute myocardial infarction in cardiac rehabilitation centers. *Curr Med Res Opin*. 2015;31(2):211–219. doi:10.1185/03007995.2014.977854
229. Miche E, Roelleke E, Zoller B, et al. A longitudinal study of quality of life in patients with chronic heart failure following an exercise training program. *Eur J Cardiovasc Nurs*. 2009;8(4):281–287. doi:10.1016/j.ejcnurse.2009.03.001
230. Vasic D, Novakovic M, Bozic Mijovski M, Barbic Zagar B, Jug B. Short-term water- and land-based exercise training comparably improve exercise capacity and vascular function in patients after a recent coronary event: a pilot randomized controlled trial. *Front Physiol*. 2019;10:903. doi:10.3389/fphys.2019.00903
231. Dehbarez NT, Lynggaard V, May O, Sogaard R. Learning and coping strategies versus standard education in cardiac rehabilitation: a cost-utility analysis alongside a randomised controlled trial. *BMC Health Serv Res*. 2015;15(1):1–10. doi:10.1186/s12913-015-1072-0
232. Marcassa C, Giordano A, Giannuzzi P. Five-year hospitalisations and survival in patients admitted to inpatient cardiac rehabilitation after cardiac surgery. *Eur J Prev Cardiol*. 2016;23(15):1609–1617. doi:10.1177/2047487316655452
233. Choo J, Burke LE, Pyo Hong K. Improved quality of life with cardiac rehabilitation for post-myocardial infarction patients in Korea. *Eur J Cardiovasc Nurs*. 2007;6(3):166–171. doi:10.1016/j.ejcnurse.2006.07.004
234. Russo N, Compostella L, Fadini G, et al. Prediabetes influences cardiac rehabilitation in coronary artery disease patients. *Eur J Prev Cardiol*. 2012;19(3):382–388. doi:10.1177/1741826711404503
235. Wichrowski M, Whiteson J, Haas F, Mola A, Rey MJ. Effects of horticultural therapy on mood and heart rate in patients participating in an inpatient cardiopulmonary rehabilitation program. *J Cardiopulm Rehabil*. 2005;25(5):270–274. doi:10.1097/00008483-200509000-00008

236. Spencer FA, Salami B, Yarzebski J, Lessard D, Gore JM, Goldberg RJ. Temporal trends and associated factors of inpatient cardiac rehabilitation in patients with acute myocardial infarction: a community-wide perspective. *J Cardiopulm Rehabil.* 2001;21(6):377–384. doi:10.1097/00008483-200111000-00006
237. Aldana SG, Whitmer WR, Greenlaw R, et al. Cardiovascular risk reductions associated with aggressive lifestyle modification and cardiac rehabilitation. *Heart Lung.* 2003;32(6):374–382. doi:10.1016/S0147-9563(03)00106-7
238. Malfatto G, Revera M, Branzi G, et al. A brief period of intensive cardiac rehabilitation improves global longitudinal strain and diastolic function after a first uncomplicated myocardial infarction. *Acta Cardiol.* 2017;72(3):284–291. doi:10.1080/00015385.2017.1305196
239. Reibis R, Voller H, Gitt A, et al. Management of patients with ST-segment elevation or non-ST-segment elevation acute coronary syndromes in cardiac rehabilitation centers. *Clin Cardiol.* 2014;37(4):213–221. doi:10.1002/clc.22241
240. Carless D, Douglas K, Fox K, McKenna J. An alternative view of psychological well-being in cardiac rehabilitation: considering temperament and character. *Eur J Cardiovasc Nurs.* 2006;5(3):237–243. doi:10.1016/j.ejcnurse.2006.03.004
241. Benzer W, Platter M, Oldridge NB, et al. Short-term patient-reported outcomes after different exercise-based cardiac rehabilitation programmes. *Eur J Cardiovasc Prev Rehabil.* 2007;14(3):441–447. doi:10.1097/HJR.0b013e32802bf7ae
242. Schwaab B, Zeymer U, Jannowitz C, Pittrow D, Gitt A. Improvement of low-density lipoprotein cholesterol target achievement rates through cardiac rehabilitation for patients after ST elevation myocardial infarction or non-ST elevation myocardial infarction in Germany: results of the PATIENT CARE registry. *Eur J Prev Cardiol.* 2019;26(3):249–258. doi:10.1177/2047487318817082
243. Russo N, Compostella L, Tarantini G, et al. Cardiac rehabilitation after transcatheter versus surgical prosthetic valve implantation for aortic stenosis in the elderly. *Eur J Prev Cardiol.* 2014;21(11):1341–1348. doi:10.1177/2047487313494029
244. Motohiro M, Yuasa F, Hattori T, et al. Cardiovascular adaptations to exercise training after uncomplicated acute myocardial infarction. *Am J Phys Med Rehabil.* 2005;84(9):684–691. doi:10.1097/01.phm.0000171167.31010.f4
245. Xu L, Zhao H, Qiu J, et al. The different effects of BMI and WC on organ damage in patients from a cardiac rehabilitation program after acute coronary syndrome. *Biomed Res Int.* 2015;2015:942695. doi:10.1155/2015/942695
246. Boitor M, Martorella G, Arbour C, Michaud C, Gélinas C. Evaluation of the preliminary effectiveness of hand massage therapy on post-operative pain of adults in the intensive care unit after cardiac surgery: a pilot randomized controlled trial. *Pain Manag Nurs.* 2015;16(3):354–366. doi:10.1016/j.pmn.2014.08.014
247. El-Kader SMA. Blood gases response to different breathing modalities in phase I of cardiac rehabilitation program after coronary artery bypass graft. *Eur J Gen Med.* 2011;8(2):85–91. doi:10.29333/ejgm/82706
248. Parsa A, Sadeghi M, Roghani F, Golshani J, Khani A, Yazdekhashti S. Effects of changes in myocardial dysfunction on quality of life in patients undergoing coronary angioplasty after cardiac rehabilitation. *Iran Heart J.* 2018;19(1):52–60.
249. Voller H, Gitt A, Jannowitz C, et al. Treatment patterns, risk factor control and functional capacity in patients with cardiovascular and chronic kidney disease in the cardiac rehabilitation setting. *Eur J Prev Cardiol.* 2014;21(9):1125–1133. doi:10.1177/2047487313482285
250. Gecaite J, Burkauskas J, Bunevicius A, Brozaitiene J, Kazukauskienė N, Mickuviene N. The association of cardiovascular reactivity during the Trier Social Stress Test with quality of life in coronary artery disease patients. *J Psychosom Res.* 2019;126:N.PAG–N.PAG. doi:10.1016/j.jpsychores.2019.109824
251. Twardella D, Kupper-Nybelen J, Rothenbacher D, Hahmann H, Wusten B, Brenner H. Short-term benefit of smoking cessation in patients with coronary heart disease: estimates based on self-reported smoking data and serum cotinine measurements. *Eur Heart J.* 2004;25(23):2101–2108. doi:10.1016/j.ehj.2004.08.017
252. Vasheghani-Farahani A, Asef-Kabiri L, Masoudkabir F, et al. Effect of exercise-based cardiac rehabilitation following coronary artery bypass surgery on ventricular repolarization indices. *J Cardiopulm Rehabil Prev.* 2011;31(4):239–244. doi:10.1097/hcr.0b013e318211e3c0
253. Bestehorn K, Wegscheider K, Voller H. Contemporary trends in cardiac rehabilitation in Germany: patient characteristics, drug treatment, and risk-factor management from 2000 to 2005. *Eur J Cardiovasc Prev Rehabil.* 2008;15(3):312–318. doi:10.1097/HJR.0b013e3282f40e14