

Oviducal gland transcriptomics of *Octopus maya* through physiological stages and the negative effects of temperature on fertilization

Oscar E. Juárez^{1,*}, Lousiana Arreola-Meraz^{1,*},
Edna Sánchez-Castrejón¹, Omar Hernando Avila-Poveda^{2,3},
Laura L. López-Galindo⁴, Carlos Rosas⁵ and Clara E. Galindo-Sánchez¹

¹ Departamento de Biotecnología Marina, Centro de Investigación Científica y de Educación Superior de Ensenada, Baja California, Ensenada, Baja California, México

² Facultad de Ciencias del Mar, Universidad Autónoma de Sinaloa, Mazatlán, Sinaloa, México

³ Programa Investigadoras e Investigadores por México, Consejo Nacional de Ciencia y Tecnología, Ciudad de México, México

⁴ Instituto de Investigaciones Oceanológicas, Universidad Autónoma de Baja California, Ensenada, Baja California, México

⁵ Unidad Multidisciplinaria de Docencia e Investigación - Sisal, Facultad de Ciencias, Universidad Nacional Autónoma de México, Sisal, Yucatán, México

* These authors contributed equally to this work.

ABSTRACT

Background: Elevated temperatures reduce fertilization and egg-laying rates in the octopus species. However, the molecular mechanisms that control the onset of fertilization and egg-laying in the octopus' oviducal gland are still unclear; and the effect of temperature on the expression of key reproductive genes is unknown. This study aims to better understand the molecular bases of octopus fertilization and egg-laying, and how they are affected by elevated temperatures.

Method: RNA-seq of oviducal glands was performed for samples before, during, and after fertilization and their transcriptomic profiles were compared. Also, at the fertilization stage, the optimal and thermal-stress conditions were contrasted.

Expression levels of key reproductive genes were validated *via* RT-qPCR.

Results: In mated females before egg-laying, genes required for the synthesis of spermine, spermidine, which may prevent premature fertilization, and the myomodulin neuropeptide were upregulated. Among the genes with higher expression at the fertilization stage, we found those encoding the receptors of serotonin, dopamine, and progesterone; genes involved in the assembly and motility of the sperm flagellum; genes that participate in the interaction between male and female gametes; and genes associated with the synthesis of eggshell mucoproteins. At temperatures above the optimal range for reproduction, mated females reduced the fertilization rate. This response coincided with the upregulation of myomodulin and APGW-amide neuropeptides. Also, genes associated with fertilization like LGALS3, VWC2, and Pcsk1 were downregulated at elevated temperatures. Similarly, in senescent females, genes involved in fertilization were downregulated but those involved in the metabolism of steroid hormones like SRD5A1 were highly expressed.

Submitted 8 October 2021

Accepted 16 January 2022

Published 30 March 2022

Corresponding author

Clara E. Galindo-Sánchez,

cgalindo@cicese.mx

Academic editor

Khor Waiho

Additional Information and
Declarations can be found on
page 25

DOI 10.7717/peerj.12895

© Copyright

2022 Juárez et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Genomics, Marine Biology, Molecular Biology, Zoology, Climate Change Biology
Keywords Egg-laying, Fertilization, Gamete recognition proteins, RNA-Seq, Thermal stress

INTRODUCTION

The endemic *Octopus maya* from the Yucatan Continental Shelf is becoming a good model to study the eco-physiological adaptations to environmental challenges, such as ocean warming (Juárez *et al.*, 2015, 2016; Sanchez-García *et al.*, 2017; López-Galindo *et al.*, 2019a; Meza-Buendía *et al.*, 2021). This ectothermic species is adapted to a narrow temperature range: reproductive events and embryonic development have an upper thermal limit around 27 °C; below that temperature—and optimally around 24 °C—these processes take place successfully. Therefore, reproductive events occur typically on winter and during upwelling pulses when sea temperatures are cooler on this region (Juárez *et al.*, 2015; Caamal-Monsreal *et al.*, 2016; Avila-Poveda *et al.*, 2016; Angeles-Gonzalez *et al.*, 2017; Sanchez-García *et al.*, 2017; Pascual *et al.*, 2019; López-Galindo *et al.*, 2019a, 2019b). However, higher temperatures have been reported in its distribution area in recent years, mainly in summer, which may negatively affect the physiological condition and reproductive success in *O. maya* males and females (Juárez *et al.*, 2015, 2016; Angeles-Gonzalez *et al.*, 2017; Pascual *et al.*, 2019; López-Galindo *et al.*, 2019a, 2019b). For instance, gene expression patterns in the testis of thermally stressed *O. maya* males were associated with physiological deficiencies and low motility of the spermatozoa (López-Galindo *et al.*, 2019b). In *O. maya* females, the exposure to temperatures above 27 °C significantly reduced the ova production, yolk amount, fertilization, and egg-laying rates (Juárez *et al.*, 2015). Moreover, the effect of thermal stress on mated females substantially reduced the hatchling survival, and hatchling growth rate. Therefore, the ability of *O. maya* females to store spermatozoa and delay fertilization until thermal conditions become favorable may be an adaptation that prevents low survival of the offspring at high temperatures (Juárez *et al.*, 2015, 2016). However, the molecular mechanisms controlling the onset of fertilization and egg-laying are still unclear.

Octopus maya is a semelparous species with only one reproductive event (egg-laying) in its lifetime and near the end of life when the female dies after the eggs hatch (Van Heukelem, 1983). In *Octopus*, the reproductive system consists of a gonad oval with two tubular oviducts and one oviducal gland (OvG) arranged halfway along each oviduct (Wells & Wells, 1977; Arkhipkin, 1992). In general terms, the OvG is involved in spermatozoa storage in spermathecae (Mangold, 1987; Marian, 2011, 2015), ova fertilization, and two critical activities: the production of cement by the peripheral gland and the cement polymerization in the central gland (Mangold, von Boletzky & Frösch, 1971; Froesch & Marthy, 1975; Wells & Wells, 1977). This cement is used to stick the eggs together in strings and attach them to the walls or roof of the female's shelter (Froesch & Marthy, 1975; Wells, 1978).

During mating, the passage of spermatozoa into the OvG towards the spermathecae does not imply fertilization; spermatozoa are stored in the spermathecae and used later in batches (Mangold, 1987; Marian, 2015; Sato, 2021). In wild and laboratory conditions, females continue mating and storing spermatozoa as the OvG enlarges (Froesch & Marthy,

1975; López-Galindo *et al.*, 2019a). Fertilization is internal: mature ova descend one by one from their follicular sheath to the proximal oviduct; simultaneously, spermatozoa are released from the spermathecae into the oviduct of the OvG for fertilization to occur (Mangold, 1987). Then, the egg-laying female stops hunting and feeding, to dedicate herself exclusively to the care and protection of her egg mass; therefore, the onset of egg-laying depends on the energetic budget of the female, because she needs enough energy reserves to perform the oviposition and egg incubation—which takes around 45 days—without food (O’Dor & Wells, 1978; Di Cristo, 2013; Juárez *et al.*, 2015; Meza-Buendía *et al.*, 2021). Egg-incubation coincides with the onset of senescence, where a general physiological decline and diseases occur, as part of a natural process that precedes the octopus death; females die just after the newborns’ hatch (Anderson, Wood & Byrne, 2002). Physical deterioration of distinct organs occurs in the senescent female’s body, including the depletion of reproductive organs, where the OvG drastically shrinks (Olivares *et al.*, 2017).

The OvG plays a crucial role in the reproductive strategy, thus deserves more consideration in the study of octopus reproductive success (Olivares *et al.*, 2017; Sato, 2021). In the present study, the transcriptomic profiles of OvG at different physiological stages were analyzed to understand the molecular mechanisms controlling the onset of fertilization and egg-laying in *O. maya*. We also evaluated the effect of temperature on the expression of key reproductive genes at the fertilization stage.

MATERIALS AND METHODS

Ethics statement

In this study, octopuses were anesthetized with ethanol 3% in seawater at experimental temperatures (Estefanell *et al.*, 2011; Gleadall, 2013) to enable humane killing (Andrews *et al.*, 2013) in consideration of ethical protocols (Mather & Anderson, 2007), and the animals’ welfare during manipulations (Moltschaniwskyj *et al.*, 2007; Winlow *et al.*, 2018), all previous takes into account the nociception in aquatic invertebrates (Sneddon, 2015). Our protocols were approved by the experimental Animal Ethics Committee of the Faculty of Chemistry at Universidad Nacional Autónoma de México (Permit number: Oficio/FQ/CICUAL/099/15) and following the ethical recommendations for the humanitarian killing of animals as established under Mexican law (NOM-033-SAG/ZOO-2014 derogating to NOM-033-ZOO-1995) and following as closely as possible the five R’s Principle in invertebrates (Crespi-Abril & Rubilar, 2021).

Capture and acclimation of octopuses

Female ($n = 30$) and male ($n = 30$) octopuses with a body mass that ranged between 400–600 g were captured off the coast of Sisal Yucatán, México, by an artisanal fishing fleet. This *O. maya* size range assumes that they are reproductively mature, and even some females have already collected sperm in their OvG during the process of maturation (Avila-Poveda *et al.*, 2016; Angeles-Gonzalez *et al.*, 2017; Markaida, Méndez-Loeza & Rosales-Raya, 2017). Captured octopuses were transported to the Experimental Cephalopod Production Unit at the Unidad Multidisciplinaria de Docencia e Investigación

(UMDI-UNAM), Sisal, Yucatan, Mexico. [Figure 1](#) (on top) shows how the octopuses were acclimated for 10 d in three outdoor ponds (6 m diameter) containing aerated natural seawater (oxygen level maintained to $5.5 \pm 0.5 \text{ mg L}^{-1}$) at $25 \pm 1 \text{ }^\circ\text{C}$ and a rate of 20 individuals per pond with a sex ratio of 1:1 ([Rosas et al., 2014](#)) following the acclimation protocol of [López-Galindo et al. \(2019a\)](#). Octopuses were observed daily at 6:00 AM, 12:00 PM, and 6:00 PM for 1 h. Mating was observed since the first day of acclimation which was confirmed by observing a male inserting his hectocotylus (mating arm) into the female mantle cavity to deposit spermatophores. During the 10-d acclimation period, all the females mated at least once.

Experimental design and thermal conditions

[Figure 1](#) illustrates the thermal experimental design and oviducal gland dissection after the acclimation, which are subsequently described in detail: on day 11, mated females were distributed randomly (coin toss) into two recirculating aquaculture systems, one for the Control condition ($n = 15$) and another for the thermal challenge ($n = 15$). Females were individually reared in 80 L tanks for 40 d in both Control and Treatment conditions (from day 11 to 51). Each tank contained a fiberglass box that served as a refuge and for egg-laying. Physio-chemical parameters (dissolved oxygen, temperature, salinity, and pH) of individual tanks were monitored twice daily to ensure homogeneous conditions within each treatment and minimize potential confounders. In the Control condition, females were reared at $24 \text{ }^\circ\text{C}$, which is the best condition for egg-laying ([Rosas et al., 2014](#); [Juárez et al., 2015](#)). The thermal challenge consisted of a high-temperature shock treatment with a temperature decrease (HTST-TD), which was performed following the protocol of [Juárez et al. \(2015\)](#), where mated females were initially exposed to a stress temperature of $30 \text{ }^\circ\text{C}$ for 10 d (from day 11 to day 21), then they were exposed 5 d at temperatures from 29 to 24 (decreasing of $1 \text{ }^\circ\text{C}$ every 5 days) ([Fig. 1](#)). After the heat shock of $30 \text{ }^\circ\text{C}$, the system was cooled down to induce the egg-laying ([Juárez et al., 2015](#)).

Egg-laying and fertilization rates

Through the experiment, the average number of eggs per spawn was calculated and the fertilization was visually inspected in each treatment following the protocol described in [Juárez et al. \(2015\)](#) including all the females that laid eggs in each experimental condition. The egg-laying rates were normalized by dividing the number of eggs per spawn by the females' weight in grams. Fertilization rates were expressed as the proportion of eggs containing embryos at the end of the incubation process, from the total eggs laid per female. Statistical differences in egg-laying rates and fertilization rates between the Control and the HTST-TD were evaluated by implementing Student t-tests with a statistical significance of $P < 0.05$.

Sampling and dissection of oviducal glands

We replicated the sampling schedule implemented by [Juárez et al. \(2015\)](#): sampling began after 2 weeks of exposure to the experimental conditions, on day 26. Females that started the egg-laying before that day were not considered for further analysis. In the Control




Figure 1 Schematic drawing of the acclimation, thermal experimental design, and oviducal gland dissection in *O. maya*. MAT24 for mated females at 24 °C, FER24 for fertilization in females at 24 °C, SEN24 for senescent females at 24 °C and, FER-TD for fertilization in females from the high-temperature shock treatment with temperature decrease (HTST-TD).

Full-size DOI: 10.7717/peerj.12895/fig-1

condition, four females in the mated stage (MAT24, $n = 4$) were sampled on day 26. The mated stage was characterized by an OvG that received spermatophores and stored spermatozoa in the spermathecae, according to the previous observation of the male(s)

inserting the hectocotylus arm into the female's mantle cavity. The onset of egg-laying—which coincides with fertilization and precedes the senescence—was unpredictable for each female. Therefore, a female was considered at the fertilization stage as soon as the egg-laying was detected during the daily monitoring. At that moment, females were sheltered in their nests and stopped feeding, whereas the OvG was presumably releasing sperm from the spermatheca and producing secretions from its glands. The next egg-laying female detected was sampled 1 week after she started laying her eggs, which represented the senescence stage. This stage was characterized by an OvG that entered in physiological (without evident secretions) and anatomical deterioration due to a phase of programmed cell death; besides, females were also near the end of their life. This was repeated until obtaining four samples for each stage (FER24, $n = 4$; SEN24, $n = 4$).

In the HTST-TD, females observed in the fertilization stage (FER-TD, $n = 4$) were sampled from day 26 to day 51 (Fig. 1). Before OvG dissection, animals were anesthetized by keeping them in ethanol-seawater (3%) solution for up to 4 min as indicated in the Ethics statement. Subsequently, the reproductive system was dissected from each octopus for each experimental condition (MAT24, FER24, SEN24, and FER-TD) and OvG samples were cut from the peripheral gland, central gland, and spermatheca area, which were mixed trying to obtain representative samples of the whole oviducal gland (Fig. 1 at the bottom), and finally preserved in RNAlater solution (Thermo Fisher) at 4 °C. Once all the samples were obtained, they were sent to the Laboratorio de Genómica Funcional de Organismos Marinos of CICESE in Ensenada, BC, Mexico where they were stored at -70 °C until RNA extraction.

RNA sequencing

RNA extraction, quantitation, and quality check were performed using the protocol, reagents, and instruments described in [López-Galindo et al. \(2019b\)](#) and [Juárez et al. \(2019\)](#), starting with 20–30 mg of every whole gland sample. For each experimental condition (MAT24, FER24, SEN24, and FER-TD), a pooled sample was prepared which consisted of 100 ng of RNA from four different individuals ($n = 4$). Paired-end libraries of complementary DNA (cDNA) were prepared for each pool using the TruSeq DNA Sample Preparation Kit v2 (Illumina, San Diego, CA, USA), following the manufacturer's protocol. Libraries were sequenced in the MiSeq system (Illumina, San Diego, CA, USA) to obtain reads of 150 bp long. Libraries and sequencing were conducted without the knowledge of treatment allocation of the pooled samples (blinding).

Bioinformatic workflow

The quality reports of raw sequence data were obtained with FastQC v0.11.6 (Babraham Bioinformatics, Babraham, UK). Low-quality reads, ambiguous nucleotides, and sequencing adaptors were removed using Trimmomatic v0.35 software ([Bolger, Lohse & Usadel, 2014](#)). The transcriptome was *de novo* assembled using Trinity v2.4.0 ([Grabherr et al., 2011](#)) and was deposited at DDBJ/EMBL/GenBank database under the accession [GJEO00000000](#). TransDecoder ([Haas et al., 2013](#)) was implemented to predict the open reading frames (ORFs) and coding sequences (CDS) of each transcript (with a minimum

length of 50 amino acids), which were annotated using BLASTx searches ([Camacho et al., 2009](#)) in the UniProt database. Differential gene expression was analyzed implementing pairwise comparisons among the physiological stages in the Control condition (*i.e.*, MAT24 to FER24, MAT24 to SEN24, FER24 to SEN24), then the representative DEGs for each stage included those obtained against the other two stages; for example, representative DEGs of MAT24 included those obtained against FER24 plus those against SEN24, removing the redundancy. The fertilization stages of the Control and the HTST-TD were compared (FER24 against FER-TD); for this, Bowtie2 v2.3.2 ([Langmead & Salzberg, 2012](#)) was used to align the reads of each library on the assembled transcriptome; RSEM v1.3.0 ([Li & Dewey, 2011](#)) to quantify transcript abundance of each library; and edgeR (Bioconductor) for identification of differential expression at isoform level ([Robinson, McCarthy & Smyth, 2009](#)). UniProt IDs of differentially expressed genes (DEGs) were used for enrichment analysis of gene ontology (GO) terms and metabolic pathways from the Kyoto Encyclopedia of Genes and Genomes (KEGG) ([Kanehisa et al., 2012](#)) with DAVID v6.8 ([Huang, Sherman & Lempicki, 2009](#)). DEGs included in the best-represented GO and KEGG categories and those associated with the regulation of OvG activity ([Di Cosmo, Di Cristo & Paolucci, 2001](#); [Iwakoshi-Ukena et al., 2004](#); [Di Cristo & Di Cosmo, 2007](#); [Minakata et al., 2009](#)) were selected for cluster analysis and plotted in heatmaps using R software ([R Development Core Team, 2010](#)). The analyses were performed following the scripts and parameters set and statistical significance implemented by [Juárez et al. \(2019\)](#) and [López-Galindo et al. \(2019b\)](#).

Validation of gene expression *via* RT-qPCR

To validate expression levels found by bioinformatic methods, DEGs with relevant reproductive functions were selected to estimate their expression by using RT-qPCR. Five potential reference genes were evaluated: V-type proton ATPase subunit d (VATD), Elongation factor 1-beta (EF1 β), Gelsolin-like protein 2, Heterogeneous nuclear ribonucleoprotein D (HNR), and Ribosomal Protein L6 (RPL6). Specific primers for the target and reference genes were designed using Primer3 ([Untergasser et al., 2012](#)). For each stage, additional replicate samples were obtained from animals under the same experimental conditions MAT24 ($n = 4$), FER24 ($n = 3$), SEN24 ($n = 3$), FER-TD ($n = 3$), including MAT-TD ($n = 5$) and SEN-TD ($n = 3$). RNAs were extracted as mentioned previously and treated with the RQ1 RNase-free DNase (Promega, Madison, WI, USA) according to the manufacturer's protocol. The cDNA was synthesized using the Improm II Reverse Transcription System (Promega, Madison, WI, USA) following the manufacturer's instructions starting with 2 μ g of RNA of each sample. The primer amplification efficiency was estimated following the procedure of [Bustin et al. \(2009\)](#). The RT-qPCR reactions were performed by triplicate following the protocol of [López-Galindo et al. \(2019b\)](#) without knowledge of group allocation of the samples (blind analysis). The stability of reference genes was evaluated using RefFinder ([Xie et al., 2012](#)). The relative expression of target genes was estimated following the method proposed by [Hellemans et al. \(2007\)](#). To assess if gene expression levels were significantly affected by the OvG physiological stages and the HTST-TD condition a two-way ANOVA was performed

Table 1 Sequencing and reads assembly of cDNA libraries from oviducal glands of *Octopus maya* at different conditions.

Libraries	Number of paired reads
MAT24	5,872,593
FER24	4,931,726
SEN24	5,507,769
FER-TD	5,352,396
Average	5,416,121
Total	21,664,484
Passed QC filter	19,969,819
Transcriptome assembly	quantity
Contigs	61,575
Coding sequences	36,136
Contig length N50, nucleotides	593
Total assembled nucleotides	32,348,437

Note:

MAT, mated; FER, fertilization; SEN, senescence; number 24 indicates 24 °C; TD, heat shock treatment with temperature decrease.

followed by Tukey's HSD test. A Spearman correlation for RNA-seq and RT-qPCR expression values was performed. Analyses were performed using STATISTICA 8.0 (StatSoft, Tulsa, OK, USA), and a statistically significant difference was accepted at $P < 0.05$.

RESULTS

In the Control condition, all females laid eggs at a temperature of 24 ± 1 °C, while in the HTST-TD, females did not lay eggs at temperatures around 30 °C, they did it until the rearing system reached temperatures below 28 °C at an average temperature of 26.1 °C. Significant differences in the egg-laying rates ($P = 8.12E-5$) and fertilization rates ($P = 0.0012$) were obtained between the Control and the HTST-TD. In the Control condition, the average number of eggs laid was three times higher and the average number of eggs laid per female's mass (grams) was two times higher than those observed in the HTST-TD. The average fertilization rate in the HTST-TD was 40.6 with an SD of 26.7, while in the Control condition the average fertilization rate was 72.5 with an SD of 4.0.

Sequencing and transcriptome assembly

The RNA sequencing generated a total of 21,664,484 paired reads, with an average of 5,416,121 paired reads per library. The raw sequence reads were deposited in the NCBI-SRA database (accession numbers: [SRR13512014](#)–[SRR13512017](#)). After the quality filter, 19,969,819 paired reads survived (92.17%), which were utilized for transcriptome assembly. The transcriptome consisted of 61,575 contigs with N50 of 593 and 32,348,437 assembled bases. A total of 36,136 coding sequences (CDS) were detected (Table 1).




Figure 2 Diagrams representing the number of upregulated transcripts in pairwise comparisons among (A) physiological stages; and (B) between thermal conditions of *O. maya* oviducal glands. Full-size DOI: 10.7717/peerj.12895/fig-2

Differential expression analysis

In the Control condition, a total of 1,719 transcripts were differentially expressed including all the pairwise comparisons among the physiological stages, of which, 210 showed the highest expression in MAT24, 626 in FER24, and 633 in SEN24. Gene upregulation was shared between stage pairs: 209 transcripts by the MAT24-FER24 pair, 11 transcripts by MAT24 and SEN24, and 30 transcripts by the FER24-SEN24 pair. In the comparison between FER24 and FER-TD, 93 transcripts showed differential expression, of which 47 showed higher expression in FER24 and 46 in FER-TD (Fig. 2).

Functional annotation and enrichment analysis

A total of 15,145 transcripts showed significant BLASTx-hit with the UniProt database, of which 12,678 obtained GO annotations. These annotations were analyzed to find enriched biological processes in the transcriptome (Table 2). The higher enrichment was

Table 2 Functional enrichment of biological processes for the whole oviducal gland transcriptome of *Octopus maya*.

GO	Term	P-value	FE
GO:0060285	cilium-dependent cell motility	1.39E-04	5.51
GO:0035082	axoneme assembly	7.24E-04	4.01
GO:0000722	telomere maintenance <i>via</i> recombination	4.69E-04	3.08
GO:0006383	transcription from RNA polymerase III promoter	6.50E-04	2.99
GO:0075733	intracellular transport of virus	5.96E-04	2.53
GO:0006396	RNA processing	4.18E-06	2.42
GO:0090630	activation of GTPase activity	1.50E-04	2.30
GO:0006283	transcription-coupled nucleotide-excision repair	4.20E-04	2.25
GO:0016925	protein sumoylation	2.09E-04	2.01
GO:0060271	cilium morphogenesis	5.97E-05	2.01

Note:

GO, gene ontology; FE, fold enrichment.

obtained for cilium-dependent cell motility and axoneme assembly, including genes like SPEF2, TTLL1, TTLL3, DRC1, CCDC39, RSPH4A, DNAH1, SPAG16, and LRGUK.

In mated females of the Control condition (MAT24), DEGs enriched biological processes like the fat-soluble vitamin metabolic process, IMP biosynthetic process, regulation of mRNA splicing *via* spliceosome, and cellular biogenic amine metabolic process. Upregulation was detected on genes encoding proteins like spermidine synthase, diamine acetyltransferase 2, kynurenine formamidase, low-density lipoprotein receptor-related protein 2, and the myomodulin neuropeptide; and downregulation was observed on genes encoding the disks large-associated protein 1, brorin, and the 5-hydroxytryptamine receptor 1 (Table 3).

DEGs obtained for FER24 enriched biological processes like the regulation of calcineurin-NFAT signaling cascade, cellular water homeostasis, regulation of male germ cell proliferation, cellular hormone metabolic process, regulation of reproductive process, extracellular matrix organization, and glycosylation. Upregulation was detected on genes encoding aquaporin, calcium and integrin-binding protein 1, chorion peroxidase, dopamine receptor 1, FMRamide-activated amiloride-sensitive sodium channel, peroxiredoxin-4, protein catecholamines up, 5-hydroxytryptamine receptor 1, and 16 genes included in the glycoprotein biosynthetic process GO term (Table 4), like the beta-1,4-N-acetylgalactosaminyltransferase bre-4; by contrast, downregulation was observed on genes encoding the 3-oxo-5-alpha-steroid 4-dehydrogenase 1, polypyrimidine tract-binding protein 1, and the myomodulin neuropeptide.

In SEN24, DEGs enriched biological processes like the cellular hormone metabolic process, iron ion homeostasis, cellular response to growth factor stimulus, lipid metabolic process, and oxidation-reduction process. Upregulation was observed on the retinoid-inducible serine carboxypeptidase, retinol dehydrogenase 14, 3-oxo-5-alpha-steroid 4-dehydrogenase 1, and matrix metalloproteinase-19; while downregulation was observed on genes encoding the chorion peroxidase, galectin-3, protein catecholamines up,

Table 3 Functional enrichment for differentially expressed genes in oviducal glands of *Octopus maya* at each condition.

GO term	FE	P-value
MAT24		
fat-soluble vitamin metabolic process	30.28	3.83E-03
IMP biosynthetic process	20.18	8.90E-03
negative regulation of RNA splicing	13.76	4.01E-04
cellular biogenic amine metabolic process	12.75	3.44E-03
cellular amine metabolic process	9.18	1.95E-03
regulation of mRNA splicing, <i>via</i> spliceosome	6.49	2.15E-03
extracellular structure organization	4.07	3.32E-03
alpha-amino acid metabolic process	3.41	4.68E-03
nucleotide biosynthetic process	3.32	1.00E-02
DNA replication	3.06	8.77E-03
FER24		
positive regulation of calcineurin-NFAT signaling cascade	31.33	2.96E-03
regulation of male germ cell proliferation	18.80	9.45E-03
positive regulation of cholesterol efflux	18.80	9.45E-03
cell volume homeostasis	13.05	4.04E-04
water homeostasis	12.53	3.23E-03
hydrogen peroxide catabolic process	11.39	4.34E-03
phagocytosis, engulfment	9.64	7.17E-03
protein hydroxylation	9.64	7.17E-03
serine family amino acid biosynthetic process	8.95	8.92E-03
retinoid metabolic process	7.23	1.19E-03
muscle organ morphogenesis	6.53	6.42E-03
cellular hormone metabolic process	4.95	6.65E-03
positive regulation of cell-substrate adhesion	4.70	8.28E-03
regulation of reproductive process	4.14	6.40E-03
cellular amino acid catabolic process	3.86	2.11E-03
extracellular structure organization	3.69	9.70E-05
extracellular matrix organization	3.30	9.64E-04
reactive oxygen species metabolic process	3.03	9.28E-03
glycosylation	2.83	7.98E-04
glycoprotein metabolic process	2.42	1.70E-03
SEN24		
regulation of microvillus organization	58.80	8.40E-04
cellular hormone metabolic process	9.28	4.09E-04
retinoid metabolic process	9.05	9.22E-03
iron ion homeostasis	8.91	2.18E-03
transition metal ion transport	7.06	1.47E-03
hormone metabolic process	5.35	1.84E-03
cellular response to growth factor stimulus	3.27	3.18E-04
lipid metabolic process	1.99	3.69E-03

(Continued)

Table 3 (continued)

GO term	FE	P-value
FER24 vs FER-TD		
extracellular matrix disassembly	26.13	7.08E-02
plasma membrane organization	7.56	5.45E-02
regulation of cell adhesion	5.75	8.79E-02
single-organism membrane organization	3.45	9.63E-02
signaling	1.72	7.87E-02

Note:

GO, gene ontology; FE, fold enrichment; MAT24, mated; FER24, fertilized; SEN24, senescence; all at 24 °C. FER-TD, fertilized and exposed to the heat shock treatment with temperature decrease.

Table 4 Transcripts corresponding to the glycoprotein biosynthesis process upregulated in the FER24 condition.

Transcript ID	UniProt ID	Gene name	Protein name
TRINITY_DN43413_c0_g1_i1	ALG8_HUMAN	ALG8	Probable dolichyl pyrophosphate Glc1Man9GlcNAc2 alpha-1,3-glycosyltransferase
TRINITY_DN13830_c0_g1_i5	B3GN5_PIG	B3GNT5	Lactosylceramide 1,3-N-acetyl-beta-D-glucosaminyltransferase
TRINITY_DN13397_c0_g2_i3	BRE4_CAEBR	bre-4	Beta-1,4-N-acetylgalactosaminyltransferase bre-4
TRINITY_DN5237_c0_g1_i1	CANT1_HUMAN	CANT1	Soluble calcium-activated nucleotidase 1
TRINITY_DN16173_c0_g2_i10	D19L1_HUMAN	DPY19L1	Probable C-mannosyltransferase DPY19L1
TRINITY_DN16072_c0_g1_i2	EDEM2_HUMAN	EDEM2	ER degradation-enhancing alpha-mannosidase-like protein 2
TRINITY_DN16442_c0_g2_i1	FUCTA_DROME	FucTA	Glycoprotein 3-alpha-L-fucosyltransferase A
TRINITY_DN14002_c0_g1_i1	G3ST2_MOUSE	Gal3st2	Galactose-3-O-sulfotransferase 2
TRINITY_DN11335_c0_g1_i1	GALT9_CAEEL	gly-9	Probable N-acetylgalactosaminyltransferase 9
TRINITY_DN3280_c0_g1_i1	GCNT1_MOUSE	Gcnt1	Beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase
TRINITY_DN10779_c0_g1_i1	GOGA2_RAT	Golga2	Golgin subfamily A member 2
TRINITY_DN16242_c0_g1_i8	LRP2_HUMAN	LRP2	Low-density lipoprotein receptor-related protein 2
TRINITY_DN29933_c0_g1_i1	MGT4B_DANRE	mgat4b	Alpha-1,3-mannosyl-glycoprotein 4-beta-N-acetylglucosaminyltransferase B
TRINITY_DN11201_c0_g1_i1	PMGT1_HUMAN	POMGNT1	Protein O-linked-mannose beta-1,2-N-acetylglucosaminyltransferase 1
TRINITY_DN10762_c0_g1_i1	STT3A_BOVIN	STT3A	Dolichyl-diphosphooligosaccharide-protein glycosyltransferase subunit STT3A
TRINITY_DN7997_c0_g1_i1	TMM59_MOUSE	Tmem59	Transmembrane protein 59

Note:

These transcripts were upregulated in oviducal glands of fertilized *Octopus maya* females in the Control condition (24 °C).

FMRamide-activated amiloride-sensitive sodium channel, and the cAMP-responsive element modulator. The top differentially expressed genes (DEGs) among the different physiological stages were plotted and clustered in a heatmap (Fig. 3). In this heatmap, the conditions MAT24 and FER-TD were grouped due to similar gene expression patterns.

In the comparison between FER24 vs FER-TD, DEGs enriched biological processes like the extracellular matrix disassembly, plasma membrane organization, regulation of cell adhesion, signaling, and cell communication. In FER-TD, we detected upregulation on genes encoding the CAD protein, matrix metalloproteinase-19, myomodulin neuropeptide, and tetraspanin-18; whereas strong downregulation was detected on the gene encoding the lactosylceramide 1,3-N-acetyl-beta-D-glucosaminyltransferase.




Figure 3 Heatmap of top DEGs among the physiological stages of oviducal glands of *O. maya*. Row labels indicate the UniProt IDs of the best BLASTx hit for each transcript. MAT24 for mated females at 24 °C, FER24 for fertilization in females at 24 °C, SEN24 for senescent females at 24 °C and, FERTD for fertilization in females from the HTST-TD. Values in $\log_2(\text{TPM}+1)$.

Full-size DOI: 10.7717/peerj.12895/fig-3

By contrast, FER24 showed higher expression of genes encoding brorin, galectin-3, epidermal retinol dehydrogenase 2, neuroendocrine convertase 1, and spectrin beta chain. The top DEGs between the Control and the HTST-TD conditions were plotted and clustered in a heatmap (Fig. 4). In this heatmap, the conditions SEN24 and FER-TD showed similar expression patterns and were grouped, another group consisted of MAT24 and FER24.




Figure 4 Heatmap of top DEGs from the comparison between *O. maya* oviducal gland samples of the Control condition against samples of the HTST-TD. Row labels indicate the UniProt IDs of the best BLASTx hit for each transcript. MAT24 for mated females at 24 °C, FER24 for fertilization in females at 24 °C, SEN24 for senescent females at 24 °C and, FERTD for fertilization in females from the HTST-TD. Values in $\log_2(\text{TPM}+1)$. [Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90_img.jpg\) DOI: 10.7717/peerj.12895/fig-4](https://doi.org/10.7717/peerj.12895/fig-4)

A third heatmap was constructed for transcripts related to the regulation of the OvG activity (Fig. 5). In this heatmap, stronger expression differences were evident in genes encoding the 5-hydroxytryptamine receptor 1 (5HT-7), cAMP-dependent protein kinase regulatory subunit (PKAR), cAMP-responsive element modulator (CREM), dopamine receptor 1 (Dop1R1), FMRFamide-activated amiloride-sensitive sodium channel (FanaCh), protein catecholamines up (Catsup), and steroid 17-alpha-hydroxylase/17,20 lyase (CYP17A1) showing all an expression peak at the fertilization stage (FER24); and




Figure 5 Expression heatmap for regulatory genes of oviducal glands' activity in *O. maya* at different conditions. Row labels indicate the UniProt IDs of the best BLASTx hit for each transcript. MAT24 for mated females at 24 °C, FER24 for fertilization in females at 24 °C, SEN24 for senescent females at 24 °C and, FERTD for fertilization in females from the HTST-TD. Values in $\log_2(\text{TPM}+1)$, ***DEGs (fold change > 2, false discovery rate < 0.05). [Full-size !\[\]\(fd7fe780e8fd8eece60268c87d0c3e04_img.jpg\) DOI: 10.7717/peerj.12895/fig-5](https://doi.org/10.7717/peerj.12895/fig-5)

myomodulin (MYOM) which was upregulated in MAT24 and FER-TD. In this heatmap, the conditions MAT24 and FER-TD were grouped due to similar expression patterns. Annotations of upregulated transcripts in each reproductive stage are shown in [Table S1](#). The list of all transcripts used in heatmaps including their gene products is shown in [Table S2](#).

Validation of gene expression by RT-qPCR

Three different gene expression patterns were represented: the upregulation in MAT24 was represented by the genes MYOM and SAT2; the upregulation in FER24 by Ptx, Dop1R1, PRDX4, and Catsup; and the upregulation in SEN24 by the genes MMP-19 and SRD5A1. Primer efficiency ranged from 92.1 to 109.8, the annealing temperatures,

Table 5 Primers for target and reference genes of *Octopus maya* oviducal glands used in RT-qPCR.

Transcript	Protein name	E	Tm	Size	Primers
TRINITY_DN22438_c0_g1_i1	3-oxo-5-alpha-steroid 4-dehydrogenase 1	109.8	60	119	CGGAAACCTAACGAAACAGG GACCAGCATGCAATAGCAAA
TRINITY_DN16362_c4_g6_i1	Chorion peroxidase	97.3	60	116	CGGCTTATCACGACGGTTAT GCATTTTGC GTTGAAAGGTT
TRINITY_DN14910_c0_g1_i2	Diamine acetyltransferase 2	90.5	60	127	ACCCACCTTCTGTTGATGATCT ACCAATGGTCCTGTGCTTAGT
TRINITY_DN12903_c1_g1_i1	Dopamine receptor 1	91.4	60	120	GGCTGTGACCTCGACATCAA GTGTTGCTGAGCCGTA CTGT
TRINITY_DN12984_c0_g1_i1	Matrix metalloproteinase-19	94	60	150	TGACGAAGAATGGACTGCAA TGAAATCCTTCACGAAACC
TRINITY_DN32740_c0_g1_i1	Myomodulin neuropeptides 1	97.2	62	118	GCAGTGGACCATTTCCTTGAT TTTTCGAAGCCATTTTGTCC
TRINITY_DN16150_c1_g1_i4	Peroxiredoxin-4	100.9	60	107	ATGGCCAAGATTCTGAAGGA CACCACGAAACAAAGGAGGT
TRINITY_DN15757_c0_g1_i2	Protein catecholamines up	98.4	60	114	TTGGGTCTGCGAGTCTTCTT AGCCATTCTCACAGCGAAGT
Reference (Xu & Zheng, 2018)	60S ribosomal protein L6	92.1	60	171	GGAAGGCACAAGGGAAAGCG CCTGGCTGGGATCTGAACCT
Reference (Juárez, 2016)	Elongation factor 1B	95.3	60	108	TGATGTCAAACCATGGGACG AGAGGTGCTAACTTGGACGC

Note:

E, primer efficiency; Tm, annealing temperature; Size, product size in base pairs.

product sizes, and sequences of each primer pair are shown in Table 5. The elongation factor 1 beta (EF1B) and the ribosomal protein L6 (RPL6) showed higher stability and were used as reference genes. Gene expression values estimated by RNA-Seq and RT-qPCR methods showed a significant Spearman correlation (P -value = 0.009). A scatter plot with linear regression representing the correlation between RNA-Seq and RT-qPCR gene expression values is provided as Fig. S1.

In the RT-qPCR estimation, the gene SAT2 showed a higher expression in the mated stage (MAT24) but the differences were not significant. The MYOM gene showed a significant effect for the interaction of factors ($P = 0.0304$) with higher expression in MAT24 and FER-TD (Fig. 6A). The genes with an expected peak in the fertilization stage showed significant effect for the “stage” factor (Catsup, $P = 0.0029$; Dop1R1, $P = 0.0064$; Ptx, $P = 0.0161$; PRDX4, $P = 0.0002$) with higher expression in fertilized females. However, the gene Catsup also showed a significant effect of the “treatment” factor ($P = 0.0219$) with higher expression in the Control (Fig. 6B). The genes with a bioinformatic peak in the senescence stage (SEN24) MMP-19 and SRD5A1 showed in the RT-qPCR a significant effect for the “stage” factor with higher expression in senescent females (both $P < 0.0001$); they also showed significant effect due to the interaction of factors ($P = 0.0284$ and 0.0001




Figure 6 Relative expression of target genes (RT-qPCR, average and standard deviation) at different conditions of oviducal glands of *O. maya*. Comparison between the Control temperature (24 °C) and HTST-TD. MAT, mated; FER, fertilization; SEN, senescence. Relative expression (fold change, FC) was calculated using the 60S ribosomal protein L6 and Elongation factor 1 B as reference genes (values in Log₁₀ scale). The expected gene expression patterns (A, B, and C) were estimated in the Control condition by bioinformatic methods.

Full-size DOI: [10.7717/peerj.12895/fig-6](https://doi.org/10.7717/peerj.12895/fig-6)

respectively), and SRD5A1 showed significant effect due to the “treatment” factor ($P = 0.0104$) with the highest expression in SEN24 (Fig. 6C).

DISCUSSION

Mated stage

Before the fertilization, the spermatozoa stored in the OvG must be reactivated (*De Lisa et al., 2013*). Important genes involved in the axoneme assembly and cilium motility such as SPEF2, TLL1, TLL3, DRC1, CCDC39, RSPH4A, DNAH1, SPAG16, and LRGUK were detected in the OvG transcriptome. Mutations in these genes have been associated with sperm immobility and male infertility, which suggests that they are essential for fertilization (*Leigh et al., 2009; Berg et al., 2011; Onoufriadis et al., 2014; Pereira et al., 2017; Wu et al., 2016; zur Lage, Newton & Jarman, 2019; Takeuchi et al., 2020*).

The expression of this set of genes in the OvG suggests that spermatozoa are stored in spermathecae without functional tails or disassembled and that flagellar components involved in sperm motility are reassembled there just before fertilization. Interestingly, in the spermathecae epithelium, spermatozoa are immobilized with their heads inserted within the mucosa, while the tails are not visible (*Di Cosmo, Di Cristo & Paolucci, 2001; Olivares et al., 2017*).

Females of *Octopus vulgaris* can store sperm in their OvG for up to 10 months (*Mangold, 1987*), which implies that during this prolonged period, the spermatozoa integrity depends on the protective molecular mechanisms of the female. For example, the upregulation of the AFMID gene, encoding the kynurenine formamidase, in mated females of *O. maya* may contribute to the protection of spermatozoa since it is important to eliminate toxic metabolites (*Dobrovolsky et al., 2005*).

In mated *O. maya* females, genes associated with the synthesis of the biogenic amines spermine and spermidine like SAT2 and SRM (*Chen et al., 2003; Wu et al., 2007*) were upregulated. In mammals, spermine and spermidine play a protective role for spermatozoa since they inhibit the glycation and fragmentation of sperm DNA in the epididymis (*Méndez & Sauer-Ramírez, 2018*). These biogenic amines also have a key role in the precise timing for successful fertilization; seminal spermine prevents premature capacitation and acrosome reaction (*Rubinstein & Breitbart, 1991*). Moreover, *in vitro* studies showed that spermine enhances the activity of seminal maltase, which increases glucose utilization by spermatozoa (*Sheth & Moodbidri, 1977*). High spermine and spermidine levels may be required in the OvG of mated females to maintain the integrity of spermatozoa and to inhibit fertilization until it is induced by environmental and metabolic signals. In this species, such a mechanism could guarantee a precise temporality for sperm activation and fertilization of eggs (*Wells & Wells, 1959; O’Dor & Wells, 1978; Di Cristo, 2013*). Interestingly, genes encoding the 5-hydroxytryptamine receptor 1 (serotonin receptor) and the gonadotropin-releasing hormone receptor (GnRHR) showed a lower expression at the mated stage (MAT24), suggesting that these receptors are not required before fertilization in the OvG under optimal thermal conditions.

The lifecycle of octopuses is characterized by a physiological transition from the growth to the reproductive phase: after reaching enough energy reserves during the growth phase,

the female stops feeding and dedicates exclusively to egg-laying and egg-care (O'Dor & Wells, 1978; Di Cristo, 2013). Such transition seems to be mediated by regulatory neuropeptides and catecholamines (Di Cristo, 2013; Wang & Ragsdale, 2018). These regulatory peptides are typically abundant in non-mated (growing) individuals which display hunting and active feeding (Wang & Ragsdale, 2018). Interestingly, the high expression of specific neuropeptides opposes certain reproductive events. In the OvG of *O. maya*, the myomodulin neuropeptide was upregulated in the MAT24 condition. The RT-qPCR estimation confirmed that, in optimal thermal conditions, a decrease in the expression of the MYOM gene coincided with the onset of fertilization. Myomodulin has been detected in mollusk of genera like *Aplysia* (Cropper et al., 1987, 1991; Brezina et al., 1995), *Lymnea* (Kellett et al., 1996), *Haliothis* (York et al., 2012), *Helix* (Greenberg et al., 1997), and *Sepia* (Zatylny-Gaudin et al., 2016). This neuropeptide has been associated with the modulation of feeding rates and muscle contractions (Cropper et al., 1987, 1991; Brezina et al., 1995; Kellett et al., 1996; Greenberg et al., 1997). More recently, a role of myomodulin in the regulation of egg-laying was suggested, since it was detected in the OvG of *Sepia officinalis* (Zatylny-Gaudin et al., 2016) and showed an upregulation before the spawning in *Haliothis asinina* females (York et al., 2012), which coincides with was found in the present study. Structurally, the oviducal glands consist of mixed layers of muscular and connective tissue vascularized and innervated (Peterson, 1959; Budelmann, Schipp & von Boletzky, 1997; Olivares et al., 2017; Anadón, 2019). Myomodulin is an important regulatory neuropeptide in the multi-messenger innervation of the sexual organs with muscle tissue in mollusks, including the glands (De Lange, Joosse & Van Minnen, 1998; Koene, 2010). Thus, it is possible that during the mated stage the OvG has greater vascularization and innervation, and therefore a higher expression of myomodulin, as was observed.

Fertilization stage

Downregulation of the myomodulin gene was observed in FER24 and strengthens the idea that the downregulation of some neuropeptides precedes reproductive events in *Octopus* (Wang & Ragsdale, 2018). In the OvG of *O. maya*, it seems that myomodulin must be downregulated for the onset of fertilization.

In the *O. vulgaris* female, fertilization is partially controlled by steroid hormones like progesterone and 17 β -estradiol, whose levels fluctuate through the reproductive phase (Di Cosmo et al., 1998; Di Cosmo, Di Cristo & Paolucci, 2001; Tosti et al., 2001). These hormones are associated with the growth and differentiation of the reproductive system, including the OvG, and the remobilization of spermatozoa (Di Cosmo, Di Cristo & Paolucci, 2001; Tosti et al., 2001). In the present study, a high expression of the Mpra gene was detected in the OvG of fertilized females (FER24); this gene encodes the membrane progesterin receptor alpha, which binds to progesterone (Zhu et al., 2003); this coincides with was found in *O. vulgaris* (Di Cosmo et al., 1998) and suggests that the role of sex steroids in the control of reproduction could be a more generalized adaptation among octopus species. Another endocrine gene upregulated in the fertilization stage was the CYP17A1, encoding the steroid 17-alpha-hydroxylase/17,20 lyase, which participates in

the metabolism and synthesis of steroid hormones (Auchus, Lee & Miller, 1998; Strauss, Modi & McAllister, 2014; Petrunak et al., 2014; Yoshimoto et al., 2016). These results suggest that OvGs metabolize and secrete steroid hormones to coordinate the activity of different organs for successful fertilization and egg-laying. Alternatively, Di Cristo & Di Cosmo (2007) proposed that sex steroids in the OvG may play a role in sustaining the production and secretion of the mucoproteins and mucopolysaccharides that coat the eggs. Authors also suggested that cyclic AMP (cAMP) modulates the secretory activity of the OvG (Di Cristo & Di Cosmo, 2007). In the present study, we detected upregulation on genes encoding the cAMP-dependent protein kinase regulatory subunit (PKAR) and cAMP-responsive element modulator (CREM) at the fertilization stage. This suggests a high secretory activity of the OvG during the fertilization stage. Moreover, the CREM gene has been associated with male fertility (Pati, Meistrich & Plon, 1999; Yanagimachi et al., 2004).

The expression of putative sperm genes was conspicuous during the fertilization stage. For instance, the Catsup gene encoding the protein catecholamines-up showed an upregulation in FER24. This protein is a zinc ion transmembrane transporter; involved in the zinc ion influx required for sperm capacitation and fertilization in mammals (Kerns, Zigo & Sutovsky, 2018; Kerns et al., 2018). Likewise, CIB1 and PRDX4 which are key genes for male fertility (Yuan et al., 2006; Iuchi et al., 2009), were upregulated at the fertilization stage. Furthermore, transcripts of the Drip gene, encoding aquaporin also showed upregulation in FER24. It is well known that aquaporins are essential for male fertility since they are responsible for the regulation of sperm volume, which is crucial for fertilization. They are also important for the osmotic adaptation of the spermatozoa to different microenvironments until reaching the egg, and for maintaining the osmotic homeostasis for both male and female gametes during fertilization (Delgado-Bermúdez, Ribas-Maynou & Yeste, 2022; Yeung et al., 2009; Ribeiro et al., 2021). The high expression of sperm genes during the fertilization stage is evidence of the synchronized reactivation of the sperm that was stored in the spermathecae.

A key step in fertilization (internal or external) is the interaction between male and female gametes (*i.e.*, gametic compatibility). The union of the spermatozoon to the extracellular matrix of the egg is mediated by gamete recognition proteins (GRPs) which have an important influence on the reproductive success of taxa with external or internal fertilization (Panhuis, Clark & Swanson, 2006; Kosman & Levitan, 2014). In our study, the ZAN gene encoding zonadhesin showed upregulation in the fertilization stage. Zonadhesin is a sperm protein that binds to the pellucid zone in a species-specific manner in mammals (Hardy & Garbers, 1995; Gao & Garbers, 1998; Bi et al., 2003; Tardif et al., 2010; Springate & Frasier, 2017). At the same time, LGALS3 which encodes galectin-3 was upregulated. Galectin-3 is a lectin that also participates in the binding of spermatozoa to the pellucid zone (Mei et al., 2019); therefore, in an analogous approach, ZAN and LGALS3 may cooperate in the union of male and female gametes in this octopus species.

After the fertilization, the egg envelope (chorion) hardens (Wells, 1978); this process is crucial to block polyspermy in internal fertilization, protecting the embryo from

mechanical damage and preventing bacterial infections ([Wang et al., 2021](#)). The hardening of the egg envelope requires the activity of the chorion peroxidase, encoded by the Ptx gene. This peroxidase is essential for the cross-link of chorion proteins and participates in chorion melanization ([Mindrinos et al., 1980](#); [Margaritis, 1985](#); [Li, Hodgeman & Christensen, 1996](#); [Han, Li & Li, 2000](#); [Konstandi et al., 2005, 2006](#); [Li & Li, 2006](#); [Wang et al., 2021](#)). In the present study, the Ptx gene peaked at the fertilization stage of *O. maya*.

Later, eggs are coated with a slime that is composed of mucoproteins secreted from the peripheral gland and a sulfonated mucopolysaccharide from the central gland ([Froesch & Marthy, 1975](#)). Mucoproteins are heavily glycosylated containing O-linked oligosaccharide chains that are covalently attached to serine or threonine residues of their polypeptide backbones ([Brockhausen & Stanley, 2015](#)). In these proteins, glycosylation begins with the addition of N-acetylgalactosamine by a large family of UDP-GalNac: polypeptide N-acetylgalactosaminyltransferases ([Clausen & Bennett, 1996](#)). In the present study, it was detected a set of 16 genes involved in the glycoprotein biosynthetic process and glycosylation process (ALG8, B3GNT5, bre-4, CANT1, DPY19L1, EDEM2, FucTA, Gal3st2, gly-9, Gcnt1, Golga2, LRP2, mgat4b, POMGNT1, STT3A, and Tmem59) including N-acetylgalactosaminyltransferases, showing upregulation in FER24. Therefore, this set of genes may be crucial for slime synthesis and important for normal egg-laying in *O. maya*.

Finally, the eggs enter the distal oviduct and are transported by peristalsis, one behind the other. In the Control condition, the receptors of serotonin (5HT-7), and dopamine (Dop1R1) were upregulated during the egg-laying in the OvG of *O. maya* females. Interestingly, serotonin and dopamine were identified as stimulating neurotransmitters that induce spawning in bivalves ([Arendse, Pitcher & Griffiths, 2018](#); [Gibbons & Castagna, 1984](#); [Braley, 1985](#); [Osada, Matsutani & Nomura, 1987](#); [Deguchi & Osada, 2020](#)). In this sense, the high expression of these receptors at the fertilization stage suggests that serotonin and dopamine are required to induce egg-laying also in *O. maya* females. Clusters of eggs and slime are then released; the female molds an egg string and fixes the end of the string to a suitable substrate ([Froesch & Marthy, 1975](#)).

Senescence stage

Once fertilization and egg-laying were finalized, there was a downregulation of important reproductive genes mentioned above, including Ptx, Catsup, FanaCh, CREM, and LGALS3. On the other hand, one of the genes with the highest expression in SEN24 was the SRD5A1. This gene encodes the 3-oxo-5-alpha-steroid 4-dehydrogenase 1, which participates in the metabolism of steroid hormones like progesterone. The expression of SRD5A1 highly increased from the fertilization to senescence stage suggesting that fertilization and egg-laying may terminate due to an enzymatic depletion of progesterone. On the other hand, this enzyme participates in the synthesis of allopregnanolone, a neuroactive metabolite of progesterone that acts in the brain ([Tsutsui & Haraguchi, 2016](#)). In senescent *O. maya* females, the OvGs may release this steroid into the bloodstream to reach the central nervous system.

Another gene highly expressed in senescent females was the MMP-19 encoding the matrix metalloproteinase-19, which participates in the degradation of the extracellular matrix. This gene has been associated with wound healing and tissue remodeling (Nagase & Murphy, 2013), which may be part of the natural senescence process. In senescent females of *O. vulgaris*, *O. mimus*, and *O. maya* (present study), the OvG notably shrank compared to the previous fertilization stage (Cuccu et al., 2013; Olivares et al., 2017), which may imply a tissue remodeling process.

Effect of high temperatures on fertilization stage

FER-TD and MAT24 were grouped in the cluster analyses for DEGs among the stages, and a set of regulatory genes. In these genes, the expression levels in FER-TD resemble those of the previous stage (MAT24) of the Control, suggesting that certain processes were delayed due to the HTST-TD. This supports the idea that octopus females under thermal stress can delay certain reproductive processes until temperatures become favorable (Juárez et al., 2015).

In the cluster analysis of DEGs between the Control and the HTST-TD, which are those directly affected by the thermal stress, FER-TD was grouped with the senescence condition SEN24 of the Control. This grouping may be related to a high physiological deterioration of the gland in FER-TD, which resembles that of the senescence stage. There was no grouping between FER24 and FER-TD in any dendrogram, confirming that thermal stress drastically affected gene expression in the oviducal glands during the fertilization stage.

In *O. maya*, the temperature is an important modulator of fertilization and egg-laying rates. These rates are significantly reduced by temperatures above 27 °C (Juárez et al., 2015), and in the males, temperatures around 30 °C provoke damages in testis and spermatozoa (López-Galindo et al., 2019a, 2019b). Gene expression changes in the OvG caused by temperature may be the cause of the reduced fertilization and egg-laying rates observed in this species.

Previously, we discussed the inhibitory role for fertilization of the myomodulin neuropeptide in MAT24, and that the MYOM gene must be downregulated to initiate fertilization. However, under thermal stress this gene was not downregulated during the fertilization stage, instead, it was highly expressed. In gastropods, this bioactive neuropeptide potentiates muscular contractions (Cropper et al., 1987; Kellett et al., 1996; Greenberg et al., 1997). Similarly, the gene encoding the APGW-amide neuropeptide (cerebral peptide 1) significantly increased its expression under thermal stress. This neuropeptide also potentiates muscle contractions in bivalves and gastropods (Minakata et al., 1991; Henry, Zatylny & Favrel, 2000), and in the OvG of *O. vulgaris*, where a function in the oviduct contractility was proposed (Di Cristo & Di Cosmo, 2007). Therefore, we can hypothesize that in *O. maya* the coordinated release of myomodulin and APGW-amide may keep the proximal oviduct contracted for longer periods, limiting the passage of eggs from the ovary to the OvG, thus reducing the fertilization rate under high temperatures as was observed in this octopus species (Juárez et al., 2015). This hypothesis should be evaluated in future research.

Essential genes for successful fertilization were downregulated in the HTST-TD; for instance, the gene LGALS3, which participates in the union of the spermatozoon to the extracellular matrix of the egg (Mei et al., 2019) showed high levels in the Control condition but was downregulated in the HTST-TD. Similarly, the gene VWC2, encoding the brorin protein, was highly expressed in the optimal temperature and downregulated in the HTST-TD; brorin is also involved in cell adhesion (Manabe et al., 2008).

Although the activity of OvG is partially regulated by steroid hormones (Di Cosmo et al., 1998; Di Cosmo, Di Cristo & Paolucci, 2001), our results suggest that these glands also play a role in the biosynthesis and release of hormones. In this regard, under the optimal temperature, an upregulation of the Pcsk1 gene was detected, while it was downregulated in the HTST-TD. This gene encodes the neuroendocrine convertase 1 and is involved in the conversion of secretory precursor proteins to bioactive polypeptides (Morash, Soanes & Anini, 2011). According to our results, the processing of prohormones in the OvG of *O. maya* females is related to fertilization, and the downregulation of the Pcsk1 gene could modulate this process under thermal stress. Another gene with strong downregulation in FER-TD was the B3GNT5, which encodes the Lactosylceramide 1,3-N-acetyl-beta-D-glucosaminyltransferase. In mice, the knock-out of this gene provokes a series of reproductive defects, therefore it is a key gene for successful reproduction (Kuan et al., 2010).

Ecological implications of a temperature-driven inhibition of fertilization in *O. maya*

Elevated temperatures caused gene expression changes in the oviducal gland that may be associated with the low fertilization and egg-laying rates observed in this octopus species (Juárez et al., 2015). The negative regulation of fertilization rates under elevated temperatures may be a strategy that prevents excessive thermal stress for embryos and hatchlings, which can improve populations' fitness (Juárez et al., 2015, 2016; Caamal-Monsreal et al., 2016); but at the same time, such a mechanism makes this species vulnerable to ocean warming and interannual thermal anomalies. Stational upwelling pulses modulate the thermal conditions at the Northeast region of the peninsula, maintaining suitable temperatures for octopus reproduction all year long, but its influence weakens towards the Western region, where sea temperature rises especially during thermal anomaly events (Angeles-Gonzalez et al., 2017). The incidence of thermal anomalies has been associated with a decrease in yields of the octopus' fishery in the Yucatan Peninsula, especially in the Western region (Noyola Regil et al., 2015; Angeles-Gonzalez et al., 2017). In an ocean warming scenario or during thermal anomalies, octopuses from Western Yucatan Peninsula may move away into deeper environments or towards the upwelling zone, looking for cooler waters. This may alter ecological interactions and increase the mortality rate if the species finds additional predators or competitors while reaching such environments. In this scenario, the *O. maya* fishery may decline due to a reduction in fertilization and egg-laying rates, but also because the population could migrate to deeper waters reducing the species catchability, or because of a

higher mortality rate (Ángeles-González *et al.*, 2021). In this regard, the octopus aquaculture—with stringent temperature control—can emerge as the best alternative for octopus' production.

Limitations of the study and recommendations for further research

In the sequencing step, it is recommended to use at least three biological replicates per condition (or more, if the funding allows it) to enhance the statistical power of the analysis, especially to detect DEGs with low expression (Williams *et al.*, 2014; Honaas, Altman & Krzywinski, 2016). However, due to funding constraints, we implemented the biological averaging approach by using pooled samples per condition in the sequencing step (Honaas, Altman & Krzywinski, 2016) and utilized biological replicates to validate the expression of key genes through RT-qPCR analysis. This low-cost strategy provided valuable insights into the physiology and thermal stress response in the oviducal gland of *O. maya*.

On the other hand, considering that the oviducal gland consists of multiple sections with specialized tissues (Froesch & Marthy, 1975; Olivares *et al.*, 2017; Anadón, 2019), we recommend, in further research, the use of single-cell transcriptomics to better understand the role of each section in the glands' physiology.

CONCLUSIONS

At optimal temperatures, key reproductive genes in the OvG control the onset of fertilization and egg-laying: Before fertilization, the upregulation of genes encoding the myomodulin neuropeptide and enzymes for the synthesis of spermine and spermidine may prevent premature fertilization. In the OvG of mated and fertilized females, upregulation of genes related to the assembly and motility of the spermatozoa flagellum indicates the metabolic and transcriptomic reactivation of sperm. During fertilization, genes encoding the receptors of serotonin, dopamine, and progesterone were highly expressed. Likewise, fertilization was favored by the expression of genes that play a role in the interaction of male and female gametes. Other important genes for the reproductive process like Ptx, Catsup, FanaCh, and CYP171A, and those involved in the synthesis of eggshell mucoproteins were conspicuous at this stage under optimal thermal conditions. In senescent females, genes involved in fertilization were downregulated and those involved in the metabolism of steroid hormones like the SRD5A1 were highly expressed.

The fertilization rate decreased in the HTST-TD; this coincided with the upregulation of neuropeptides like myomodulin and APGW-amide, downregulation of genes involved in the adhesion between spermatozoa and eggs like galectin-3 and brorin, and downregulation of the neuroendocrine convertase 1 gene. This regulation may be required to reduce the fertilization rate under high temperatures which are stressful for embryos and hatchlings of this species, therefore the species may be highly vulnerable to ocean warming.

Finally, some proteins associated with the fertilization process in *O. maya* have not been previously detected in other invertebrates, to our knowledge. Although, they have been found in vertebrate taxa, where they exhibit homologies in the molecular mechanisms to

achieve fertilization and leave offspring. The reproductive proteins involved in the fertilization process evolve rapidly, partly due to the type of fertilization (internal or external) and the reproductive system (see Table 1 in *Turner & Hoekstra (2008)*). Thus, the adaptive evolution of functionally reproductive proteins is contrasting between the external fertilization species (mediated by simple gamete proteins) and internal fertilization species (involving complex interactions between multiple proteins) (*Turner & Hoekstra, 2008*).

ACKNOWLEDGEMENTS

The results presented here are part of Louisiana Arreola-Meraz's MS Thesis. We also thank Claudia Ventura-López for her technical support during the RT-qPCR procedure and Claudia Caamal-Monsreal for her support with biological material during the experiments and sampling.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This research was funded by the project SEP-CONACYT-CB-2014-01/241690, and CICESE internal project 682-136 awarded to Clara E. Galindo-Sanchez, also by projects DGAPA-UNAM, PAPIIT IN203022, and CONACYT 61503 awarded to Carlos Rosas. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
SEP-CONACYT-CB-2014-01/241690.
CICESE internal project 682-136.
DGAPA-UNAM, PAPIIT IN203022, and CONACYT 61503.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Oscar E. Juárez conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Louisiana Arreola-Meraz performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Edna Sánchez-Castrejón analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Omar Hernando Avila-Poveda analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Laura L. López-Galindo performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

- Carlos Rosas conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Clara E. Galindo-Sánchez conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (*i.e.*, approving body and any reference numbers):

Our protocols were approved by the experimental Animal Ethics Committee of the Faculty of Chemistry at Universidad Nacional Autónoma de México (Permit number: Oficio/FQ/CICUAL/099/15).

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

The transcriptome assembly is available at DDBJ/EMBL/GenBank: [GJEO000000001](https://www.ncbi.nlm.nih.gov/nuccore/GJEO000000001).

Data Availability

The following information was supplied regarding data availability:

The RT-qPCR raw data is available in the [Supplementary File](#).

The raw sequence reads are available at NCBI-SRA: [SRR13512014](https://www.ncbi.nlm.nih.gov/sra/SRR13512014)–[SRR13512017](https://www.ncbi.nlm.nih.gov/sra/SRR13512017), BioProject [PRJNA544090](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA544090).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.12895#supplemental-information>.

REFERENCES

- Anadón R. 2019.** Functional histology: the tissues of common coleoid cephalopods. In: *Handbook of Pathogens and Diseases in Cephalopods*, Cham: Springer International Publishing, 39–85.
- Anderson RC, Wood JB, Byrne RA. 2002.** Octopus senescence: the beginning of the end. *Journal of Applied Animal Welfare Science* 5(4):275–283 DOI [10.1207/S15327604JAWS0504_02](https://doi.org/10.1207/S15327604JAWS0504_02).
- Andrews PLR, Darmaillacq A-S, Dennison N, Gleadall IG, Hawkins P, Messenger JB, Osorio D, Smith VJ, Smith JA. 2013.** The identification and management of pain, suffering and distress in cephalopods, including anaesthesia, analgesia and humane killing. *Journal of Experimental Marine Biology and Ecology* 447(13):46–64 DOI [10.1016/j.jembe.2013.02.010](https://doi.org/10.1016/j.jembe.2013.02.010).
- Angeles-Gonzalez LE, Calva R, Santos-Valencia J, Avila-Poveda OH, Olivares A, Diaz F, Rosas C. 2017.** Temperature modulates spatio-temporal variability of the functional reproductive maturation of *Octopus maya* (Cephalopoda) on the shelf of the Yucatan Peninsula, Mexico. *Journal of Molluscan Studies* 83(3):280–288 DOI [10.1093/mollus/eyx013](https://doi.org/10.1093/mollus/eyx013).
- Ángeles-González LE, Martínez-Meyer E, Yañez-Arenas C, Velázquez-Abunader I, López-Rocha JA, Torrejón-Magallanes J, Rosas C. 2021.** Climate change effect on *Octopus maya* (Voss and Solís-Ramírez, 1966) suitability and distribution in the Yucatan Peninsula, Gulf of Mexico: A correlative and mechanistic approach. *Estuarine, Coastal and Shelf Science* 260 DOI [10.1016/j.ecss.2021.107502](https://doi.org/10.1016/j.ecss.2021.107502).

- Arendse DCZ, Pitcher GC, Griffiths C. 2018.** Reproductive condition and spawning induction in the South African scallop *pecten sulcicostatus*. *Journal of Shellfish Research* **37**:1005–1011 DOI [10.2983/035.037.0511](https://doi.org/10.2983/035.037.0511).
- Arkhipkin AI. 1992.** Reproductive system structure, development and function in cephalopods with a new general scale for maturity stages. *Journal of Northwest Atlantic Fishery Science* **12**:63–74 DOI [10.2960/J.v12.a7](https://doi.org/10.2960/J.v12.a7).
- Auchus RJ, Lee TC, Miller WL. 1998.** Cytochrome b5 augments the 17,20-lyase activity of human P450c17 without direct electron transfer. *The Journal of Biological Chemistry* **273**(6):3158–3165 DOI [10.1074/jbc.273.6.3158](https://doi.org/10.1074/jbc.273.6.3158).
- Avila-Poveda OH, Koueta N, Benítez-Villalobos F, Santos-Valencia J, Rosas C. 2016.** Reproductive traits of *Octopus maya* (Cephalopoda: Octopoda) with implications for fisheries management. *Molluscan Research* **36**(1):29–44 DOI [10.1080/13235818.2015.1072912](https://doi.org/10.1080/13235818.2015.1072912).
- Berg JS, Evans JP, Leigh MW, Omran H, Bizon C, Mane K, Knowles MR, Weck KE, Zariwala MA. 2011.** Next generation massively parallel sequencing of targeted exomes to identify genetic mutations in primary ciliary dyskinesia: implications for application to clinical testing. *Genetics in Medicine* **13**(3):218–229 DOI [10.1097/GIM.0b013e318203cff2](https://doi.org/10.1097/GIM.0b013e318203cff2).
- Bi M, Hickox JR, Winfrey VP, Olson GE, Hardy DM. 2003.** Processing, localization and binding activity of zonadhesin suggest a function in sperm adhesion to the zona pellucida during exocytosis of the acrosome. *The Biochemical Journal* **375**(2):477–488 DOI [10.1042/BJ20030753](https://doi.org/10.1042/BJ20030753).
- Bolger AM, Lohse M, Usadel B. 2014.** Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**(15):2114–2120 DOI [10.1093/bioinformatics/btu170](https://doi.org/10.1093/bioinformatics/btu170).
- Braley RD. 1985.** Serotonin-induced spawning in giant clams (Bivalvia: Tridacnidae). *Aquaculture* **47**(4):321–325 DOI [10.1016/0044-8486\(85\)90217-0](https://doi.org/10.1016/0044-8486(85)90217-0).
- Brezina V, Bank B, Cropper EC, Rosen S, Vilim FS, Kupfermann I, Weiss KR. 1995.** Nine members of the myomodulin family of peptide cotransmitters at the B16-ARC neuromuscular junction of *Aplysia*. *Journal of Neurophysiology* **74**(1):54–72 DOI [10.1152/jn.1995.74.1.54](https://doi.org/10.1152/jn.1995.74.1.54).
- Brockhausen I, Stanley P. 2015.** *O-GalNAc glycans*. New York: Cold Spring Harbor Laboratory Press.
- Budelmann BU, Schipp R, von Boletzky S. 1997.** Cephalopoda. In: Harrison FW, Kohn AJ, eds. *Microscopic Anatomy of Invertebrates. Mollusca II*. New York: Wiley-Liss, 119–414.
- Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. 2009.** The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clinical Chemistry* **55**(4):611–622 DOI [10.1373/clinchem.2008.112797](https://doi.org/10.1373/clinchem.2008.112797).
- Caamal-Monsreal C, Uriarte I, Farias A, Díaz F, Sánchez A, Re D, Rosas C. 2016.** Effects of temperature on embryo development and metabolism of *O. maya*. *Aquaculture* **451**:156–162 DOI [10.1016/j.aquaculture.2015.09.011](https://doi.org/10.1016/j.aquaculture.2015.09.011).
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.** BLAST+: architecture and applications. *BMC Bioinformatics* **10**(1):421 DOI [10.1186/1471-2105-10-421](https://doi.org/10.1186/1471-2105-10-421).
- Chen Y, Vujcic S, Liang P, Diegelman P, Kramer DL, Porter CW. 2003.** Genomic identification and biochemical characterization of a second spermidine/spermine N1-acetyltransferase. *The Biochemical Journal* **373**(3):661–667 DOI [10.1042/BJ20030734](https://doi.org/10.1042/BJ20030734).
- Clausen H, Bennett EP. 1996.** A family of UDP-GalNAc: polypeptide N-acetylgalactosaminyltransferases control the initiation of mucin-type O-linked glycosylation. *Glycobiology* **6**(6):635–646 DOI [10.1093/glycob/6.6.635](https://doi.org/10.1093/glycob/6.6.635).

- Crespi-Abril AC, Rubilar T. 2021.** Moving forward in the ethical consideration of invertebrates in experimentation: beyond the Three R's Principle. *Revista de Biología Tropical* **69**(Suppl.1):346–357 DOI [10.15517/rbt.v69isuppl.1.46366](https://doi.org/10.15517/rbt.v69isuppl.1.46366).
- Cropper EC, Tenenbaum R, Kolks MA, Kupfermann I, Weiss KR. 1987.** Myomodulin: a bioactive neuropeptide present in an identified cholinergic buccal motor neuron of *Aplysia*. *Proceedings of the National Academy of Sciences of the United States of America* **84**(15):5483–5486 DOI [10.1073/pnas.84.15.5483](https://doi.org/10.1073/pnas.84.15.5483).
- Cropper EC, Vilim FS, Alevizos A, Tenenbaum R, Kolks MAG, Rosen S, Kupfermann I, Weiss KR. 1991.** Structure, bioactivity, and cellular localization of myomodulin B: a novel *Aplysia* peptide. *Peptides* **12**(4):683–690 DOI [10.1016/0196-9781\(91\)90120-E](https://doi.org/10.1016/0196-9781(91)90120-E).
- Cuccu D, Mereu M, Porcu C, Follesa MC, Cau AL, Cau A. 2013.** Development of sexual organs and fecundity in *Octopus vulgaris* Cuvier, 1797 from the Sardinian waters (Mediterranean Sea). *Mediterranean Marine Science* **14**(2):270–277 DOI [10.12681/mms.412](https://doi.org/10.12681/mms.412).
- De Lange RPJ, Joosse J, Van Minnen J. 1998.** Multi-messenger innervation of the male sexual system of *Lymnaea stagnalis*. *The Journal of Comparative Neurology* **390**:564–577 DOI [10.1002/\(SICI\)1096-9861\(19980126\)390:4<564::AID-CNE8>3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1096-9861(19980126)390:4<564::AID-CNE8>3.0.CO;2-Z).
- De Lisa E, Salzano AM, Moccia F, Scaloni A, Di Cosmo A. 2013.** Sperm-attractant peptide influences the spermatozoa swimming behavior in internal fertilization in *Octopus vulgaris*. *Journal of Experimental Biology* **216**(12):2229–2237 DOI [10.1242/jeb.081885](https://doi.org/10.1242/jeb.081885).
- Deguchi R, Osada M. 2020.** Chapter 7 gametogenesis, spawning, and fertilization in bivalves and other protostomes. In: *Reproduction in Aquatic Animals*, Singapore, Springer Singapore, 113–165.
- Delgado-Bermúdez A, Ribas-Maynou J, Yeste M. 2022.** Relevance of aquaporins for gamete function and cryopreservation. *Animals* **12**:573 DOI [10.3390/ani12050573](https://doi.org/10.3390/ani12050573).
- Di Cosmo A, Di Cristo C, Paolucci M. 2001.** Sex steroid hormone fluctuations and morphological changes of the reproductive system of the female of *Octopus vulgaris* throughout the annual cycle. *Journal of Experimental Zoology* **289**:33–47 DOI [10.1002/1097-010X\(20010101/31\)289:1<33::AID-JEZ4>3.0.CO;2-A](https://doi.org/10.1002/1097-010X(20010101/31)289:1<33::AID-JEZ4>3.0.CO;2-A).
- Di Cosmo A, Paolucci M, Di Cristo C, Botte V, Ciarcia G. 1998.** Progesterone receptor in the reproductive system of the female of *Octopus vulgaris*: characterization and immunolocalization. *Molecular Reproduction and Development* **50**:451–460 DOI [10.1002/\(SICI\)1098-2795\(199808\)50:4<451::AID-MRD9>3.0.CO;2-H](https://doi.org/10.1002/(SICI)1098-2795(199808)50:4<451::AID-MRD9>3.0.CO;2-H).
- Di Cristo C. 2013.** Nervous control of reproduction in *Octopus vulgaris*: a new model. *Invertebrate Neuroscience* **13**(1):27–34 DOI [10.1007/s10158-013-0149-x](https://doi.org/10.1007/s10158-013-0149-x).
- Di Cristo C, Di Cosmo A. 2007.** Neuropeptidergic control of *Octopus* oviducal gland. *Peptides* **28**(1):163–168 DOI [10.1016/j.peptides.2006.09.016](https://doi.org/10.1016/j.peptides.2006.09.016).
- Dobrovolsky VN, Bowyer JF, Pabarcus MK, Heflich RH, Williams LD, Doerge DR, Arvidsson B, Bergquist J, Casida JE. 2005.** Effect of arylformamidase (kynurenine formamidase) gene inactivation in mice on enzymatic activity, kynurenine pathway metabolites and phenotype. *Biochimica et Biophysica Acta (BBA)-General Subjects* **1724**(1–2):163–172 DOI [10.1016/j.bbagen.2005.03.010](https://doi.org/10.1016/j.bbagen.2005.03.010).
- Estefanell J, Socorro J, Afonso JM, Roo J, Fernandez-Palacios H, Izquierdo MS. 2011.** Evaluation of two anaesthetic agents and the passive integrated transponder tagging system in *Octopus vulgaris* (Cuvier 1797). *Aquaculture Research* **42**(3):399–406 DOI [10.1111/j.1365-2109.2010.02634.x](https://doi.org/10.1111/j.1365-2109.2010.02634.x).

- Froesch D, Marthy HJ. 1975.** The structure and function of the oviducal gland in octopods (Cephalopoda). *Proceedings of the Royal Society of London. Series B, Biological Sciences* **188(1090)**:95–101 DOI [10.1098/rspb.1975.0005](https://doi.org/10.1098/rspb.1975.0005).
- Gao Z, Garbers DL. 1998.** Species diversity in the structure of zonadhesin, a sperm-specific membrane protein containing multiple cell adhesion molecule-like domains. *Journal of Biological Chemistry* **273(6)**:3415–3421 DOI [10.1074/jbc.273.6.3415](https://doi.org/10.1074/jbc.273.6.3415).
- Gibbons MC, Castagna M. 1984.** Serotonin as an inducer of spawning in six bivalve species. *Aquaculture* **40(2)**:189–191 DOI [10.1016/0044-8486\(84\)90356-9](https://doi.org/10.1016/0044-8486(84)90356-9).
- Gleadall IG. 2013.** The effects of prospective anaesthetic substances on cephalopods: summary of original data and a brief review of studies over the last two decades. *Journal of Experimental Marine Biology and Ecology* **447**:23–30 DOI [10.1016/j.jembe.2013.02.008](https://doi.org/10.1016/j.jembe.2013.02.008).
- Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, Di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. 2011.** Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* **29(7)**:644–652 DOI [10.1038/nbt.1883](https://doi.org/10.1038/nbt.1883).
- Greenberg MJ, Doble KE, Lesser W, Lee TD, Pennell NA, Morgan CG, Price DA. 1997.** Characterization of myomodulin-related peptides from the pulmonate snail *Helix aspersa*. *Peptides* **18(8)**:1099–1106 DOI [10.1016/S0196-9781\(97\)00149-6](https://doi.org/10.1016/S0196-9781(97)00149-6).
- Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, Couger MB, Eccles D, Li B, Lieber M, Macmanes MD, Ott M, Orvis J, Pochet N, Strozzi F, Weeks N, Westerman R, William T, Dewey CN, Henschel R, Leduc RD, Friedman N, Regev A. 2013.** De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols* **8(8)**:1494–1512 DOI [10.1038/nprot.2013.084](https://doi.org/10.1038/nprot.2013.084).
- Han Q, Li G, Li J. 2000.** Purification and characterization of chorion peroxidase from *Aedes aegypti* eggs. *Archives of Biochemistry and Biophysics* **378(1)**:107–115 DOI [10.1006/abbi.2000.1821](https://doi.org/10.1006/abbi.2000.1821).
- Hardy DM, Garbers DL. 1995.** A sperm membrane protein that binds in a species-specific manner to the egg extracellular matrix is homologous to von willebrand factor. *Journal of Biological Chemistry* **270(44)**:26025–26028 DOI [10.1074/jbc.270.44.26025](https://doi.org/10.1074/jbc.270.44.26025).
- Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. 2007.** qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biology* **8(2)**:R19 DOI [10.1186/gb-2007-8-2-r19](https://doi.org/10.1186/gb-2007-8-2-r19).
- Henry J, Zatylny C, Favrel P. 2000.** HPLC and electrospray ionization mass spectrometry as tools for the identification of APGWamide-related peptides in gastropod and bivalve mollusks: comparative activities on *Mytilus* muscles. *Brain Research* **862(1–2)**:162–170 DOI [10.1016/S0006-8993\(00\)02108-9](https://doi.org/10.1016/S0006-8993(00)02108-9).
- Honaas LA, Altman NS, Krzywinski M. 2016.** Study design for sequencing studies. In: *Methods in Molecular Biology (Clifton, N.J.)*, New York: Humana Press, 39–66.
- Huang DW, Sherman BT, Lempicki RA. 2009.** Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* **4(1)**:44–57 DOI [10.1038/nprot.2008.211](https://doi.org/10.1038/nprot.2008.211).
- Iuchi Y, Okada F, Tsunoda S, Kibe N, Shirasawa N, Ikawa M, Okabe M, Ikeda Y, Fujii J. 2009.** Peroxiredoxin 4 knockout results in elevated spermatogenic cell death via oxidative stress. *The Biochemical Journal* **419(1)**:149–158 DOI [10.1042/BJ20081526](https://doi.org/10.1042/BJ20081526).
- Iwakoshi-Ukena E, Ukena K, Takuwa-Kuroda K, Kanda A, Tsutsui K, Minakata H. 2004.** Expression and distribution of octopus gonadotropin-releasing hormone in the central nervous system and peripheral organs of the octopus (*Octopus vulgaris*) by in situ hybridization and

immunohistochemistry. *Journal of Comparative Neurology* 477(3):310–323
DOI 10.1002/cne.20260.

- Juárez OE. 2016. *Evaluación a nivel fisiológico y transcriptómico del efecto de la temperatura sobre la regulación y el éxito reproductivo en hembras Octopus maya* [Doctoral Thesis, Centro de Investigación Científica y Educación Superior de Ensenada, Baja California]. Available at <http://cicese.repositorioinstitucional.mx/jspui/handle/1007/1351>.
- Juárez OE, Galindo-Sánchez CE, Díaz F, Re D, Sánchez-García AM, Camaal-Monsreal C, Rosas C. 2015. Is temperature conditioning *Octopus maya* fitness? *Journal of Experimental Marine Biology and Ecology* 467:71–76 DOI 10.1016/j.jembe.2015.02.020.
- Juárez OE, Hau V, Camaal-Monsreal C, Galindo-Sánchez CE, Díaz F, Re D, Rosas C. 2016. Effect of maternal temperature stress before spawning over the energetic balance of *Octopus maya* juveniles exposed to a gradual temperature change. *Journal of Experimental Marine Biology and Ecology* 474:39–45 DOI 10.1016/j.jembe.2015.10.002.
- Juárez OE, López-Galindo L, Pérez-Carrasco L, Lago-Lestón A, Rosas C, Di Cosmo A, Galindo-Sánchez CE. 2019. *Octopus maya* white body show sex-specific transcriptomic profiles during the reproductive phase, with high differentiation in signaling pathways. *PLOS ONE* 14(5):e0216982 DOI 10.1371/journal.pone.0216982.
- Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. 2012. KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Research* 40(D1):D109–D114 DOI 10.1093/nar/gkr988.
- Kellett E, Perry SJ, Santama N, Worster BM, Benjamin PR, Burke JF. 1996. Myomodulin gene of *Lymnaea*: structure, expression, and analysis of neuropeptides. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* 16(16):4949–4957 DOI 10.1523/jneurosci.16-16-04949.1996.
- Kerns K, Zigo M, Drobnis EZ, Sutovsky M, Sutovsky P. 2018. Zinc ion flux during mammalian sperm capacitation. *Nature Communications* 9:2061 DOI 10.1038/s41467-018-04523-y.
- Kerns K, Zigo M, Sutovsky P. 2018. Zinc: a necessary ion for mammalian sperm fertilization competency. *International Journal of Molecular Sciences* 19(12):4097 DOI 10.3390/ijms19124097.
- Koene JM. 2010. Neuro-endocrine control of reproduction in hermaphroditic freshwater snails: mechanisms and evolution. *Frontiers in Behavioral Neuroscience* 4:1–17 DOI 10.3389/fnbeh.2010.00167.
- Konstandi OA, Papassideri IS, Stravopodis DJ, Antonelou MH, Kenoutis CA, Stefanidou DC, Margaritis LH. 2006. The dual role of chorion peroxidase in *Bactrocera oleae* chorion assembly. *International Journal of Developmental Biology* 50:543–552 DOI 10.1387/ijdb.052122ok.
- Konstandi OA, Papassideri IS, Stravopodis DJ, Kenoutis CA, Hasan Z, Katsorchis T, Wever R, Margaritis LH. 2005. The enzymatic component of *Drosophila melanogaster* chorion is the Pxd peroxidase. *Insect Biochemistry and Molecular Biology* 35(9):1043–1057 DOI 10.1016/j.ibmb.2005.04.005.
- Kosman ET, Levitan DR. 2014. Sperm competition and the evolution of gametic compatibility in externally fertilizing taxa. *Molecular Human Reproduction* 20(12):1190–1197 DOI 10.1093/molehr/gau069.
- Kuan C-T, Chang J, Mansson J-E, Li J, Pegram C, Fredman P, McLendon RE, Bigner DD. 2010. Multiple phenotypic changes in mice after knockout of the B3gnt5 gene, encoding Lc3 synthase—a key enzyme in lacto-neolacto ganglioside synthesis. *BMC Developmental Biology* 10:114 DOI 10.1186/1471-213X-10-114.

- Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 9(4):357–359 DOI 10.1038/nmeth.1923.
- Leigh MW, Pittman JE, Carson JL, Ferkol TW, Dell SD, Davis SD, Knowles MR, Zariwala MA. 2009. Clinical and genetic aspects of primary ciliary dyskinesia/Kartagener syndrome. *Genetics in Medicine* 11(7):473–487 DOI 10.1097/GIM.0b013e3181a53562.
- Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 12:323 DOI 10.1186/1471-2105-12-323.
- Li J, Hodgeman BA, Christensen BM. 1996. Involvement of peroxidase in chorion hardening in *Aedes aegypti*. *Insect Biochemistry and Molecular Biology* 26(3):309–317 DOI 10.1016/0965-1748(95)00099-2.
- Li JS, Li J. 2006. Major chorion proteins and their crosslinking during chorion hardening in *Aedes aegypti* mosquitoes. *Insect Biochemistry and Molecular Biology* 36(12):954–964 DOI 10.1016/j.ibmb.2006.09.006.
- López-Galindo L, Galindo-Sánchez C, Olivares A, Avila-Poveda OH, Díaz F, Juárez OE, Lafarga F, Pantoja-Pérez J, Caamal-Monsreal C, Rosas C. 2019a. Reproductive performance of *Octopus maya* males conditioned by thermal stress. *Ecological Indicators* 96:437–447 DOI 10.1016/j.ecolind.2018.09.036.
- López-Galindo L, Juárez OE, Larios-Soriano E, Del Vecchio G, Ventura-López C, Lago-Lestón A, Galindo-Sánchez C. 2019b. Transcriptomic analysis reveals insights on male infertility in *Octopus maya* under chronic thermal stress. *Frontiers in Physiology* 9:1920 DOI 10.3389/fphys.2018.01920.
- Manabe R, Tsutsui K, Yamada T, Kimura M, Nakano I, Shimono C, Sanzen N, Furutani Y, Fukuda T, Oguri Y, Shimamoto K, Kiyozumi D, Sato Y, Sado Y, Senoo H, Yamashina S, Fukuda S, Kawai J, Sugiura N, Kimata K, Hayashizaki Y, Sekiguchi K. 2008. Transcriptome-based systematic identification of extracellular matrix proteins. *Proceedings of the National Academy of Sciences of the United States of America* 105(35):12849–12854 DOI 10.1073/pnas.0803640105.
- Mangold K. 1987. Reproduction. In: *Cephalopod Life Cycles Vol. 2 Comparative Reviews*, Cambridge: Academic Press, 157–200.
- Mangold K, von Boletzky S, Frösch D. 1971. Reproductive biology and embryonic development of *Eledone cirrosa* (Cephalopoda: Octopoda). *Marine Biology* 8(2):109–117 DOI 10.1007/BF00350926.
- Margaritis LH. 1985. The egg-shell of *Drosophila melanogaster* III. Covalent crosslinking of the chorion proteins involves endogenous hydrogen peroxide. *Tissue and Cell* 17(4):553–559 DOI 10.1016/0040-8166(85)90031-X.
- Marian JEAR. 2011. O enigma da reação espermatofórica: breve síntese do conhecimento sobre a estrutura e o funcionamento dos espermátóforos dos cefalópodes (Mollusca: Cephalopoda). *Papéis Avulsos de Zoologia* 51(13):207–219 DOI 10.1590/S0031-10492011001300001.
- Marian JEAR. 2015. Evolution of spermatophore transfer mechanisms in cephalopods. *Journal of Natural History* 49(21–24):1423–1455 DOI 10.1080/00222933.2013.825026.
- Markaida U, Méndez-Loeza I, Rosales-Raya ML. 2017. Seasonal and spatial trends of Mayan octopus, *Octopus maya*, population dynamics from Campeche, Mexico. *Journal of the Marine Biological Association of the United Kingdom* 97(8):1663–1673 DOI 10.1017/S0025315416001132.
- Mather JA, Anderson RC. 2007. Ethics and invertebrates: a cephalopod perspective. *Diseases of Aquatic Organisms* 75:119–129 DOI 10.3354/dao075119.

- Mei S, Chen P, Lee C-L, Zhao W, Wang Y, Lam KKW, Ho P-C, Yeung WSB, Fang C, Chiu PCN. 2019. The role of galectin-3 in spermatozoa-zona pellucida binding and its association with fertilization in vitro. *Molecular Human Reproduction* 25(8):458–470 DOI 10.1093/molehr/gaz030.
- Meza-Buendía AK, Trejo-Escamilla I, Piu M, Caamal-Monsreal C, Rodríguez-Fuentes G, Díaz F, Re D, Galindo-Sánchez CE, Rosas C. 2021. Why high temperatures limit reproduction in cephalopods? The case of *Octopus maya*. *Aquaculture Research* 52(11):5111–5123 DOI 10.1111/are.15387.
- Minakata H, Kuroki Y, Ikeda T, Fujisawa Y, Nomoto K, Kubota I, Muneoka Y. 1991. Effects of the neuropeptide APGW-amide and related compounds on molluscan muscles-GW-amide shows potent modulatory effects. *Comparative Biochemistry and Physiology. Part C, Comparative Pharmacology* 100(3):565–571 DOI 10.1016/0742-8413(91)90041-Q.
- Minakata H, Shigeno S, Kano N, Haraguchi S, Osugi T, Tsutsui K. 2009. Octopus gonadotrophin-releasing hormone: a multifunctional peptide in the endocrine and nervous systems of the cephalopod. *Journal of Neuroendocrinology* 21(4):322–326 DOI 10.1111/j.1365-2826.2009.01852.x.
- Mindrinos MN, Petri WH, Galanopoulos VK, Lombard MF, Margaritis LH. 1980. Crosslinking of the *Drosophila* chorion involves a peroxidase. *Wilhelm Roux's Archives of Developmental Biology* 189:187–196 DOI 10.1007/BF00868677.
- Moltschanivskyj NA, Hall K, Lipinski MR, Marian JEAR, Nishiguchi M, Sakai M, Shulman DJ, Sinclair B, Sinn DL, Staudinger M, Van Gelderen R, Villanueva R, Warnke K. 2007. Ethical and welfare considerations when using cephalopods as experimental animals. *Reviews in Fish Biology and Fisheries* 17:455–476 DOI 10.1007/s11160-007-9056-8.
- Morash MG, Soanes K, Anini Y. 2011. Prohormone processing in zebrafish. *Methods in Molecular Biology (Clifton, N.J.)* 768:257–271 DOI 10.1007/978-1-61779-204-5_14.
- Méndez J, Sauer-Ramírez J. 2018. Ram Sperm DNA fragmentation under hyperglycemia conditions is inhibited by polyamines. *Journal of Clinical and Molecular Endocrinology* 3(1):2–26 DOI 10.21767/2572-5432.100046.
- Nagase H, Murphy G. 2013. Metalloproteinases, matrix. In: *Encyclopedia of Biological Chemistry: Second Edition*, Cambridge: Academic Press, 90–97.
- Noyola Regil J, Mascaro M, Díaz F, Re AD, Sánchez-Zamora A, Caamal-Monsreal C, Rosas C. 2015. Thermal biology of prey (*Melongena corona bispinosa*, *Strombus pugilis*, *Callinectes similis*, *Libinia dubia*) and predators (*Ocyurus chrysurus*, *Centropomus undecimalis*) of *Octopus maya* from the Yucatan Peninsula. *Journal of Thermal Biology* 53:151–161 DOI 10.1016/j.jtherbio.2015.11.001.
- Olivares A, Avila-Poveda OH, Leyton V, Zuñiga O, Rosas C, Northland-Leppe I. 2017. Oviducal glands throughout the gonad development stages: a case study of *Octopus mimus* (Cephalopoda). *Molluscan Research* 37(4):229–241 DOI 10.1080/13235818.2017.1334275.
- Onoufriadis A, Shoemark A, Schmidts M, Patel M, Jimenez G, Liu H, Thomas B, Dixon M, Hirst RA, Rutman A, Burgoyne T, Williams C, Scully J, Bolard F, Lafitte JJ, Beales PL, Hogg C, Yang P, Chung EMK, Emes RD, O'Callaghan C, Bouvagnet P, Mitchison HM. 2014. Targeted NGS gene panel identifies mutations in RSPH1 causing primary ciliary dyskinesia and a common mechanism for ciliary central pair agenesis due to radial spoke defects. *Human Molecular Genetics* 23(13):3362–3374 DOI 10.1093/hmg/ddu046.
- Osada M, Matsutani T, Nomura T. 1987. Implication of catecholamines during spawning in marine bivalve molluscs. *International Journal of Invertebrate Reproduction and Development* 12(3):241–251 DOI 10.1080/01688170.1987.10510324.

- O'Dor RK, Wells MJ. 1978. Reproduction versus somatic growth: hormonal control in *Octopus vulgaris*. *Journal of Experimental Biology* 77:15–31 DOI 10.1242/jeb.77.1.15.
- Panhuis TM, Clark NL, Swanson WJ. 2006. Rapid evolution of reproductive proteins in abalone and *Drosophila*. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 361(1466):261–268 DOI 10.1098/rstb.2005.1793.
- Pascual C, Mascaro M, Rodríguez-Canul R, Gallardo P, Sánchez AA, Rosas C, Cruz-López H. 2019. Sea surface temperature modulates physiological and immunological condition of *Octopus maya*. *Frontiers in Physiology* 10:1–11 DOI 10.3389/fphys.2019.00739.
- Pati D, Meistrich ML, Plon SE. 1999. Human Cdc34 and Rad6B ubiquitin-conjugating enzymes target repressors of cyclic AMP-induced transcription for proteolysis. *Molecular and Cellular Biology* 19(7):5001–5013 DOI 10.1128/mcb.19.7.5001.
- Pereira R, Sá R, Barros A, Sousa M. 2017. Major regulatory mechanisms involved in sperm motility. *Asian Journal of Andrology* 19:5–14 DOI 10.4103/1008-682X.167716.
- Peterson RP. 1959. The anatomy and histology of the reproductive systems of *Octopus bimaculoides*. *Journal of Morphology* 104(1):61–87 DOI 10.1002/jmor.1051040103.
- Petrunk EM, DeVore NM, Porubsky PR, Scott EE. 2014. Structures of human steroidogenic cytochrome P450 17A1 with substrates. *Journal of Biological Chemistry* 289(47):32952–32964 DOI 10.1074/jbc.M114.610998.
- R Development Core Team. 2010. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Available at <http://www.R-project.org/>.
- Ribeiro JC, Alves MG, Yeste M, Cho YS, Calamita G, Oliveira PF. 2021. Aquaporins and (in) fertility: more than just water transport. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1867(3):166039 DOI 10.1016/j.bbadis.2020.166039.
- Robinson MD, McCarthy DJ, Smyth GK. 2009. edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26(1):139–140 DOI 10.1093/bioinformatics/btp616.
- Rosas C, Gallardo P, Mascaró M, Caamal-Monsreal C, Pascual C. 2014. *Octopus maya*. In: *Cephalopod Culture*, Dordrecht: Springer Netherlands, 383–396.
- Rubinstein S, Breitbart H. 1991. Role of spermine in mammalian sperm capacitation and acrosome reaction. *Biochemical Journal* 278(1):25–28 DOI 10.1042/bj2780025.
- Sanchez-García A, Rodríguez-Fuentes G, Díaz F, Galindo-Sánchez CE, Ortega K, Mascaró M, López E, Caamal-Monsreal C, Juárez O, Noreña-Barroso E, Re D, Rosas C. 2017. Thermal sensitivity of *O. maya* embryos as a tool for monitoring the effects of environmental warming in the Southern of Gulf of Mexico. *Ecological Indicators* 72(3):574–585 DOI 10.1016/j.ecolind.2016.08.043.
- Sato N. 2021. A review of sperm storage methods and post-copulatory sexual selection in the Cephalopoda. *Biological Journal of the Linnean Society* 134(2):285–302 DOI 10.1093/biolinnean/blab096.
- Sheth AR, Moodbidri SB. 1977. Significance of polyamines in reproduction. *Advances in Sex Hormone Research* 3:51–74.
- Sneddon LU. 2015. Pain in aquatic animals. *The Journal of Experimental Biology* 218(7):967–976 DOI 10.1242/jeb.088823.
- Springate L, Frasier TR. 2017. Gamete compatibility genes in mammals: candidates, applications and a potential path forward. *Royal Society Open Science* 4(8):170577 DOI 10.1098/rsos.170577.
- Strauss JF, Modi B, McAllister JM. 2014. Defects in ovarian steroid hormone biosynthesis. In: *Cellular Endocrinology in Health and Disease*, Amsterdam: Elsevier Inc, 285–309.

- Takeuchi K, Xu Y, Kitano M, Chiyonobu K, Abo M, Ikegami K, Ogawa S, Ikejiri M, Kondo M, Gotoh S, Nagao M, Fujisawa T, Nakatani K. 2020. Copy number variation in DRC1 is the major cause of primary ciliary dyskinesia in the Japanese population. *Molecular Genetics and Genomic Medicine* 8(3):3943 DOI 10.1002/mgg3.1137.
- Tardif S, Wilson MD, Wagner R, Hunt P, Gertsenstein M, Nagy A, Lobe C, Koop BF, Hardy DM. 2010. Zonadhesin is essential for species specificity of sperm adhesion to the egg zona pellucida. *Journal of Biological Chemistry* 285(32):24863–24870 DOI 10.1074/jbc.M110.123125.
- Tosti E, Di Cosmo A, Cuomo A, Di Cristo C, Gragnaniello G. 2001. Progesterone induces activation in *Octopus vulgaris* spermatozoa. *Molecular Reproduction and Development* 59(1):97–105 DOI 10.1002/mrd.1011.
- Tsutsui K, Haraguchi S. 2016. Allopregnanolone. In: *Handbook of Hormones*, Amsterdam: Elsevier, 544-e96C-3.
- Turner LM, Hoekstra HE. 2008. Causes and consequences of the evolution of reproductive proteins. *The International Journal of Developmental Biology* 52:769–780 DOI 10.1387/ijdb.082577lt.
- Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG. 2012. Primer3-new capabilities and interfaces. *Nucleic Acids Research* 40(15):e115 DOI 10.1093/nar/gks596.
- Van Heukelem WF. 1983. Octopus maya. In: Boyle PR, ed. *Cephalopod Life Cycles. Vol. 1. Species Accounts*. London, UK: Academic Press, 311–323.
- Wang Y, Chen F, He J, Xue G, Chen J, Xie P. 2021. Cellular and molecular modification of egg envelope hardening in fertilization. *Biochimie* 181(16):134–144 DOI 10.1016/j.biochi.2020.12.007.
- Wang ZY, Ragsdale CW. 2018. Multiple optic gland signaling pathways implicated in octopus maternal behaviors and death. *Journal of Experimental Biology* 221(19):jeb185751 DOI 10.1242/jeb.185751.
- Wells MJ. 1978. *Octopus*. Dordrecht, Netherlands: Springer.
- Wells MJ, Wells J. 1959. Hormonal control of sexual maturity in Octopus. *Journal of Experimental Biology* 36:1–33 DOI 10.1242/jeb.36.1.1.
- Wells MJ, Wells J. 1977. Cephalopoda. In: Giese A, Pearse JS, eds. *Reproduction of Marine Invertebrates. Molluscs: Gastropods and Cephalopods*. Vol. 4. London, UK: Academic Press Inc, 291–337.
- Williams AG, Thomas S, Wyman SK, Holloway AK. 2014. RNA-seq data: challenges in and recommendations for experimental design and analysis. *Current Protocols in Human Genetics* 83(1):11.13.1–11.13.120 DOI 10.1002/0471142905.hg1113s83.
- Winlow W, Polese G, Moghadam H-F, Ahmed IA, Di Cosmo A. 2018. Sense and insensibility—an appraisal of the effects of clinical anesthetics on gastropod and cephalopod molluscs as a step to improved welfare of cephalopods. *Frontiers in Physiology* 9:1147 DOI 10.3389/fphys.2018.01147.
- Wu Y, Hu X, Li Z, Wang M, Li S, Wang X, Lin X, Liao S, Zhang Z, Feng X, Wang S, Cui X, Wang Y, Gao F, Hess RA, Han C. 2016. Transcription factor RFX2 is a key regulator of mouse spermiogenesis. *Scientific Reports* 6(1):20435 DOI 10.1038/srep20435.
- Wu H, Min J, Ikeguchi Y, Zeng H, Dong A, Loppnau P, Pegg AE, Plotnikov AN. 2007. Structure and mechanism of spermidine synthases †. *Biochemistry* 46(28):8331–8339 DOI 10.1021/bi602498k.

- Xie F, Xiao P, Chen D, Xu L, Zhang B. 2012.** miRDeepFinder: a miRNA analysis tool for deep sequencing of plant small RNAs. *Plant Molecular Biology* **80**(1):75–84
DOI [10.1007/s11103-012-9885-2](https://doi.org/10.1007/s11103-012-9885-2).
- Xu R, Zheng X. 2018.** Selection of reference genes for quantitative real-time PCR in *Octopus minor* (Cephalopoda: Octopoda) under acute ammonia stress. *Environmental Toxicology and Pharmacology* **60**(1–2):76–81 DOI [10.1016/j.etap.2018.04.010](https://doi.org/10.1016/j.etap.2018.04.010).
- Yanagimachi R, Wakayama T, Kishikawa H, Fimia GM, Monaco L, Sassone-Corsi P. 2004.** Production of fertile offspring from genetically infertile male mice. *Proceedings of the National Academy of Sciences of the United States of America* **101**(6):1691–1695
DOI [10.1073/pnas.0307832100](https://doi.org/10.1073/pnas.0307832100).
- Yeung CH, Callies C, Rojek A, Nielsen S, Cooper TG. 2009.** Aquaporin isoforms involved in physiological volume regulation of murine spermatozoa. *Biology of Reproduction* **80**(2):350–357
DOI [10.1095/biolreprod.108.071928](https://doi.org/10.1095/biolreprod.108.071928).
- York PS, Cummins SF, Degan SM, Woodcroft BJ, Degan BM. 2012.** Marked changes in neuropeptide expression accompany broadcast spawnings in the gastropod *Haliotis asinina*. *Frontiers in Zoology* **9**(1):9 DOI [10.1186/1742-9994-9-9](https://doi.org/10.1186/1742-9994-9-9).
- Yoshimoto FK, Gonzalez E, Auchus RJ, Guengerich FP. 2016.** Mechanism of 17 α ,20-lyase and new hydroxylation reactions of human cytochrome P450 17A1 18O labeling and oxygen surrogate evidence for a role of a perferryl oxygen. *Journal of Biological Chemistry* **291**(33):17143–17164 DOI [10.1074/jbc.M116.732966](https://doi.org/10.1074/jbc.M116.732966).
- Yuan W, Leisner TM, McFadden AW, Clark S, Hiller S, Maeda N, O'Brien DA, Parise LV. 2006.** CIB1 is essential for mouse spermatogenesis. *Molecular and Cellular Biology* **26**(22):8507–8514
DOI [10.1128/MCB.01488-06](https://doi.org/10.1128/MCB.01488-06).
- Zatylny-Gaudin C, Cornet V, Leduc A, Zanuttini B, Corre E, Le Corguillé G, Bernay B, Garderes J, Kraut A, Couté Y, Henry J. 2016.** Neuropeptidome of the cephalopod *Sepia officinalis*: identification, tissue mapping, and expression pattern of neuropeptides and neurohormones during egg laying. *Journal of Proteome Research* **15**(1):48–67
DOI [10.1021/acs.jproteome.5b00463](https://doi.org/10.1021/acs.jproteome.5b00463).
- Zhu Y, Rice CD, Pang Y, Pace M, Thomas P. 2003.** Cloning, expression, and characterization of a membrane progesterin receptor and evidence it is an intermediary in meiotic maturation of fish oocytes. *Proceedings of the National Academy of Sciences of the United States of America* **100**(5):2231–2236 DOI [10.1073/pnas.0336132100](https://doi.org/10.1073/pnas.0336132100).
- zur Lage P, Newton FG, Jarman AP. 2019.** Survey of the ciliary motility machinery of *Drosophila* sperm and ciliated mechanosensory neurons reveals unexpected cell-type specific variations: a model for motile ciliopathies. *Frontiers in Genetics* **10**:229 DOI [10.3389/fgene.2019.00024](https://doi.org/10.3389/fgene.2019.00024).