

## POSTER PRESENTATION

## Open Access

# Pericentrin interacts with KASH domain-containing protein Syne-2

N Falk<sup>1\*</sup>, K Kessler<sup>2</sup>, J Glöckner<sup>3</sup>, K Boldt<sup>3</sup>, M Ueffing<sup>3</sup>, R Roepman<sup>4</sup>, C Thiel<sup>2</sup>, JH Brandstätter<sup>1</sup>, A Gießl<sup>1</sup>

From Cilia 2014 - Second International Conference  
Paris, France. 18-21 November 2014

### Objective

Pericentrin, a highly conserved protein of the pericentriolar material, serves as a multifunctional scaffold for numerous proteins and plays an important role in microtubule organization. Mutations in the human PCNT gene are associated with a range of diseases including primordial dwarfism and ciliopathies. In the mouse retina Pericentrin colocalizes with several proteins responsible for transport processes at the connecting cilium between the two photoreceptor compartments. In order to get more insights on the function of Pericentrin in the retina we try to identify new ciliary as well as centrosomal interaction partners.

### Methods

Identification of Pericentrin interaction partners was done by Tandem Affinity Purification, Yeast two-hybrid with a self-constructed cDNA library from mouse retina and GST pull-down.

### Results

We were able to show that Pericentrin interacts with Klarsicht/ANC-1/Syne-homologue (KASH) domain-containing protein Syne-2. Furthermore, we found a partial colocalization of Pericentrin and Syne-2 in the ciliary region of mouse photoreceptors. Pericentrin is localized at the basal body complex of the connecting cilium while Syne-2 seems to be localized in the whole inner segment of photoreceptors.

### Conclusion

Yu et al. (2010) suggested an interaction between Syne-2 and dynein/dynactin as well as kinesin complexes as the molecular motor of nuclear migration in the mouse

retina. The interaction between Pericentrin and Syne-2 could play an essential role in interkinetic nuclear migration and may provide us new insights in the photoreceptor cell migration progress in general.

**Support:** DFG (GI770/1-1), Schmauser-Stiftung, Universitätsbund Erlangen-Nürnberg e.V.

### Authors' details

<sup>1</sup>Department of Biology, Division of Animal Physiology, FAU Erlangen-Nürnberg, Erlangen, Germany. <sup>2</sup>Institute of Human Genetics, FAU Erlangen-Nürnberg, Erlangen, Germany. <sup>3</sup>Division of Experimental Ophthalmology and Medical Proteome Center, University of Tübingen, Tübingen, Germany.

<sup>4</sup>Department of Human Genetics, Radboud University Nijmegen, Nijmegen, The Netherlands.

Published: 13 July 2015

doi:10.1186/2046-2530-4-S1-P19

Cite this article as: Falk et al.: Pericentrin interacts with KASH domain-containing protein Syne-2. *Cilia* 2015 **4**(Suppl 1):P19.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Department of Biology, Division of Animal Physiology, FAU Erlangen-Nürnberg, Erlangen, Germany

Full list of author information is available at the end of the article