



POSTER PRESENTATION

Open Access

Liposomal formulation of Gp41 derivate with adjuvant MPLA: vaccine design, immunogenicity in animals and safety in humans

D Katinger^{1*}, A Wagner¹, I Luque², S Crespillo², F Conejero-Lara², M Roger³, C Martin³, N Mouz³, S Mourao⁴, A Farsang⁵, F Notka⁶, K Malcolm⁷, N Bosquet⁸, R Le Grand⁸, C Moog⁹, A Cope¹⁰, R Shattock¹⁰, DJ Lewis¹¹, R El Habib¹²

From AIDS Vaccine 2012
Boston, MA, USA. 9-12 September 2012

Background

Gp41 of the HIV envelope, especially the MPER, contains highly conserved epitopes recognized by neutralizing monoclonal antibodies such as D5, 2F5 and 4E10. The correct presentation of the antigens is considered to be key for eliciting neutralizing immune response. The lipid bilayer of liposomes can mimic the virus surface and therefore provides the appropriate presentation of this membrane protein. In addition, liposomes can integrate the adjuvant monophosphoryl lipid A (MPLA).

Methods

The gp41 derivate FPA2 was mutated to increase its solubility and modify its structure for exposure of neutralizing epitopes. The protein was expressed in *E. coli*. The purified protein was solubilized with detergent. Liposomes were prepared containing antigen FPA2 and MPLA, a Toll-like receptor 4 agonist, using a continuous ethanol crossflow injection technology. Immunogenicity and safety was determined in rabbits and macaques. The liposomes formulated in HEC gel were applied i.n. followed by two intramuscular boosts in healthy female volunteers in a phase I study of safety and immunogenicity.

Results

Liposomes containing FPA2 and MPLA manufactured under GMP conditions were uniform and stable at 2-8°C. Immunogenicity was demonstrated in rabbits and monkey after mucosal (i.vag., i.n.) prime, followed by

intramuscular boosts. Immune response was induced systemically and in mucosal surfaces. Preliminary data from the clinical phase 1 study in healthy volunteers demonstrated good safety with three nasal applications of 200 µg of the formulated protein in HEC gel and of two i.m. boosts.

Conclusion

The liposomal formulation of the gp41 subunit of HIV envelope together with MPLA as adjuvant proved to be well tolerated in animals and human. Liposome-formulated FPA2 was able to induce binding antibodies measured by ELISA and neutralizing activity in both blood and vaginal secretions in animals. Immunogenicity testing in humans is ongoing.

Author details

¹Polymun Scientific GmbH, Klosterneuburg, Austria. ²Universidad de Granada, Granada, Spain. ³PXTherapeutics, Grenoble, France. ⁴I.M. PROJET, Miribel, France. ⁵National Food Chain Safety Office, Budapest, Hungary. ⁶GeneArt AG / Life Technologies, Regensburg, Germany. ⁷Queen's University of Belfast, Belfast, UK. ⁸CEA, Fontenay-aux-Roses, France. ⁹Université de Strasbourg, Strasbourg, France. ¹⁰Imperial College, London, UK. ¹¹University of Surrey, Guildford, UK. ¹²Sanofi Pasteur, Marcy l'Etoile, France.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P354

Cite this article as: Katinger et al.: Liposomal formulation of Gp41 derivate with adjuvant MPLA: vaccine design, immunogenicity in animals and safety in humans. *Retrovirology* 2012 **9**(Suppl 2):P354.

¹Polymun Scientific GmbH, Klosterneuburg, Austria
Full list of author information is available at the end of the article