

The Arianna thread: the matching of S-100 family with the RyR's muscle receptor

Stefania Fulle (1,2), Silvia Belia (2,3), Giorgio Fanò Illic (2,4)

(1) Department of Neuroscience, Imaging and Clinical Science, University "G. d'Annunzio" Chieti-Pescara, Italy; (2) Interuniversity Institute of Myology (IIM); (3) Department of Chemistry, Biology and Biotechnology, University of Perugia, Italy; (4) Free University of Alcatraz, Santa Cristina of Gubbio, Perugia, Italy

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Abstract

The functional state of RyR depends on the intracellular calcium concentration and on the oxidation state of its protein components in some particular sites and of some sentinel amino acids. In addition to the regulation of the RyR channel by exogenous substances (caffeine, ryanodine), ions environmental situations (oxidative state), other components, such as some endogenous proteins present in the sarcoplasm and/or in muscle membranes that are able to determine changes in Ca^{2+} channel activity. Among these, calmodulin and S-100A could determine modifications in the status of RyR channel in the skeletal muscle. The currently available data can be justified the use of a simplified S-100/CaM and RyR interaction model for the regulation of Ca^{2+} release in skeletal muscle. Under resting conditions, the CaM/S100A1 binding domain on RyR1 is predominantly dependent on S100A1. Vice versa when the intracellular Ca^{2+} concentration becomes high as well as during repetitive (tetanus) stimulation, the Ca-CaM bond becomes dominant, shifting S100A1 from RyR1 and promoting channel inactivation. This may be one of the mechanism of muscle fatigue.

Key Words: Ca^{2+} release, RyR modulation, S-100 proteins, calmodulin.

Eur J Transl Myol 30 (1): 48-52, 2020

In the mammalian skeletal muscle the contraction occurs because the intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) increases from 10^{-7} M until 10^{-5} M. The junctional Sarcoplasmic Reticulum (jSR) is involved in the release of this ion; may contains three isoforms (RyR₁, RyR₂ and RyR₃) of a particular calcium operated channel, the ryanodine Ca-channels called "feet" (juxtaposed to the voltage-dependent channels of type L-DHPR- present in the T tubule). The channels for Ca^{2+} DHPR and RyRs, form groups (4:1) called Ca^{2+} release units¹ As a result of the depolarization brought by the action potential, the voltage-sensitive L-channels cause, with a different mechanism in the skeletal and cardiac muscle, RyR channels to open, causing the release of massive amounts of Ca^{2+} into the sarcoplasm.² The functional state of RyR (closed-open-inactivated) essentially depends on the intracellular calcium concentration and on the oxidation state of its protein components in some particular sites and of some sentinel amino acids.³ Many pharmacological substances are able to modify the functional status of the channel and its sensitivity to

Ca^{2+} .⁴ Among these, the best known are ryanodine and a trimethylxanthine such as caffeine. Ryanodine added in micromolar concentrations, modifies the gating and conductance of RyR in a characteristic way bringing it into a state of subconductivity. Vice versa, if the concentration of the drug reaches the millimolar level, the channel switches into a closed state. A characteristic property of ryanodine-modified channel states is insensitivity to regulation by Ca^{2+} , Mg^{2+} , and ATP.⁵ Caffeine is bound to a specific site of the channel. In contrast to ryanodine, caffeine always has an activating action on the purified channel increasing the probability of opening and without loss of sensitivity to regulation by Ca^{2+} , Mg^{2+} and ATP.⁶ In addition to the regulation of the RYR channel by exogenous substances (caffeine, ryanodine), ions (Ca^{2+} , Mg^{2+}) environmental conditions (oxidative state), other components, such as some endogenous proteins present in the sarcoplasm and/or muscle membranes are able to directly determine changes in the Ca^{2+} channel activity state. Among these, we will consider only calmodulin and the group of proteins belonging to the S-100 family.⁷ Sarcoplasmic CaM is a 16.7 kD protein that regulates the

Matching of S-100 family with the RyR's muscle receptor

Eur J Transl Myol 30 (1): 48-52 2020




Fig 1. A, S-100 protein domains. The AA sequence and subunits (alfa and beta) of the protein share a large degree of homology. Some differences, which are used as immunological markers, are present in the hydrophilic region. The presence of two cysteine residues in the subunit can confer to this subunit some particular feature (data derived from Fanò *et al.* 1995).⁷

B, The monomeric structure (MW 10.4 Kd) consists of a repetitive EF-hand motif, whereas each Ca²⁺-binding Loop (Loop I and II) is flanked by α -helices. The N-terminal and the C-terminal EF hands are connected by a linker region (hinge region). The data reported derived from Völkers.¹²

release of SR Ca²⁺ through direct binding to RyRs or through other proteins that interact with RyRs and bind CaM. As with ryanodine, the effect of CaM on the release of Ca by the RyR channel depends on the concentration of Ca-free. For [Ca²⁺] > 1 μ M, CaM inhibits all three mammalian RyR isoforms. Vice versa, for Ca-free concentrations < 1 μ M, the activity of RyR₁ and RyR₃ channels is increased, while RyR₂ is inhibited. From this comes the hypothesis of a physiological role for CaM which, by prolonging the channel closure state at [Ca²⁺] greater than 10⁻⁶ M, would allow the ion to spread in the sarcomere, away from the release sites.⁸

Studies with CaM-free mutants indicate that if CaM's binding to RyR₂ is altered, cardiac function is impaired.⁹ The S100 protein was named S100 because of its solubility in a 100% saturated solution with ammonium sulphate.¹⁰ The term S100 is used to embrace a multigenic family of mostly dimeric calcium-binding proteins.¹¹ In nature can be found both as a low molecular weight monomer (alpha or beta) and as an alpha-alpha (S100A) and beta-beta (S-100B) homodimer. Each chain has a molecular weight of 10-12 kDa.

The S100 proteins are highly conserved in amino acid composition among vertebrate species, suggesting that they may have crucially conserved biological roles.⁷

S100 proteins are small, acidic proteins and contain two

distinct EF-hands, 4 α -helical segments, a central hinge region of variable length and the N- and C-terminal variable domains (Figure 1). At present, at least 25 proteins have been identified as belonging to the S100 protein family, 21 of them having genes clustered at chromosome locus 1q21.¹² S100 proteins do not have intrinsic catalytic activity. They are generally thought to be that, in a similar manner to calmodulin and troponin C, undergo conformational changes and modulate biological activity via calcium binding. The two subunits have many aspects in common and each of them is characterized by two Ca²⁺ site binding: one at high affinity (KD= 10⁻⁵ M) on the C-terminal portion and another unconventional with low affinity for the ion.¹³ Even though the members of the S100 family share a remarkable sequence homology, they express themselves in specific tissue patterns. S100A is the most highly expressed member of the family in striated muscle, where the protein shows the highest presence in the heart muscle, followed by slow contraction red fibres and finally white, fast fibers.¹⁴

To date, a considerable amount of data has accumulated to hypothesize a functional role for S100A in cardiac muscle, but research on a possible role for S100 in skeletal muscle is still incomplete.¹⁵ The first experiments involving protein intervention in the

Matching of S-100 family with the RyR's muscle receptor

Eur J Transl Myol 30 (1): 48-52 2020




Fig 2. (a) Effect of S-100 on Ca^{2+} uptake and release in vesicles derived from SR fragmentation of skeletal muscle of *Rana esculenta*. The uptake was induced by 5mM ATP plus 5.0 μM S100; (b) Specific binding of 3-5 nM ryanodine in presence of 1 μM S100. (c) S100A1 modulation of skeletal muscle EC coupling in genetic (S100^{-/-}) ablated fibers. Data reported in part (a) and (b) derived from Fanò et al.¹⁷ while part (c) represents the result of observations carried out from Prosser et al.¹⁸

excitation-contraction cycle in skeletal muscle, were performed by our group and showed that S100A improved the release of RyR₁ Ca^{2+} in SR terminal cisternae preparations (Figure 2). The most interesting fact of these experiments was to highlight how, despite belonging to the same superfamily as the EF-hand calcium binding protein, CaM and S100A seemed to have contrasting effects on the activation of Ca^{2+} release via RyR₁.^{16,17}

Several years later Prosser and collaborators have shown that the effective concentration of S100A available locally for modulation of RyR₁ is probably higher. However, as has been demonstrated, both ultrastructurally and immunofluorescently, S-100 is located at the overlap area of the A/I band and at the

tetrads, where the release of Ca^{2+} into the skeletal muscle occurs.¹⁸ Therefore, it should come as no surprise that S-100 can act as a regulator of the functional status of RyR channels as it has been assumed for another protein with similar characteristics, the calmodulin.¹⁹ In experiments using vesicles derived from the SR fraction containing the RyR channels preloaded with Ca^{2+} , the addition of S-100 in mM concentration increased the release of ion induced by the presence of [Ca^{2+}] mM in the experimental medium. This may be interpreted as an effect of S-100 on the activation time of the channel, which becomes, thus, longer. In the same experimental model, CaM induces inhibition of Ca release while the use of S100 inactivated is not able to increase the release of the ion from the vesicles previously loaded with Ca in

Matching of S-100 family with the RyR's muscle receptor

Eur J Transl Myol 30 (1): 48-52 2020

the presence of ATP.²⁰

The currently available data can be justified if one uses a simplified S-100/CaM and RyR interaction model for the regulation of Ca²⁺ release in skeletal muscle. Under this assumption, and in resting conditions, the CaM/S100A1 binding domain on RyR₁ is predominantly dependent on S100A1, which increases Ca²⁺ release and force generation when the muscle is stimulated. When the [Ca²⁺]_i becomes high during repetitive (tetanus) stimulation, the Ca-CaM bond becomes dominant, shifting S100A1 from RyR₁ and promoting channel inactivation.¹⁸

In this way, CaM and S100A1 compete to fine-tune the release of Ca²⁺ from RyR during normal muscle activation. The interruption of this system leads to a decrease in force production and a reduction in muscle performance. However, the molecular details of these interactions and the consequences of their dysregulation on channel activity are not yet known.¹⁹ Nevertheless, recent studies based on the FRET technique, have suggested that CaM and S100A can simultaneously, but separately, modulate the RyR function without S100A competing for CaM on its specific channel site.²⁰

In conclusion, the data that are available today indicate that there is a mechanism, in the skeletal muscle, able to modulate finely the functional state of the RyR channel and therefore the available Ca²⁺ necessary for the contraction. This mechanism is represented by the joint interaction of CaM and S100A on specific common RyR sites. However, fundamental information is still missing to understand the molecular dynamics of this interaction. Furthermore, there are conditions not fully studied, in which the competition between the two calcium binding proteins for the same site changes (during exercise, oxidative stress, fatigue, etc.). Nevertheless, there is strong evidence that the interaction of CaM and S100A can regulate skeletal muscle EC coupling.²⁰⁻²² This kind of balanced control could also be at the basis of the onset of the phenomenon of fatigue, which does not yet have a precise determinant because the accumulation of lactate does not seem to be sufficient to explain its occurrence. If this hypothesis will be confirmed it could represent a turning point in the definition of the functional role for the protein. S100 could finally emerge from the labyrinth in which it has been confined since its discovery. Of course, no one wants it to end up like Theseus.

List of acronyms

ATP - Adenosine Triphosphate

CaM - Calmodulin

EC - Excitation Contraction

FRET - Fluorescence Resonance Energy Transfer.

RyR - Ryanodine Receptor

Authors contributions

Giorgio Fanò-Illic played a main role in the writing of the manuscript, Silvia Belia in the drafting and finalizing the

manuscript, Stefania Fulle in the extensive manuscript revision.

Funding

The A&C M-C Foundation for Translational Myology, Padova, Italy sponsored the publication of this manuscript.

Conflict of Interest

The authors declare they have no financial, personal, or other conflicts of interest.

Ethical Publication Statement

We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

Corresponding Author

Giorgio Fanò-Illic, Interuniversity Institute of Myology (IIM) and Free University of Alcatraz, Santa Cristina of Gubbio, Perugia, Italy.

ORCID iD: 0000-0002-8753-5707

Email: fanoillic@gmail.com

E-mail and ORCID iD of co-authors

Silvia Belia: silvia.belia@unipg.it

ORCID iD: 0000-0002-0323-4959

Stefania Fulle: stefania.fulle@unich.it

ORCID iD: 0000-0003-4557-9127

References

1. Franzini-Armstrong C, and Protasi F. Ryanodine receptors of striated muscles: A complex channel capable of multiple interactions. *Physiol Rev* 1997;77:699-729. doi: 10.1152/physrev.1997.77.3.699.
2. Fabiato A. Calcium-induced release of calcium from the cardiac sarcoplasmic reticulum. *Am J Physiol* 1983;245:C1-C14. doi: 10.1152/ajpcell.1983.245.1.C1.
3. Sun QA, Wang B, Miyagi M, et al. Oxygen-coupled redox regulation of the skeletal muscle ryanodine receptor/Ca²⁺ release channel (RyR1): Sites and nature of oxidative modification. *J Biol Chem* 2013;288:22961-71. doi: 10.1074/jbc.M113.48022 8.
4. Meissner G. Ryanodine receptor/Ca²⁺ release channels and their regulation by endogenous effectors. *Annu Rev Physiol* 1994;56:485-508. doi: 10.1146/annurev.ph.56.030194.002413.
5. Ngo VA, Perissinotti LL, Miranda W, et al. Mapping ryanodine binding sites in the pore cavity of ryanodine receptors. *Biophys J* 2017;112:1645-53. doi: 10.1016/j.bpj.2017.03.014.
6. Liu W, Meissner G. Structure-activity relationship of xanthines and skeletal muscle ryanodine receptor/Ca²⁺ channel release. *Pharmacology* 1997;54:35-43. doi: 10.1159/000139480.
7. Fanò G, Biocca S, Fulle S, et al. The S-100: a protein family in search of a function. *Prog*

Matching of S-100 family with the RyR's muscle receptor

Eur J Transl Myol 30 (1): 48-52 2020

- Neurobiol 1995;46:71-82. doi: 10.1016/0301-0082(94)00062-m.
8. Xu L, Meissner G. Mechanism of calmodulin inhibition of cardiac sarcoplasmic reticulum Ca²⁺ release channel (ryanodine receptor). *Biophys J* 2004;86:797-804. doi: 10.1016/S0006-3495(04)74155-7.
 9. Yamaguchi N, Chakraborty A, Huang TQ, et al. Cardiac hypertrophy associated with impaired regulation of cardiac ryanodine receptor by calmodulin and S100A1. *Am J Physiol Heart Circ Physiol* 2013;305:H86-H94. doi: 10.1152/ajpheart.00144.2013.
 10. Moore BW. A soluble protein characteristic of the nervous system. *Biochem Biophys Res Commun* 1965;19:739-744. doi: 10.1016/0006-291x(65)90320-7.
 11. Baudier J, Gerard D. Ions binding to S100 proteins. Conformational studies and calcium-induced conformational changes in S100 alpha protein: the effect of acidic pH and calcium incubation on subunit exchange in S100a (alpha beta) protein. *J Biol Chem* 1986;261:8204-12.
 12. Volkens M, Rohde D, Goodman C, et al. S100: a regulator of striated muscle sarcoplasmic reticulum Ca²⁺ handling, sarcomeric and mitochondrial function. *J Biomed Biotechnol* 2010;2010:178614. doi: 10.1155/2010/178614.
 13. Ikura M. Calcium binding and conformational response in EFhand proteins. *Trends Biochem Sci* 1996;21:14-7.
 14. Adhikari BB, Wang K. S100A1 modulates skeletal muscle contraction by desensitizing calcium activation of isometric tension, stiffness and ATPase. *FEBS Lett* 2001;497:95-8. doi: 10.1016/s0014-5793(01)02444-9.
 15. Riuzzi F, Sorci G, Arcuri C, et al. Cellular and molecular mechanisms of sarcopenia: the S100B perspective. *J Cachexia Sarcopenia Muscle* 2018;9(7):1255-1268. doi: 10.1002/jcsm.12363.
 16. Fanò G, Marsili V, Angelella P, et al. S-100a0 protein stimulates Ca²⁺-induced Ca²⁺ release from isolated sarcoplasmic reticulum vesicles. *FEBS Lett* 1989;255:381-4. doi: 10.1016/0014-5793(89)81127-5.
 17. Fanò G, Marsili V, Protasi F, et al. Relationship between an endogenous calcium binding protein (S-100ab) and Ca²⁺ transport in frog skeletal muscle. *Basic Applied Myol (BAM) - Eur J Transl Myol* 1992;2:309-15.
 18. Prosser BL, Hernandez-Ochoa EO, Schneider MF. S100A1 and calmodulin regulation of ryanodine receptor in striated muscle. *Cell Calcium* 2011;50:323-3131. doi: 10.1016/j.ceca.2011.06.001.
 19. Yamaguchi N, Prosser BL, Ghassemi F, et al. Modulation of sarcoplasmic reticulum Ca²⁺ release in skeletal muscle expressing ryanodine receptor impaired in regulation by calmodulin and S100A1. *Am J Physiol Cell Physiol* 2011;300:C998-C1012. doi: 10.1152/ajpcell.00370.2010.
 20. Marsili V, Mancinelli L, Menchetti G, et al. S-100ab increases Ca²⁺ release in purified sarcoplasmic reticulum vesicles of frog skeletal muscle. *J Muscle Res Cell Motil* 1992;13:511-55. doi: 10.1007/bf01737993.
 21. Bers D. Calmodulin and S100A1 fine tune skeletal muscle E-C coupling (editorial) *Am J Physiol Cell Physiol* 2011;300:C974-5.
 22. Rebbeck RT, Nitu FR, Rohde P, et al. S100A1 protein does not compete with calmodulin for ryanodine receptor binding but structurally alters the ryanodine receptor-calmodulin complex. *J Biol Chem* 2016;291:15896-907. doi: 10.1074/jbc.M115.713107.

Submitted: January 20, 2020

Accepted for publication: January 31, 2020