


## Fast endoscopic rendezvous to the rescue after maldeployment of lumen-apposing metal stent during endoscopic ultrasound-guided choledochoduodenostomy

OPEN  
ACCESS



**▶ Video 1** Maldeployment of lumen-apposing metal stent during endoscopic ultrasound choledochoduodenostomy managed by advancing a guidewire directly through the choledochal breach followed by endoscopic retrograde cholangiopancreatography rendezvous with transpapillary biliary stent placement and clips to seal the perforation.



**▶ Fig. 1** Evidence of gross biliary leak and bile duct opacification.



**▶ Fig. 2** Biliary stent and clip closure of the duodenal defect.



**▶ Fig. 3** Final occlusion cholangiogram showed no calculi or biliary leak.

A 70-year-old woman with acute cholangitis was referred to our unit for treatment. Complex choledocholithiasis with impacted stones prevented endoscopic retrograde cholangiopancreatography (ERCP) and the patient underwent biliary drainage by endoscopic ultrasound (EUS)-guided choledochoduode-

nostomy (CDS) using a cautery-enhanced 8×8-mm lumen-apposing metal stent (LAMS) over a guidewire. After LAMS release, EUS imaging revealed maldeployment of the distal flange of the LAMS between the common bile duct (CBD) and duodenal wall. This was probably due to an angled scope tip and diffi-

culty in manipulating the guidewire (stripping of the guidewire occurred) (**▶ Video 1**).

Passage of purulent fluid was visible through the LAMS endoscopically. The EUS scope was replaced with a gastro-scope to easily inject contrast through the LAMS. The fluoroscopic view showed a subhepatic leak and quick opacification of the biliary tree (**▶ Fig. 1**).

Careful inspection through the LAMS lumen using the front view gastro-scope enabled us to identify the breach in the CBD: a guidewire was advanced antegrade through a 5-Fr biliary catheter into the CBD to the duodenum. This allowed transpapillary insertion of a guidewire with a sphincterotome using a duodenoscopy. Then a fully covered metal stent was inserted to enable biliary drainage and closure of the perforation. The LAMS was removed, and the duodenal perforation was closed using two clips (**▶ Fig. 2**). The post-procedure period was uneventful and the patient was discharged after 6 days of hospitalization. Four weeks later the biliary stent was removed, choledo-

chal stone clearance was completed, and the occlusion cholangiogram confirmed that the biliary leak was sealed (► **Fig. 3**). Maldeployment of a LAMS during EUS-CDS can lead to a double perforation and requires prompt recognition and management [1]. When feasible, endoscopic rescue therapies are effective in managing this serious adverse event [2–5]; endoscopists should be aware of possible solutions to a serious complication that would otherwise require a surgical approach.

Endoscopy\_UCTN\_Code\_CPL\_1AK\_2AD

### Competing interests

The authors declare that they have no conflict of interest.

### The authors

**Elia Armellini<sup>1</sup>**, **Flavio Metelli<sup>1</sup>**, **Antonio Piazzini Albani<sup>2</sup>**, **Lorenzo Vescovi<sup>2</sup>**, **Luisella Spinelli<sup>2</sup>**, **Fabio Pace<sup>1</sup>**

- 1 ASST Bergamo Est, Gastroenterology Department, Seriate, Bergamo, Italy
- 2 ASST Bergamo Est, Surgery Department, Seriate, Bergamo, Italy

### Corresponding author

**Flavio Metelli, MD**  
ASST Bergamo Est, Gastroenterology  
Department, Via Paderno, 21, 24068 Seriate,  
BG, Italy  
flaviomet90@gmail.com

### References

- [1] Fabbri C, Coluccio C, Binda C et al. Lumen-apposing metal stents: How far are we from standardization? An Italian survey. *Endosc Ultrasound* 2022; 11: 59–67
- [2] Di Mitri R, Amata M, Mocciano F et al. EUS-guided biliary drainage with LAMS for distal malignant biliary obstruction when ERCP fails: single-center retrospective study and maldeployment management. *Surg Endosc* 2022; 36: 4553–4569
- [3] Sanchez-Ocana R, de Benito Sanz M, Nájera-Muñoz R et al. Rendezvous ERCP via endoscopic ultrasound-guided gallbladder drainage to salvage a dislodged lumen-apposing metal stent during choledochoduodenostomy. *Endoscopy* 2022; 54: 65–67
- [4] Fugazza A, Fabbri C, Di Mitri R et al. EUS-guided choledochoduodenostomy for malignant distal biliary obstruction after failed ERCP: a retrospective nationwide analysis. *Gastrointest Endosc* 2022; 95: 896–904
- [5] Graves M, Krafft M, Nasr J. EUS-guided choledochoduodenostomy creation using a biliary fully covered self-expanding metal stent after maldeployment of lumen-apposing metal stent. *VideoGIE* 2021; 6: 234–235

### Bibliography

*Endoscopy* 2023; 55: E37–E38  
DOI 10.1055/a-1929-9148  
ISSN 0013-726X  
published online 22.9.2022  
© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,  
70469 Stuttgart, Germany



### ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is an open access online section, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>