

# Older Adults and Immune Thrombocytopenia: Considerations for the Clinician

Etienne Crickx<sup>1,2</sup>, Matthieu Mahévas<sup>1,3,4</sup>, Marc Michel<sup>1</sup>, Bertrand Godeau<sup>1</sup>

<sup>1</sup>Service de Médecine Interne, Centre national de référence des cytopénies auto-immunes de l'adulte, Hôpital Henri Mondor, Fédération Hospitalo-Universitaire TRUE InnovaTive theRapy for immUne disordErs, Assistance Publique Hôpitaux de Paris (AP-HP), Université Paris Est Créteil, Créteil, France; <sup>2</sup>Université de Paris, Imagine Institute, Laboratory of Immunogenetics of Pediatric Autoimmune Diseases, Paris, F-75015, France; <sup>3</sup>Institut Necker Enfants Malades (INEM), INSERM U1151/CNRS UMS 8253, ATIP-Avenir Team AI2B, Université de Paris, Université Paris-Est-Créteil, Paris, France; <sup>4</sup>INSERM U955, équipe 2, Université Paris-Est Créteil (UPEC), Créteil, France

Correspondence: Bertrand Godeau, Service de Médecine Interne, Hôpital Henri Mondor, Assistance Publique Hôpitaux de Paris (AP-HP), 51 avenue du maréchal de Lattre de Tassigny, Créteil, 94000, France, Tel +331 49 81 29 05, Fax +331 49 81 29 02, Email bertrand.godeau@aphp.fr

**Abstract:** Many epidemiological studies have shown that the incidence of immune thrombocytopenia (ITP) increases after age 60 years and peaks in patients over age 80 years. Therefore, ITP is a concern for physicians taking care of older patients, especially regarding its diagnosis and management. The diagnostic work-up should exclude other causes of thrombocytopenia and secondary ITP, including myelodysplastic syndrome and drug-induced ITP. The treatment decision is influenced by an increased risk of bleeding, infectious diseases and thrombosis in this population and should take into account comorbidities and concomitant medications such as anticoagulant drugs. First-line treatment is based on short corticosteroids courses and intravenous immunoglobulin, which should be reserved for patients with more severe bleeding complications, with their higher risk of toxic effects as compared with younger patients. Second-line treatment should be tailored to the patient's history, comorbidities and preferences. Preferred second-line treatments are thrombopoietin receptor agonists for most groups and guidelines given their good efficacy/tolerance ratio, but the thrombotic risk is increased in older people. Other second-line options that can be good alternatives depending on the clinical context include rituximab, dapsone, fostamatinib or immunosuppressive drugs. Splenectomy is less often performed but remains an option for fit patients with chronic refractory disease. Emerging treatments such as Syk or Bruton tyrosine kinase inhibitors and FcRn antagonists are becoming available for ITP and may modify the treatment algorithm in the near future. The aim of this review is to describe the particularities of the diagnosis and treatment of ITP in older people, including the response and tolerance to the currently available drugs. We also discuss some situations related to co-morbidities that can frequently lead to adapt the management strategy in older patients.

**Keywords:** immune thrombocytopenia, ITP, elderly, intravenous immunoglobulin, IVIg, thrombopoietin receptor agonists, splenectomy, rituximab

## Introduction

Immune thrombocytopenia (ITP) is an autoimmune disease characterized by antibody-mediated platelet destruction and impaired platelet production resulting in bleeding symptoms.<sup>1</sup> Although it can affect individuals of all age categories, the disease incidence peaks in older patients.<sup>2,3</sup> Hence, combined with the worldwide trend in the ageing of the population, ITP is of particular interest for physicians taking care of older patients. ITP management is challenging in older versus younger patients given the frequent comorbidities and increased risk of bleeding, infections and thrombosis of the former group.<sup>4,5</sup>

A growing number of studies focusing on older patients with ITP are now available, as are new treatments for ITP. In this review, we provide an update on the diagnosis, prognosis and treatment of older patients with ITP in light of these recent data. We also discuss some situations related to co-morbidities that can frequently lead to adapt the management strategy in older patients. To date, no prospective study focusing on this population has been conducted and therefore most recommendations presented here are not evidence-based but rather extrapolated from observational and retrospective studies as well as our own experience.

# ITP Diagnosis and Epidemiology

## Epidemiology

Several large epidemiological studies have shown that the ITP epidemiology is influenced by sex and age,<sup>2,3,6</sup> with peaks in young women and old men. ITP is also a geriatric disease, with incidence rates reaching 23.9/100 000 in men >80 years old in the United Kingdom<sup>3</sup> and 9/100 000 person-years in men >75 years old in France,<sup>2</sup> that is, an approximately 10-fold increase as compared with men aged 30 to 39 years in both studies. In a recent French study including 541 adults with incident ITP included in a prospective national registry, 251 (46%) were  $\geq 65$  years and among them, 47% were  $\geq 80$  years. In this later group of very old patients, 37.9% were exposed to antiplatelet drugs and 18.4% to anticoagulants.<sup>7</sup>

## Diagnosis

According to international guidelines, primary ITP is defined by isolated thrombocytopenia  $< 100 \times 10^9/L$  of an autoimmune origin in the absence of any underlying cause or disorder.<sup>8</sup> ITP usually presents as isolated thrombocytopenia, and the diagnostic work-up mainly focuses on eliminating other etiologies because of no gold-standard diagnostic test. Secondary ITP refers to immune thrombocytopenia associated with other conditions (eg, hematological malignancies, systemic lupus, primary immunodeficiencies) at diagnosis. The main differential diagnoses of thrombocytopenia and causes of secondary ITP are shown in Table 1.

Patient history, physical examination, complete blood count and peripheral blood smear examination are the cornerstone of ITP diagnosis. In addition, some exams of interest to identify particular situations or secondary ITP include serologies for HIV, hepatitis C and B virus, antinuclear antibodies and protein serum electrophoresis. Other blood tests should be oriented to the clinical context. Because of lack of sensitivity and specificity, the search for antibodies against platelet antigens is not recommended in routine practice.<sup>8,9</sup>

However, the diagnostic approach proposed in the international guidelines is a global approach that addresses adults without taking into account age. In older patients, some points deserve consideration in the ITP diagnostic work-up. First,

**Table 1** Other Main Causes of Thrombocytopenia

Secondary ITP	Non-Immune Thrombocytopenia
<p><b>Associated auto-immune disorder</b></p> <ul style="list-style-type: none"> <li>Evans syndrome</li> <li>Systemic lupus erythematosus</li> <li>Antiphospholipid syndrome</li> <li>Other auto-immune disorders</li> </ul> <p><b>Drug-induced ITP and vaccine-induced ITP</b></p> <p><b>Infectious diseases</b></p> <ul style="list-style-type: none"> <li>HIV</li> <li>HCV</li> <li>Helicobacter pylori</li> </ul> <p><b>Lymphoproliferative disorders</b></p> <ul style="list-style-type: none"> <li>Chronic lymphocytic leukemia</li> <li>Indolent B-cell lymphoma</li> <li>Angioimmunoblastic T-cell lymphoma</li> </ul> <p><b>Primary immune deficiencies/inborn error of immunity</b></p> <ul style="list-style-type: none"> <li>Common variable immune deficiency</li> <li>Autoimmune lymphoproliferative syndrome</li> </ul>	<p><b>Splenic sequestration</b></p> <ul style="list-style-type: none"> <li>Portal hypertension</li> <li>Splenomegaly</li> <li>Splenic infiltrative diseases</li> </ul> <p><b>Decreased production</b></p> <ul style="list-style-type: none"> <li>Inherited thrombocytopenia</li> <li>Primary and secondary bone-marrow failures</li> <li>Hematological drug toxicity, alcohol toxicity</li> <li>Folate and/or B12 deficiency</li> <li>Viral infections: HIV, HBV, HCV, EBV, CMV, parvovirus B19</li> </ul> <p><b>Peripheral platelet consumption</b></p> <ul style="list-style-type: none"> <li>Disseminated intravascular coagulation</li> <li>Thrombotic microangiopathy</li> <li>Sepsis</li> </ul>
	<b>Alloimmune thrombocytopenia</b>

**Abbreviations:** ITP, immune thrombocytopenia; CMV, cytomegalovirus; EBV, Epstein-Barr virus; HBV, hepatitis B virus; HCV, hepatitis C virus; HIV, human immunodeficiency virus.

drug-induced ITP should be considered because exposure to treatments increases with age. Several mechanisms can be responsible for drug-induced ITP, but most commonly, platelet destruction is mediated by a drug-dependent antibody.<sup>10</sup> Patients usually present profound thrombocytopenia and bleeding symptoms soon (in a few hours to a few days) after drug introduction. Because *in vitro* confirmation of drug-induced ITP is usually unavailable, the diagnosis relies on a complete history of drug intake, the relative chronology with thrombocytopenia, and spontaneous recovery upon drug discontinuation. Many drugs but also vaccines have been implicated in drug-induced ITP and updated reviews have been published recently.<sup>10–12</sup>

## Interest of Bone-Marrow Evaluation (BME) in Older Patients

Another differential diagnosis of thrombocytopenia in older patients is myelodysplastic syndrome (MDS) affecting megakaryocyte lineage. Although most MDS cases exhibit associated thrombocytopenia (65%),<sup>13</sup> isolated thrombocytopenia is less frequent (12% of cases).<sup>14</sup> Therefore, the presence of other cytopenia or macrocytosis on complete blood count and signs of dyserythropoiesis and/or dysgranulopoiesis on peripheral blood smear examination should prompt a BME to rule out a myelodysplastic syndrome. Notably, patients with early ITP can present some abnormalities in megakaryocyte morphology as well as some reticulin fiber deposits on bone-marrow biopsy.<sup>15,16</sup> The coexistence of MDS and ITP has been described.<sup>17</sup>

We suggest to use a pragmatic approach and to propose a short course of corticosteroids or intravenous immunoglobulin (IVIg) when suspecting ITP. Although not validated in prospective studies, a response to these treatments suggests an immunological contribution to thrombocytopenia but does not rule out MDS.<sup>5</sup>

Performing systematic BME in patients >60 years presenting isolated thrombocytopenia has been debated in the past years. However, growing evidence suggests that the rate of abnormality on bone-marrow smear in this context is very low. In the prospective French registry, only one in 197 (0.8%) patients with ITP and older than 60 years had an abnormal bone-marrow smear result.<sup>18</sup> In a recent retrospective study of 324 patients with relapsed/refractory ITP, 56% had a BME, which resulted in eight other diagnoses (8% of tested patients). Of note, age was not associated with abnormal findings on BME in this study.<sup>19</sup>

Current international guidelines recommend BME only in patients with systemic symptoms, abnormal signs, or a suspected different diagnosis independent of age. BME can also be performed before splenectomy or in case of treatment failure.<sup>8,20</sup> If performed in older patients to rule out MDS, a cytogenetic study and next-generation sequencing should be performed at the same time; flow cytometry analysis and bone-marrow biopsy should be performed in case of suspected lymphoma.<sup>8</sup>

## ITP Prognosis: What Particularities in Older Patients?

Given the increased frequency of ITP in older patients, interest has been growing in determining whether the disease course, severity and complications are similar to those of younger patients. Infectious and thrombotic risk are also relevant when considering therapeutic options, especially in older patients.

## Disease Course

ITP can be classified according to disease duration as newly diagnosed (0–3 months), persistent (3–12 months), or chronic (>12 months).<sup>8</sup> In adults, newly diagnosed ITP evolves toward chronic disease in about 70% of cases. Chronic ITP has a very low probability of spontaneous recovery.<sup>21</sup> Whether older patients diagnosed with ITP are at increased risk of chronic disease is unknown.

## Bleeding Risk

Data from the literature are conflicting about bleeding risk in older patients. Some studies did not find an association between age and bleeding,<sup>22,23</sup> and others found more bleeding in older patients with similar platelet counts.<sup>2,24–31</sup> In a recent series comparing 311 patients aged <65 years to 154 aged ≥65 years, Palandri et al found no differences in platelet count at diagnosis or in bleeding symptoms but more grade 3/4 bleeding and thrombosis in older patients, with a first-line therapy started for higher platelet counts.<sup>32</sup> Several other factors such as comorbidities and drug exposure could influence

the bleeding risk, particularly in older patients. Comorbidities such as cardiovascular diseases, kidney or liver failure, or gastrointestinal diseases increase in number with age.<sup>4,7,29</sup> In a French cross-sectional study, age was significantly associated with severe bleeding on univariate analysis but not after adjustment for other covariates, including anti-coagulant drugs. In this study, platelet count, female sex and exposure to non-steroidal anti-inflammatory drugs were associated with risk of any bleeding, and exposure to anticoagulant drugs was a major risk factor for severe bleeding.<sup>33</sup>

The definition of “older” has also varied among studies and may explain some discrepancies. In a recent study from our group, the disease course of very old patients (aged  $\geq 80$  years) was compared to those of old patients (aged 65–79 years). Although platelet counts at ITP diagnosis were similar, severe bleeding and mortality were more frequent in very old patients although not significantly. Comorbidities were also more frequent, and exposure to anticoagulant drugs was strongly associated with severe bleeding.<sup>7</sup>

Overall, these data suggest that platelet count and bleeding symptoms at ITP onset are probably independent of age as well as disease course but that older patients probably are at increased risk of severe bleeding, especially when exposed to anticoagulant drugs. In other words, the care of a patient over age 65 years without co-morbidities and not exposed to anticoagulants can probably be modelled on that of younger patients. However, a more aggressive strategy and a higher platelet threshold to decide to treat should probably be considered in very old patients, especially in the presence of a co-morbidity and exposure to anticoagulants because of greater risk of bleeding complications.

## Thrombosis and Infection

Multiple factors can contribute to arterial and venous thrombosis observed during the ITP course.<sup>34</sup> Paradoxically, ITP could be a prothrombotic disease, and thrombosis can occur even with low platelet counts.<sup>35,36</sup> Thrombotic risk also increases with age,<sup>37,38</sup> and cardiovascular risk factors are more frequent in older patients. Some ITP treatments that could favor thrombosis include IVIg,<sup>39</sup> splenectomy<sup>40</sup> or thrombopoietin receptor agonists (TPO-RAs).<sup>41–45</sup> In this context, the addition of treatment-related prothrombotic factors to patient- and ITP-related prothrombotic factors should be weighed carefully.

Older patients may be more sensitive to infections than younger patients. Infectious risk is also increased in ITP probably because of immunosuppressive drug use and splenectomy,<sup>50</sup> and contribute to mortality.<sup>51,52</sup> This is important to consider for ITP management and treatment decisions in older patients with a possibly increased risk of infection observed with some treatments such as corticosteroids, immunosuppressive drugs and rituximab.

## ITP Treatments: What Tolerance and What Efficacy in Older Patients?

### Treatment Decision

The treatment decision is mainly based on bleeding symptoms and platelet count, although other factors important to consider include disease duration, other medications, comorbidities, expected tolerance, accessibility of care, quality of life and patient expectations. Age but more importantly frailty also influence the treatment choice, but in any case, treatment should always be tailored to the patient, and guidelines highlight that ITP treatment should be personalized. This is especially true for older people because for the same number of platelets, many individual factors can influence the decision. Also, patients' concerns and fears and disease burden are often misjudged by the physician.<sup>53,54</sup> Therefore, the decision to treat must always be shared with the patient, as it has been shown in other hematologic diseases.<sup>55</sup> If the patient has high functional impairment, a fair discussion with the caregivers outlining the benefits, limitations and side effects of treatment is essential.

International guidelines have long proposed a threshold of  $30 \times 10^9/L$  platelets to indicate treatment, even in a patient without bleeding.<sup>56</sup> We now have considerable evidence to suggest that this threshold can be lowered to  $20 \times 10^9/L$ .<sup>33</sup> Because bleeding risk clearly increases with platelet count  $< 20 \times 10^9/L$ , particularly in older patients (see above), the recommendation is to treat older patients with platelet count below this threshold.<sup>8</sup> However, bleeding symptoms secondary to thrombocytopenia are unusual with platelet count  $> 50 \times 10^9/L$ , and a watch-and-wait approach should be proposed in this case unless a higher platelet count is required for an intervention.<sup>8</sup>

When platelet counts are 20 to 50 x 10<sup>9</sup>/L, no evidence-based medicine is available and the treatment decision should be based on the presence of additional risk factors of bleeding such as a history of bleeding; co-morbidities (eg, severe hypertension, renal or liver failure, peptic ulcer); age >80 years and frailty, including risk of falls; and drug exposure. In this setting, a watch-and-wait approach is easier when a rapid response to corticosteroids or IVIg is known from previous treatment courses, allowing for safe management of the potential occurrence of a bleeding complication.

When the patient is taking drugs interfering with hemostasis, the referring physician should be contacted to assess the risk/benefit balance. These drugs should not eventually be discontinued when indicated (except transiently when platelet counts are very low [ie, <10 x 10<sup>9</sup>/L] or in case of bleeding) because a low platelet count does not preclude thrombotic risk.<sup>35,36</sup> ITP treatment should be considered to maintain a safe platelet count. Anticoagulant drugs have been associated with increased risk of severe bleeding, and some experts and epidemiological studies argue that maintaining the platelet count above 50 x 10<sup>9</sup>/L is safer.<sup>7,33,57</sup> Although anti-platelet drugs interfere with primary hemostasis, the bleeding risk seems lower with these agents,<sup>7,33</sup> perhaps because platelet turnover is increased in ITP. Therefore, maintaining a higher platelet count is probably safer when the patient takes anti-platelet drugs, although the optimal threshold is unknown, but a threshold of 30 x 10<sup>9</sup>/L could probably be tolerated.


If a watch-and-wait approach is favored, patients should have a close monitoring of platelet count and bleeding symptoms. Our current strategy for ITP treatment decisions in older patients is summarized in Figure 1.

## First-Line Treatments

### Corticosteroids

ITP first-line therapy relies on a short course of corticosteroids.<sup>8</sup> Long-term corticosteroids should be avoided in most cases because they are poorly effective in maintaining remission at a low dose and do not affect the ITP natural history while generating side effects.<sup>58,59</sup> Prednisone 1 mg/kg for 3 weeks or dexamethasone 40 mg/day for 4 days (repeated up to three times if necessary) can be used. A number of controlled studies have suggested a faster and greater platelet response with dexamethasone over prednisolone, but the rate of long-term response was similar in both groups.<sup>60,61</sup>

Age does not affect the response to corticosteroids, but this treatment generates more adverse events in older patients.<sup>27,29</sup> Tolerance remains acceptable with short courses of corticosteroids: in a retrospective study of 440 patients aged >60 years who received corticosteroids (prednisone or dexamethasone) as first-line ITP therapy, only 16 (3.7%)



**Figure 1** Proposed algorithm for treatment decision. In patients with immune thrombocytopenia (ITP) without anticoagulant drugs and with platelet counts of 20 to 50 x 10<sup>9</sup>/L, treatment can be delayed if there are no bleeding risk factors but should be considered if there are more than 2 bleeding risk factors.

discontinued treatment because of toxic effects (mainly diabetes and psychiatric symptoms).<sup>62</sup> The risk of infection is also increased with corticosteroids but mainly with prolonged treatment, which should be strongly avoided in ITP.<sup>63</sup>

Several studies have suggested lower than standard corticosteroids dose regimens as an alternative strategy.<sup>26,64</sup> We frequently use a reduced dexamethasone dose of 20 mg/day (instead of 40 mg/day) in frail patients. Although particularly relevant in older patients to limit toxic effects, this dose remains to be validated in this particular population.

### Intravenous Immunoglobulin (IVIg)

A combined course of IVIg and corticosteroids has been found more effective than corticosteroids alone for time to response, platelet count and response duration.<sup>65</sup> Therefore, IVIg is a treatment of choice for patients with a high bleeding score,<sup>66</sup> although its cost, availability, transient response (less than 1 month) and potential adverse events limits its more general use.<sup>8</sup> The recommended dose is 1 g/kg/day, which can be repeated on day 2 in case of severe bleeding with a life-threatening situation or on day 3 with less severe bleeding and in case of lack of response after the first infusion.<sup>67</sup> Side effects include anaphylaxis, hemolytic reactions, and minor symptoms such as headache, nausea, or fever. Other adverse events such as renal failure, fluid overload and thrombosis seem more frequent in older patients and justify a lower daily dosage (0.5 g/kg days 1–4 or 0.4 g/kg days 1–5), with careful monitoring of hydration and renal function.<sup>39,68–70</sup>

### Anti-Rho(D) Drugs

Intravenous anti-Rho(D) drugs are not licensed in Europe but have been proposed to treat Rho(D)-positive patients who have not undergone splenectomy, with high response rates at the cost of rare but life-threatening side effects.<sup>71</sup> Fatal disseminated intravascular coagulation associated with acute hemoglobinemia or hemoglobinuria after Rho(D) intravenous administration has been reported, particularly in older people in whom this treatment should be avoided.<sup>72</sup>

### Management of Life-Threatening Bleeding and Refractory Disease

In case of life-threatening bleeding situations such as visceral hemorrhage with decreased hemoglobin level or intracranial bleeding, the addition of platelet transfusions to IVIg and dexamethasone should be administered with a proven beneficial hemostatic effect.<sup>73,74</sup> Drugs interfering with hemostasis should be stopped immediately. In patients with active bleeding and with disease refractory to first-line therapy, vinca alkaloids such as vinblastin (weekly injection of 5 mg/m<sup>2</sup> with a maximum of 10 mg) rather than vincristine due to better digestive tolerance can provide rapid response but should not be used for long-term treatment because of neurological toxic effects, particularly in older people.<sup>8</sup> TPO-RAs can also be used as an off-label salvage therapy in our experience but with a risk of thrombosis.<sup>75</sup> Until these encouraging results are confirmed by other groups, the use of high-dose TPO-RAs should be reserved for patients with very severe bleeding resistant to total therapy (IVIg + high-dose corticosteroids + multiple and repeated platelet transfusions).

## Second-Line Treatments

Most adults with ITP have a chronic disease course. When disease is refractory to or relapses after first-line treatment, an increasing number of therapeutic options are available for second-line treatment. Each has an expected pattern of response, but to date, there are no accurate predictors to help select the ITP treatment. The treatment choice also relies on caveats specific to each molecule that are described below and in [Table 2](#). Co-morbidities and patient preference are also important in therapeutic decisions. According to the context, treatment should aim to avoid bleeding by maintaining a safe platelet count while limiting short- and long-term toxic effects. An overview of our current treatment strategy is proposed in [Figure 2](#).

### Thrombopoietin Receptor Agonists (TPO-RAs)

TPO-RAs are drugs increasing platelet production without immunosuppression that have a demonstrated good efficacy/safety profile, which explains why they have become the preferred second-line treatment in older patients with ITP by most groups and guidelines ([Figure 2](#)).<sup>4,5,46</sup> Two TPO-RAs, namely eltrombopag and romiplostim, are currently licensed for chronic ITP in Europe, and a third one, avatrombopag, is also available in some countries. Although no head-to-head comparison is available, response rates seem similar between both drugs<sup>76</sup> and close to 80% within 2 weeks.<sup>41,42</sup> The molecule choice mostly relies on the administration route preference: romiplostim requires a weekly subcutaneous


**Table 2** Factors influencing second-line treatment choices for ITP

	Avoid if Increased Thrombotic Risk	Avoid if Increased Infectious Risk	Avoid if Rapid Response Needed	Avoid if ITP is Secondary to Lymphoproliferative Disorder	Avoid if Severe Co-Morbidities	Avoid in Non-Chronic Patients
TPO-RA	x					
Rituximab		x				
Immunosuppressive drugs		x	x	x	x	
Dapsone			x			
Danazole	x		x			
Splenectomy	x	x		x	x	x
Fostamatinib				?		

**Abbreviations:** ITP, immune thrombocytopenia; TPO-RA, thrombopoietin receptor agonist.

injection, whereas eltrombopag is administered orally but at the cost of food or drug interactions and potential poor compliance in patients with cognitive disorders. In case of non-response or intolerance to a TPO-RA, a switch to the other drug can provide a better clinical result.<sup>77</sup> In case of failure to eltrombopag and romiplostim, response can be obtained with a switch to avatrombopag.<sup>78</sup> Because they increase platelet production, TPO-RAs have long been considered to very rarely lead to sustained remission off-treatment in ITP, but we now have evidence for sustained remission after their discontinuation in 10% to 30% of patients.<sup>79,80</sup> Therefore, after a response is obtained, the treatment should be tapered in order to use the lowest dose to maintain platelet counts.<sup>8</sup>

Several studies have suggested that TPO-RA efficacy is similar in older and younger patients.<sup>47,48,81</sup> Although the overall tolerance of TPO-RA is good, there are some concerns about long- and short-term adverse events. The risk of



**Figure 2** Proposed therapeutic strategy for ITP in older patients.

**Abbreviations:** ITP, immune thrombocytopenia; IVIg, intravenous immunoglobulin; RTX, rituximab; TPO-RA, thrombopoietin receptor agonist.

reticulin deposits in bone marrow has been raised in patients with long-term use of TPO-RAs but does not lead to bone-marrow failure and is reversible after drug discontinuation.<sup>82,83</sup>

Although not evidenced in pivotal trials, some studies have suggested a two- to three-fold increase in thrombosis risk with TPO-RA use as compared with untreated ITP patients.<sup>41–45</sup> This risk is particularly relevant in older patients because age is an independent risk factor for thrombosis,<sup>37,38</sup> in addition to the risks with other ITP treatments such as IgIV<sup>39</sup> or splenectomy<sup>40</sup> and the disease itself.<sup>35,36</sup> Despite these multiple confounding factors, a higher rate of thrombotic events has been described in older patients particularly those exposed to TPO-RAs.<sup>46–49</sup>

### Anti-CD20 Monoclonal Antibodies

The efficacy of rituximab is well documented in ITP, with initial response rates of about 60%, decreasing to 40% at 2 years and to almost 30% after 5 years.<sup>84–90</sup> Although low, these long-term responses suggest that rituximab can be a curative treatment for ITP, and re-treatment was found effective in most relapsing patients.<sup>88,89,91</sup>

Rituximab is usually administered as four weekly infusions of 375 mg/m<sup>2</sup>, but a more convenient dosage of 1000 mg on days 1 and 15 also used in other autoimmune diseases is considered equivalent.<sup>88</sup> Treatment with lower doses (100 mg weekly for 4 weeks) has comparable short-term response but with slower time to response and shorter response duration.<sup>92</sup> Several retrospective studies suggested a lower response rate in older patients<sup>85,86</sup> and in patients with longer disease duration.<sup>88</sup> However, a large prospective register including almost 250 patients with 5 years of follow-up did not show a difference in long-term response according to age.<sup>89</sup>

The tolerance profile of rituximab is good, but some points are important to consider when treating older patients. The infectious risk is only moderately increased,<sup>93</sup> although severe infections are predominantly observed in older patients, without excluding that concomitant prolonged corticosteroids and/or immunosuppressive drugs could have been the cause of these severe infectious events.<sup>88</sup> Hypogammaglobulinemia is uncommon after one course of rituximab, and the risk of severe hypogammaglobulinemia after repeated courses and/or the addition of immunosuppressive agents is unknown in ITP but seems rare when anti-CD20 antibodies are used to treat other autoimmune diseases such as rheumatoid arthritis.<sup>94</sup> Rituximab is associated with a risk of hepatitis B virus reactivation that is preventable.<sup>95</sup> Progressive multifocal leukoencephalopathy is exceptional.<sup>96</sup> Most importantly, an impaired vaccine response in the months following rituximab treatment has been well documented<sup>97</sup> and is particularly relevant for older patients requiring yearly influenza and/or SARS-CoV-2 vaccination. Anti-pneumococcal vaccination should also be performed before rituximab administration in older patients. During the COVID-19 pandemic, most groups avoided rituximab use in older patients because of the fear that rituximab would increase the risk of fatal infections, even if this risk has not been confirmed in the setting of ITP, for which rituximab is rarely associated with a strong immunosuppressive treatment.

Despite these limitations, rituximab remains a valid option for older patients and should be assessed on an individual basis. Anti-CD20 treatment is also a good option in the particular case of ITP associated with B-cell lymphoid malignancies or with associated systemic autoimmunity. Associating rituximab with other agents such as dexamethasone and/or ciclosporin has also been proposed, with higher response rates but at the cost of greater toxic effects, including infections and decreased gamma globulin levels.<sup>98–101</sup> Promising results were obtained by combining belimumab with rituximab in a pilot study.<sup>102</sup> The efficacy and tolerance of these combined strategies in older patients are unknown.

### Splenectomy

Splenectomy has been used to treat ITP for decades and provides a high rate of long-term response with an overall long-term remission rate of 60% to 70%.<sup>103</sup> It is now admitted that splenectomy should be deferred for at least 12 months to allow for spontaneous remission.<sup>8</sup> Given the availability of new ITP treatments, the rate of splenectomy is decreasing, particularly in older patients.<sup>32,104</sup> Response rates in older patients could be lower with increased surgical risk. Although laparoscopic surgery can reduce the risk of post-operative complications,<sup>103,105</sup> some studies reported a higher rate of immediate surgical complications such as severe bleeding<sup>105,106</sup> in older patients, which could be explained by the frequency of co-morbidities in this population.<sup>107</sup> Some studies have suggested a lower response rate<sup>105</sup> and higher relapse rate<sup>104,105,108</sup> in older patients. In a recent multicenter retrospective study after TPO-RA introduction in France, lack of sustained response after splenectomy was associated with older age on multivariable analysis (60–75 years: odds



ratio 0.39 [95% confidence interval 0.17–0.86],  $p = .02$ ; >75 years: 0.28 [0.10–0.75],  $p = .013$ ).<sup>104</sup> Splenectomy also exposes to increased risk of thrombosis, which should be prevented by prophylactic use of an anticoagulant immediately after the procedure, but given that this risk persists over time, it should be taken into account in the treatment decision.<sup>40,109</sup> Lastly, splenectomy also favors infection, particularly infection to encapsulated germs, and prior immunization against influenza and *Streptococcus pneumoniae* is recommended, as is patient education to take immediate antibiotics in case of fever.<sup>40,109</sup>

Nonetheless, our current opinion is that splenectomy can still be considered a third-line treatment in older patients, particularly for those with few comorbidities and low risk of thrombosis, and should not be contra-indicated solely on an age basis.<sup>29</sup> Some groups suggest that a platelet isotopic study could be useful to select the best patient candidates for splenectomy because in case of a predominant splenic sequestration pattern, most studies reported an excellent positive predictive value of response to splenectomy.<sup>110</sup> However, conflicting results were published in case of a hepatic or mixed sequestration pattern. A large UK study showed that most patients with mixed platelet destruction experienced clinical benefit after splenectomy despite being classified as non-responding.<sup>111</sup>

Partial splenic embolization was recently reported as a safe and effective treatment by Japanese colleagues.<sup>112</sup> If these good results are confirmed, this technique could be an interesting alternative in older patients with disease refractory to medical treatment and in whom splenectomy appears dangerous because of comorbidities.

### Immunosuppressive Agents

Immunosuppressive drugs have been used in ITP with various response rates. Mycophenolate mofetil<sup>113–116</sup> and azathioprine<sup>117,118</sup> are the most widely used, but data on ciclosporin,<sup>119–121</sup> rapamycin<sup>122–124</sup> or cyclophosphamide<sup>125,126</sup> efficacy have also been published. The treatment usually takes several months to obtain a response and requires long-term use with a high risk of relapse in case of stopping and with increased infectious risk that can be problematic in older patients. Despite these limitations, mycophenolate mofetil is widely prescribed as second-line therapy in the United Kingdom and has even been proposed as first-line therapy in a prospective controlled study.<sup>116</sup> This attitude is not widely adopted, and in France, our strategy is to reserve immunosuppressive drugs for patients with failure to respond to TPO-RAs and rituximab. We have also observed good responses in refractory patients when combining immunosuppressive drugs with TPO-RAs (unpublished observations).

### Dapsone

Dapsone is an effective treatment for ITP that can be used in older patients with moderate response rates (20–60%).<sup>127–130</sup> Previous splenectomy is associated with low efficacy, and it should not be used when a rapid response is needed given that the median time to response is close to 1 month. This treatment induces hemolysis, requiring a close monitoring of hemoglobin level especially in the first weeks of treatment, but in many patients, the decrease in hemoglobin level is transient (a few weeks) and limited (1 to 2 g/dL) and does not require discontinuation of the treatment. In contrast, the patient should be aware of the risk of dapsone-induced hypersensitivity syndrome that combines a generalized skin eruption with fever, with lymphadenopathy and hepatitis in the more severe forms. The syndrome always occurs in the first 4 weeks after the beginning of treatment. In our experience, skin eruption is observed in 7% of patients, and its severity is probably diminished when dapsone is associated with a short course of prednisone.<sup>131</sup>

### Danazol

Danazol has been used for years as second-line treatment for ITP and is considered efficient, with 40% to 70% response rates, even in older patients.<sup>27,132,133</sup> Its use is limited by its long time to response and its toxic effects (androgenic effects in women, risk of accelerated prostate cancer in men, liver cytotoxicity and increased thrombotic risk) but also by the concurrence of alternative second-line treatments for ITP.

### Fostamatinib and Emerging Agents

Fostamatinib, a Syk inhibitor, has been approved recently in the United States and the European Medicines Agency for treating chronic ITP. Both pivotal trials included older adults,<sup>134</sup> but to date there are no data regarding its specific use in

older people. Common adverse events reported in a pivotal study, such as diarrhea, hypertension, nausea, and increased transaminase levels, should be considered in the therapeutic decision. Other treatments such as anti-FcRn, rizalbrutinib (which inhibits Bruton tyrosine kinase) or daratumumab (an anti-CD38 monoclonal antibody) are currently under development for chronic ITP, but their respective efficacy/tolerance ratio in older patients remains to be determined.<sup>135</sup>

## Management of Particular Situations

### B-Cell Malignancies

The frequency of clonal B-cell disorders increases with age, and ITP is a well-known complication of B-cell lymphoid malignancies such as indolent lymphomas or chronic lymphoid leukemias. In this particular setting, lymphoma treatment should prevail over ITP treatment. With no indication for treating the B-cell disorder, rituximab<sup>136–138</sup> and TPO-RAs<sup>139</sup> can be used, and splenectomy should be avoided.

### Myelodysplastic Syndrome (MDS)

In older people, MDS can coexist with ITP or complicate the ITP course and is associated with increased bleeding risk.<sup>17</sup> The overall therapeutic strategy is unchanged, but the risk of acute myeloid leukemia transformation favored by TPO-RAs has been a concern. However, the risk seems limited and reassuring data have been produced recently.<sup>17,140–143</sup> In a randomized double-blind trial, the risk of leukemic progression in thrombocytopenic patients with low-risk MDS was similar between romiplostim and placebo,<sup>143</sup> although there are fewer data available for the use of TPO-RAs in high-risk MDS.

### Management of Thrombosis and TPO-RAs

Thrombotic events have been described in older patients receiving TPO-RAs.<sup>46–49</sup> In a real-life study of 384 patients >60 years and receiving TPO-RAs with a median follow-up of 2.7 years, Palandri et al observed 43 thromboses in 35 patients (including 22 arterial thromboses), corresponding to a cumulative incidence of 6.2% at 12 months.<sup>48</sup> The authors found a significant association between thrombosis history and thrombosis. Median platelet count was  $127 \times 10^9/L$  at the time of thrombosis event, which confirms that thrombosis can occur even with low/moderate platelet counts. Of note, thrombosis recurrence was observed mostly in patients who continued TPO-RAs, particularly those not receiving long-term anti-thrombotic treatment, which suggests that it could be used as a secondary prophylaxis if TPO-RA continuation is needed. These data strongly suggest that TPO-RAs should be used with caution in older patients with a history of thrombosis, especially if they are no longer receiving anticoagulants. In this situation, we suggest considering another therapeutic strategy.

### Refractory Patients

Despite the increasing therapeutic options to treat ITP, some patients do not respond to or lose response after the initiation of second-line treatment.<sup>144</sup> In these cases, a differential diagnosis should be ruled out, if necessary, by performing BME. When the disease is refractory to several second-line treatments such as both TPO-RAs and rituximab, the treatment choice should be personalized according to the context and the expected benefit/risk ratio. A better knowledge of the pathophysiology of ITP suggested that a combination of treatments could be synergic.<sup>135</sup> According to this hypothesis, we propose that the combination of immunosuppressive drugs and TPO-RA could produce a significant response in multirefractory ITP with previous failure to respond to TPO-RAs and immunosuppressive drugs alone.<sup>140</sup> Splenectomy can also be discussed for patients with few comorbidities and low risk of thrombosis.

## Conclusion

ITP is increasingly frequent in older patients, and the diagnostic work-up is mainly the same as in younger patients, with special attention to ruling out a differential diagnosis. The underlying mechanisms responsible for disease may differ and are probably a hot topic for research because it could help tailor therapy in the future. One of the limits of this review is that evidence-based medicine is lacking in older patients, although there is a growing number of retrospective studies focusing on this particular population. Age is associated with the accumulation of chronic conditions but is probably not a prognostic factor in ITP after controlling for other variables. Thus, the therapeutic decision should be patient-tailored according to the patient's preferences, co-morbidities and fitness rather than age itself. The efficacy of second-line

treatments has relegated splenectomy to patients with refractory disease only, and the good efficacy/safety profile of TPO-RAs has changed the therapeutic landscape of ITP in older people at the cost of increased thrombosis risk. The availability of new treatments will likely modify our therapeutic strategy in the next years.

## Acknowledgment

We thank Laura Smales for English editing.

## Funding

EC was supported by a poste d'accueil AP-HP.

## Disclosure

Mahévas received funds for research from GSK and fees from Amgen and Novartis for lectures. Bertrand Godeau served as an expert for Amgen, Novartis, Grifols and Sobi. Marc Michel received honoraria (advisory boards, speaker fees) from Novartis, Amgen, UCB, Argenx, Alexion and Sanofi, and personal fees from Sobi. Etienne Crickx received honoraria (advisory boards, speaker fees) from Novartis, UCB, and Sanofi. The authors report no other potential conflicts of interest in this work.

## References

1. Audia S, Mahévas M, Samson M, Godeau B, Bonnotte B. Pathogenesis of immune thrombocytopenia. *Autoimmun Rev*. 2017;16(6):620–632. doi:10.1016/j.autrev.2017.04.012
2. Moulis G, Palmaro A, Montastruc JL, Godeau B, Lapeyre-Mestre M, Sailler L. Epidemiology of incident immune thrombocytopenia: a nationwide population-based study in France. *Blood*. 2014;124(22):3308–3315. doi:10.1182/blood-2014-05-578336
3. Doobaree IU, Conway K, Miah H, et al. Incidence of adult primary immune thrombocytopenia in England—an update. *Eur J Haematol*. 2022;109(3):238–249. doi:10.1111/ejh.13803
4. Lucchini E, Fanin R, Cooper N, Zaja F. Management of immune thrombocytopenia in elderly patients. *Eur J Intern Med*. 2018;58:70–76. doi:10.1016/j.ejim.2018.09.005
5. Mahévas M, Michel M, Godeau B. How we manage immune thrombocytopenia in the elderly. *Br J Haematol*. 2016;173(6):844–856. doi:10.1111/bjh.14067
6. Terrell DR, Beebe LA, Neas BR, Vesely SK, Segal JB, George JN. Prevalence of primary immune thrombocytopenia in Oklahoma. *Am J Hematol*. 2012;87(9):848–852. doi:10.1002/ajh.23262
7. Sokal A, de Nadaï T, Maquet J, et al. Primary immune thrombocytopenia in very elderly patients: particularities in presentation and management: results from the prospective CARMEN-France registry. *Br J Haematol*. 2022;196(5):1262–1270. doi:10.1111/bjh.17935
8. Provan D, Arnold DM, Bussell JB, et al. Updated international consensus report on the investigation and management of primary immune thrombocytopenia. *Blood Adv*. 2019;3(22):3780–3817. doi:10.1182/bloodadvances.2019000812
9. McMillan R. Antiplatelet antibodies in chronic immune thrombocytopenia and their role in platelet destruction and defective platelet production. *Hematol Oncol Clin North Am*. 2009;23(6):1163–1175. doi:10.1016/j.hoc.2009.08.008
10. Vayne C, Guéry EA, Rollin J, Baglo T, Petermann R, Gruel Y. Pathophysiology and diagnosis of drug-induced immune thrombocytopenia. *J Clin Med*. 2020;9(7):E2212. doi:10.3390/jcm9072212
11. Crickx E, Moulis G, Ebbo M, et al. Safety of anti-SARS-CoV-2 vaccination for patients with immune thrombocytopenia. *Br J Haematol*. 2021. doi:10.1111/bjh.17813
12. Fuentes S, Chrétien B, Dolladille C, et al. An updated list of drugs suspected to be associated with immune thrombocytopenia based on the WHO pharmacovigilance database. *Blood*. 2022;140(8):922–927. doi:10.1182/blood.2022015936
13. Kantarjian H, Giles F, List A, et al. The incidence and impact of thrombocytopenia in myelodysplastic syndromes. *Cancer*. 2007;109(9):1705–1714. doi:10.1002/cncr.22602
14. Waisbren J, Dinner S, Altman J, et al. Disease characteristics and prognosis of myelodysplastic syndrome presenting with isolated thrombocytopenia. *Int J Hematol*. 2017;105(1):44–51. doi:10.1007/s12185-016-2081-4
15. Brynes RK, Orazi A, Theodore D, et al. Evaluation of bone marrow reticulin in patients with chronic immune thrombocytopenia treated with eltrombopag: data from the EXTEND study. *Am J Hematol*. 2015;90(7):598–601. doi:10.1002/ajh.24011
16. Rivière É, Viallard JF, Guy A, et al. Intrinsically impaired platelet production in some patients with persistent or chronic immune thrombocytopenia. *Br J Haematol*. 2015;170(3):408–415. doi:10.1111/bjh.13444
17. Jachiet V, Moulis G, Hadjadj J, et al. Clinical spectrum, outcome and management of immune thrombocytopenia associated with myelodysplastic syndromes and chronic myelomonocytic leukemia. *Haematologica*. 2021;106(5):1414–1422. doi:10.3324/haematol.2020.272559
18. Comont T, Germain J, Beyne-Rauzy O, Adoue D, Moulis G; Group and the C investigators. Positivity rate of systematic bone marrow smear in patients over 60 years old with newly diagnosed immune thrombocytopenia. *Blood Adv*. 2020;4(10):2136–2138. doi:10.1182/bloodadvances.2020001654
19. Taparia K, Wall E, Arnold DM, Sun HL. Frequency and utility of bone marrow examination in relapsed/refractory immune thrombocytopenia. *J Thromb Haemost*. 2022;20(9):2119–2126. doi:10.1111/jth.15802
20. Neunert C, Terrell DR, Arnold DM, et al. American Society of Hematology 2019 guidelines for immune thrombocytopenia. *Blood Adv*. 2019;3(23):3829–3866. doi:10.1182/bloodadvances.2019000966

21. Stasi R, Stipa E, Masi M, et al. Long-term observation of 208 adults with chronic idiopathic thrombocytopenic purpura. *Am J Med.* 1995;98(5):436–442. doi:10.1016/s0002-9343(99)80342-8
22. Frederiksen H, Schmidt K. The incidence of idiopathic thrombocytopenic purpura in adults increases with age. *Blood.* 1999;94(3):909–913. doi:10.1182/blood.V94.3.909.415k02\_909\_913
23. Vianelli N, Valdrè L, Fiacchini M, et al. Long-term follow-up of idiopathic thrombocytopenic purpura in 310 patients. *Haematologica.* 2001;86(5):504–509.
24. Cortelazzo S, Finazzi G, Buelli M, Molteni A, Viero P, Barbui T. High risk of severe bleeding in aged patients with chronic idiopathic thrombocytopenic purpura. *Blood.* 1991;77(1):31–33. doi:10.1182/blood.V77.1.31.31
25. Bourgeois E, Caulier MT, Delarozee C, Brouillard M, Bauters F, Fenaux P. Long-term follow-up of chronic autoimmune thrombocytopenic purpura refractory to splenectomy: a prospective analysis. *Br J Haematol.* 2003;120(6):1079–1088. doi:10.1046/j.1365-2141.2003.04211.x
26. Bizzoni L, Mazzucconi MG, Gentile M, et al. Idiopathic thrombocytopenic purpura (ITP) in the elderly: clinical course in 178 patients. *Eur J Haematol.* 2006;76(3):210–216. doi:10.1111/j.1600-0609.2005.00602.x
27. Daou S, Federici L, Zimmer J, Maloïsel F, Serraj K, André E. Idiopathic thrombocytopenic purpura in elderly patients: a study of 47 cases from a single reference center. *Eur J Intern Med.* 2008;19(6):447–451. doi:10.1016/j.ejim.2007.07.006
28. Cohen YC, Djulbegovic B, Shamai-Lubovitz O, Mozes B. The bleeding risk and natural history of idiopathic thrombocytopenic purpura in patients with persistent low platelet counts. *Arch Intern Med.* 2000;160(11):1630–1638. doi:10.1001/archinte.160.11.1630
29. Michel M, Rauzy OB, Thoraval FR, et al. Characteristics and outcome of immune thrombocytopenia in elderly: results from a single center case-controlled study. *Am J Hematol.* 2011;86(12):980–984. doi:10.1002/ajh.22170
30. Mithoowani S, Cervi A, Shah N, et al. Management of major bleeds in patients with immune thrombocytopenia. *J Thromb Haemost.* 2020;18(7):1783–1790. doi:10.1111/jth.14809
31. Zhou H, Fu R, Wang H, et al. Immune thrombocytopenia in the elderly: clinical course in 525 patients from a single center in China. *Ann Hematol.* 2013;92(1):79–87. doi:10.1007/s00277-012-1567-2
32. Palandri F, Catani L, Auteri G, et al. Understanding how older age drives decision-making and outcome in immune thrombocytopenia. A single centre study on 465 adult patients. *Br J Haematol.* 2019;184(3):424–430. doi:10.1111/bjh.15668
33. Piel-Julian ML, Mahévas M, Germain J, et al. Risk factors for bleeding, including platelet count threshold, in newly diagnosed immune thrombocytopenia adults. *J Thromb Haemost.* 2018;16(9):1830–1842. doi:10.1111/jth.14227
34. Swan D, Newland A, Rodeghiero F, Thachil J. Thrombosis in immune thrombocytopenia - current status and future perspectives. *Br J Haematol.* 2021;194(5):822–834. doi:10.1111/bjh.17390
35. Sarpatwari A, Bennett D, Logie JW, et al. Thromboembolic events among adult patients with primary immune thrombocytopenia in the United Kingdom general practice research database. *Haematologica.* 2010;95(7):1167–1175. doi:10.3324/haematol.2009.018390
36. Doobaree IU, Nandigam R, Bennett D, Newland A, Provan D. Thromboembolism in adults with primary immune thrombocytopenia: a systematic literature review and meta-analysis. *Eur J Haematol.* 2016;97(4):321–330. doi:10.1111/ejh.12777
37. Ekstrand C, Linder M, Baricault B, et al. Impact of risk factors on the occurrence of arterial thrombosis and venous thromboembolism in adults with primary immune thrombocytopenia – results from two nationwide cohorts. *Thromb Res.* 2019;178:124–131. doi:10.1016/j.thromres.2019.04.016
38. Naess IA, Christiansen SC, Romundstad P, Cannegieter SC, Rosendaal FR, Hammerstrøm J. Incidence and mortality of venous thrombosis: a population-based study. *J Thromb Haemost.* 2007;5(4):692–699. doi:10.1111/j.1538-7836.2007.02450.x
39. Paran D, Herishanu Y, Elkayam O, Shopin L, Ben-Ami R. Venous and arterial thrombosis following administration of intravenous immunoglobulins. *Blood Coagul Fibrinolysis.* 2005;16(5):313–318. doi:10.1097/01.mbc.0000172694.85233.a8
40. Thai LH, Mahévas M, Roudot-Thoraval F, et al. Long-term complications of splenectomy in adult immune thrombocytopenia. *Medicine.* 2016;95(48):e5098. doi:10.1097/MD.0000000000005098
41. Wong RSM, Saleh MN, Khelif A, et al. Safety and efficacy of long-term treatment of chronic/persistent ITP with eltrombopag: final results of the EXTEND study. *Blood.* 2017;130(23):2527–2536. doi:10.1182/blood-2017-04-748707
42. Kuter DJ, Bussel JB, Newland A, et al. Long-term treatment with romiplostim in patients with chronic immune thrombocytopenia: safety and efficacy. *Br J Haematol.* 2013;161(3):411–423. doi:10.1111/bjh.12260
43. Catalá-López F, Corrales I, de la Fuente-Honrubia C, et al. Risk of thromboembolism with thrombopoietin receptor agonists in adult patients with thrombocytopenia: systematic review and meta-analysis of randomized controlled trials. *Med Clin.* 2015;145(12):511–519. doi:10.1016/j.medcli.2015.03.014
44. Ghanima W, Cooper N, Rodeghiero F, Godeau B, Bussel JB. Thrombopoietin receptor agonists: ten years later. *Haematologica.* 2019;104(6):1112–1123. doi:10.3324/haematol.2018.212845
45. Bussel JB, Kuter DJ, Pullarkat V, Lyons RM, Guo M, Nichol JL. Safety and efficacy of long-term treatment with romiplostim in thrombocytopenic patients with chronic ITP. *Blood.* 2009;113(10):2161–2171. doi:10.1182/blood-2008-04-150078
46. Lozano ML, Mingot-Castellano ME, Perera MM, et al. A decade of changes in management of immune thrombocytopenia, with special focus on elderly patients. *Blood Cells Mol Dis.* 2021;86:102505. doi:10.1016/j.bcmd.2020.102505
47. Michel M, Wasser J, Godeau B, et al. Efficacy and safety of the thrombopoietin receptor agonist romiplostim in patients aged ≥ 65 years with immune thrombocytopenia. *Ann Hematol.* 2015;94(12):1973–1980. doi:10.1007/s00277-015-2485-x
48. Palandri F, Rossi E, Bartoletti D, et al. Real-world use of thrombopoietin receptor agonists in older patients with primary immune thrombocytopenia. *Blood.* 2021;138(7):571–583. doi:10.1182/blood.2021010735
49. Ruggeri M, Tosetto A, Palandri F, et al. Thrombotic risk in patients with primary immune thrombocytopenia is only mildly increased and explained by personal and treatment-related risk factors. *J Thromb Haemost.* 2014;12(8):1266–1273. doi:10.1111/jth.12636
50. Nørgaard M, Jensen AØ, Engebjerg MC, et al. Long-term clinical outcomes of patients with primary chronic immune thrombocytopenia: a Danish population-based cohort study. *Blood.* 2011;117(13):3514–3520. doi:10.1182/blood-2010-10-312819
51. Portielje JE, Westendorp RG, Kluin-Nelemans HC, Brand A. Morbidity and mortality in adults with idiopathic thrombocytopenic purpura. *Blood.* 2001;97(9):2549–2554. doi:10.1182/blood.v97.9.2549
52. Schoonen WM, Kucera G, Coalson J, et al. Epidemiology of immune thrombocytopenic purpura in the general practice research database. *Br J Haematol.* 2009;145(2):235–244. doi:10.1111/j.1365-2141.2009.07615.x

53. Cooper N, Kruse A, Kruse C, et al. Immune thrombocytopenia (ITP) World Impact Survey (iWISh): patient and physician perceptions of diagnosis, signs and symptoms, and treatment. *Am J Hematol.* 2021;96(2):188–198. doi:10.1002/ajh.26045
54. Cooper N, Kruse A, Kruse C, et al. Immune thrombocytopenia (ITP) World Impact Survey (I-WISh): impact of ITP on health-related quality of life. *Am J Hematol.* 2021;96(2):199–207. doi:10.1002/ajh.26036
55. Giordano P, Lassandro G, Valente M, Molinari AC, Ieranò P, Coppola A. Current management of the hemophilic child: a demanding interlocutor. Quality of life and adequate cost-efficacy analysis. *Pediatr Hematol Oncol.* 2014;31(8):687–702. doi:10.3109/08880018.2014.930768
56. Provan D, Stasi R, Newland AC, et al. International consensus report on the investigation and management of primary immune thrombocytopenia. *Blood.* 2010;115(2):168–186. doi:10.1182/blood-2009-06-225565
57. Matzdorff A, Beer JH. Immune thrombocytopenia patients requiring anticoagulation—maneuvering between scylla and charybdis. *Semin Hematol.* 2013;50(Suppl 1):S83–S88. doi:10.1053/j.seminhematol.2013.03.020
58. Godeau B, Bierling P. High-dose dexamethasone as initial treatment for immune thrombocytopenic purpura. *N Engl J Med.* 2003;349(23):2267–2268;author reply 2267–2268. doi:10.1056/NEJM200312043492318
59. Pirunsarn A, Kijrattanakul P, Chamnanchanunt S, Polprasert C, Rojnuckarin P. A randomized multicenter trial comparing low-dose prednisolone versus observation for prevention of recurrences in adult immune thrombocytopenia. *Clin Appl Thromb Hemost.* 2018;24(6):867–873. doi:10.1177/1076029618764843
60. Wei Y, Ji XB, Wang YW, et al. High-dose dexamethasone vs prednisone for treatment of adult immune thrombocytopenia: a prospective multicenter randomized trial. *Blood.* 2016;127(3):296–302;quiz 370. doi:10.1182/blood-2015-07-659656
61. Mithoowani S, Gregory-Miller K, Goy J, et al. High-dose dexamethasone compared with prednisone for previously untreated primary immune thrombocytopenia: a systematic review and meta-analysis. *Lancet Haematol.* 2016;3(10):e489–e496. doi:10.1016/S2352-3026(16)30109-0
62. Palandri F, Santoro C, Carpenedo M, et al. Management of elderly patients with immune thrombocytopenia: real-world evidence from 451 patients older than 60 years. *Thromb Res.* 2020;185:88–95. doi:10.1016/j.thromres.2019.11.026
63. Moulis G, Palmaro A, Sailler L, Lapeyre-Mestre M, Latus J. Corticosteroid risk function of severe infection in primary immune thrombocytopenia adults. A nationwide nested case-control study. *PLoS One.* 2015;10(11):e0142217. doi:10.1371/journal.pone.0142217
64. Bellucci S, Charpak Y, Chastang C, Tobelem G. Low doses v conventional doses of corticoids in immune thrombocytopenic purpura (ITP): results of a randomized clinical trial in 160 children, 223 adults. *Blood.* 1988;71(4):1165–1169. doi:10.1182/blood.V71.4.1165.1165
65. Godeau B, Chevret S, Varet B, et al. Intravenous immunoglobulin or high-dose methylprednisolone, with or without oral prednisone, for adults with untreated severe autoimmune thrombocytopenic purpura: a randomised, multicentre trial. *Lancet.* 2002;359(9300):23–29. doi:10.1016/S0140-6736(02)07275-6
66. Khellaf M, Michel M, Schaeffer A, Bierling P, Godeau B. Assessment of a therapeutic strategy for adults with severe autoimmune thrombocytopenic purpura based on a bleeding score rather than platelet count. *Haematologica.* 2005;90(6):829–832.
67. Godeau B, Caulier MT, Decuyper L, et al. Intravenous immunoglobulin for adults with autoimmune thrombocytopenic purpura: results of a randomized trial comparing 0.5 and 1 g/kg b.w. *Br J Haematol.* 1999;107(4). doi:10.1046/j.1365-2141.1999.01766.x
68. Caress JB, Kennedy BL, Eickman KD. Safety of intravenous immunoglobulin treatment. *Expert Opin Drug Saf.* 2010;9(6):971–979. doi:10.1517/14740338.2010.484419
69. Marie I, Maurey G, Hervé F, Hellot MF, Levesque H. Intravenous immunoglobulin-associated arterial and venous thrombosis; report of a series and review of the literature. *Br J Dermatol.* 2006;155(4):714–721. doi:10.1111/j.1365-2133.2006.07390.x
70. Stiehm ER. Adverse effects of human immunoglobulin therapy. *Transfus Med Rev.* 2013;27(3):171–178. doi:10.1016/j.tmr.2013.05.004
71. Gaines AR. Acute onset hemoglobinemia and/or hemoglobinuria and sequelae following Rh(o)(D) immune globulin intravenous administration in immune thrombocytopenic purpura patients. *Blood.* 2000;95(8):2523–2529. doi:10.1182/blood.V95.8.2523
72. Ar G. Disseminated intravascular coagulation associated with acute hemoglobinemia or hemoglobinuria following Rh(0)(D) immune globulin intravenous administration for immune thrombocytopenic purpura. *Blood.* 2005;106:5. doi:10.1182/blood-2004-11-4303
73. Salama A, Kiesewetter H, Kalus U, Movassaghi K, Meyer O. Massive platelet transfusion is a rapidly effective emergency treatment in patients with refractory autoimmune thrombocytopenia. *Thromb Haemost.* 2008;100(5):762–765. doi:10.1160/TH08-06-0418
74. Spahr JE, Rodgers GM. Treatment of immune-mediated thrombocytopenia purpura with concurrent intravenous immunoglobulin and platelet transfusion: a retrospective review of 40 patients. *Am J Hematol.* 2008;83(2):122–125. doi:10.1002/ajh.21060
75. Roumier M, Le Burel S, Audia S, et al. High dose romiplostim as a rescue therapy for adults with severe bleeding and refractory immune thrombocytopenia. *Am J Hematol.* 2021;96(2):E43–E46. doi:10.1002/ajh.26040
76. Zhang J, Liang Y, Ai Y, et al. Eltrombopag versus romiplostim in treatment of children with persistent or chronic immune thrombocytopenia: a systematic review incorporating an indirect-comparison meta-analysis. *Sci Rep.* 2018;8(1):576. doi:10.1038/s41598-017-19099-8
77. Kuter DJ, Macahilig C, Grotzinger KM, et al. Treatment patterns and clinical outcomes in patients with chronic immune thrombocytopenia (ITP) switched to eltrombopag or romiplostim. *Int J Hematol.* 2015;101(3):255–263. doi:10.1007/s12185-014-1731-7
78. Al-Samkari H, Jiang D, Gernsheimer T, et al. Adults with immune thrombocytopenia who switched to avatrombopag following prior treatment with eltrombopag or romiplostim: a multicentre US study. *Br J Haematol.* 2022;197(3):359–366. doi:10.1111/bjh.18081
79. Bussel JB, Wang X, Lopez A, Eisen M. Case study of remission in adults with immune thrombocytopenia following cessation of treatment with the thrombopoietin mimetic romiplostim. *Hematology.* 2016;21(4):257–262. doi:10.1179/1607845415Y.0000000041
80. Mahévas M, Fain O, Ebbo M, et al. The temporary use of thrombopoietin-receptor agonists may induce a prolonged remission in adult chronic immune thrombocytopenia. Results of a French observational study. *Br J Haematol.* 2014;165(6):865–869. doi:10.1111/bjh.12888
81. González-López TJ, Sánchez-González B, Jarque I, et al. Use of eltrombopag for patients 65 years old or older with immune thrombocytopenia. *Eur J Haematol.* 2020;104(3):259–270. doi:10.1111/ejh.13370
82. Ghanima W, Geyer JT, Lee CS, et al. Bone marrow fibrosis in 66 patients with immune thrombocytopenia treated with thrombopoietin-receptor agonists: a single-center, long-term follow-up. *Haematologica.* 2014;99(5):937–944. doi:10.3324/haematol.2013.098921
83. Rizvi H, Butler T, Calaminici M, et al. United Kingdom immune thrombocytopenia registry: retrospective evaluation of bone marrow fibrosis in adult patients with primary immune thrombocytopenia and correlation with clinical findings. *Br J Haematol.* 2015;169(4):590–594. doi:10.1111/bjh.13330
84. Godeau B, Porcher R, Fain O, et al. Rituximab efficacy and safety in adult splenectomy candidates with chronic immune thrombocytopenic purpura: results of a prospective multicenter Phase 2 study. *Blood.* 2008;112(4):999–1004. doi:10.1182/blood-2008-01-131029

85. Auger S, Duny Y, Rossi JF, Quittet P. Rituximab before splenectomy in adults with primary idiopathic thrombocytopenic purpura: a meta-analysis. *Br J Haematol.* 2012;158(3):386–398. doi:10.1111/j.1365-2141.2012.09169.x
86. Marangon M, Vianelli N, Palandri F, et al. Rituximab in immune thrombocytopenia: gender, age, and response as predictors of long-term response. *Eur J Haematol.* 2017;98(4):371–377. doi:10.1111/ejh.12839
87. Arnold DM, Dentali F, Crowther MA, et al. Systematic review: efficacy and safety of rituximab for adults with idiopathic thrombocytopenic purpura. *Ann Intern Med.* 2007;146(1):25–33. doi:10.7326/0003-4819-146-1-200701020-00006
88. Khellaf M, Charles-Nelson A, Fain O, et al. Safety and efficacy of rituximab in adult immune thrombocytopenia: results from a prospective registry including 248 patients. *Blood.* 2014;124(22):3228–3236. doi:10.1182/blood-2014-06-582346
89. Deshayes S, Khellaf M, Zarour A, et al. Long-term safety and efficacy of rituximab in 248 adults with immune thrombocytopenia: results at 5 years from the French prospective registry ITP-ritux. *Am J Hematol.* 2019;94(12):1314–1324. doi:10.1002/ajh.25632
90. Patel VL, Mahévas M, Lee SY, et al. Outcomes 5 years after response to rituximab therapy in children and adults with immune thrombocytopenia. *Blood.* 2012;119(25):5989–5995. doi:10.1182/blood-2011-11-393975
91. Hasan A, Michel M, Patel V, et al. Repeated courses of rituximab in chronic ITP: three different regimens. *Am J Hematol.* 2009;84(10):661–665. doi:10.1002/ajh.21512
92. Zaja F, Vianelli N, Volpetti S, et al. Low-dose rituximab in adult patients with primary immune thrombocytopenia. *Eur J Haematol.* 2010;85(4):329–334. doi:10.1111/j.1600-0609.2010.01486.x
93. Chugh S, Darvish-Kazem S, Lim W, et al. Rituximab plus standard of care for treatment of primary immune thrombocytopenia: a systematic review and meta-analysis. *Lancet Haematol.* 2015;2(2):e75–e81. doi:10.1016/S2352-3026(15)00003-4
94. van Vollenhoven RF, Emery P, Bingham CO, et al. Long-term safety of rituximab in rheumatoid arthritis: 9.5-year follow-up of the global clinical trial programme with a focus on adverse events of interest in RA patients. *Ann Rheum Dis.* 2013;72(9):1496–1502. doi:10.1136/annrheumdis-2012-201956
95. Cho CH, Hwang WL, Cheng SB, Lee TY, Teng CL. Hepatitis B reactivation induced by Rituximab maintenance therapy for lymphoma. *Ann Hematol.* 2011;90(1):111–112. doi:10.1007/s00277-010-0962-9
96. Carson KR, Evens AM, Richey EA, et al. Progressive multifocal leukoencephalopathy after rituximab therapy in HIV-negative patients: a report of 57 cases from the research on adverse drug events and reports project. *Blood.* 2009;113(20):4834–4840. doi:10.1182/blood-2008-10-186999
97. Nazi I, Kelton JG, Larché M, et al. The effect of rituximab on vaccine responses in patients with immune thrombocytopenia. *Blood.* 2013;122(11):1946–1953. doi:10.1182/blood-2013-04-494096
98. Bussel JB, Lee CS, Seery C, et al. Rituximab and three dexamethasone cycles provide responses similar to splenectomy in women and those with immune thrombocytopenia of less than two years duration. *Haematologica.* 2014;99(7):1264–1271. doi:10.3324/haematol.2013.103291
99. Choi PYI, Roncolato F, Badoux X, Ramanathan S, Ho SJ, Chong BH. A novel triple therapy for ITP using high-dose dexamethasone, low-dose rituximab, and cyclosporine (TT4). *Blood.* 2015;126(4):500–503. doi:10.1182/blood-2015-03-631937
100. Zaja F, Baccarani M, Mazza P, et al. Dexamethasone plus rituximab yields higher sustained response rates than dexamethasone monotherapy in adults with primary immune thrombocytopenia. *Blood.* 2010;115(14):2755–2762. doi:10.1182/blood-2009-07-229815
101. Gudbrandsdottir S, Birgens HS, Frederiksen H, et al. Rituximab and dexamethasone vs dexamethasone monotherapy in newly diagnosed patients with primary immune thrombocytopenia. *Blood.* 2013;121(11):1976–1981. doi:10.1182/blood-2012-09-455691
102. Mahévas M, Azzaoui I, Crickx E, et al. Efficacy, safety and immunological profile of combining rituximab with belimumab for adults with persistent or chronic immune thrombocytopenia: results from a prospective phase 2b trial. *Haematologica.* 2021;106(9):2449–2457. doi:10.3324/haematol.2020.259481
103. Kojouri K, Vesely SK, Terrell DR, George JN. Splenectomy for adult patients with idiopathic thrombocytopenic purpura: a systematic review to assess long-term platelet count responses, prediction of response, and surgical complications. *Blood.* 2004;104(9):2623–2634. doi:10.1182/blood-2004-03-1168
104. Mageau A, Terriou L, Ebbo M, et al. Splenectomy for primary immune thrombocytopenia revisited in the era of thrombopoietin receptor agonists: new insights for an old treatment. *Am J Hematol.* 2022;97(1):10–17. doi:10.1002/ajh.26378
105. Gonzalez-Porras JR, Escalante F, Pardal E, et al. Safety and efficacy of splenectomy in over 65-yr-old patients with immune thrombocytopenia. *Eur J Haematol.* 2013;91(3):236–241. doi:10.1111/ejh.12146
106. Andrés E, Zimmer J, Noel E, Kaltenbach G, Koumariou A, Maloisel F. Idiopathic thrombocytopenic purpura: a retrospective analysis in 139 patients of the influence of age on the response to corticosteroids, splenectomy and danazol. *Drugs Aging.* 2003;20(11):841–846. doi:10.2165/00002512-200320110-00005
107. Kavic SM, Segan RD, Park AE. Laparoscopic splenectomy in the elderly: a morbid procedure? *Surg Endosc.* 2005;19(12):1561–1564. doi:10.1007/s00464-005-0125-6
108. Park YH, Yi HG, Kim CS, et al. Clinical outcome and predictive factors in the response to splenectomy in elderly patients with primary immune thrombocytopenia: a multicenter retrospective study. *Acta Haematol.* 2016;135(3):162–171. doi:10.1159/000442703
109. Vianelli N, Palandri F, Polverelli N, et al. Splenectomy as a curative treatment for immune thrombocytopenia: a retrospective analysis of 233 patients with a minimum follow up of 10 years. *Haematologica.* 2013;98(6):875–880. doi:10.3324/haematol.2012.075648
110. Cuker A, Cines DB. Evidence-based mini-review: is indium-labeled autologous platelet scanning predictive of response to splenectomy in patients with chronic immune thrombocytopenia? *Hematol Am Soc Hematol Educ Program.* 2010;2010(1):385–386. doi:10.1182/asheducation-2010.1.385
111. Sarpatwari A, Provan D, Erqou S, Sobnack R, David Tai FW, Newland AC. Autologous 111 in-labelled platelet sequestration studies in patients with primary immune thrombocytopenia (ITP) prior to splenectomy: a report from the United Kingdom ITP registry. *Br J Haematol.* 2010;151(5):477–487. doi:10.1111/j.1365-2141.2010.08377.x
112. Togasaki E, Shimizu N, Nagao Y, et al. Long-term efficacy of partial splenic embolization for the treatment of steroid-resistant chronic immune thrombocytopenia. *Ann Hematol.* 2018;97(4):655–662. doi:10.1007/s00277-018-3232-x
113. Hou M, Peng J, Shi Y, et al. Mycophenolate mofetil (MMF) for the treatment of steroid-resistant idiopathic thrombocytopenic purpura. *Eur J Haematol.* 2003;70(6):353–357. doi:10.1034/j.1600-0609.2003.00076.x
114. Taylor A, Neave L, Solanki S, et al. Mycophenolate mofetil therapy for severe immune thrombocytopenia. *Br J Haematol.* 2015;171(4):625–630. doi:10.1111/bjh.13622

115. Provan D, Moss AJ, Newland AC, Bussel JB. Efficacy of mycophenolate mofetil as single-agent therapy for refractory immune thrombocytopenic purpura. *Am J Hematol.* 2006;81(1):19–25. doi:10.1002/ajh.20515
116. Bradbury CA, Pell J, Hill Q, et al. Mycophenolate mofetil for first-line treatment of immune thrombocytopenia. *N Engl J Med.* 2021;385(10):885–895. doi:10.1056/NEJMoa2100596
117. Vesely SK, Perdue JJ, Rizvi MA, Terrell DR, George JN. Management of adult patients with persistent idiopathic thrombocytopenic purpura following splenectomy: a systematic review. *Ann Intern Med.* 2004;140(2):112–120. doi:10.7326/0003-4819-140-3-200402030-00012
118. Mishra K, Pramanik S, Sandal R, et al. Safety and efficacy of azathioprine in immune thrombocytopenia. *Am J Blood Res.* 2021;11(3):217–226.
119. Choudhary DR, Naithani R, Mahapatra M, Kumar R, Mishra P, Saxena R. Efficacy of cyclosporine as a single agent therapy in chronic idiopathic thrombocytopenic purpura. *Haematologica.* 2008;93(10):e61–e62;discussion e63. doi:10.3324/haematol.13481
120. Kappers-Klunne MC, Van't Veer MB. Cyclosporin A for the treatment of patients with chronic idiopathic thrombocytopenic purpura refractory to corticosteroids or splenectomy. *Br J Haematol.* 2001;114(1):121–125. doi:10.1046/j.1365-2141.2001.02893.x
121. Emilia G, Morselli M, Luppi M, et al. Long-term salvage therapy with cyclosporin A in refractory idiopathic thrombocytopenic purpura. *Blood.* 2002;99(4):1482–1485. doi:10.1182/blood.V99.4.1482
122. Li H, Ji J, Du Y, et al. Sirolimus is effective for primary relapsed/refractory autoimmune cytopenia: a multicenter study. *Exp Hematol.* 2020;89:87–95. doi:10.1016/j.exphem.2020.08.001
123. Feng Y, Xiao Y, Yan H, et al. Sirolimus as rescue therapy for refractory/relapsed immune thrombocytopenia: results of a single-center, prospective, single-arm study. *Front Med.* 2020;7:110. doi:10.3389/fmed.2020.00110
124. Li J, Wang Z, Dai L, et al. Effects of rapamycin combined with low dose prednisone in patients with chronic immune thrombocytopenia. *Clin Dev Immunol.* 2013;2013:548085. doi:10.1155/2013/548085
125. Verlin M, Laros RK, Penner JA. Treatment of refractory thrombocytopenic purpura with cyclophosphamide. *Am J Hematol.* 1976;1(1):97–104. doi:10.1002/ajh.2830010111
126. Pizzuto J, Ambriz R. Therapeutic experience on 934 adults with idiopathic thrombocytopenic purpura: multicentric trial of the cooperative Latin American group on hemostasis and thrombosis. *Blood.* 1984;64(6):1179–1183. doi:10.1182/blood.V64.6.1179.1179
127. Godeau B, Oksenhendler E, Bierling P. Dapsone for autoimmune thrombocytopenic purpura. *Am J Hematol.* 1993;44(1):70–72. doi:10.1002/ajh.2830440117
128. Godeau B, Durand JM, Roudot-Thoraval F, et al. Dapsone for chronic autoimmune thrombocytopenic purpura: a report of 66 cases. *Br J Haematol.* 1997;97(2):336–339. doi:10.1046/j.1365-2141.1997.412687.x
129. Zaja F, Marin L, Chiozzotto M, Puglisi S, Volpetti S, Fanin R. Dapsone salvage therapy for adult patients with immune thrombocytopenia relapsed or refractory to steroid and rituximab. *Am J Hematol.* 2012;87(3):321–323. doi:10.1002/ajh.22266
130. Durand JM, Lefèvre P, Hovette P, Mongin M, Soubeyrand J. Dapsone for idiopathic autoimmune thrombocytopenic purpura in elderly patients. *Br J Haematol.* 1991;78(3):459–460. doi:10.1111/j.1365-2141.1991.tb04467.x
131. Sauvetre G, MahÉvas M, Limal N, et al. Cutaneous rash and dapsone-induced hypersensitivity syndrome a common manifestation in adult immune thrombocytopenia. Presentation and outcome in 16 cases. *Am J Hematol.* 2015;90(10):E201–E202. doi:10.1002/ajh.24068
132. Maloisel F, Andrès E, Zimmer J, et al. Danazol therapy in patients with chronic idiopathic thrombocytopenic purpura: long-term results. *Am J Med.* 2004;116(9):590–594. doi:10.1016/j.amjmed.2003.12.024
133. Liu W, Gu X, Fu R, et al. The effect of danazol in primary immune thrombocytopenia: an analysis of a large cohort from a single center in China. *Clin Appl Thromb Hemost.* 2016;22(8):727–733. doi:10.1177/1076029615622002
134. Bussel J, Arnold DM, Grossbard E, et al. Fostamatinib for the treatment of adult persistent and chronic immune thrombocytopenia: results of two Phase 3, randomized, placebo-controlled trials. *Am J Hematol.* 2018;93(7):921–930. doi:10.1002/ajh.25125
135. Audia S, MahÉvas M, Nivet M, Ouandji S, Ciudad M, Bonnotte B. Immune thrombocytopenia: recent advances in pathogenesis and treatments. *Hemasphere.* 2021;5(6):e574. doi:10.1097/HS9.0000000000000574
136. Zent CS, Ding W, Reinalda MS, et al. Autoimmune cytopenia in chronic lymphocytic leukemia/small lymphocytic lymphoma: changes in clinical presentation and prognosis. *Leuk Lymphoma.* 2009;50(8):1261–1268. doi:10.1080/10428190903026492
137. Rogers KA, Woyach JA. Secondary autoimmune cytopenias in chronic lymphocytic leukemia. *Seminars in Oncology.* 2016;43(2):300–310. doi:10.1053/j.seminoncol.2016.02.011
138. Hauswirth AW, Skrabs C, Schützing C, et al. Autoimmune thrombocytopenia in non-Hodgkin's lymphomas. *Haematologica.* 2008;93(3):447–450. doi:10.3324/haematol.11934
139. Ferretti A, Baldacci E, Miulli E, et al. Thrombopoietin receptor agonists to control immune thrombocytopenia in patients with active lymphoma. *Br J Haematol.* 2019;186(6):e217–e219. doi:10.1111/bjh.16114
140. Kantarjian HM, Giles FJ, Greenberg PL, et al. Phase 2 study of romiplostim in patients with low- or intermediate-risk myelodysplastic syndrome receiving azacitidine therapy. *Blood.* 2010;116(17):3163–3170. doi:10.1182/blood-2010-03-274753
141. Giagounidis A, Mufti GJ, Fenaux P, et al. Results of a randomized, double-blind study of romiplostim versus placebo in patients with low/intermediate-risk myelodysplastic syndrome and thrombocytopenia. *Cancer.* 2014;120(12):1838–1846. doi:10.1002/cncr.28663
142. Fenaux P, Muus P, Kantarjian H, et al. Romiplostim monotherapy in thrombocytopenic patients with myelodysplastic syndromes: long-term safety and efficacy. *Br J Haematol.* 2017;178(6):906–913. doi:10.1111/bjh.14792
143. Kantarjian HM, Fenaux P, Sekeres MA, et al. Long-term follow-up for up to 5 years on the risk of leukaemic progression in thrombocytopenic patients with lower-risk myelodysplastic syndromes treated with romiplostim or placebo in a randomised double-blind trial. *Lancet Haematol.* 2018;5(3):e117–e126. doi:10.1016/S2352-3026(18)30016-4
144. MahÉvas M, Gerfaud-Valentin M, Moulis G, et al. Characteristics, outcome, and response to therapy of multirefractory chronic immune thrombocytopenia. *Blood.* 2016;128(12):1625–1630. doi:10.1182/blood-2016-03-704734

Clinical Interventions in Aging

Dovepress

## Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>