

## REVIEW ARTICLE

# The community nurse in Australia. Who are they? A rapid systematic review

Nicole Blay RN, PhD, BHA, Adjunct Fellow<sup>1,2,3</sup>  |

Mariana S. Sousa MD, PhD, Postdoctoral Research Fellow<sup>1,3,4</sup>  |

Mick Rowles RN MHSM, Director of Nursing Primary and Community Health<sup>3,5</sup> |

Pauline Murray-Parahi RN, RM, MEd, PhD<sup>1,3,4</sup> 

<sup>1</sup>Centre for Applied Nursing Research, School of Nursing and Midwifery, Western Sydney University, Penrith, New South Wales, Australia

<sup>2</sup>Centre for Applied Nursing Research (CANR), Ingham Institute for Applied Medical Research, Liverpool, New South Wales, Australia

<sup>3</sup>Nursing and Midwifery, South Western Sydney Local Health District, Liverpool, New South Wales, Australia

<sup>4</sup>Faculty of Health, University of Technology Sydney, Ultimo, New South Wales, Australia

<sup>5</sup>Primary and Community Health Division, South Western Sydney Local Health District, Liverpool, New South Wales, Australia

## Correspondence

Nicole Blay, Centre for Applied Nursing Research (CANR), School of Nursing and Midwifery, Western Sydney University, Locked Bag 1797, Penrith, New South Wales 2751, Australia.

Email: nicole.blay@westernsydney.edu.au

## Funding information

South Western Sydney Local Health District; Western Sydney University

## Abstract

**Aim:** This study aimed to profile the community nurse in Australia.

**Background:** The need for nurses in the community health care sector is increasing in response to shorter hospital stays, an aging population and chronic disease. The increase in demand has not been followed by appropriate workforce planning, leading to structural issues and lack of qualified nursing workforce in the community sector.

**Evaluation:** MEDLINE and ProQuest Public Health and grey literature were searched for records published between 2010 and 2020 relative to the profile of the community nurse in Australia. Twenty-five records (21 publications, 2 databases and 2 reports) were included in the review. Abstracted data followed the principles of workforce planning and included demographics, qualifications and roles.

**Key Issues:** Inconsistent definitions, self-reported data and a focus on practice nurses have contributed to data irregularities. Little is known about the specific aspects of community nursing work.

**Conclusion:** A lack of concrete data has overshadowed a community nursing workforce crisis with implications for patients' health and safety across the lifespan.

**Implications for Nursing Management:** There is urgent need for nurse managers globally to refocus nursing recruitment to the community sector to maintain quality and ensure sustainability of the nursing workforce.

## KEYWORDS

community health, nursing, primary health care, public health, workforce

## 1 | INTRODUCTION

The primary and community (P&C) health care system provides prevention, treatment and rehabilitative services outside the hospital system. In Australia, as globally, populations are living longer, often

with chronic disease (Australian Bureau of Statistics [ABS], 2018; Kyu et al., 2018), and lengths of hospital stays have decreased (Australian Institute of Health and Welfare [AIHW], 2017). Hospital in the Home programmes have exponentially increased (Montalto et al., 2020) as recovery from surgery or illness has moved from the acute sector to

[Correction added on 20 November 2021, after first online publication: The ORCID has been added to the fourth author's (Pauline Murray-Parahi) name in this version.]

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2021 The Authors. *Journal of Nursing Management* published by John Wiley & Sons Ltd.

the home, with support from community-based health care practitioners.

The integration of nurses into P&C care services is deemed to improve access, continuity and quality of care. The World Health Organization (WHO, 2017) considers the primary health care nursing role to be integral for health promotion, disease prevention and management. However, there are some challenges to maximizing nurses' contributions in the primary care capacity. There is a global shortage of nurses and midwives (Drennan & Ross, 2019) corresponding to over 50% of the total deficit in health care workers. The WHO estimates that an additional 9 million nurses are needed by 2030 to meet sustainable health and well-being goals (WHO, 2020). In addition, there has not been an alignment of skilled and qualified nursing supply with demand required at the community sector, with evidence of recruitment of nurses from acute care to meet the shortages in the primary setting (Ashley, 2016). With changing models of care from the acute to the community, it is necessary that workforce planning encompasses the community nursing sector to ensure sustainability and retention of skilled nursing staff in this setting.

## 2 | BACKGROUND

Multiple terms, definitions, governance and funding mechanisms of P&C-based services exist around the globe that have contributed to confusion and debate around what is primary and/or community health (Awofeso, 2004; Goodman et al., 2014; Muldoon et al., 2006; Phillips & Bazemore, 2010). This ambiguity extends to P&C nurse roles as titles are often used interchangeably (Drennan, 2019), and in Australia, health workforce data rely on self-reports at annual registration (AIHW, 2018).

P&C nurses fall into two main groups: the practice nurse, generally accepted to refer to nurses employed by privately run general practice (GP) (Jolly, 2007) and publicly funded community nurses attached to a health centre or clinic. The roles of P&C nurses are diverse as they are responsible for a population of 26 million from birth through to aged care in metropolitan, rural and remote regions (defined simply as outside Australia's major cities), across eight states and territories (ABS, 2021a; Royal Flying Doctor Service, n.d. [see map for detail]). The health system is complex, with the Federal Government funding primary health care and the states/territories responsible for targeted community health services (Productivity Commission, 2019; Swerissen et al., 2018) based on population health, geographical and socio-economic indicators. Aboriginal health services tend to be run by state/territory governments in metropolitan regions and the local community in rural and remote regions (Fitts et al., 2021).

Nurses employed within both groups include nurse practitioners (NPs), endorsed independent practitioners with master-level qualifications; registered nurses (RNs) who may be degree qualified (from 1985 to 1993 depending upon state/territory) or be certificated (hospital trained) (Jolly, 2007; The Department of Health, 2013); certificated or diploma educated (since 2014) enrolled nurses (ENs) (Blay &

Smith, 2020; Jolly, 2007); and increasingly, unregulated nurse assistants who may receive little or no training (Blay & Roche, 2020).


In light of the increasing need for P&C nurses as highlighted, it is imperative that nurse managers have a comprehensive understanding of the current workforce. As little is known about the community nurse workforce, a rapid systematic review was planned to profile the nurse working in the P&C health care sector in Australia and to ascertain future workforce needs. This review was guided by the fundamental principles of workforce planning that advocate the exploration of demographics, qualifications, activities and skills (De Bruecker et al., 2015) to answer the question who is the community nurse in Australia? Rapid reviews are vital in terms of response time for policymakers. Findings relative to employment data, role and career perceptions are important antecedents for nurse recruitment and therefore of benefit to nurse managers in the many countries with community nursing workforce shortages.

## 3 | METHODS

Rapid reviews are a method of systematically searching literature to inform policy in a more timely manner than can be achieved with the typical systematic review (Alliance for Health Policy and Systems Research, & World Health Organisation, 2017; Haby et al., 2016). Methods are many and remain under debate (Tricco et al., 2015), but include limiting databases, publication dates or language, excluding grey literature or a quality assessment, data abstraction by a single reviewer with verification by a second reviewer (Alliance for Health Policy and Systems Research, & World Health Organisation, 2017; Haby et al., 2016). We limited databases and grey literature searching and excluded a quality assessment as suggested above.

Two databases (MEDLINE and ProQuest Public Health) and Google's search engine using the MeSH terms Community 'AND' Nurse 'OR' Workforce 'AND' Australia were used to locate empirical research, workforce data and reports, and grey literature that described the community nurse in Australia. Searches conducted by the first author between March and July 2020 were limited to peer-reviewed journals, published in the English language between 2010 and 2020 (for decadal trend(s)) and availability of full text. Grey literature, accepted as print and electronic data, theses, and government, academic and business records that are not controlled by commercial publishers, helps to provide a more balanced review (Adams et al., 2016; Paez, 2017; Woods et al., 2020). Hand-searching and Really Simple Syndication (RSS) feeds were used to alert newly published manuscripts from selected journals. Opinion pieces, editorials and research where Australian data could not be extracted were excluded (see Figure 1).

The database search yielded 4801 published articles, whereas the yield from the grey literature was extensive ( $n = 8,940,000$ ); therefore, only the first five pages of results ( $n = 50$ ) were reviewed. After the removal of duplicates and following the screening of titles and abstracts, 32 published articles were retained for full-text review and two websites were retained from the grey literature.



**FIGURE 1** PRISMA published and unpublished studies included in the review

Record verification was conducted by a second author (P. M.-P.) with expertise in community nursing, and any differences of opinion were resolved by discussion and consensus. Following the method as outlined here, 21 published articles, 2 national datasets (comprising multiple tables) (AIHW, 2013b, 2015) and 2 workforce reports (Australian Primary Health Care Nurses Association [APNA], 2017, 2020) accessed from the respective websites were included in this review. The final review contained 25 records relative to the P&C health nurse (refer to Figure 1).

### 3.1 | Data abstraction and synthesis

Data on study design, aims and outcomes in terms of demographics, qualifications, work experience, activities and skills were extracted from the national databases, workforce reports and literature and entered into Microsoft Excel for integration. Descriptive statistics were applied for reporting purposes and percentage difference calculated for trend data. No assumptions were made for missing data. A

risk of bias assessment of included studies was not undertaken due to their descriptive nature, and any elimination may have compromised the description of the current workforce profile.

## 4 | RESULTS

The 21 journal publications reported findings relative to 18 research studies focussing on P&C nurses within metropolitan, rural and remote Australia. The national workforce datasets were constructed from self-reported data at nurse registration and annual renewal, and the APNA website reported findings from membership surveys (APNA, 2017, 2020).

The majority 47.6% ( $n = 10$ ) of publications described findings relative to the GP nurse, four publications respectively focussed on the community nurse (Aggar et al., 2018; Duiveman & Bonner, 2012; Happell et al., 2013; Terry et al., 2015) or P&C health workforce (Ashley, Brown, et al., 2018; Ashley, Halcomb, et al., 2018; Friesen & Comino, 2017; Oliver-Baxter et al., 2017), and the remainder the

remote area nurse (Lenthall et al., 2011; Zhao et al., 2017) or child and family health nurse (Borrow et al., 2011). Surveys were the preferred data collection method ( $n = 9$ ), followed by interviews and/or focus groups ( $n = 8$ ), secondary analyses of state and national datasets ( $n = 4$ ) and a self-reported activity log (diary) ( $n = 1$ ) (refer to Table 1).

#### 4.1 | P&C workforce

As shown in Table 2, there was a 5.4% increase in the employment of P&C nurses from 2012 to 2015. (Midwives were included in 2015 data, but as they could not be extrapolated and the ratio of RNs to registered midwives was 9.2:1, the term 'nurses' has been used hereafter [AIHW, 2015].) The increase in the P&C workforce is led by a 17.3% increase in nurses working in GPs, 'locum' or 'other private practice' (not defined) and to a lesser extent (10.3%) in Aboriginal health services. A concurrent decline (4.7%) occurred in community nurses (AIHW, 2015). Heywood and Laurence (2018a, 2018b) reported a 64% increase in practice nurses over 8 years, and workforce surveys indicated that 69% of respondents worked in GP (APNA, 2020) despite practice nurse leaders describing recruitment and retention difficulties (McKenna et al., 2015). Further analyses were unable to be conducted as turnover data, vacancy rates, population and full-time equivalent nursing data were not available.

Data relative to the clinical speciality in which P&C nurses primarily worked are conflicting. Of employed nurses nationally ( $n = 307,104$ ), fewer worked in community nursing (4.0%,  $n = 12,380$ ) and a greater number worked in GP nursing (4.2%,  $n = 12,821$ ) compared with stated employment area (AIHW, 2015) (refer to Table 2). Community nursing specialties included child and family health ( $n = 5444$ , 1.8%), Aboriginal health ( $n = 1500$ ), 'health promotion' ( $n = 1195$ , 0.4%) (AIHW, 2015), rural and remote health ( $n = 911$ ) (Heywood & Laurence, 2018b; Lenthall et al., 2011; Terry et al., 2015; Zhao et al., 2017), mining ( $n = 51$ ), tourist facilities ( $n = 19$ ) (Lenthall et al., 2011), research (Oliver-Baxter et al., 2017), corrective services, aged care and school health (APNA, 2020), although national data were not always available.

#### 4.2 | Nurse demographics

Literature supports national data in that P&C nurses are overwhelmingly female (89.3%), RNs (87.3%) and with a mean age of 44.4 years who work part-time (mean 30.9 h/week) (AIHW, 2015). GP nurses are younger (mean 46.6 years) than community (mean 48.2 years) and child and family health nurses (mean 49.9 years) following the employment of graduate nurses in the practice environment (Aggar et al., 2017; Heywood & Laurence, 2018a; Thomas et al., 2018).

Data on NPs or ENs are limited. Data from 2012 indicated that 8.6% ( $n = 4414$ ) of P&C nurses were enrolled; most worked in GP (4.6%,  $n = 2399$ ), 3.2% ( $n = 1630$ ) worked in community health and the remainder (0.7%,  $n = 385$ ) worked in child and family health or

health promotion (AIHW, 2013a). Similarly, 9% of APNA respondents were ENs and 2% were NPs (APNA, 2020).

#### 4.3 | Qualifications and skills

The majority (55–80%) of P&C nurses were trained in the hospital system (Terry et al., 2015; Thomas et al., 2018) and had on average 17 years of nursing experience and 5 years of P&C experience (AIHW, 2015; APNA, 2020; Ashley, Brown, et al., 2018; Borrow et al., 2011; Thomas et al., 2018). Nurses educated to bachelor's degree level ranged from 20% to 70% (Aggar et al., 2018; APNA, 2020; Parker et al., 2011; Thomas et al., 2018), 2–43% were working towards or had attained postgraduate qualifications (Aggar et al., 2018; APNA, 2020; Friesen & Comino, 2017; Lenthall et al., 2011; Parker et al., 2011), and less than 15% were qualified to manage chronic diseases, including asthma, diabetes and mental health issues (APNA, 2020) (refer to Table 1).

#### 4.4 | Nursing activities

Activities performed by P&C nurses as identified in the published literature were categorized into six recognized nursing workforce categories, namely: administration (general and administrative activities); direct care (activities directly related to patient care); indirect care (activities indirectly related to patient care); communication (communication with other health professionals, patients and/or carers); documentation (update or complete nursing or unit-related documentation by any medium); or other (activities not previously identified) (Blay et al., 2017; Blay & Roche, 2020).

As shown in Table 3, nursing activities ( $n = 63$ ) were diverse. The majority of identified activities were direct care ( $n = 39$ ), followed by administration ( $n = 6$ ), communication ( $n = 5$ ), indirect care ( $n = 5$ ), other ( $n = 5$ ) and documentation ( $n = 3$ ). Wound care was the nursing activity most frequently identified (APNA, 2020; Ashley, Brown, et al., 2018; Halcomb & Ashley, 2019; McInnes et al., 2019; Terry et al., 2015; Thomas et al., 2018), followed by immunization (APNA, 2020; Halcomb & Ashley, 2019; McInnes et al., 2019; Thomas et al., 2018), mental health management (Borrow et al., 2011; Halcomb & Ashley, 2019; Happell et al., 2013; Thomas et al., 2018), home visits (APNA, 2020; Borrow et al., 2011; Duiveman & Bonner, 2012; Halcomb & Ashley, 2019) and health services communication (Borrow et al., 2011; Halcomb & Ashley, 2019; Happell et al., 2013; Terry et al., 2015). The majority of listed direct care activities are associated with the practice environment indicating the research focus on this population.

### 5 | DISCUSSION

This review endeavoured to profile the community nurse in Australia. It could be argued that due to the focus on the GP nurse, the

TABLE 1 Summary of included publications, datasets and reports

Citation	Aim	Design and method	Sample, region and workplace	Demographics	Qualifications, skills and roles
Aggar et al. (2017)	To assess graduate nurse competency in a general practice transition to practice programme	Longitudinal exploratory mixed methods Survey at three monthly intervals and semistructured interviews	Graduate nurses in two metropolitan practices: Commencement ( $n = 6$ ) Completion ( $n = 4$ ) Programme preceptors ( $n = 7$ )	Community graduates: Female gender 83% ( $n = 10$ ) Mean age 33 years (SD 11.7) Previous nursing experience 33% ( $n = 4$ ) Community preceptors: Female gender 100% ( $n = 18$ ) Mean age 49 years (SD 8.6) Experience mean 24 years (SD 11.4)	Skill development—practice dependent Competency assessment as per national standards Career opportunities limited Graduates: Bachelor's degree 75% ( $n = 9$ ) Graduate certificate 17% ( $n = 2$ ) Master's degree 8% ( $n = 1$ ) Preceptors: Bachelor's degree 28% ( $n = 5$ ) Graduate certificate/diploma 86% ( $n = 12$ )
Aggar et al. (2018)	To compare competencies and experiences between graduates in a community, subacute and acute sector programme	Cohort study Survey at 6 and 12 months	Graduate nurses ( $n = 12$ ) Community preceptors ( $n = 18$ )		
Ashley, Brown, et al. (2018) and Ashley, Halcomb, et al. (2018)	To describe experiences of nurses who moved from the acute to primary or community sector	Sequential mixed methods Electronic survey Semistructured phone or face-to-face interviews	Nurse survey ( $n = 111$ ): Metropolitan (61%, $n = 67$ ) Rural (24%, $n = 26$ ) Remote (15.5%, $n = 17$ ) General practice 65% Interviewees: General practice ( $n = 6$ ) Schools ( $n = 3$ ) Community health, remote area, sexual health and refugee nursing ( $n = 1$ , respectively)	Survey: Female gender 96% Registered nurse 80% Mean age 45.4 years (SD 10.45) Experience: mean 18.9 years Primary or community: mean 3.4 years	Orientation period 81% Supernumerary period 49.5% Access to preceptor/mentor 35% Education-related leave and financial support—practice dependent Role ambiguity Role autonomy—practice dependent
Borrow et al. (2011)	To describe the community-based child and family health nurse in Western Australia	Qualitative Content analysis from 2-week self-reported diary of work activities Focus groups ( $n = 3$ )	Nurse diarists ( $n = 51$ ) Interviewees ( $n = 24$ ) from metropolitan and regional centres	Mean age 48.2 years (SD 7.9) Majority (>50%) had extensive nursing experience and 5 years child and family health nursing experience Part-time 64.4%	Postgraduate child and family health (57%) Midwifery qualification <61%
Duiveman and Bonner (2012)	To explore community nurses' experiences of negotiating client's care contracts	Qualitative descriptive: thematic analysis Focus groups ( $n = 2$ )	Two community health centres in New South Wales	Registered nurses ( $n = 14$ ) Female gender 86%	

(Continues)

TABLE 1 (Continued)

Citation	Aim	Design and method	Sample, region and workplace	Demographics	Qualifications, skills and roles
Friesen and Comino (2017)	To explore facilitators and barriers to research engagement by primary and community health staff	Exploratory Research culture in context tool distributed by email or hard copy	Multidisciplinary primary and community health staff within a New South Wales health district	Survey (n = 109) Female gender 87% (all respondents) Nurses 66% (n = 71) Child and family health 64% (n = 69) Primary and community health nurses 25% (n = 27)	Graduate certificate (nursing) 17.6% (n = 19) Tertiary-level postgraduate qualifications 27% (n = 29) (all respondents)
Halcomb and Ashley (2019) and Halcomb and Bird (2020)	To explore general practice nurses' roles, satisfaction and turnover intent	Mixed methods Cross-sectional nationwide electronic survey (n = 1166)	General practice 81.4% (n = 950) Other primary and community health settings 18.5% (n = 216) Metropolitan 56%, (n = 536) Rural 38% (n = 360) Remote 5% (n = 53) Nursing experience >20 years 69% (n = 657) Primary and community health experience 6+ years 58% (n = 553)	Female gender 98% (n = 771) Registered nurse 98% (n = 930) Enrolled nurse 9% (n = 89) Nurse practitioner 2% (n = 17) Based on completed surveys (n = 786) Mean age 49.9 years (SD 10.1) Part-time 56.8% (n = 441)	Satisfied with professional development opportunities 55% Intent to remain in general practice nursing 77% Satisfied with role 82% General practice role focussed on direct care 74% (n = 684) Regularly practised to full skill and knowledge 29% (n = 274) Majority able to practice skills
Happell et al. (2013)	To identify activities performed by community mental health nurses	Descriptive Quantitative analysis of 39 nursing activities grouped into four categories (clinical care, clinical organisation, practice administration and [service] integration)	Nursing dataset (n = 48,322) from community and ambulatory units (n = 252) in metropolitan, regional and rural Queensland over a 12-month period		Nursing activities: Clinical care 58% Clinical organisation 33% Administration 5% Integration 3%
Heywood and Laurence (2018a, 2018b)	To describe, compare and estimate the future supply of the general practice nursing workforce	Descriptive Secondary analysis of self-reported national workforce data Simulation model	General practice nurses (n = 12,746) Metropolitan 62.5% (n = 7966) Regional 35% (n = 4469) Remote 1.5% (n = 194) locations (2015 data)	Female gender 97% Registered nurse 80% Age 45 years or older 60% Nursing experience: mean 18.6 years Part-time work 65%	Intent to resign within 10 years 48% (n = 6093)

(Continues)

TABLE 1 (Continued)

Citation	Aim	Design and method	Sample, region and workplace	Demographics	Qualifications, skills and roles
Lenthall et al. (2011)	To describe the remote area nursing workforce	Descriptive Secondary analysis and comparison of population and remote area nursing data and surveys from 1995 to 2008	Identified remote area nurse, primary and community health positions ( $n = 469$ ) Survey respondents: ( $n = 349$ , 34.6% response rate) from seven states/territories Workplace: clinics, community health, mining and tourist facilities	Female gender 89% Mean age 44 years (median 46) Mean hours 47.6 per week Employers: state/territory or Aboriginal communities	Bachelor's level 55% Postgraduate rural-remote nursing 5% Significant decline in midwifery and child health qualifications Clinics: Single nurse 15% 2–5 nurses 69% 6–13 nurses 16%
McInnes et al. (2019)	Explore nurse and mentor experiences of a graduate general practice programme	Longitudinal qualitative Thematic analysis of semistructured telephone interviews prior, during and on completion of programme	Nurse graduates: Commencement ( $n = 8$ ) Completion ( $n = 4$ ) Nurse mentors ( $n = 9$ )	12-month programme incorporating two general practices	Career opportunities limited Able to practice university-acquired skills Expectation that graduates possess specialist skills
McKenna et al. (2015)	Exploration of facilitators and barriers to advanced practice in general practice environments	Descriptive Modified Delphi: thematic analysis from multidisciplinary semistructured telephone ( $n = 17$ ) or face-to-face interviews ( $n = 5$ )	General practice nurses ( $n = 4$ ) Nursing academics ( $n = 3$ ) Primary and community health decision makers <sup>a</sup> ( $n = 5$ ) Stakeholders ( $n = 11$ ) from across Australia <sup>a</sup>		Nurse education focusses on acute sector Time and financial constraints limit education No clear career pathway Recruitment and retention of skilled nurses problematic Salary less compared with acute sector Practice and culture Role can be isolating with little peer support Perception that role focusses on chronic disease and aged care Scope of practice limited by management and time constraints Role ambiguity
Oliver-Baxter et al. (2017)	To explore primary and community health higher degree research workforce	Descriptive Multidisciplinary cross-sectional: electronic survey of former higher degree research students	Survey respondents ( $n = 37$ ) Nursing background 16% ( $n = 6$ ) Workplace: University-based (74%) Metropolitan regions (58%)	Respondents in clinical practice ( $n = 2$ ) Nursing respondents currently working in P&C research 67% ( $n = 4$ )	Research career pathway unclear 78% ( $n = 29$ )

(Continues)

TABLE 1 (Continued)

Citation	Aim	Design and method	Sample, region and workplace	Demographics	Qualifications, skills and roles
Parker et al. (2011)	To explore the educational background of general practice nurses	Quantitative: electronic mail distributed survey in response to general practice network advertising	Survey respondents ( $n = 58$ , 74% response rate) Registered and enrolled nurses working in general practice environments	Female gender 96.5% Registered nurses 84% ( $n = 49$ ) Mean age 46 years (range 22–60) Mean 4.6 years in P&C environment Mean hours 28.2 per week (range 8–66)	Education perceived an important mechanism to raise status Hospital-based training 55% ( $n = 32$ ) Bachelor's degree 29% ( $n = 17$ ) Postgraduate P&C qualifications 17% ( $n = 10$ ) Preference for P&C short courses Staffing, time and financial constraints barriers to further education
Terry et al. (2015)	To explore any work, health and safety issues experienced by community nurses	Phenomenological Thematic analysis from semistructured telephone ( $n = 10$ ) or face-to-face interviews ( $n = 4$ ) with Tasmanian rural and remote community nurses	Registered nurses ( $n = 15$ ) working in 13 state-funded community centres	Female gender 87% ( $n = 13$ ) Age range 40–60 years Community nursing experience mean 8.8 years (range 1–31)	Community model ranged from single nurse to small teams working in clinics and/or homes Limited support mechanisms Work, health and safety issues: Fears for personal safety Manual handling Travel (roads, wildlife and weather) Home environment (cleanliness, passive smoking, uneven surfaces and pets)
Thomas et al. (2018)	To explore nurse and preceptor experiences of a graduate general practice programme	Qualitative: thematic analysis of semistructured interviews	Graduate nurses ( $n = 4$ , 67%) Preceptors ( $n = 5$ , 55.5%)	Graduates: Mean age 26 years (SD 8) Preceptors: Mean age 54 years (SD 13) Nursing experience mean 10 years (SD 7)	Limited financial support for education Acute sector perceived P&C experience to be of limited value P&C sector value acute experience Perception P&C environment more suitable for older nurses Good environment for skill development and unique skills, e.g., immunization Person-centred care, but autonomy is limited Career opportunities limited Preceptors: hospital-based training 80% Bachelor's degree 20% ( $n = 1$ )

(Continues)



TABLE 1 (Continued)

Citation	Aim	Design and method	Sample, region and workplace	Demographics	Qualifications, skills and roles
Walters et al. (2012)	To explore the possibility of telemonitoring by general practice nurses to supporting chronic disease	Mixed methods Survey of nurse participants trained in health mentoring (intervention) for a randomized controlled trial	General practice nurses ( <i>n</i> = 5) trained in health mentoring, Tasmania	Female gender 80% ( <i>n</i> = 4) Age range 36–60 years Median hours 24 per week (IQR 6) Primary and community experience median 4 years (IQR 1)	Self-management support training ( <i>n</i> = 1) Challenging work in a supportive environment Patient partnership important although carer involvement limited Performed health assessments 60% ( <i>n</i> = 3) Confidence in capacity to devise a patient action plan 40% ( <i>n</i> = 2)
Zhao et al. (2017)	To explore remote area practitioner workforce changes, turnover and costs between 2004 and 2015 <sup>b</sup>	Retrospective cross-sectional Secondary analysis of government payroll, personnel and financial activity	Remote area nurses, midwives, Aboriginal health practitioners and administrative staff from 54 remote area clinics in the Northern Territory	Remote area nurses increased from 120 to 135 (headcount) over 12 years Female gender 77% Non-Aboriginal 14% Significant increase in nurses aged >50 years Agency employees 15–20%	
Databases and websites	Aim	Design and method	Sample, region and workplace	Demographics	Qualifications, skills and roles
Australian Institute of Health and Welfare (AIHW, 2013a, 2013b, 2015)	To profile the nursing and midwifery workforce	Microsoft Excel data tables: self-reported survey data collated at nurse registration and renewal	2015: registered, enrolled nurses and midwives in Australia ( <i>n</i> = 360,008) Practice nursing ( <i>n</i> = 18,290) Community nursing ( <i>n</i> = 22,310)	Major work setting: Community health service ( <i>n</i> = 22,310) General practice ( <i>n</i> = 11,040) Locum or other privatepractice ( <i>n</i> = 7250) Aboriginal health service ( <i>n</i> = 1500)	
Australian Primary Health Care Nurses Association (APNA, 2017, 2020)	Annual membership survey	Brief reports	Survey respondents 2019 ( <i>n</i> = 1678)	General practice 69% Aboriginal health <sup>a</sup> Aged care <sup>a</sup> School health <sup>a</sup> Corrective services <sup>a</sup>	Bachelor's degree 70% Working towards or holding a postgraduate qualification 43% Immunization accredited 48% <15% qualified to manage chronic diseases, including asthma, diabetes and mental health issues

<sup>a</sup>No further details provided.<sup>b</sup>Longitudinal study: latest data reported.

**TABLE 2** Primary and community nursing workforce 2012 and 2015

Employment area of main job	2012 <sup>a</sup>	2015 <sup>b</sup>	% difference
Community	23,362	22,310	-4.7
Community aged care	5215		
Community mental health	4833		
Other community health service	13,314		
Practice	15,117	18,290	17.3
General practice	9165	11,040	17.0
Locum or other private practice	5952	7250	17.9
Aboriginal health service	1345	1500	10.3
Total community, practice and Aboriginal health	39,824	42,100	5.4
Employment specialty	2012 <sup>a</sup>	2015 <sup>b</sup>	% difference
Community	23,362	12,380	-88.7
General practice	9165	12,821	28.5
Child and family health		5444	
Health promotion		1195	
Total	32,527	31,840	-2.2

Note: At the time of searching, data from 2010–2011 and 2016–2020 were not available.

<sup>a</sup>Registered nurses only.

<sup>b</sup>Nurses and midwives.

Source: Adapted from AIHW (2015).

community nurse role remains largely undefined. The lack of research into the community nurse is surprising considering the importance placed by state/territory governments on targeting services to population needs and the incremental rise in Hospital in the Home programmes.

Findings highlight the need for consistent terminology and definitional specificity at a global and national level, to ensure accuracy with data reporting and for comprehensive nursing workforce planning (Drennan, 2019; Weller-Newton et al., 2020). In this instance, national data relied heavily on self-reported data (AIHW, 2018) whereas other data crucial for workforce planning such as full-time equivalents and turnover rates were not available affecting result reporting. Despite these statistical artefacts, the study has highlighted that the community nursing sector is facing a severe nursing workforce crisis relative to increased demand from an aging population, chronic disease and shorter hospitalizations. The trends identified an increase in the number of nurses in GP with a parallel decline in nursing staff working in the community. Although the rise in practice nurses is encouraging, it could be argued that with emphasis on recovery at home (Montalto et al., 2020), the need for community nurses is paramount, and patient care will be compromised if the negative trend continues (Parliament of Australia, 2002). The low numbers, older age and (part-time) employment patterns of P&C nurses, particularly in child and family health, are alarming. To attain full-time equivalence, a higher headcount (of part-time employees) is needed, and considering that almost 50% of P&C nurses are intending to resign (Heywood & Laurence, 2018a, 2018b) or likely retire within the next decade, workforce shortages are set to escalate. Moreover, if it is considered that Australia in 2017 had over 309,000 births (ABS, 2019) and that

approximately 6% of the population are aged under 5 (ABS, 2021b), it is probable that many infants and children are not being assessed by a child and family health nurse. The future health and developmental checks of babies and children are at risk.

Variances were found between studies and nurses' qualifications. Specifically, the APNA survey indicated that the majority of nurses were tertiary educated, whereas other studies showed that less than one third of P&C nurses had tertiary-level qualifications (Friesen & Comino, 2017; Parker et al., 2011; Thomas et al., 2018). Further research is needed, but it is possible that nurses with professional memberships are more keen to further their education or vice versa.

An astounding finding considering the rising incidence of chronic disease, and because mental health support was the third most frequently listed direct care activity, is that few nurses were accredited to manage these common reasons for GP visits (Finley et al., 2018; Kimble et al., 2020). Some have argued that postgraduate courses are limited (McKenna et al., 2015; Parliament of Australia, 2002) and that minimal attention is paid to P&C health in the undergraduate curriculum (Keleher et al., 2010; Murray-Parahi et al., 2020). However, it is almost 30 years since nursing education transitioned to the tertiary sector, and although P&C nurses acknowledged the importance of education for professional status, they also have a preference for short courses (Parker et al., 2011). This no doubt has limited capacity for P&C curriculum reform.

To ensure a sustainable workforce in line with population needs and changing models of care from hospital to the home, newly graduated and mid-level clinicians must be recruited. Little clarity is provided around the P&C role, scope of practice or a career pathway—factors that are known to influence nurse recruitment

**TABLE 3** Identified nurse activities and reported frequency (*n*)

Citation(s)	Categories ( <i>n</i> )	Activities ( <i>n</i> )
APNA (2020) Borrow et al. (2011) Halcomb and Ashley (2019) Happell et al. (2013)	Administration (1)	Arranging transport (1) Data processing and computer work (1) Organizing health promotion (1) Photocopying, faxing and scanning (1) Reception duties (1) Scheduling appointments, visits, recalls and reminders (3)
Borrow et al. (2011) Halcomb and Ashley (2019) Happell et al. (2013) Terry et al. (2015)	Communication	Case conferencing (1) Health professional (3) Health services and facilities (4) Telephone calls (1) Telehealth consultations (1)
APNA (2020) Ashley, Brown, et al. (2018) Borrow et al. (2011) Duiveman and Bonner (2012) Halcomb and Ashley (2019) Happell et al. (2013) McInnes et al. (2019) Thomas et al. (2018) Terry et al. (2015)	Direct care	Assisting with activities of daily living (1) Assisting with surgical procedures (2) Arthritis management (1) Basic nursing care (1) Case or care management (1) Child and family support (2) Complex and chronic disease management (1) Ear syringing (2) End-of-life care (1) Health assessment (2) <ul style="list-style-type: none"> <li>• Aged-related (1)</li> <li>• Antenatal and postnatal (2)</li> <li>• Child health (3)</li> <li>• Cardiac or respiratory (1)</li> <li>• Diabetes (1)</li> <li>• Domestic violence (1)</li> <li>• Mental health and cognition (1)</li> <li>• Men's health (1)</li> <li>• Smoking, nutrition, alcohol, physical activity and other risk factors (1)Home visits (4)</li> </ul> Immunization (5) Medication prompts and administration including insulin and intravenous medications (3) Mental health management or support (4) Patient education, health promotion and disease prevention (1) <ul style="list-style-type: none"> <li>• Mothercraft or breastfeeding (1)</li> <li>• Child health and immunization (2)</li> <li>• Chronic disease (1)</li> <li>• Community nurse role (1)</li> <li>• Drug, alcohol and smoking (2)</li> <li>• Goals for self-care or service provision (1)</li> <li>• Mental health (1)</li> <li>• Wound management (1)Plaster application and removal (1)</li> </ul> Suturing (1) Triage (2) Venepuncture and cannulation (2) Vital signs, blood sugar levels, ECG, peak flow and spirometry (2) Wart treatment (1) Wound care (6)

(Continues)

**TABLE 3** (Continued)

Citation(s)	Categories (n)	Activities (n)
Ashley, Brown, et al. (2018) Borrow et al. (2011) Halcomb and Ashley (2019) Happell et al. (2013) Terry et al. (2015)	Documentation (1)	Care plans, reports and patient records (4) Policy review (1) Service/practitioner referrals (1)
APNA (2020) Ashley, Brown, et al. (2018) Borrow et al. (2011) Duiveman and Bonner (2012) Halcomb and Ashley (2019) Terry et al. (2015)	Indirect care	Cold chain management (immunization transport and storage) (2) Establishing/directing play groups (1) Home assessment (2) Infection control and sterilizing (2) Review blood test results (1)
Borrow et al. (2011) Friesen and Comino (2017) Halcomb and Ashley (2019) Happell et al. (2013) Halcomb and Ashley (2019) Thomas et al. (2018) Terry et al. (2015)	Other	Management, leadership and mentoring (2) Ordering, restocking and cleaning (3) Research, audits and accreditation (2) Travel (1) Women's health (1) <sup>a</sup>

<sup>a</sup>No further details provided.

(Blay & Smith, 2020; Godsey et al., 2020). The APNA (2017) emphasizes, in line with global recommendations (WHO, 2017), that the P&C role is to promote health and prevent illness, yet with the exception of nurses in rural and remote locations (Roden et al., 2015), these activities are seldom realized (Ball et al., 2014; Sworn & Booth, 2020). Indeed, many of the nursing activities identified in this review are fundamental, could be performed by others and are associated with the practice environment. With a need for skilled nurses comparative with demand (WHO, 2017, 2020) nurse leaders and professional bodies can help address workforce shortages and perceived career limitations by encouraging education, upskilling and the NP role. An area for concern is the potential for vacant community nurse positions to be filled by an unregulated workforce, with negative consequences for skilled nurses' workload and patient safety. Employing more ENs would enable RNs to focus on preventative health care and help bridge the gap between fundamental care and expectations around scope of practice (Murray-Parahi et al., 2017).

Although Australia's community workforce figures are deeply concerning, they are not unique. International research has demonstrated that the majority of nurses in high-income countries work within the hospital sector (Drennan & Ross, 2019) and that globally, P&C nurses are in short supply (Buerhaus et al., 2015; WHO, 2020). Until such time that nursing research comprehensively explores and quantifies the activities undertaken by the P&C nurse, the existing confusion around role expectations (Ashley, Brown, et al., 2018; Ashley, Halcomb, et al., 2018) will continue, and current nursing workforce shortages will only worsen.

## 5.1 | Limitations

As a rapid review, only two databases were searched, data were cross-sectional and reported data were often generic restricting

comparisons. Caution should be taken when interpreting results, as changes in classification categories used in the different sources can impact workforce data. This review has identified that data on the P&C workforce are scant. Despite these limitations and resultant knowledge gaps, the review has provided a basis for our understanding of the P&C workforce and has highlighted a looming crisis in the community nursing sector.

## 6 | CONCLUSION

This review has highlighted a workforce crisis in the state and territory community nursing sector. Importantly, it has determined that definitional ambiguity has impacted self-reported data and national workforce statistics. The need for rigorous research exploring the community nurse role is of paramount importance for workforce sustainability and to ensure the health and safety of the Australian population from birth through to older age.

## 7 | IMPLICATIONS FOR NURSING MANAGEMENT

With changing population demographics, and models of care shifting from hospital to the home, nurse managers need to refocus nurse recruitment to the community sector. Profiling and developing the community nurse role to reduce negative role perceptions and expand scope of practice is a step towards the promotion of a sustainable community workforce, consistent with WHO recommendations.

## ACKNOWLEDGEMENTS

Acknowledgements are extended to Assoc. Prof. Lucie Ramjan, Assoc. Prof. Laretta Luck, Ms Amanda Sykes and Ms Karen Steel for

assistance with a preliminary exploration of contemporary literature or feedback on draft manuscripts. This manuscript is a precursor to an observational study of community health nurses funded by a partnership grant between Western Sydney University and South Western Sydney Local Health District.

## ETHICS STATEMENT

As a rapid systematic review, ethical approval was not required.

## CONFLICT OF INTERESTS

No conflicts of interest.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available at <https://researchdirect.westernsydney.edu.au/>. These data were derived from the following resources available in the public domain: AIHW (2013a, 2013b, 2015, 2017, 2018) and APNA (2017, 2020).

## ORCID

Nicole Blay  <https://orcid.org/0000-0001-8515-4209>

Mariana S. Sousa  <https://orcid.org/0000-0001-5034-9946>

Pauline Murray-Parahi  <https://orcid.org/0000-0002-6444-1150>

## REFERENCES

- Adams, J., Hillier-Brown, F. C., Moore, H. J., Lake, A. A., Araujo-Soares, V., White, M., & Summerbell, C. (2016). Searching and synthesising 'grey literature' and 'grey information' in public health: Critical reflections on three case studies. *Systematic Reviews*, 5, 164. Retrieved 7 December 2020 from <https://doi.org/10.1186/s13643-016-0337-y>
- Aggar, C., Bloomfield, J., Thomas, T. H., & Gordon, C. J. (2017). Australia's first transition to professional practice in primary care program for graduate registered nurses: A pilot study. *BMC Nursing*, 16, 14. Retrieved 18 November 2020, from [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5363053/pdf/12912\\_2017\\_Article\\_207.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5363053/pdf/12912_2017_Article_207.pdf), <https://doi.org/10.1186/s12912-017-0207-5>
- Aggar, C., Gordon, C. J., Thomas, T. H. T., Wadsworth, L., & Bloomfield, J. (2018). Evaluation of a community transition to professional practice program for graduate registered nurses in Australia. *Nurse Education in Practice*, 32, 101–107. <https://doi.org/10.1016/j.nepr.2018.03.005>
- Alliance for Health Policy and Systems Research, & World Health Organisation. (2017). Rapid reviews to strengthen health policy and systems: a practical guide; Tricco, A. C., Langlois, E. V. & Straus, S. E. (Eds); World Health Organization; Geneva. <https://www.who.int/alliance-hpsr/resources/publications/rapid-review-guide/en/>
- Ashley, C. (2016). Making the move to primary healthcare—Unearthing the experiences of acute care nurses transitioning to PHC employment. *Australian Nursing and Midwifery Journal*, 24(2), 34. <https://doi.org/10.3316/informit.223221439898833>
- Ashley, C., Brown, A., Halcomb, E., & Peters, K. (2018). Registered nurses transitioning from acute care to primary healthcare employment: A qualitative insight into nurses' experiences. *Journal of Clinical Nursing*, 27(3–4), 661–668. <https://doi.org/10.1111/jocn.13984>
- Ashley, C., Halcomb, E., Brown, A., & Peters, K. (2018). Experiences of registered nurses transitioning from employment in acute care to primary health care—Quantitative findings from a mixed-methods study. *Journal of Clinical Nursing*, 27(1–2), 355–362. <https://doi.org/10.1111/jocn.13930>
- Australian Bureau of Statistics. (2018). Chronic conditions. Australian Bureau of Statistics. Retrieved 2 June 2020 from <https://www.abs.gov.au/statistics/health/health-conditions-and-risks/national-health-survey-first-results/latest-release>
- Australian Bureau of Statistics. (2019). Births, Australia, 2018. Australian Bureau of Statistics. Retrieved 13 July 2020 from <https://www.abs.gov.au/ausstats/abs@.nsf/mf/3301.0>
- Australian Bureau of Statistics. (2021a). Population. Australian Bureau of Statistics. Retrieved 23 September from <https://www.abs.gov.au/statistics/people/population>
- Australian Bureau of Statistics. (2021b). Regional population. Australian Bureau of Statistics. Retrieved 23 June 2021 from <https://www.abs.gov.au/statistics/people/population/regional-population/2019-20#new-south-wales>
- Australian Institute of Health and Welfare. (2013a). Enrolled nurses principle area tables 1–9. Australian Institute of Health and Welfare. <https://www.aihw.gov.au/reports/workforce/nursing-and-midwifery-workforce-2015/data>
- Australian Institute of Health and Welfare. (2013b). Registered and enrolled nurses work setting tables 1–11. Australian Institute of Health and Welfare. <https://www.aihw.gov.au/reports-data/health-welfare-services/workforce/data>
- Australian Institute of Health and Welfare. (2015). Data tables: Nurses and midwives, overview tables 1–25. Australian Institute of Health and Welfare. <https://www.aihw.gov.au/reports/workforce/nursing-and-midwifery-workforce-2015/data>
- Australian Institute of Health and Welfare. (2017). Admitted patient care 2015–16: Australian hospital statistics (Health Services Series No. 75). Australian Institute of Health and Welfare. <https://www.aihw.gov.au/reports/hospitals/ahs-2015-16-admitted-patient-care/contents/table-of-contents>
- Australian Institute of Health and Welfare. (2018). Workforce. Australian Institute of Health and Welfare. Retrieved 24 September from <https://www.aihw.gov.au/reports-data/health-welfare-services/workforce/about>
- Australian Primary Health Care Nurses Association. (2017). Community health nursing. Australian Primary Health Care Nurses Association. Retrieved 29 November 2019 from <https://www.apna.asn.au/profession/what-is-primary-health-care-nursing/community-health-nursing>
- Australian Primary Health Care Nurses Association. (2020). Workforce summary report. Australian Primary Health Care Nurses Association. Retrieved 17 July 2020 from <https://www.apna.asn.au/profession/apna-workforce-survey>
- Awofeso, N. (2004). What is the difference between 'primary care' and 'primary healthcare'? [Guest editorial]. *Quality in Primary Care*, 12(2), 93–94. <https://primarycare.imedpub.com/what-is-the-difference-between-primary-care-and-primary-healthcare.php?aid=2237>
- Ball, J., Philippou, J., Pike, G., & Sethi, J. (2014). Survey of district and community nurses in 2013 (Report to the Royal College of Nursing). National Nursing Research Unit (NNRU). <https://www.kcl.ac.uk/nmpc/research/nrru/publications/reports/dn-community-rcn-survey-report—updated-27-05-14.pdf>
- Blay, N., & Roche, M. A. (2020). A systematic review of activities undertaken by the unregulated Nursing Assistant. *Journal of Advanced Nursing*, 76(7), 1538–1551. <https://doi.org/10.1111/jan.14354>
- Blay, N., Roche, M. A., Duffield, C., & Gallagher, R. (2017). Intra-hospital transfers and the impact on nursing workload. *Journal of Clinical Nursing*, 26(23–24), 4822–4829. <https://doi.org/10.1111/jocn.13838>
- Blay, N., & Smith, L. E. (2020). An integrative review of Enrolled Nurse recruitment and retention. *Collegian*, 27(1), 89–94. [https://www.collegianjournal.com/article/S1322-7696\(19\)30061-7/fulltext](https://www.collegianjournal.com/article/S1322-7696(19)30061-7/fulltext), <https://doi.org/10.1016/j.jcolegn.2019.06.005>

- Borrow, S., Munns, A., & Henderson, S. (2011). Community-based child health nurses: An exploration of current practice. *Contemporary Nurse: A Journal for the Australian Nursing Profession*, 40(1), 71–86. <https://doi.org/10.5172/conu.2011.40.1.71>
- Buerhaus, P. I., DesRoches, C. M., Dittus, R., & Donelan, K. (2015). Practice characteristics of primary care nurse practitioners and physicians. *Nursing Outlook*, 63(2), 144–153. <https://doi.org/10.1016/j.outlook.2014.08.008>
- De Bruecker, P., Van den Bergh, J., Beliën, J., & Demeulemeester, E. (2015). Workforce planning incorporating skills: State of the art. *European Journal of Operational Research*, 243(1), 1–16. <https://doi.org/10.1016/j.ejor.2014.10.038>
- Drennan, V. (2019). More care out of hospital? A qualitative exploration of the factors influencing the development of the district nursing workforce in England. *Journal of Health Services Research & Policy*, 24(1), 11–18. <https://doi.org/10.1177/1355819618769082>
- Drennan, V. M., & Ross, F. (2019). Global nurse shortages—The facts, the impact and action for change. *British Medical Bulletin*, 130(1), 1–13. <https://doi.org/10.1093/bmb/ldz014>
- Duiveman, T., & Bonner, A. (2012). Negotiating: Experiences of community nurses when contracting with clients. *Contemporary Nurse: A Journal for the Australian Nursing Profession*, 41(1), 120–125. <https://doi.org/10.5172/conu.2012.41.1.120>
- Finley, C. R., Chan, D. S., Garrison, S., Korownyk, C., Kolber, M. R., Campbell, S., Eurich, D. T., Lindblad, A. J., Vandermeer, B., & Allan, G. M. (2018). What are the most common conditions in primary care?: Systematic review. *Canadian family physician Medecin de famille canadien*, 64(11), 832–840. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6234945/>
- Fitts, M. S., Humphreys, J., Dunbar, T., Bourke, L., Mulholland, E., Guthridge, S., Zhao, Y., Jones, M. P., Boffa, J., Ramjan, M., Murakami-Gold, L., Tangey, A., Comerford, C., Schultz, R., Campbell, N., Mathew, S., Liddle, Z., Russell, D., & Wakerman, J. (2021). Understanding and responding to the cost and health impact of short-term health staffing in remote and rural Aboriginal and Torres Strait Islander community-controlled health services: A mixed methods study protocol. *BMJ Open*, 11(8), e043902. <https://doi.org/10.1136/bmjopen-2020-043902>
- Friesen, E. L., & Comino, E. J. (2017). Research culture and capacity in community health services: Results of a structured survey of staff. *Australian Journal of Primary Health*, 23(2), 123–131. <https://doi.org/10.1071/PY15131>
- Godsey, J. A., Houghton, D. M., & Hayes, T. (2020). Registered nurse perceptions of factors contributing to the inconsistent brand image of the nursing profession. *Nursing Outlook*, 68(6), 808–821. <https://doi.org/10.1016/j.outlook.2020.06.005>
- Goodman, R. A., Bunnell, R., & Posner, S. F. (2014). What is “community health”? Examining the meaning of an evolving field in public health. *Preventive Medicine*, 67(Suppl 1), S58–S61. <https://doi.org/10.1016/j.yjmed.2014.07.028>
- Haby, M. M., Chapman, E., Clark, R., Barreto, J., Reveiz, L., & Lavis, J. N. (2016). What are the best methodologies for rapid reviews of the research evidence for evidence-informed decision making in health policy and practice: A rapid review. *Health Research Policy and Systems*, 14(1), 83. <https://doi.org/10.1186/s12961-016-0155-7>
- Halcomb, E., & Ashley, C. (2019). Are Australian general practice nurses underutilised?: An examination of current roles and task satisfaction. *Collegian*, 26, 522–527. <https://doi.org/10.1016/j.colegn.2019.02.005>
- Halcomb, E., & Bird, S. (2020). Job Satisfaction and Career Intention of Australian General Practice Nurses: A Cross-Sectional Survey. *Journal of Nursing Scholarship*, 52(3), 270–280. <https://doi.org/10.1111/jnu.12548>
- Happell, B., Gaskin, C., Hoey, W., Nizette, D., & Veach, K. (2013). The activities that nurses working in community mental health perform: A geographical comparison. *Australian Health Review*, 37(4), 1–7. <https://doi.org/10.1071/AH13045>
- Heywood, T., & Laurence, C. (2018a). The general practice nurse workforce: Estimating future supply. *Australian Journal of General Practice*, 47(11), 788–795. <https://doi.org/10.31128/AJGP-01-18-4461>
- Heywood, T., & Laurence, C. (2018b). An overview of the general practice nurse workforce in Australia, 2012–15. *Australian Journal of Primary Health*, 24(3), 227–232. <https://doi.org/10.1071/PY17048>
- Jolly, R. (2007). Practice nursing in Australia (Research Paper no. 10 2007–08). Parliament of Australia. [https://www.aph.gov.au/About-Parliament/Parliamentary\\_Departments/Parliamentary\\_Library/pubs/rp/RP0708/08rp10#Endnotes](https://www.aph.gov.au/About-Parliament/Parliamentary_Departments/Parliamentary_Library/pubs/rp/RP0708/08rp10#Endnotes)
- Keleher, H., Parker, R., & Francis, K. (2010). Preparing nurses for primary health care futures: How well do Australian nursing courses perform? *Australian Journal of Primary Health*, 16(3), 211–216. <https://doi.org/10.1071/py09064>
- Kimble, L. P., Phan, Q., Hillman, J. L., Blackman, J., Shore, C., Swainson, N., & Amobi, C. N. (2020). The CAPACITY professional development model for community-based primary care nurses: Needs assessment and curriculum planning. *Nursing Economics*, 38(3), 110–120,148.
- Kyu, H. H., Abate, D., Abate, K. H., & GBD 2017 DALYs and HALE Collaborators. (2018). Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*, 392(10159), 1859–1922. Retrieved 10 November 2018 from. <https://www.sciencedirect.com/science/article/pii/S0140673618323353>
- Lenthall, S., Wakerman, J., Opie, T., Dunn, S., Macleod, M., Dollard, M., Rickard, G., & Knight, S. (2011). Nursing workforce in very remote Australia, characteristics and key issues. *The Australian Journal of Rural Health*, 19(1), 32–37. <https://doi.org/10.1111/j.1440-1584.2010.01174.x>
- McInnes, S., Halcomb, E., Huckel, K., & Ashley, C. (2019). Experiences of registered nurses in a general practice-based new graduate program: A qualitative study. *Australian Journal of Primary Health*, 25(4), 366–373. <https://doi.org/10.1071/PY19089>
- McKenna, L., Halcomb, E., Lane, R., Zwar, N., & Russell, G. (2015). An investigation of barriers and enablers to advanced nursing roles in Australian general practice. *Collegian*, 22(2), 183–189. <https://doi.org/10.1016/j.colegn.2015.02.003>
- Montalto, M., McElduff, P., & Hardy, K. (2020). Home ward bound: Features of hospital in the home use by major Australian hospitals, 2011–2017. *Medical Journal of Australia*, 213(1), 22–27. Retrieved 24 June 2021, from. <https://www.mja.com.au/journal/2020/213/1/home-ward-bound-features-hospital-home-use-major-australian-hospitals-2011-2017>, <https://doi.org/10.5694/mja2.50599>
- Muldoon, L. K., Hogg, W. E., & Levitt, M. (2006). Primary care (PC) and primary health care (PHC). What is the difference? *Canadian Journal of Public Health*, 97(5), 409–411. <https://doi.org/10.1007/BF03405354>
- Murray-Parahi, P., DiGiacomo, M., Jackson, D., Phillips, J., & Davidson, P. M. (2020). Primary health care content in Australian undergraduate nursing curricula. *Collegian*, 27(3), 271–280. <https://doi.org/10.1016/j.colegn.2019.08.008>
- Murray-Parahi, P., Edgar, V., Descallar, J., Comino, E., & Johnson, M. (2017). ENSCOPE: Scoping the practice of enrolled nurses in an Australian community health setting. *International Nursing Review*, 64(1), 59–68. <https://doi.org/10.1111/inr.12305>
- Oliver-Baxter, J., Brown, L., & McIntyre, E. (2017). Surviving or thriving in the primary health care research workforce: The Australian experience. *Australian Journal of Primary Health*, 23(2), 183–188. <https://doi.org/10.1071/PY15190>

- Paez, A. (2017). Gray literature: An important resource in systematic reviews. *Journal of Evidence-Based Medicine*, 10(3), 233–240. <https://doi.org/10.1111/jebm.12266>
- Parker, R., Keleher, H., & Forrest, L. (2011). The work, education and career pathways of nurses in Australian general practice. *Australian Journal of Primary Health*, 17(3), 227–232. <https://doi.org/10.1071/PY10074>
- Parliament of Australia. (2002). Report on the inquiry into nursing—The patient profession: Time for action. [https://www.aph.gov.au/Parliamentary\\_Business/Committees/Senate/Community\\_Affairs/Completed\\_inquiries/2002-04/nursing/report/index](https://www.aph.gov.au/Parliamentary_Business/Committees/Senate/Community_Affairs/Completed_inquiries/2002-04/nursing/report/index)
- Phillips, R. L. J., & Bazemore, A. W. (2010). Primary care and why it matters for U.S. health system reform. *Health Affairs*, 29(5), 806–810. <https://doi.org/10.1377/hlthaff.2010.0020>
- Productivity Commission. (2019). Primary and community health (Report on Government Services 2015). Australian Government. <https://www.pc.gov.au/research/ongoing/report-on-government-services>
- Roden, J., Jarvis, L., Campbell-Crofts, S., & Whitehead, D. (2015). Australian rural, remote and urban community nurses' health promotion role and function. *Health Promotion International*, 31(3), 704–714. <https://doi.org/10.1093/heapro/dav018>
- Royal Flying Doctor Service. (n.d.). Defining rural and remote. RDFS. Retrieved 23 September from <https://www.flyingdoctor.org.au/what-we-do/research/defining-rural-remote/>
- Swerissen, H., Duckett, S., & Moran, G. (2018). Mapping primary care in Australia. Grattan Institute. <https://grattan.edu.au/wp-content/uploads/2018/07/906-Mapping-primary-care.pdf>
- Sworn, K., & Booth, A. (2020). A systematic review of the impact of 'mised care' in primary, community and nursing home settings. *Journal of Nursing Management*, 28(8), 1805–1829. <https://doi.org/10.1111/jonm.12969>
- Terry, D., Lê, Q., Nguyen, U., & Hoang, H. (2015). Workplace health and safety issues among community nurses: A study regarding the impact on providing care to rural consumers. *BMJ Open*, 5(8), e008306. <https://doi.org/10.1136/bmjopen-2015-008306>
- The Department of Health. (2013). Appendix iv: History of Commonwealth involvement in the nursing and midwifery workforce. Australian Government. Retrieved 22 September from <https://www1.health.gov.au/internet/publications/publishing.nsf/Content/work-review-australian-government-health-workforce-programs-toc~appendices~appendix-iv-history-commonwealth-involvement-nursing-midwifery-workforce>
- Thomas, T. H. T., Bloomfield, J. G., Gordon, C. J., & Aggar, C. (2018). Australia's first transition to professional practice in primary care program: Qualitative findings from a mixed-method evaluation. *Collegian*, 25(2), 201–208. <https://doi.org/10.1016/j.colegn.2017.03.009>
- Tricco, A. C., Antony, J., Zarin, W., Striffler, L., Ghassemi, M., Ivory, J., Perrier, L., Hutton, B., Moher, D., & Straus, S. E. (2015). A scoping review of rapid review methods. *BMC Medicine*, 13(1), 224. <https://doi.org/10.1186/s12916-015-0465-6>
- Walters, J. A. E., Courtney-Pratt, H., Cameron-Tucker, H., Nelson, M., Robinson, A., Scott, J., Turner, P., Walters, E. H., & Wood-Baker, R. (2012). Engaging general practice nurses in chronic disease self-management support in Australia: insights from a controlled trial in chronic obstructive pulmonary disease. *Australian Journal of Primary Health*, 18(1), 74–79. <https://doi.org/10.1071/PY10072>
- Weller-Newton, J. M., Phillips, C., Roche, M. A., McGillion, A., Mapes, J., Dufty, T., Schlieff, J., Boyd, L., Geary, A., & Haines, S. (2020). Datasets to support workforce planning in nursing: A scoping review. *Collegian*, 28(3), 324–332. <https://doi.org/10.1016/j.colegn.2020.09.001>
- Woods, S., Phillips, K., & Dudash, A. (2020). Grey literature citations in top nursing journals: A bibliometric study. *Journal of the Medical Library Association*, 108(2), 262–269. <https://doi.org/10.5195/jmla.2020.760>
- World Health Organization. (2017). Enhancing the role of community health nursing for universal health coverage (ISBN: 978-92-4-151189-6). (Human Resources for Health Observer Series No. 18).
- World Health Organization. (2020). Nursing and midwifery. World Health Organization. Retrieved 20 September from <https://www.who.int/news-room/fact-sheets/detail/nursing-and-midwifery>
- Zhao, Y., Russell, D. J., Guthridge, S., Ramjan, M., Jones, M. P., Humphreys, J. S., Carey, T. A., & Wakerman, J. (2017). Long-term trends in supply and sustainability of the health workforce in remote Aboriginal communities in the Northern Territory of Australia. *BMC Health Services Research*, 17(1), 836. <https://doi.org/10.1186/s12913-017-2803-1>

**How to cite this article:** Blay, N., Sousa, M. S., Rowles, M., & Murray-Parahi, P. (2022). The community nurse in Australia. Who are they? A rapid systematic review. *Journal of Nursing Management*, 30(1), 154–168. <https://doi.org/10.1111/jonm.13493>