

RESEARCH

Open Access



# A combination of the percentages of IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and granzyme B<sup>+</sup>CD19<sup>+</sup>B cells is associated with acute hepatic rejection: a case control study

Ji-Qiao Zhu<sup>†</sup>, Jing Wang<sup>†</sup>, Xian-Liang Li<sup>†</sup>, Wen-Li Xu, Shao-cheng Lv, Xin Zhao, Ren Lang<sup>\*</sup> and Qiang He<sup>\*</sup>

## Abstract

**Background:** T cells and B cells play a key role in alloimmune responses. We aimed to characterize the shift of T cell subsets and B cell subsets during acute hepatic rejection, and further determine whether they could serve as a prognostic marker.

**Methods:** Blood samples together with the clinical data from liver transplant recipients with and without acute hepatic rejection were collected and analyzed as well as from a validation cohort.

**Results:** Upon activation the expression of TGF- $\beta$  and granzyme B in CD19<sup>+</sup>B cells, and the expression of IL-2 and IFN- $\gamma$  in CD4<sup>+</sup>T cells were higher in acute hepatic rejection. However, only the frequencies of granzyme B<sup>+</sup>CD19<sup>+</sup>B cells and IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells correlated with liver function in addition to with each other. A combination of the two cell subsets as a novel marker could classify rejection versus non-rejection (area under the curve 0.811,  $p=0.001$ ) with the cut-off value of 62.93%, which was more sensitive for worse histological changes ( $p=0.027$ ). Moreover, the occurrence rate of acute rejection was higher in the group with the novel marker  $>62.93\%$  ( $p=0.000$ ). The role of the novel marker was further confirmed in a validation cohort, which was identified to be the only significant independent risk factor for acute rejection (odds ratio: 0.923; 95% CI confidence interval: 0.885–0.964;  $p=0.000$ ).

**Conclusions:** A combination of the percentages of IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and granzyme B<sup>+</sup>CD19<sup>+</sup>B cells can distinguish rejection from non-rejection, which can be used as a potential prognostic marker for acute rejection in liver transplant recipients.

**Keywords:** Acute rejection, Liver transplantation, B lymphocyte subsets, T lymphocyte subsets

## Introduction

Liver transplantation is a major therapeutic approach in patients with end-stage liver diseases [1]. Following transplantation acute rejection remains a major challenge despite the development of immunosuppressive drugs [2]. Although abnormalities in liver function may raise concerns about acute rejection, invasive liver biopsies are still the gold standard for definitive diagnoses [3]. Therefore, a potential biomarker with minimal invasion for the diagnosis of acute rejection after liver transplantation has

\*Correspondence: dr-langren@hotmail.com; heqiangsurg@163.com

<sup>†</sup>Ji-Qiao Zhu, Jing Wang and Xian-Liang Li contributed equally to this manuscript

Department of Hepatobiliary and Pancreaticosplenic Surgery, Medical Research Center, Beijing Organ Transplant Center, Beijing Chaoyang Hospital, Capital Medical University, No. 8 Gongtinan Road, Chaoyang District, Beijing 100020, China



been required. Actually, acute rejection is realized via the immune cells, thus, blood-derived biomarkers are under intensive research.

By means of producing different cytokines and exerting different biological functions, CD4<sup>+</sup>T cells are thought to play a crucial role during the rejection process [4, 5]. Tang et al. reported IL-2 and IFN- $\gamma$  could be produced by memory CD4<sup>+</sup>T-helper cells to mediate vigorous allograft rejection [6]; Sawitzki B et al. found IFN- $\gamma$  secreted by CD4<sup>+</sup>T cells was responsible for transplantation tolerance in a skin transplantation model [7]. In another study transplantation of older organs triggered more potent alloimmune responses via proinflammatory cytokine production of IL-17 [8]. Immature dendritic cells-derived exosomes improved the percent of survival and suppressed rejection associated cytokines IFN- $\gamma$ , IL-2, IL-17 production [9]. Hence, CD4<sup>+</sup>T cells can promote the rejection process.

In the meanwhile, B cells were once considered to act as antigen-presenting cells and provide co-stimulatory signals for T cell activation in addition to antibody production. Recent studies have demonstrated the production of multiple cytokines can enable B cells to regulate the immune response [10, 11]. IL-10 is a typical cytokine produced by regulatory B cells, which suppresses the proliferation and inflammatory cytokine productions of effector T cells [12]. In our previous study, Granzyme B (GrB) produced by B cells was found to maintain allospecific tolerance in patients with renal rejection [13]. Moreover, low mRNA expression of TGF- $\beta$  in the biopsy could reveal early acute rejection implying an increased risk for renal graft failure [14]. Therefore, regulatory B cells are characterized by the immunosuppressive function.

However, the dynamic changes of T cell subsets and B cell subsets during acute rejection have not been well elucidated. Furthermore, there has not been a widely accepted biomarker for predicting acute rejection. Thus, we performed the study to investigate whether the T cell subsets and B cell subsets could serve as a prognostic marker in liver transplant recipients (LTR) with acute rejection.

## Materials and methods

### Study design

LTR with acute rejection and with stable liver function were enrolled in this study, who underwent a first single liver transplant or were followed up at Beijing Chaoyang Hospital. There were not any signs of post-operative complications in all cases at the time of blood sampling except acute rejection. The controls did not have an episode of acute rejection if the follow-up period was  $\leq 6$  months; otherwise, the period between acute rejection and sampling was  $> 6$  months. A validation

cohort was further enrolled to confirm the observations. The study was approved by the Institutional Review Board of Beijing Chaoyang Hospital (No. 2016-2-19-38) in accordance with the Helsinki declaration of 1975, as revised in 1983. Written informed consent was obtained from all participants.

### Immunosuppressive management

Immunosuppressive therapy contained induction with basiliximab (20 mg on day 0 and day 4) and maintenance on calcineurin inhibitors. Acute rejection was diagnosed using clinical and laboratory parameters and graft biopsy assessed according to the Banff schema [15]. LTR with acute rejection were treated by adding the dose of immunosuppressants or with a round of steroids (Methylprednisolone given 500 mg on day 1, 240 mg on day 2, then daily reduced by 40 mg till the 8th day, finally changed to prednisolone 20 mg/day for  $\sim 1$  month).

### Cell culture

Lymphocyte subset analysis was performed at the time of biopsy before the therapeutic intervention. Peripheral blood mononuclear cells (PBMC) were isolated by ficoll density gradient centrifugation and suspended ( $1 \times 10^6$ /ml) in RPMI-1640 medium supplemented with 10% heat inactivated fetal calf serum, 100 U/ml penicillin, 100  $\mu$ g/ml Streptomycin (Invitrogen). For cytokines produced by B cells, PBMC were stimulated with IgG + IgM (5.4  $\mu$ g/ml) in the presence of IL-21 (50 ng/ml), seeded in a 24-well flat-bottom plate, and incubated for 20 h at 37  $^{\circ}$ C, 5% CO<sub>2</sub>. Brefeldin A (1  $\mu$ g/ml) was added for the last 4 h of incubation. For cytokines produced by T cells, PBMC ( $1 \times 10^6$ /ml) were stimulated with cell stimulation cocktail (eBioscience) and seeded in a 15-ml tube and incubated for 4 h at 37  $^{\circ}$ C, 5% CO<sub>2</sub>.

### Antibodies and flow cytometric measurement

Anti-human mAbs included PE-Cy7-CD3 (Biolegend), PerCP-Cy5.5-CD8 (Biolegend), APC-IL-17 (eBioscience), PE-IFN- $\gamma$  (Biolegend), APC-IL-2 (Biolegend), PE-Cy7-CD19 (Biolegend), FITC-Zombie (Biolegend), APC-fire750-CD69 (Biolegend), PerCP-Cy5.5-TGF- $\beta$  (Biolegend), APC-IL-10 (Biolegend), PE-GrB (eBioscience). After stimulation, PBMC were harvested and first stained with surface antibodies followed by fixation/permeabilization (Cytotfix/Cytoperm kit, BD Biosciences) and subsequent intracellular staining, as previously described [13].

Flow cytometry was performed in NovoCyte D2060R (ACEA Biosciences Inc). NovoEXpress software (San Diego, CA, USA) was used for analysis. Flow cytometry characterization of lymphocyte subsets is presented in Additional file 1: Figure S1.

### Statistical analysis

Data analyses were carried out by using SPSS 19.0 computer software (IBM Corp., Armonk, NY, USA) and GraphPad Prism 5 software (GraphPad Software Inc., La Jolla, CA, USA). Values were expressed as mean  $\pm$  standard deviation. The Kolmogorov–Smirnov test was used for the normal distribution of continuous variables. The independent samples t-test was employed for quantitative variables. The Mann–Whitney U test was selected due to non-normal distribution. The Chi-square or Fisher's exact test was used to compare nominal variables. Receiver operating characteristic curve (ROC) analysis and comparison of the area under the curve (AUC) was performed. The cut-off value for positive parameters was further determined by optimal sensitivity and specificity on ROC curve analysis. Multivariate Cox analysis was employed to determine the predictors. Relative risk was expressed as an odds ratio (OR) with a 95% confidence interval (CI). A p-value  $< 0.05$  was considered statistically significant.

## Results

### Characteristics of LTR

A total of 15 LTR with acute rejection and 30 paired controls with stable liver function (1:2) were enrolled in this study. They were matched by gender, age ( $\pm 3$  years), primary diseases for transplantation (benign or malignant), main immunosuppressants, and follow-up periods ( $\pm 7$  days). Patients were all first deceased donor allograft recipients. The characteristics of LTR are listed in Additional file 1: Table S1. Of the 15 patients, 13 had T cell-mediated rejection proven by histological evidence and two mixed T cell and antibody-mediated rejection. Six patients were treated with adding the dose of immunosuppressants alone and nine followed by a round of steroids. All patients recovered with stable liver function.

### The percentages of IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and IL-2<sup>+</sup>CD4<sup>+</sup>T cells could differentiate rejection from non-rejection upon activation

CD4<sup>+</sup>T cells have been reported to play an important role in allograft rejection [16, 17] while the dynamic changes of T lymphocyte subsets have not been well investigated in LTR. Therefore, we wanted to check the shift of circulating T cells. After analysis, we found the production of IFN- $\gamma$ , IL-2, and IL-17 in resting CD4<sup>+</sup>T cells increased in LTR with rejection but did not reach statistical significance between the two groups ( $P > 0.05$ , Fig. 1a–c). When activated with stimulation cocktail, CD4<sup>+</sup>T cell derived IL-2 production ( $p = 0.042$ , Fig. 1d) and IFN- $\gamma$  production ( $p = 0.007$ , Fig. 1e) rose significantly in LTR


with rejection; meanwhile, the percentages of IL-17<sup>+</sup>CD4<sup>+</sup>T cells remained similar ( $P > 0.05$ , Fig. 1f) between the two groups (Additional file 1: Table S2).

### The percentages of GrB<sup>+</sup>CD19<sup>+</sup>B cells and TGF- $\beta$ <sup>+</sup>CD19<sup>+</sup>B cells could differentiate rejection from non-rejection upon activation

We previously showed that the frequency of GrB<sup>+</sup>CD19<sup>+</sup>B cells increased in patients with renal rejection [13]. In this study, we extended our research to determine how various B cell subsets would shift in LTR with acute rejection. First, we detected the expression of IL-10, TGF- $\beta$ , and GrB in circulating B cells. The results showed the production of IL-10, TGF- $\beta$  and GrB in resting CD19<sup>+</sup>B cells was similar between the two groups, respectively ( $P > 0.05$ , Fig. 2a–c). Next, B cells were stimulated with anti-IgG/IgM in the presence of IL-21. We observed an obvious increase in the percentages of TGF- $\beta$ <sup>+</sup>CD19<sup>+</sup>B cells ( $p = 0.049$ , Fig. 2d) and GrB<sup>+</sup>CD19<sup>+</sup>B cells ( $p = 0.005$ , Fig. 2e) from LTR with acute rejection while the percentage of IL-10<sup>+</sup>CD19<sup>+</sup>B cells ( $P > 0.05$ , Fig. 2f) was comparable upon stimulation between two groups (Additional file 1: Table S3).

### The percentages of GrB<sup>+</sup>CD19<sup>+</sup>B cells and IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells correlate with liver function

Since activated cell subsets could differentiate rejection from non-rejection, we wanted to know whether a sharp increase in the subset percentages was in close relation to liver function. First, clinical data from LTR with and without rejection were collected and compared (Additional file 1: Table S4). LTR with acute rejection had higher levels of aspartate transaminase (AST,  $p = 0.001$ , Additional file 1: Figure S2A) and alanine aminotransferase (ALT,  $p = 0.002$ , Additional file 1: Figure S2B). Surprisingly, levels of TBIL were comparable (Figure S2C). We attributed the TBIL change to the diagnosis of acute rejection at an early stage as the rise of transaminases was more rapid. Notably, tacrolimus levels were also similar between the two groups (Additional file 1: Figure S2D), implying the similar immunosuppressive therapy of the two groups. Then, we performed the correlation analysis between activated cell subsets and the transaminases. We found the percentages of GrB<sup>+</sup>CD19<sup>+</sup>B cells correlated positively with AST ( $p = 0.010$ , Fig. 3a) and ALT ( $p = 0.002$ , Fig. 3b) in LTR with acute rejection, and with ALT ( $p = 0.035$ , Fig. 3f) in LTR without rejection. Such a correlation was not found between AST and ALT, and the percentages of TGF- $\beta$ <sup>+</sup>CD19<sup>+</sup>B cells either in LTR with rejection (Fig. 3c, d) or in LTR without rejection (Fig. 3g, h) as well as the percentages of GrB<sup>+</sup>CD19<sup>+</sup>B cells and AST in LTR without rejection (Fig. 3e). Similarly, the percentages of IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells correlated positively



with AST ( $p=0.014$ , Fig. 4a) and ALT ( $p=0.011$ , Fig. 4b) in LTR with acute rejection and with ALT ( $p=0.040$ , Fig. 4f) in LTR without rejection. In contrast, the percentages of IL-2<sup>+</sup>CD4<sup>+</sup>T cells did not correlate with AST and ALT either in LTR with rejection (Fig. 4c, d) or in LTR without rejection (Fig. 4g, h). Neither did the percentages of IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells correlate with AST in LTR without rejection (Fig. 4e).

#### A combination of the percentages of IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and GrB<sup>+</sup>CD19<sup>+</sup>B cells as a novel marker can identify acute rejection



From the above analysis, the percentages of activated IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and activated GrB<sup>+</sup>CD19<sup>+</sup>B cells were determined to have a strong association with acute rejection. Besides, we also observed a positive correlation between the percentages of activated IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and the percentages of activated GrB<sup>+</sup>CD19<sup>+</sup>B cells after further analysis (Fig. 5a). Based on their relationship we proposed a novel marker of a combination of the two cell subsets and tested its efficacy.

First, we performed the analysis of AUC-ROC for the percentages of activated IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and the percentages of activated GrB<sup>+</sup>CD19<sup>+</sup>B cells and the

new marker (Fig. 5b), which revealed the latter strongly classified rejection versus non-rejection (AUC 0.811,  $p=0.001$ ) with the cut-off value of 62.93%. The corresponding sensitivity and specificity were 93.3% and 66.7%, respectively. Then, the LTR were regrouped according to the cut-off value. We found the occurrence rate of acute rejection was higher in the group with the new marker > 62.93% ( $p=0.000$ ; Fig. 5c). Finally, the association between three parameters and the severity of the Banff histological grades (mild group and non-mild group) was analyzed (Fig. 5d, e). Of note, LTR with mild inflammation and damage had low values of the new marker ( $p=0.027$ , Fig. 5f). While the percentages of activated IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and activated GrB<sup>+</sup>CD19<sup>+</sup>B cells failed to identify the LTR (Table S5, Fig. 5g, h). Thus, the novel marker not only identified LTR with acute rejection but also was more sensitive for the worse acute histological changes.


#### The novel marker can predict acute rejection in a validation cohort

In an attempt to confirm the ability of the novel marker to identify LTR with rejection, we performed an independent study. In this validation cohort, a total of 101



LTR were enrolled, who accepted liver transplantation or were followed up at our center within one year after surgery. Blood samples were obtained to analyze

the lymphocyte subsets. These patients were followed up for at least one year to observe the occurrence of acute rejection as most acute rejections occur within



the first year following liver transplantation [18–20]. 23 LTR had a rejection episode during the follow-up period and were treated by adding the dose of immunosuppressants or with a round of steroids.


In this validation cohort, LTR with acute rejection had the higher percentages of activated IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells ( $p=0.000$ , Fig. 6a) and activated GrB<sup>+</sup>CD19<sup>+</sup>B cells ( $p=0.002$ , Fig. 6b). Again, the percentages of activated IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and the percentages of activated GrB<sup>+</sup>CD19<sup>+</sup>B cells correlated with each other ( $p=0.047$ , Fig. 6c). We considered that might explain the effect of the new marker, which showed higher values in LTR with rejection ( $p=0.000$ , Fig. 6d, Additional file 1: Table S6). Subsequent ROC analysis further confirmed that the new marker was a strong predictor of acute rejection with an AUC of 0.810 (Fig. 6e). When the LTR were regrouped at the cut-off value of 62.93%, the occurrence rate of acute rejection remained higher in the group with the new marker >62.93% ( $p=0.004$ ; Fig. 6f). Finally, we asked if the new marker could independently predict acute rejection. On univariate analysis age ( $p=0.009$ ), CMV infection (0.049), postoperative lymphocyte percentage ( $p=0.020$ ), and the new marker ( $p=0.000$ ) were identified to be positive between LTR with and without rejection. Comparison of other parameters did not reach significance ( $p>0.05$ ). Using multiple logistic regression analysis, the new marker (OR: 0.923; 95%

CI: 0.885–0.964;  $p=0.000$ ) was the only significant independent risk factor (Table 1).

## Discussion

In this study, we found that a combination of percentages of activated IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and activated GrB<sup>+</sup>CD19<sup>+</sup>B cells can distinguish rejection from non-rejection, which can be used as a novel biomarker for diagnosis of acute rejection in LTR.


IL-17 is a pro-inflammatory cytokine implicated in the pathogenesis of lung and renal transplantation as reduced IL-17 production was associated with attenuation of acute rejection [21, 22]. Similar results were reported in liver transplantation [23, 24]. In contrast, we found high IL-17 production in LTR with rejection upon activation, which failed to reach significance. Notably, we detected the percentages of IL-17<sup>+</sup>CD4<sup>+</sup>T cells rather than serum levels of IL-17 in this study. In addition, LTR with and without rejection were all found to have higher levels of IL-17 than healthy controls during the postoperative period [25], which might minimize the difference. IL-2 and IFN- $\gamma$  are typically secreted by activated CD4<sup>+</sup>T cells, specifically T-helper 1 cells, which play an important role in acute rejection. Our findings are in line with reported studies stressing the function of IL-2 and IFN- $\gamma$  during the rejection process [26–28]. They also showed LTR with improved liver function had low levels of IL-2 and IFN- $\gamma$ . However, the correlation analysis has not



been investigated among their studies. We first demonstrated the percentages of activated IFN- $\gamma$ +CD4 $^{+}$ T cells instead of activated IL-2 $^{+}$ CD4 $^{+}$ T cells was found to positively correlate with transaminases in both the rejection group and the non-rejection group. A possible explanation for the difference between the two cytokines might be their main functions. IL-2 promotes the proliferation and differentiation of lymphocytes [29, 30]; while IFN- $\gamma$  can induce hepatocyte apoptosis or inhibit hepatocyte cell cycle progression [31].

Recent studies have revealed that B cells can regulate the immune response via immunosuppressive cytokine secretion in addition to producing antibodies and presenting antigens to T cells [32]. Due to its immunosuppressive nature, regulatory B cells have been repeatedly reported to be involved in transplantation tolerance [33, 34]. In contrast, its regulation in acute rejection following

transplantation is poorly understood. In our study, we found LTR with acute rejection had higher percentages of TGF- $\beta$ +CD19 $^{+}$ B cells and GrB $^{+}$ CD19 $^{+}$ B cells, which might affect as feedback to main tolerance [35]. Surprisingly, the percentage of IL-10 $^{+}$ CD19 $^{+}$ B cells was similar to controls. Although TGF- $\beta$ , GrB, and IL-10 are all produced by regulatory B cells they belong to distinct cell subsets (data not shown). The most efficient way for IL-10 expression is to stimulate B cells through TLR signaling activation while maximal TGF- $\beta$  and GrB production is induced through BCR [13, 36–38]. When we further analyzed the correlations between cytokines and liver function, only the percentages of GrB $^{+}$ CD19 $^{+}$ B cells showed an association with transaminases. TGF- $\beta$  can regulate both cell differentiation and cell survival while GrB can strongly induce T cell apoptosis in addition to suppressing CD4 $^{+}$ T cell proliferation [39, 40]. Thus,



higher values of transaminases call for higher production of GrB during the rejection process.

Of note, the percentages of GrB<sup>+</sup>CD19<sup>+</sup>B cells correlated positively with the percentages of IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells. As they are pro-inflammatory cytokine and anti-inflammatory cytokine, respectively, these two kinds of cytokines might work as a loop as discussed above. Therefore, we proposed a combination of the cell subsets as a new marker based on their relationship. We also showed the new marker had strong predictive capacity after analysis of AUC. When regrouped by the cut-off value, 93.3% LTR exhibiting rejection had a high value and 6.7% a low value. Furthermore, LTR with a high value of the novel marker also had worse histological scores, confirming its association with the severity of rejection. Taken together, our data suggest that the new marker is more sensitive for the risk of acute rejection.

The function of the new marker was further validated in an independent cohort. The results from the validation cohort showed the new marker could identify

LTR with a very low risk for rejection from LTR who were at substantial risk for developing acute rejection. We found the new marker was identified to be the only independent predictor for rejection, which outperformed the other parameters. Several biomarkers have been shown to predict clinical acute rejection, including primary biliary cirrhosis, younger age, hepatitis C [41], IL-10-1082 polymorphism [42], de novo donor-specific antibody [20], and cytokine promoter polymorphisms [43]. However, all these risk factors need prospective validation. Notably, tacrolimus levels were comparable between LTR with and without acute rejection in both cohorts. Since the LTR received similar immunosuppressive therapy, then, rapid recovery of lymphocyte subsets might account for this phenomenon.

In conclusion, a combination of the percentages of IFN- $\gamma$ <sup>+</sup>CD4<sup>+</sup>T cells and granzyme B<sup>+</sup>CD19<sup>+</sup>B cells can distinguish rejection from non-rejection, which can be used as a potential prognostic marker for acute rejection in liver transplant recipients.

**Table 1** Risk factors for acute rejection in liver transplant recipients

Parameters	Univariate analysis			Multivariate analysis		
	Rejection (n = 23)	Non-rejection (n = 78)	P	OR	CI	P
Sex (male)	22	71	0.777			
Age	49.09 ± 9.91	54.19 ± 7.40	0.009	1.073	0.997–1.154	0.060
Primary disease			0.821			
Hepatitis related cirrhosis	13	42				
Hepacellular carcinoma	10	36				
Preoperative creatinine (μmol/l)	65.43 ± 17.85	73.84 ± 28.69	0.364			
Preoperative bilirubin (μmol/l)	52.80 ± 71.58	72.52 ± 142.72	0.827			
Preoperative INR	1.20 ± 0.27	1.30 ± 0.30	0.160			
Preoperative lymphocyte count (10 <sup>9</sup> /l)	1.26 ± 0.80	1.09 ± 0.71	0.331			
Preoperative lymphocyte percentage	25.49 ± 13.64	24.11 ± 11.94	0.639			
Operating time (min)	401.22 ± 282.80	312.68 ± 288.11	0.176			
Warm ischemia time (min)	2.48 ± 0.85	2.63 ± 0.69	0.521			
Cold storage time (h)	7.35 ± 1.03	7.62 ± 1.06	0.307			
Bleeding (> 800 ml)	12	39	0.855			
Transfusion (> 800 ml)	5	31	0.113			
Anhepatic phase (min)	127.74 ± 26.45	118.99 ± 35.11	0.272			
Immunosuppressants			0.484			
Tacrolimus-based	19	59				
Cyclosporin A-based	4	19				
Tacrolimus level (ng/ml)	6.04 ± 4.99	7.35 ± 5.63	0.365			
CMV	9	15	0.049	2.355	0.700–7.930	0.167
Postoperative lymphocyte count (10 <sup>9</sup> /l)	1.41 ± 1.04	1.19 ± 0.72	0.252			
Postoperative lymphocyte percentage	23.88 ± 8.23	19.10 ± 8.56	0.020	0.962	0.902–1.027	0.246
New marker	86.62 ± 16.76	66.48 ± 17.33	0.000	0.923	0.885–0.964	0.000

INR international normalized ratio, CMV cytomegalovirus, OR odds ratio, CI 95% confidence interval

### Abbreviations

GrB: Granzyme B; PBMC: Peripheral blood mononuclear cells; ROC: Receiver operating characteristic; AUC: Area under the curve; OR: Odds ratio; CI: Confidence interval; AST: Aspartate transaminase; ALT: Alanine aminotransferase; TBIL: Total bilirubin.

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12967-021-02855-w>.

**Additional file 1: Table S1.** Characteristics of liver transplant recipients with and without acute rejection. **Table S2.** Comparison of the percentages of T cell subsets between liver transplant recipients with and without rejection. **Table S3.** Comparison of the percentages of B cell subsets between liver transplant recipients with and without rejection. **Table S4.** Analysis of liver function and tacrolimus levels between liver transplant recipients with and without rejection. **Table S5.** Comparison of the Banff grades in the rejection group with the percentages of GrB+CD19+B cell and IFN-γ+CD4+T cell and the new marker. **Table S6.** Comparison of the percentages of GrB+CD19+B cell and IFN-γ+CD4+T cell and the new marker in a validation cohort. **Figure S1.** Flow cytometry characterization of T cell subsets and B cell subsets. **Figure S2.** AST and ALT deteriorate in liver transplant recipients with acute rejection. Comparison of levels of AST (A), ALT (B), TBIL (C) and FK506 (D) between liver transplant recipients with and without acute rejection. Bars represent mean and standard deviation. AST, aspartate transaminase; ALT, alanine aminotransferase; TBIL, total bilirubin.

### Acknowledgements

Not applicable.

### Authors' contributions

LR and HQ contributed conception and design of the study; ZJ, WJ and LX organized the database; XW, XZ and WQ performed the statistical analysis; ZJ, WJ and LX wrote the first draft of the manuscript; the rest of the authors wrote sections of the manuscript. All authors contributed to manuscript revision, read and approved the submitted version. All authors read and approved the final manuscript.

### Funding

This work is supported by Open Project of Beijing Key Laboratory of Tolerance Induction and Organ Protection in Transplantation (2017YZNS01) and National Natural Science Foundation of China (81601392) and Beijing Municipal Science and Technology Commission (Z181100001718164).

### Availability of data and materials

All data generated or analyzed during this study are included in this published article and its supplementary information files.

### Declarations

#### Ethics approval and consent to participate

The study was approved by the Institutional Review Board of Beijing Chaoyang Hospital (No. 2016-2-19-38) in accordance with the Helsinki declaration of 1975, as revised in 1983. Written informed consent was obtained from all participants.

**Consent for publication**

Not applicable.

**Competing interests**

The authors declare that they have no competing interests.

Received: 21 February 2021 Accepted: 21 April 2021

Published online: 01 May 2021

**References**

- Adams DH, Sanchez-Fueyo A, Samuel D. From immunosuppression to tolerance. *J Hepatol*. 2015;62(1 Suppl):S170–85.
- Rodriguez-Peralvarez M, Germani G, Papastergiou V, Tsochatzis E, Thalassinou E, Luong TV, et al. Early tacrolimus exposure after liver transplantation: relationship with moderate/severe acute rejection and long-term outcome. *J Hepatol*. 2013;58(2):262–70.
- Nilles KM, Krupp J, Lapin B, Sustento-Reodica N, Gallon L, Levitsky J. Incidence and impact of rejection following simultaneous liver-kidney transplantation. *J Hepatol*. 2015;62(2):340–5.
- Li J, Li X, Liang C, Ling L, Chen Z, Wong CK, et al. Coreceptor blockade targeting CD4 and CD8 allows acceptance of allogeneic human pluripotent stem cell grafts in humanized mice. *Biomaterials*. 2020;248:120013.
- Liu Z, Fan H, Jiang S. CD4(+) T-cell subsets in transplantation. *Immunol Rev*. 2013;252(1):183–91.
- Tang AL, Bingaman AW, Kadavil EA, Leiser DB, Farber DL. Generation and functional capacity of polyclonal alloantigen-specific memory CD4 T cells. *Am J Transplant*. 2006;6(6):1275–84.
- Sawitzki B, Kingsley CI, Oliveira V, Karim M, Herber M, Wood KJ. IFN-gamma production by alloantigen-reactive regulatory T cells is important for their regulatory function in vivo. *J Exp Med*. 2005;201(12):1925–35.
- Oberhuber R, Heinbokel T, Cetina Bieffer HR, Boenisch O, Hock K, Bronson RT, et al. CD11c+ dendritic cells accelerate the rejection of older cardiac transplants via interleukin-17A. *Circulation*. 2015;132(2):122–31.
- Pang XL, Wang ZG, Liu L, Feng YH, Wang JX, Xie HC, et al. Immature dendritic cells derived exosomes promotes immune tolerance by regulating T cell differentiation in renal transplantation. *Aging (Albany NY)*. 2019;11(20):8911–24.
- Rosser EC, Mauri C. Regulatory B cells: origin, phenotype, and function. *Immunity*. 2015;42(4):607–12.
- Oleinika K, Mauri C, Salama AD. Effector and regulatory B cells in immune-mediated kidney disease. *Nat Rev Nephrol*. 2019;15(1):11–26.
- Cherukuri A, Salama AD, Carter CR, Landsittel D, Arumugakani G, Clark B, et al. Reduced human transitional B cell T1/T2 ratio is associated with subsequent deterioration in renal allograft function. *Kidney Int*. 2017;91(1):183–95.
- Zhu J, Zeng Y, Dolff S, Bienholz A, Lindemann M, Brinkhoff A, et al. Granzyme B producing B-cells in renal transplant patients. *Clin Immunol*. 2017;184:48–53.
- Viklicky O, Hribova P, Volk HD, Slatinska J, Petrasek J, Bandur S, et al. Molecular phenotypes of acute rejection predict kidney graft prognosis. *J Am Soc Nephrol*. 2010;21(1):173–80.
- Banff schema for grading liver allograft rejection: an international consensus document. *Hepatology*. 1997;25(3):658–663.
- Gerlach UA, Vogt K, Schlickeiser S, Meisel C, Streitz M, Kunkel D, et al. Elevation of CD4+ differentiated memory T cells is associated with acute cellular and antibody-mediated rejection after liver transplantation. *Transplantation*. 2013;95(12):1512–20.
- Horne PH, Lunsford KE, Eiring AM, Wang Y, Gao D, Bumgardner GL. CD4+ T-cell-dependent immune damage of liver parenchymal cells is mediated by alloantibody. *Transplantation*. 2005;80(4):514–21.
- Volpin R, Angeli P, Galioto A, Fasolato S, Neri D, Barbazza F, et al. Comparison between two high-dose methylprednisolone schedules in the treatment of acute hepatic cellular rejection in liver transplant recipients: a controlled clinical trial. *Liver Transpl*. 2002;8(6):527–34.
- Saliba F, Duvoux C, Gugenheim J, Kamar N, Dharancy S, Salame E, et al. Efficacy and safety of everolimus and mycophenolic acid with early tacrolimus withdrawal after liver transplantation: a multicenter randomized trial. *Am J Transplant*. 2017;17(7):1843–52.
- Jucaud V, Shaked A, DesMarais M, Sayre P, Feng S, Levitsky J, et al. Prevalence and impact of de novo donor-specific antibodies during a multicenter immunosuppression withdrawal trial in adult liver transplant recipients. *Hepatology*. 2019;69(3):1273–86.
- Gupta PK, Wagner SR, Wu Q, Shilling RA. IL-17A blockade attenuates obliterative bronchiolitis and IFN-gamma cellular immune response in lung allografts. *Am J Respir Cell Mol Biol*. 2017;56(6):708–15.
- Kwan T, Chadban SJ, Ma J, Bao S, Alexander SI, Wu H. IL-17 deficiency attenuates allograft injury and prolongs survival in a murine model of fully MHC-mismatched renal allograft transplantation. *Am J Transplant*. 2015;15(6):1555–67.
- Zhou Y, Yang X, Zhang H, Jiang J. The roles of T helper type 17/regulatory T cells in acute rejection after liver transplantation in rats. *Transplantation*. 2015;99(6):1126–31.
- Li J, Lai X, Liao W, He Y, Liu Y, Gong J. The dynamic changes of Th17/Treg cytokines in rat liver transplant rejection and tolerance. *Int Immunopharmacol*. 2011;11(8):962–7.
- Fabrega E, Lopez-Hoyos M, San Segundo D, Casafont F, Pons-Romero F. Changes in the serum levels of interleukin-17/interleukin-23 during acute rejection in liver transplantation. *Liver Transpl*. 2009;15(6):629–33.
- Li DY, Xie SL, Wang GY, Dang XW. CD47 blockade alleviates acute rejection of allogeneic mouse liver transplantation by reducing ischemia/reperfusion injury. *Biomed Pharmacother*. 2020;123:109793.
- Jiang ZJ, Shen QH, Chen HY, Yang Z, Shuai MQ, Zheng S. Galectin-1 restores immune tolerance to liver transplantation through activation of hepatic stellate cells. *Cell Physiol Biochem*. 2018;48(3):863–79.
- Li P, Zhang YY, Deng J. PDL1 gene-loaded BMSCs induce liver transplantation immune tolerance. *Eur Rev Med Pharmacol Sci*. 2018;22(10):3214–23.
- Liao W, Lin JX, Leonard WJ. IL-2 family cytokines: new insights into the complex roles of IL-2 as a broad regulator of T helper cell differentiation. *Curr Opin Immunol*. 2011;23(5):598–604.
- Williams MA, Tzgnik AJ, Bevan MJ. Interleukin-2 signals during priming are required for secondary expansion of CD8+ memory T cells. *Nature*. 2006;441(7095):890–3.
- Horras CJ, Lamb CL, Mitchell KA. Regulation of hepatocyte fate by interferon-gamma. *Cytokine Growth Factor Rev*. 2011;22(1):35–43.
- Mann MK, Maresz K, Shriver LP, Tan Y, Dittel BN. B cell regulation of CD4+CD25+ T regulatory cells and IL-10 via B7 is essential for recovery from experimental autoimmune encephalomyelitis. *J Immunol*. 2007;178(6):3447–56.
- Lai G, Nakayama Y, Sethi A, Singh AK, Burrell BE, Kulkarni N, et al. Interleukin-10 from marginal zone precursor B-cell subset is required for costimulatory blockade-induced transplantation tolerance. *Transplantation*. 2015;99(9):1817–28.
- Cugini D, Noris M. Toward a B-cell signature of tolerance? *Kidney Int*. 2010;78(5):435–7.
- Chen M, Mohtize M, Mattei MF, Villemot JP, Kohler C, Faure GC, et al. Reduced levels of both circulating CD4+ CD25+ CD127(low/neg) and CD4+ CD8(neg) invariant natural killer regulatory T cells in stable heart transplant recipients. *Clin Exp Immunol*. 2011;163(1):104–12.
- Lampropoulou V, Hoehlig K, Roch T, Neves P, Calderon Gomez E, Sweeney CH, et al. TLR-activated B cells suppress T cell-mediated autoimmunity. *J Immunol*. 2008;180(7):4763–73.
- Lampropoulou V, Calderon-Gomez E, Roch T, Neves P, Shen P, Stervbo U, et al. Suppressive functions of activated B cells in autoimmune diseases reveal the dual roles of Toll-like receptors in immunity. *Immunol Rev*. 2010;233(1):146–61.
- Dedobbeleer O, Stockis J, van der Woning B, Coulie PG, Lucas S. Cutting edge: active TGF-beta1 released from GARP/TGF-beta1 complexes on the surface of stimulated human B lymphocytes increases class-switch recombination and production of IgA. *J Immunol*. 2017;199(2):391–6.
- Kriegel MA, Li MO, Sanjabi S, Wan YY, Flavell RA. Transforming growth factor-beta: recent advances on its role in immune tolerance. *Curr Rheumatol Rep*. 2006;8(2):138–44.

40. Chesneau M, Michel L, Dugast E, Chenouard A, Baron D, Pallier A, et al. Tolerant kidney transplant patients produce B cells with regulatory properties. *J Am Soc Nephrol*. 2015;26(10):2588–98.
41. Levitsky J, Goldberg D, Smith AR, Mansfield SA, Gillespie BW, Merion RM, et al. Acute rejection increases risk of graft failure and death in recent liver transplant recipients. *Clin Gastroenterol Hepatol*. 2017;15(4):584–932.
42. Warle MC, Metselaar HJ, Hop WC, Tilanus HW. Cytokine gene polymorphisms and acute liver graft rejection: a meta-analysis. *Liver Transpl*. 2005;11(1):19–26.
43. Fernandes H, Koneru B, Fernandes N, Hameed M, Cohen MC, Raveche E, et al. Investigation of promoter polymorphisms in the tumor necrosis factor-alpha and interleukin-10 genes in liver transplant patients. *Transplantation*. 2002;73(12):1886–91.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

