



# Managing general anesthesia for low invasive dental procedures while maintaining spontaneous respiration with low concentration remifentanyl: a cross-sectional study

Daijiro Ogumi, Shota Abe, Hikaru Sato, Fumihiko Suzuki, Hiroyoshi Kawaai, Shinya Yamazaki

Department of Dental Anesthesiology, Ohu University, School of Dentistry, Fukushima, Japan

**Background:** We assessed the relationship between patient age and remifentanyl dosing rate in patients managed under general anesthesia with spontaneous breathing using low-dose remifentanyl in sevoflurane.

**Methods:** The participants were patients with an American Society of Anesthesiologists Physical Status of 1 or 2 maintained under general anesthesia with low-dose remifentanyl in 1.5-2.0% sevoflurane. The infusion rate of remifentanyl was adjusted so that the spontaneous respiratory rate was half the rate prior to the induction of anesthesia, and  $\gamma^{\text{H}}$  ( $\mu\text{g}/\text{kg}/\text{min}$ ) was defined as the infusion rate of remifentanyl under stable conditions where the respiratory rate was half the rate prior to the induction of anesthesia for  $\geq 15$  minutes. The relationship between  $\gamma^{\text{H}}$  and patient age was analyzed statistically by Spearman's correlation analysis.

**Results:** During dental treatment under general anesthesia using low-dose remifentanyl in sevoflurane, a significant correlation was detected between  $\gamma^{\text{H}}$  and patient age. The regression line of  $y = -0.00079x + 0.066$  (y-axis;  $\gamma^{\text{H}}$ , x-axis; patient's age) was provided. The values of  $\gamma^{\text{H}}$  provide 0.064  $\mu\text{g}/\text{kg}/\text{min}$  at 2 years and 0.0186  $\mu\text{g}/\text{kg}/\text{min}$  at 60 years. Therefore, as age increases, the dosing rate exhibits a declining trend. Furthermore, in the dosing rate of remifentanyl when the patient's respiratory rate was reduced by half from the preanesthetic respiratory rate, the dosing rate provided was around 0.88 mL/h in all ages if the remifentanyl was diluted as 0.1 mg/mL. EtCO<sub>2</sub> showed  $51.0 \pm 5.7$  mmHg, and SpO<sub>2</sub> was controlled within the normal range by this method. In addition, all dental treatments were performed without major problems, such as awakening and body movement during general anesthesia, and the post-anesthetic recovery process was stable.

**Conclusion:** General anesthesia with spontaneous breathing provides various advantages, and the present method is appropriate for minimally invasive procedures.

**Keywords:** Age; Minimally Invasive Surgical Procedures; Remifentanyl; Sevoflurane; Spontaneous Respiration.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



## INTRODUCTION

Remifentanyl is a widely used ultra-short-acting opioid analgesic that causes severe respiratory depression. However, low concentrations of remifentanyl that are used

under sedation with artificial ventilation in the ICU have been reported to improve respiratory patterns, reduce the workload on the respiratory muscles while maintaining spontaneous respiration [1], suppress stress hormone secretion, secure urine volume during perioperative management, and effectively prevent hyperglycemia [2].

Received: March 28, 2023 • Revised: June 12, 2023 • Accepted: June 21, 2023

Corresponding Author: Shinya Yamazaki, Chief Professor, Department of Dental Anesthesiology, Ohu University, School of Dentistry, 31-1 Misumido, Tomita, Koriyama, Fukushima, 963-8611 Japan

Tel: +81-24-932-9337 E mail: [zakiyama@ops.dti.ne.jp](mailto:zakiyama@ops.dti.ne.jp)

Copyright© 2023 Journal of Dental Anesthesia and Pain Medicine

Furthermore, the coadministration of remifentanyl with propofol at a dose of 0.05-0.1  $\mu\text{g}/\text{kg}/\text{min}$  allowed for intravenous sedation while maintaining spontaneous respiration and was effective in suppressing an abnormal gag reflex [3].

Low concentrations of remifentanyl are often used with sevoflurane [4,5,6] or propofol [7,8,9] to maintain general anesthesia under spontaneous respiration. Many previous case reports were limited to specific age groups, such as children [4,5,8,9] and adults [6,7], with no examination of the correlation between administration rates across all age groups (from infants to adults). Based on the above findings, we maintained general anesthesia in all age groups using a low concentration of remifentanyl, at which the spontaneous respiration rate was half of the rate prior to the induction of anesthesia. Based on this large amount of data, we hypothesized that the maintenance dose of remifentanyl at which the spontaneous respiration rate becomes half of the rate prior to the induction of anesthesia could be predicted at any age by obtaining a regression line for age and maintenance dose of remifentanyl.

## METHODS

This was a cross-sectional study (case series). The participants included patients who had been treated in the previous 5 years (2015-2020) at our university and affiliated clinics for whom dental treatment while maintaining consciousness was difficult due to dental phobia, abnormal gag reflex, intellectual disability, or refusal of treatment (in the case of children), and those classified as grades 1-2 according to the American Society of Anesthesiologists Physical Status (ASA-PS) classification system. General anesthesia was maintained with spontaneous breathing using low concentrations of remifentanyl and 1.5-2.0% sevoflurane alone. Patients undergoing highly invasive surgeries, such as those involving the jawbone, were excluded from the study. The participants were limited to those who underwent

conservative treatment, prosthetic treatment, or simple extraction under general anesthesia. For procedures accompanied by pain, infiltration anesthesia was administered as required. This study was approved by the Ethics Committee of Ohu University (approval number: 186). The informed consent, which was approved by this Committee, was obtained from all participants in this study. The authors declare no conflicts of interest.

A total of 200 patients were included in this study. General anesthesia was induced in all 200 patients at an oxygen rate of 2 L/min, nitrous oxide rate of 4 L/min, and sevoflurane at 5% and maintained using sevoflurane at 1.0-1.5% following endotracheal intubation. At the same time, the administration of remifentanyl, which was adjusted to a dosing rate of 0.01-0.1  $\mu\text{g}/\text{kg}/\text{min}$ , was started. Around 20-40 minutes later, when the spontaneous respiration rate decreased by ~30% compared with that prior to the induction of anesthesia, administration of nitrous oxide was discontinued, and only oxygen was administered at 6 L/min, whereas sevoflurane was increased by 1.5-2.0% and maintained at that value. Thereafter, the infusion rate was adjusted such that  $\gamma^{\text{H}}$  was the infusion rate of remifentanyl ( $\mu\text{g}/\text{kg}/\text{min}$ ) at which the spontaneous respiration rate was half of the rate prior to the induction of anesthesia and remained stable for  $\geq 15$  minutes. A 2 mg vial of remifentanyl was diluted with 20 mL of physiological saline and continuously administered with a syringe pump to a dose of 0.1 mg/mL. The volume of fluid that was administered was in the range 40-60 mL/h. The infusion rate of remifentanyl was calculated based on actual body weight for patients with a body mass index (BMI) of  $\leq 25 \text{ kg}/\text{m}^2$  and standard body weight for patients with a BMI of  $\geq 25 \text{ kg}/\text{m}^2$ . Pain treatment procedures were performed under local anesthesia; however, patients with obvious fluctuations in heart rate, blood pressure, and respiration were excluded because local anesthesia was not successfully administered. The respiratory rate prior to the induction of anesthesia (RRcont), ( $\gamma^{\text{H}}$ ), the respiratory rate at which the infusion rate was stable (RRremi), the amount of carbon dioxide in exhaled air

**Table 1.** Subjects background

Subjects number	200
Sex (M / F)	101 / 99
Age (yr)	29.2 ± 20.3 (3-62)
Body height (cm)	145.1 ± 23.0 (87-182)
Body weight (kg)	44.4 ± 21.1 (13-105)
ASA -PS	1 or 2

mean ± SD (Min - Max).

ASA-PS, The American Society of Anesthesiologists Physical Status. F, female; M, male. Other data during spontaneous general anesthesia.

(EtCO<sub>2</sub>), and oxygen saturation (SpO<sub>2</sub>) were measured. The Spearman's correlation test was performed to statistically analyze the relationship between patients'  $\gamma^H$  and age. For the multivariate analysis, multiple regression analysis with the dosing rate of remifentanyl at half of the preanesthetic respiratory rate (mL/h at 0.1 mg/mL) was used as the dependent variable. Variable selection was based on the forced input method. Data were analyzed using IBM SPSS version 29 for Windows (IBM, Armonk, NY, USA). Statistical significance was set at  $P < 0.05$ .

The free software G-power was used to calculate the sample size. In the t-test for Spearman's correlation, the effect size, alpha error probability, and power were set to 0.25, 0.05, and 0.95, respectively. The total sample size and power were 197 and 0.950, respectively. For the F-tests of the linear multiple regression, the effect size, alpha error probability, power, and number of predictors were set to 0.15, 0.05, 0.95, and two, respectively. The total sample size and power were 107 and 0.952, respectively. Therefore, the sample size of this study was confirmed to be sufficient.

## RESULTS

Data from 286 patients were analyzed in this study. Fifteen patients were excluded because anesthesia-related agents other than sevoflurane, nitrous oxide, and remifentanyl were administered during general anesthesia. Thirty-three patients were excluded because their respiratory and circulatory conditions could not be

**Table 2.** Other data during spontaneous general anesthesia

RRcont (breath/min)	18.9 ± 4.1 (12-30)
$\gamma^H$ ( $\mu\text{g}/\text{kg}/\text{min}$ )	0.042 ± 0.019 (0.016-0.093)
RRremi (breath/min)	8.3 ± 2.8 (4-18)
EtCO <sub>2</sub> (mmHg)	51.0 ± 5.7 (36-63)
SpO <sub>2</sub> (%)	99.4 ± 0.8 (96-100)

mean ± SD (Min - Max).

RRcont, The respiratory rate prior to the induction of anesthesia;  $\gamma^H$ , The remifentanyl dosing rate that reduced preanesthesia spontaneous respiratory rate by half; RRremi, The respiratory rate at which the infusion rate was stable; EtCO<sub>2</sub>, End tidal CO<sub>2</sub>; SpO<sub>2</sub>, Saturation of percutaneous oxygen.

stabilized for more than 15 min, the treatment time was too short, or the effect of local anesthesia was insufficient. Thirty-eight patients were excluded because the data of  $\gamma^H$  or BMI data were outside the box-and-whisker plot of the initial data. Therefore, 86 patients were excluded from the study, and quality data from 200 patients were evaluated.

Participants' backgrounds are listed in Table 1. The mean age of the patients was 29.2 (± 20.3 SD) years, ranging from 3 years 2 months to 62 years. The mean body weight was 44.4 (± 21.1 SD) kg, ranging from 13 kg to 105 kg. In all patients who were included in this study, general anesthesia was maintained with spontaneous respiration using a low-concentration remifentanyl infusion, which allowed dental treatment without any major problems, such as intraoperative awakening or body movement. No postoperative abnormalities or problems were observed in any patient after regaining consciousness. The data obtained during spontaneous general anesthesia are shown in Table 2. The SpO<sub>2</sub> level during the maintenance of general anesthesia under spontaneous respiration with low concentration remifentanyl varied between 96% and 100%, and the mean EtCO<sub>2</sub> was 51.0 mmHg (± 5.7 SD).

The Spearman's correlation test with age on the x-axis and  $\gamma^H$  expressed as  $\mu\text{g}/\text{kg}/\text{min}$  on the y-axis (Fig. 1) revealed a significant negative correlation between the two variables ( $r_s = 0.82$ ,  $P = 2.1^{-49}$ ), and a regression line of  $y = -0.00079x + 0.066$  was obtained. The  $\gamma^H$  corresponding to the age of 2 years was 0.064  $\mu\text{g}/\text{kg}/\text{min}$ , as shown by the regression line and that corresponding




Fig. 1. Relationship between age and dosing rate of remifentanyl ( $\mu\text{g}/\text{kg}/\text{min}$ )




Fig. 2. Relationship between age and dosing rate of remifentanyl ( $\text{mL}/\text{h}$  at  $0.1\text{mg}/\text{mL}$ )

**Table 3.** Multiple regression analysis on dosing rate of remifentanyl (mL/h at 0.1 mg/mL)

	$\beta$	P value	95%CI	
			Lower	Upper
Age	-0.101	0.164	< 0.001	< 0.001
Sex	-0.025	0.729	-0.082	0.058

Dependent variable was dosing rate of remifentanyl at half of preanesthetic respiratory rate (mL/h at 0.1 mg/mL).  $\beta$ , standardized partial regression coefficient; CI, confidence interval.

to 60 years was 0.0186  $\mu\text{g}/\text{kg}/\text{min}$ . This demonstrates that the remifentanyl dosing rate decreases with increasing age.

Moreover, when the infusion rate of remifentanyl diluted to 0.1 mg/mL was changed to mL/h,  $\gamma^H$  was used as the vertical axis and the patient's age was used as the horizontal axis. Spearman's correlation analysis (Fig. 2) showed no increasing or decreasing changes. Therefore, there was no significant correlation between  $\gamma^H$  expressed in mL/h and patient age ( $r_s = 0.073$ ,  $P = 0.306$ ). The obtained regression line of  $y = 0.000396x + 0.88$  showed that  $\gamma^H$  expressed as mL/h did not increase or decrease with age and that  $\gamma^H$  was around 0.88 mL/h at any age (Fig. 2).

Table 3 shows the results of multiple regression analysis with the dosing rate of remifentanyl (mL/h at 0.1 mg/mL) as the dependent variable. Age and sex were not significant independent variables for the dosing rate of remifentanyl at half the pre-anesthetic respiratory rate.

## DISCUSSION

The regression equation  $y = -0.00079x + 0.066$  (x-axis: age, y-axis:  $\gamma^H$  in  $\mu\text{g}/\text{kg}/\text{min}$ ) enabled us to calculate  $\gamma^H$  in  $\mu\text{g}/\text{kg}/\text{min}$ , which was best adjusted for the patient's age. Minimally invasive procedures, such as dental procedures, can be performed by maintaining anesthesia with the respiratory rate controlled at approximately half of that at preanesthetic levels. Moreover, when using remifentanyl, diluted to 0.1 mg/mL, the initial dosing rate was approximately 0.88 mL/h for all age groups. The dose for continuous infusion is usually expressed as  $\mu\text{g}/\text{kg}/\text{min}$  per body weight.

However, when administering drugs to patients, the diluent is placed on a syringe pump, and the dose is set to mL/h. Remifentanyl is usually administered continuously as a 0.1 mg/mL dilution using a syringe pump. Therefore, it is interesting that the remifentanyl infusion rate, which reduces the spontaneous respiration rate to half that before induction of anesthesia, can be achieved regardless of age or physique by simply setting the syringe pump to about 0.88 mL/h. Furthermore, by making small adjustments according to the degree of the patient's respiratory depression, anesthesia can be maintained with the respiratory rate controlled at approximately half of the preanesthetic respiratory rate. This maintenance method is also effective during the final stage of the procedure to reinduce spontaneous breathing in the patient; however, stabilization of the spontaneous respiratory rate after changing the dosing rate may require approximately 20-40 min.

Compared with general anesthesia with positive pressure ventilation, the maintenance of spontaneous ventilation is associated with a lesser degree of decreased venous return and unequal ventilation perfusion ratio [10,11]. Furthermore, as reported by Osaka et al. [12], the presence of spontaneous ventilation or  $\text{EtCO}_2$  values during spontaneous ventilation under general anesthesia provides valuable data regarding the depth of general anesthesia or the state of invasion. Numerous studies have recommended anesthetic management under spontaneous respiration to maintain pulmonary blood flow in patients with pulmonary hypertension. Patients with pulmonary hypertension managed with regional anesthesia and propofol-ketamine without aggravation of hemodynamics through spontaneous respiration [13], and patients managed with sevoflurane and ketamine without elevation of pulmonary vascular resistance during general anesthesia by preserving spontaneous respiration [14] were reportedly successfully managed with adequate pulmonary blood flow. Based on these findings, spontaneous ventilation during general anesthesia may have several advantages.

Our findings showed that increased age was associated

with a gradual decrease in the remifentanyl dosing rate that is required to reduce the spontaneous respiration rate to half of the preanesthetic respiratory rate ( $\gamma^H$ ). Peacock et al. [7] varied the remifentanyl and propofol infusion rates and reported adequate spontaneous respiration in 88% of the study's 64 participants who were aged 18-65 years by remifentanyl of 0.026-0.053  $\mu\text{g}/\text{kg}/\text{min}$ . Barker et al. [8] also examined the relationship between age and remifentanyl dose rate as associated with the occurrence of spontaneous respiration in propofol-remifentanyl anesthesia. According to their study, spontaneous breathing occurred when remifentanyl was administered at 0.192  $\mu\text{g}/\text{kg}/\text{min}$  for children aged 0.5-3 years, 0.095  $\mu\text{g}/\text{kg}/\text{min}$  for children aged 3-6 years, and 0.075  $\mu\text{g}/\text{kg}/\text{min}$  for children aged 6-9 years, which is generally consistent with our results for patients aged 3 years old and over. Furthermore, in a study of 32 children aged 2-7 years undergoing conservative dental treatment, Ansermino et al. [4] reported that for general anesthesia combined with the coadministration of remifentanyl and sevoflurane, the mean dosing rate before the disappearance of spontaneous breathing was 0.127  $\mu\text{g}/\text{kg}/\text{min}$ . These results are also consistent with ours, in which we determined the dosing rate of remifentanyl that would reduce the spontaneous respiration rate to half of the preanesthetic respiratory rate.

Shen et al. [5] found that for general anesthesia with spontaneous breathing in children in whom remifentanyl at a mean dosing rate of 0.036  $\mu\text{g}/\text{kg}/\text{min}$  was co-administered with sevoflurane, the mean  $\text{EtCO}_2$  was 53 mmHg. Moreover, compared with general anesthesia with sevoflurane alone, the time to awakening was 9 min shorter, and the incidence of post-extubation airway obstruction was significantly lower. Hu et al. [6] reported that the administration of general anesthesia with spontaneous breathing in adults who were co-administered remifentanyl at a mean dosing rate of 0.028  $\mu\text{g}/\text{kg}/\text{min}$  with sevoflurane resulted in a mean  $\text{EtCO}_2$  of 52 mmHg, a shorter time to awakening (4 min shorter than that in general anesthesia with sevoflurane alone), and a significantly lower incidence of post-extubation airway

obstruction. Furthermore, Shen et al. [9] reported that for tracheal foreign object removal in 65 children under general anesthesia with propofol-remifentanyl and spontaneous breathing, the mean  $\text{EtCO}_2$  was 59 mmHg but was safely controlled. As indicated by the above reports,  $\text{EtCO}_2$  tends to be higher in general anesthesia with spontaneous breathing using a low concentration of remifentanyl; however,  $\text{EtCO}_2$  of  $53 \pm 6$  mmHg can be safely managed based on the concepts of permissive hypercapnia [15,16]. Therefore, general anesthesia with spontaneous ventilation using low concentrations of remifentanyl that is co-administered with sevoflurane is considered a safe method that shortens the time to awakening and is associated with a low incidence of post-extubation airway obstruction. In contrast, conscious intubation is recommended to secure the airways in patients with difficult intubation. Appropriate sedation and preservation of spontaneous breathing are the principles of conscious intubation [17]. Among these, opioids such as remifentanyl are often used to alleviate pain [18,19]. However, excessive opioid administration can easily stop spontaneous respiration. Therefore, controlling the administration rate is important to preserve spontaneous respiration. The administration rates of remifentanyl in these studies were within the ranges specified in this report. In addition, the age-adjusted rate shown in this study would be useful for setting an appropriate opioid administration rate to preserve spontaneous breathing.

Nevertheless, a higher concentration of remifentanyl is required for some procedures, such as highly invasive surgeries, making management with spontaneous ventilation impossible. Therefore, this method of maintaining general anesthesia using a low concentration of remifentanyl that preserves spontaneous ventilation is considered feasible for minimally invasive procedures, such as dental treatments, which can be performed without inducing pain, as local anesthesia induces a sufficient response.

Multiple regression analysis adjusted for sex also showed that the dosing rate of remifentanyl at half the preanesthetic respiratory rate was not correlated with age. BMI was not used as an independent variable because



the dosing rate of remifentanyl in obese patients was calculated from the standard body weight based on a BMI of 22 kg/m<sup>2</sup>, which was not associated with the actual BMI. Moreover, this regression equation is only an estimate for setting the dosing rate, and the precise concentration or dosing rate must be adjusted for each patient according to the individual responses to sevoflurane or remifentanyl. There are various cases of clinical general anesthesia, and it is hoped that this report will be of some help in cases where the preservation of spontaneous breathing is advantageous.

In conclusion, in minimally invasive procedures, such as dental procedures, anesthesia can be maintained by the administration at a dosing rate of  $-0.00079 \times \text{patient age} + 0.066 \mu\text{g/kg/min}$  with 1.5-2.0% sevoflurane to reduce and control the spontaneous respiratory rate to approximately half of the preanesthetic respiratory rate after approximately 20-40 min. The administration of remifentanyl solution diluted to 0.1 mg/mL at approximately 0.88 mL/h allows anesthesia to be maintained while reducing the spontaneous respiratory rate to approximately half of the preanesthetic respiratory rate after approximately 20-40 min in patients of all age groups. Because general anesthesia under spontaneous ventilation has many advantages, our method is a promising and effective means of administering anesthesia in minimally invasive procedures.

#### AUTHOR ORCID*s*

**Daijiro Ogumi:** <https://orcid.org/0000-0002-3668-573X>

**Shota Abe:** <https://orcid.org/0000-0003-4597-3327>

**Hikaru Sato:** <https://orcid.org/0000-0002-0106-0211>

**Fumihiko Suzuki:** <https://orcid.org/0000-0002-5156-9729>

**Hiroyoshi Kawaai:** <https://orcid.org/0000-0002-4718-0550>

**Shinya Yamazaki:** <https://orcid.org/0000-0002-9273-3927>

#### AUTHOR CONTRIBUTIONS

**Daijiro Ogumi:** Writing - original draft

**Shota Abe:** Formal analysis

**Hikaru Sato:** Data curation

**Fumihiko Suzuki:** Data curation, Formal analysis

**Hiroyoshi Kawaai:** Supervision

**Shinya Yamazaki:** Conceptualization, Supervision, Writing - review & editing

**DECLARATION OF INTEREST:** The authors of this paper have no conflicts of interest to declare.

**FUNDING:** None

#### REFERENCES

1. Natalini G, Di Maio A, Rosano A, Ferretti P, Bertelli M, Bernardini A. Remifentanyl improves breathing pattern and reduces inspiratory workload in tachypneic patients. *Respir Care* 2011; 56: 827-33.
2. Ohara S, Nishimura A, Tachikawa S, Iijima T. Effect of remifentanyl on intraoperative fluid balance: a retrospective statistical examination of factors contributing to fluid balance. *J Dent Anesth Pain Med* 2020; 20: 129-35.
3. Ishii M, Tomoyasu Y, Arai Y, Hayashi T, Yamane A, Maeda S, et al. A case of desensitized severe gagging reflex resulting from sedation with propofol and remifentanyl during dental treatment. *J Jpn Dent Soc Anesthesiol* 2012; 40: 46-7.
4. Ansermino JM, Brooks P, Rosen D, Vandebek CA, Reichert C. Spontaneous ventilation with remifentanyl in children. *Paediatr Anaesth* 2005; 15: 115-21.
5. Shen X, Hu C, Li W. Tracheal extubation of deeply anesthetized pediatric patients: a comparison of sevoflurane and sevoflurane in combination with low-dose remifentanyl. *Paediatr Anaesth* 2012; 22: 1179-84.
6. Hu C, Yu H, Ye M, Shen X. Sevoflurane in combination with remifentanyl for tracheal extubation after otologic surgery. *Am J Health Syst Pharm* 2014; 71: 1108-11.
7. Peacock JE, Luntley JB, O'Connor B, Reilly CS, Ogg TW, Watson BJ, et al. Remifentanyl in combination with propofol for spontaneous ventilation anaesthesia. *Br J Anaesth* 1998; 80: 509-11.
8. Barker N, Lim J, Amari E, Malherbe S, Ansermino JM. Relationship between age and spontaneous ventilation during intravenous anesthesia in children. *Paediatr Anaesth* 2007; 17: 948-55.
9. Shen X, Hu CB, Ye M, Chen YZ. Propofol-remifentanyl intravenous anesthesia and spontaneous ventilation for airway foreign body removal in children with preoperative

- respiratory impairment. *Paediatr Anaesth* 2012; 22: 1166-70.
10. Brecher GA, Hubay CA. Pulmonary blood flow and venous return during spontaneous respiration. *Circ Res* 1955; 3: 210-4.
  11. Putensen C, Mutz NJ, Putensen-Himmer G, Zinserling J. Spontaneous breathing during ventilatory support improves ventilation-perfusion distributions in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1999; 159: 1241-8.
  12. Osaka Y, Koitabashi T. The relationship between the respiratory parameters under spontaneous breathing and effect site propofol concentrations. *Masui* 2004; 53: 1130-5.
  13. Koyata T, Yamamoto M, Shionoya Y, Nakamura K. General anesthesia for tooth extractions in a child with fontan circulation and a pacemaker. *J Jpn Dent Soc Anesthesiol* 2019; 47: 1-3.
  14. Williams GD, Philip BM, Chu LF, Boltz MG, Kamra K, Terwey H, et al. Ketamine does not increase pulmonary vascular resistance in children with pulmonary hypertension undergoing sevoflurane anesthesia and spontaneous ventilation. *Anesth Analg* 2007; 105: 1578-84.
  15. Bigatello LM, Patroniti N, Sangalli F. Permissive hypercapnia. *Curr Opin Crit Care* 2001; 7: 34-40.
  16. Ikegami Y, Tase C. Permissive hypercapnia. *Jpn J Reanimatol* 2002; 21: 1-8.
  17. Ahmad I, El-Boghdadly K, Bhagrath R, Hodzovic I, McNarry AF, Mir F, et al. Difficult Airway Society guidelines for awake tracheal intubation (ATT) in adults. *Anaesthesia* 2020; 75: 509-28.
  18. Donaldson AB, Meyer-Witting M, Roux A. Awake fiberoptic intubation under remifentanyl and propofol target-controlled infusion. *Anaesth Intensive Care* 2002; 30: 93-5.
  19. Santiago Martín J, Torres Fernández V, Muñoz Blanco F, Santiago Martín FM, Puente Rodero A, Barranco Armenteros F. Nasotracheal fiberoptic intubation under remifentanyl for sedation and analgesia in a boy with a difficult airway due to giant gingival hypertrophy. *Rev Esp Anesthesiol Reanim* 2005; 52: 363-6.