

mutations associated to MVC resistance were analyzed, as well as the association of the subtype with clinical, epidemiological, virological, and immunological variables. The subtypes detected using the C2V3 region were CRF20_23_24_BG (35 patients, 30.4%); Subtype B (33 patients, 28.7%); CRF19_cpx (30 patients, 26.1%); CRF18_cpx (10 patients, 8.7%); and others (7 patients, 6.1%). Overall, 60 per cent of the viruses exhibited R5 phenotype, 14.8 per cent were R5X4 and 25.2 per cent were X4. Interestingly, CRF19_cpx virus was associated with having phenotype X4 [46.7%, $P = 0.0047$, odds ratio (OR): 3.96, 95% confidence interval (95% CI): 1.59–9.84], with infection in young individuals (39.1%, $P = 0.025$, OR: 3.548; 95% CI: 1.136–11.077) and with higher values of viral load ($P \leq 0.05$). The comparison of the amino acid sequences of the V3 loop showed differences between the B and non-B subtypes ($P = 0.0001$). Mutations reported to be associated with MVC resistance, were detected in 75.7 per cent of the samples, in positions 11 (6.1%), 13 (49.6%), 25 (6.1%), 316 (7.0%), 323 (11.3%), and 319 (3.5%) of Gp120, particularly in the recombinant forms CRF19_cpx and CRF_BGs. HIV variants that use the CXCR4 co-receptor were associated with more than 10 years of diagnosis, with older individuals, in the AIDS stage, with low CD4 counts and higher viral load levels ($P < 0.05$). The results support the hypothesis previously stated that CRF19_cpx viruses could be more pathogenic and would have limitations for the use of MVC. The high rate of mutations associated to MVC among non-B Cuban subtypes should be further studied.

A8 Epidemiological study of transmission clusters in a local HIV-1 cohort

Carmen M. González Domenech,¹ Isabel Viciano-Ramos,^{2,3} Gabriel Sena-Corrales,² Laura Mora Navas,² Guillermo Ojeda,² Enrique Nuño,² Encarnación Clavijo,² Rosario Palacios,^{2,3} and Jesus Santos^{2,3}

¹Department of Microbiology, University of Granada, Granada, Spain, ²Infectious Diseases and Clinical Microbiology Unit, Virgen de la Victoria University Hospital, Malaga, Spain and ³Institute of Biomedical Investigation of Malaga (IBIMA), Malaga, Spain

Integration of molecular, clinical, and demographic data represents a powerful tool to understand the dynamics of local HIV-1 transmission chains (TCs). The aim of our study was the phylogenetic analysis of the TCs within a HIV-1 cohort and the description of the relevant patient data within a TC. We performed a phylogenetic analysis of 757 sequences from newly HIV-1 diagnosed patients in Málaga (Southern Spain) during the period 2004–15. We used partial *pol* gene sequences in a preliminary phylogenetic reconstruction using the Neighbour Joining method (MEGA v6.06 program). After eliminating branches with bootstrap values <80 per cent, we constructed a new phylogeny by Maximum likelihood method (FastTree program). We considered as TC any cluster with bootstrap values ≥ 90 per cent. Patient sequences within and outside TCs were compared. Resistance mutations in the protease (PR) and reverse transcriptase (RT) sequences were analyzed using the Stanford algorithm. Four hundred and fifty-one out of 757 patients (59.6%) were grouped into fifty-three TCs, seventeen of them with five or more subjects. The largest number of patients associated within a TC was ninety. Patients younger than 40 years [odds ratio (OR) 1.75, 95% confidence interval (95% CI) 1.2–2.4, $P = 0.002$], men who have sex with men (MSM) (OR 2.14, 95% CI 1.3–3.2, $P < 0.0001$), non-Spanish (OR 1.48, 95% CI 1.0–2.1, $P = 0.038$), with a non-B subtype HIV-1 (OR 3.12, 95% CI 2.0–4.8, $P < 0.0001$), and presenting primary resistance mutations (OR 14.1, 95% CI 3.1–62.6, $P = 0.001$) were more likely to be associated within a cluster. Ninety-four out of 118 patients (79.6%) with transmission resistance mutations were included in some TC. The most frequent mutations associated with clusters were T69D/N, L210W, and K219E/Q, for NRTIs, K103N, and G190A/S for NNRTIs, and the I54L/M and L90M mutations for PIs. The prevalence for resistance to NNRTIs in TCs was 13.7 per cent. There were two TCs of rarer non-B subtypes: CRF19_cpx, with twenty-one individuals, sixteen of them (76.2%) with mutation G190A; and CRF51_01B with thirty-nine patients, twenty of them with the K103N mutation. Approximately 60 per cent of newly HIV-1 diagnosed patients were included in a TC. Younger patients, MSM, non-Spanish, with non-B subtype HIV-1 and primary resistance mutations were more likely to belong to a cluster. NNRTI mutations were the most frequent ones among patients in TCs. We observed two TCs represented by infrequent non-B subtypes in our area—CRF19_cpx and CRF51_01B—both of which were associated to the transmission of primary resistance.

A9 A method to obtain full-length HIV proviral sequences and their sites of integration

Sean C. Patro,¹ Xiaolin Wu,² Shuang Guo,² Michael J. Bale,¹ Ann Wiegand,¹ Jonathan Spindler,¹ Christine M. Fennessey,³ Elias K. Halvas,⁴ Brandon F. Keele,³ Stephen H. Hughes,¹ John W. Mellors,⁴ John M. Coffin,⁵ Jason W. Rausch,⁶ and Mary F. Kearney⁴

¹HIV Dynamics and Replication Program, National Cancer Institute, Frederick, MD 21702, USA, ²Leidos Biomedical Research, Inc., Frederick National Laboratory for Cancer Research, Frederick, MD 21702, USA, ³AIDS and Cancer Virus Program, Leidos Biomedical Research, Inc., Frederick National Laboratory, Frederick, MD 21702, USA, ⁴Department of Medicine, University of Pittsburgh, Pittsburgh, PA 15213, USA, ⁵Department of Molecular Biology and Microbiology, Tufts University, Boston, MA 02111, USA and ⁶Basic Research Laboratory, Center for Cancer Research, National Cancer Institute, Frederick, MD 21702, USA

Accurate definition of the HIV-1 reservoir on antiretroviral therapy (ART) is of paramount importance to the development of curative strategies. Much of this reservoir is derived from clonal expansion of latently infected CD4+ T cells. Methods used to characterize the reservoir include near full-length single-genome sequencing (NFL-SGS) and integration site analysis (ISA). However, current technologies do not link the intact proviruses detected by NFL-SGS to their sites of integration. Therefore, we developed a method to obtain both near full-length single-proviral sequences and their sites of integration. We call our method full-length integrated proviral single-genome sequencing (FLIP-SGS). Genomic DNA from ACH2 and CEM cells mixed at 1:1,000, or patient samples were diluted to a single proviral endpoint. An in-house, optimized whole genome amplification (WGA) method was performed on wells at the endpoint, generating multiple copies of all DNA molecules within each well. The number of proviral copies after WGA was determined by droplet digital PCR targeting the long terminal region (LTR). Forty per cent of each WGA reaction was used to obtain the provirus–host integration sites with ISA (linker ligation, nested PCR, and illumina sequencing). The remaining fraction was used to amplify the full-length proviruses in four overlapping fragments (LTR-pol, gag-int, int-env, and env-LTR) for Sanger sequencing. WGA performed on the endpoint-diluted ACH2:CEM DNA amplified single-copy HIV-1 proviral templates greater than 500-fold, making it possible to obtain unique integration sites from single proviruses in ACH2 cells, including one that was previously reported (in the NT5C3A gene on chromosome 7) and two that were not previously reported (in the EIF4ENIF1 gene of chromosome 22 and an unknown region of chromosome 6). Near full-length PCR amplification and Sanger sequencing was performed on proviruses integrated in the NT5C3A gene. FLIP-SGS was applied to peripheral blood mononuclear cells from one HIV-1 infected donor with viremia suppressed on ART and yielded integration sites of four genomes that appear to contain large internal deletions. We report a method for near full-length HIV-1 single-genome sequencing combined with host integration site detection that we call FLIP-SGS. This assay will further define clonal expansion of infected CD4+ T cells as a mechanism that maintains the HIV-1 reservoir and as the source of identical sequences observed during therapy and rebound, rather than from ongoing replication.

A10 Presence and frequency of M184V mutation in the MOBIDIP trial

A. Armero,^{1,2} M. L. Chaix,^{1,2} M. L. Nere,^{1,2} E. Delaporte,^{3,4} M. Peeters,^{3,4} and C. Delaugerre^{1,2}

¹Virologie, Hôpital Saint-Louis, Assistance Publique Hôpitaux de Paris, Paris, France, ²INSERM UMR 941, Université de Paris Diderot, Sorbonne Paris Cité, France, ³Institut de Recherche pour le Développement UMI 233, Montpellier, France and ⁴INSERM U1175, Université de Montpellier, Montpellier, France

The MOBIDIP trial evaluated the simplification by protease (PI/r) monotherapy for HIV infection versus dual therapy and boosted protease inhibitor plus lamivudine (PI/r + 3TC) in controlled patients under second-line regimens. MOBIDIP was interrupted because of a significant number of patients with virological failure (VF) at week 48 (W48) in PI/r (33/133, ~25%) versus in PI/r + 3TC (4/132, ~3%). At the time of first-line VF, 96 per cent of patients harbored the M184V mutation. The presence of the M184V mutation was related to a protective effect against VF in the